



# An air-stable N-heterocyclic carbene iminoxyl borate radical zwitterion†

Cite this: *Chem. Commun.*, 2018, 54, 6824

Youngsuk Kim <sup>ab</sup> and Eunsung Lee <sup>\*abc</sup>

Received 18th February 2018,  
Accepted 26th March 2018

DOI: 10.1039/c8cc01399c

rsc.li/chemcomm

**A remarkably stable radical zwitterion derived from N-heterocyclic carbene nitric oxide and  $B(C_6F_5)_3$  is reported. The presented radical was generated by steric and electronic protection of the nitric oxide moiety using  $B(C_6F_5)_3$ , which secured its stability toward air and moisture. An analogous yet less stable radical derived from  $C(C_6H_5)_3$ <sup>+</sup> is also synthesized and characterized.**

Stable organic radicals are of great interest to chemists since they offer fundamental understanding of reactive intermediates, as well as they have numerous applications as functional materials.<sup>1</sup> While the majority of organic radicals are thermodynamically and kinetically unstable, there are several kinds of stable organic radicals, namely triarylmethyl, nitroxyl, and hydrazyl radicals, for example.<sup>2</sup> In the past decade, N-heterocyclic carbenes (NHCs)<sup>3</sup> have been introduced to stabilise various organic radicals and radical ions,<sup>4</sup> as a variety of previously inaccessible organic radicals have been successfully prepared and characterised with the aid of NHCs.<sup>5</sup> To date, NHC-stabilised carbonyl (A and B),<sup>6</sup> propargyl (C),<sup>7</sup> aminyl (D and E),<sup>8</sup> phosphinyl (F),<sup>9</sup> and many other organic radicals have been isolated or spectroscopically characterised by the groups of Bertrand, Roesky, Curran, and many others (Scheme 1).<sup>10</sup> These radicals were successfully stabilised due to the  $\pi$ -accepting properties of NHCs that delocalise the spin density,<sup>11</sup> as well as the steric protection of the bulky NHC substituents.<sup>12</sup>

In this context, our group recently reported the synthesis of persistent iminoxyl radicals derived from NHCs with nitric oxide.<sup>10a,13</sup> The iminoxyl radical **1** was quite stable in the solid



**Scheme 1** Selected examples of organic radicals stabilised by NHC (Dipp = 2,6-diisopropylphenyl).

state; however, in the solution phase, it slowly decomposed even under an inert atmosphere of nitrogen. Here we report the synthesis and characterisation of a remarkably stable radical zwitterion **2** obtained from the steric protection of **1** using tris(pentafluorophenyl)borane ( $B(C_6F_5)_3$ ) (Scheme 2).

The radical **2** can be synthesised in high yield by mixing equimolar amounts of **1** and  $B(C_6F_5)_3$  in diethyl ether solution under a  $N_2$  atmosphere. The reaction mixture turned dark brown immediately, and after 15 minutes, volatiles were removed under vacuum. After washing the resulting solid using *n*-pentane, **2** was isolated as a dark brown solid in 98% yield. It is notable that a similar reactivity was recently reported for the (2,2,6,6-tetramethylpiperidin-1-yl)oxyl (TEMPO) radical with  $B(C_6F_5)_3$ .<sup>14</sup> The molecular structure of **2** was unambiguously determined using single crystal X-ray crystallography (Fig. 1). The bond lengths of C1–N3 (1.339(2) Å) and N3–O1 (1.344(2) Å) indicate a bond order of 1.5. The (imidazole ring)–N3–O1–B1 group is planar, which also suggests delocalisation of the radical through  $\pi$ -conjugation. The structural parameters of **2** were well reproduced using density functional theory (DFT) calculations at the B3LYP/6-31G(d,p) level of theory. The Wiberg bond orders were calculated for the C1–N3 (1.49), N3–O1 (1.44), and O1–B1 (0.86) bonds, which were consistent with the structure obtained from single crystal X-ray analysis. The experimental electron paramagnetic resonance

<sup>a</sup> Center for Self-assembly and Complexity, Institute for Basic Science (IBS), Pohang, 37673, Republic of Korea

<sup>b</sup> Department of Chemistry, Pohang University of Science and Technology, Pohang, 37673, Republic of Korea. E-mail: eslee@postech.ac.kr

<sup>c</sup> Division of Advanced Materials Science, Pohang University of Science and Technology, Pohang, 37673, Republic of Korea

† Electronic supplementary information (ESI) available: Experimental details, EPR spectra, X-ray crystal data, DFT calculation results. CCDC 1823243 and 1823244. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c8cc01399c



**Scheme 2** Synthesis of the radical zwitterion **2** from IPr (1,3-bis-(2,6-diisopropylphenyl)imidazol-2-ylidene) and NO (nitric oxide). <sup>a</sup> Isolated yield.




**Fig. 1** Molecular structure of **2** from X-ray crystallography. The thermal ellipsoids are shown at the 50% probability level. Hydrogen atoms and solvent molecules (*n*-pentane) were omitted for clarity.


(EPR) spectrum shows a complex splitting pattern (Fig. 2), which was well reproduced in the simulated spectra. DFT calculations suggest that the N3 atom has the largest spin density (44%), consistent with the largest hyperfine coupling constant on N3 (23.4 MHz). The calculation also shows that the singly occupied molecular orbital (SOMO) of **2** is delocalised over the molecular plane (Fig. S1, ESI<sup>†</sup>). The UV-vis absorption spectrum of **2** in benzene at room temperature shows a peak at  $\lambda_{\text{max}} = 448$  nm (Fig. S8, ESI<sup>†</sup>). The cyclic voltammogram of **2** reveals one reversible redox peak at  $E_{1/2} = -0.022$  V (*versus* saturated Ag/AgCl electrode), showing that **2** is a weak oxidant (Fig. S13, ESI<sup>†</sup>).

It is notable that **2** shows remarkable stability toward air and moisture. For example, a solution of **2** in wet technical-grade benzene was monitored using UV-vis, which showed no detectable decomposition over 12 days (Fig. 3). In addition, **2** was stable even under silica chromatographic conditions as the benzene solution of **2** was still EPR active even after the filtration through silica gel under air (Fig. S6, ESI<sup>†</sup>).


After discovering the remarkable stability of the radical **2**, we were curious whether the radical cation **3** with almost equivalent steric bulk is also stable or not. When **1** was treated with a stoichiometric amount of trityl tetrakis(pentafluorophenyl)borate



**Fig. 2** Experimental (bottom) and simulated (top) EPR spectra of **2** ( $g = 2.0107$ ; hyperfine coupling constants:  $a(^{14}\text{N}) = 23.4, 9.1, 6.9$  MHz,  $a(^{11}\text{B}) = 6.7$  MHz,  $a(^1\text{H}) = 5.3, 3.6$  MHz,  $a(^{19}\text{F}) = 2.6, 1.7$  MHz).




**Fig. 3** Decay of the radicals **2** and **3** in wet benzene solution under air; monitored by UV-vis.



**Scheme 3** Synthesis of the radical cation **3**. <sup>a</sup> Isolated yield.

( $\text{Ph}_3\text{C}^+\text{B}(\text{C}_6\text{F}_5)_4^-$ ) in diethyl ether solution under a  $\text{N}_2$  atmosphere, the color of the reaction mixture changed immediately to dark brown (Scheme 3). After 5 minutes, the product was precipitated by the addition of *n*-pentane, and subsequently washed to yield **3** as a dark brown solid (96%). Single crystal X-ray crystallographic analysis revealed the molecular structure of **3** with the  $\text{B}(\text{C}_6\text{F}_5)_4^-$  counteranion (Fig. 4). The planar molecular structure and bond lengths of C1–N3 (1.339(1) Å) and N3–O1 (1.361(1) Å) of **3** were almost identical with the structure of **2**. The calculated Wiberg bond lengths of C1–N3 (1.50), N3–O1 (1.38), and O1–C2 (0.87) bonds of **3** were also similar with those of **2**. On the other hand, the experimental and simulated EPR spectra of **3** were very different from those of **2** due to the absence of nearby boron and fluorine atoms (Fig. 5). The UV-vis spectrum of **3** in benzene was recorded at room temperature, which showed a peak at  $\lambda_{\text{max}} = 458$  nm (Fig. S9, ESI<sup>†</sup>). In contrast to **2**, **3** is a quite strong oxidant as its cyclic



**Fig. 4** Molecular structure of **3** from X-ray crystallography. The thermal ellipsoids are shown at the 50% probability level. Hydrogen atoms were omitted for clarity.



Fig. 5 Experimental (bottom) and simulated (top) EPR spectra of **3** ( $g = 2.0098$ ; hyperfine coupling constants:  $a(^{14}\text{N}) = 26.6, 9.3, 7.5$  MHz,  $a(^1\text{H}) = 10.2, 6.6$  MHz).

voltammogram shows one reversible redox peak at  $E_{1/2} = 0.582$  V versus sat. Ag/AgCl (Fig. S14, ESI<sup>†</sup>). One-electron reduction of **3** using decamethylferrocene also resulted in the neutral oxime compound (see the ESI<sup>†</sup>). Interestingly, **3** was much more sensitive toward air and moisture than **2** as the half-life of **3** in wet technical-grade benzene was approximately 35 minutes (Fig. 3).

Therefore, simply protecting the radical center using a bulky substituent did not guarantee the stability of the radical. DFT calculations at the B3LYP/6-31G(d,p) level using the SMD solvation model showed that the trityl group is not binding strongly enough to the oxygen atom of **1** compared to the  $\text{B}(\text{C}_6\text{F}_5)_3$  group. The standard free energy of the dissociation of the  $\text{B}(\text{C}_6\text{F}_5)_3$  group from **2** is energetically uphill by  $12.6 \text{ kcal mol}^{-1}$  in benzene solution, while the trityl group of **3** required only  $1.4 \text{ kcal mol}^{-1}$  for dissociation (Fig. S2, ESI<sup>†</sup>). Additional calculations using  $\text{BPh}_3$  and  $\text{C}(\text{C}_6\text{F}_5)_3^+$  groups also suggested that the introduction of electron-withdrawing fluorine atoms makes huge difference in the dissociation energies, as perfluorinated substituents have much higher electrophilicity (see the ESI<sup>†</sup>). The dissociation of the trityl group from **3** was also evidenced by the crossover experiment: adding 2 equivalents of  $\text{B}(\text{C}_6\text{F}_5)_3$  to a solution of **3** successfully generated **2** along with the trityl group as observed by EPR (Fig. S7, ESI<sup>†</sup>) and UV-vis (Fig. S12, ESI<sup>†</sup>).

In summary, two different radicals **2** and **3** were synthesised from **1** and  $\text{B}(\text{C}_6\text{F}_5)_3$  or  $\text{Ph}_3\text{C}^+\text{B}(\text{C}_6\text{F}_5)_4^-$ , respectively. The radical zwitterion **2** showed remarkable stability toward air, moisture, and even toward silica. On the other hand, **3** showed limited stability upon exposure to air and moisture, showing a half-life of about 35 minutes in wet benzene solution. The difference of stability is mainly because the trityl moiety of **3** binds weaker than the  $\text{B}(\text{C}_6\text{F}_5)_3$  group of **2** to the oxygen, as analysed *via* DFT calculations. With the help from the novel properties of NHCs, this work clearly shows a successful example of designing a stable radical. Possible applications of the stable radical **2** are currently under active investigation.

This work was supported by Institute for Basic Science (IBS) [IBS-R007-D1]. The X-ray diffraction experiment with synchrotron radiation was performed at the Pohang Accelerator Laboratory (Beamline 2D).

## Conflicts of interest

There are no conflicts to declare.

## Notes and references

- 1 I. Ratera and J. Veciana, *Chem. Soc. Rev.*, 2012, **41**, 303–349.

- 2 R. G. Hicks, *Stable Radicals: Fundamentals and Applied Aspects of Odd-Electron Compounds*, John Wiley & Sons, Ltd, 2010.
- 3 (a) M. N. Hopkinson, C. Richter, M. Schedler and F. Glorius, *Nature*, 2014, **510**, 485–496; (b) D. Bourissou, O. Guerret, F. P. Gabbaï and G. Bertrand, *Chem. Rev.*, 2000, **100**, 39–92; (c) A. Igau, H. Grutzmacher, A. Baceiredo and G. Bertrand, *J. Am. Chem. Soc.*, 1988, **110**, 6463–6466; (d) A. J. Arduengo, R. L. Harlow and M. Kline, *J. Am. Chem. Soc.*, 1991, **113**, 361–363; (e) V. Lavallo, Y. Canac, C. Prasang, B. Donnadieu and G. Bertrand, *Angew. Chem., Int. Ed.*, 2005, **44**, 5705–5709; (f) T. W. Hudnall and C. W. Bielawski, *J. Am. Chem. Soc.*, 2009, **131**, 16039–16041; (g) M. Melaimi, M. Soleilhavoup and G. Bertrand, *Angew. Chem., Int. Ed.*, 2010, **49**, 8810–8849.
- 4 (a) C. D. Martin, M. Soleilhavoup and G. Bertrand, *Chem. Sci.*, 2013, **4**, 3020–3030; (b) D. P. Curran, A. Solov'yev, M. Makhlof Brahmi, L. Fensterbank, M. Malacria and E. Lacote, *Angew. Chem., Int. Ed.*, 2011, **50**, 10294–10317; (c) M. Soleilhavoup and G. Bertrand, *Acc. Chem. Res.*, 2015, **48**, 256–266; (d) J. P. Moerdyk, D. Schilter and C. W. Bielawski, *Acc. Chem. Res.*, 2016, **49**, 1458–1468; (e) M. Melaimi, R. Jazzar, M. Soleilhavoup and G. Bertrand, *Angew. Chem., Int. Ed.*, 2017, **56**, 10046–10068.
- 5 (a) O. Back, M. A. Celik, G. Frenking, M. Melaimi, B. Donnadieu and G. Bertrand, *J. Am. Chem. Soc.*, 2010, **132**, 10262–10263; (b) O. Back, B. Donnadieu, P. Parameswaran, G. Frenking and G. Bertrand, *Nat. Chem.*, 2010, **2**, 369–373; (c) R. Kinjo, B. Donnadieu and G. Bertrand, *Angew. Chem., Int. Ed.*, 2010, **49**, 5930–5933; (d) R. Kinjo, B. Donnadieu, M. A. Celik, G. Frenking and G. Bertrand, *Science*, 2011, **333**, 610–613; (e) H. Tanaka, M. Ichinohe and A. Sekiguchi, *J. Am. Chem. Soc.*, 2012, **134**, 5540–5543; (f) K. C. Mondal, H. W. Roesky, M. C. Schwarzer, G. Frenking, I. Tkach, H. Wolf, D. Kratzert, R. Herbst-Irmer, B. Niepötter and D. Stalke, *Angew. Chem., Int. Ed.*, 2013, **52**, 1801–1805.
- 6 (a) J. K. Mahoney, D. Martin, C. E. Moore, A. L. Rheingold and G. Bertrand, *J. Am. Chem. Soc.*, 2013, **135**, 18766–18769; (b) D. Martin, C. E. Moore, A. L. Rheingold and G. Bertrand, *Angew. Chem., Int. Ed.*, 2013, **52**, 7014–7017; (c) J. K. Mahoney, D. Martin, F. Thomas, C. E. Moore, A. L. Rheingold and G. Bertrand, *J. Am. Chem. Soc.*, 2015, **137**, 7519–7525; (d) C. L. Deardorff, R. Eric Sikma, C. P. Rhodes and T. W. Hudnall, *Chem. Commun.*, 2016, **52**, 9024–9027; (e) V. Regnier, F. Molton, C. Philouze and D. Martin, *Chem. Commun.*, 2016, **52**, 11422–11425; (f) J. K. Mahoney, R. Jazzar, G. Royal, D. Martin and G. Bertrand, *Chem. – Eur. J.*, 2017, **23**, 6206–6212.
- 7 (a) M. M. Hansmann, M. Melaimi and G. Bertrand, *J. Am. Chem. Soc.*, 2017, **139**, 15620–15623; (b) Y. Li, K. C. Mondal, P. P. Samuel, H. Zhu, C. M. Orben, S. Pannesar, B. Dittrich, B. Schwederski, W. Kaim, T. Mondal, D. Koley and H. W. Roesky, *Angew. Chem., Int. Ed.*, 2014, **53**, 4168–4172; (c) L. Jin, M. Melaimi, L. Liu and G. Bertrand, *Org. Chem. Front.*, 2014, **1**, 351–354; (d) M. M. Hansmann, M. Melaimi and G. Bertrand, *J. Am. Chem. Soc.*, 2018, **140**, 2206–2213; (e) M. M. Hansmann, M. Melaimi, D. Munz and G. Bertrand, *J. Am. Chem. Soc.*, 2018, **140**, 2546–2554.
- 8 (a) L. Y. Eymann, A. G. Tskhovrebov, A. Sienkiewicz, J. L. Bila, I. Zivkovic, H. M. Ronnow, M. D. Wodrich, L. Vannay, C. Corminboeuf, P. Pattison, E. Solari, R. Scopelliti and K. Severin, *J. Am. Chem. Soc.*, 2016, **138**, 15126–15129; (b) J. Back, J. Park, Y. Kim, H. Kang, Y. Kim, M. J. Park, K. Kim and E. Lee, *J. Am. Chem. Soc.*, 2017, **139**, 15300–15303.
- 9 O. Back, B. Donnadieu, M. von Hopffgarten, S. Klein, R. Tonner, G. Frenking and G. Bertrand, *Chem. Sci.*, 2011, **2**, 858.
- 10 (a) J. Park, H. Song, Y. Kim, B. Eun, Y. Kim, D. Y. Bae, S. Park, Y. M. Rhee, W. J. Kim, K. Kim and E. Lee, *J. Am. Chem. Soc.*, 2015, **137**, 4642–4645; (b) S. Roy, A. C. Stuckl, S. Demeshko, B. Dittrich, J. Meyer, B. Maity, D. Koley, B. Schwederski, W. Kaim and H. W. Roesky, *J. Am. Chem. Soc.*, 2015, **137**, 4670–4673; (c) S. Styra, M. Melaimi, C. E. Moore, A. L. Rheingold, T. Augenstein, F. Breher and G. Bertrand, *Chem. – Eur. J.*, 2015, **21**, 8441–8446; (d) A. D. Ledet and T. W. Hudnall, *Dalton Trans.*, 2016, **45**, 9820–9826; (e) S. Kundu, P. P. Samuel, S. Sinhababu, A. V. Luebben, B. Dittrich, D. M. Andrada, G. Frenking, A. C. Stuckl, B. Schwederski, A. Paretzki, W. Kaim and H. W. Roesky, *J. Am. Chem. Soc.*, 2017, **139**, 11028–11031; (f) S. Kundu, S. Sinhababu, S. Dutta, T. Mondal, D. Koley, B. Dittrich, B. Schwederski, W. Kaim, A. C. Stuckl and H. W. Roesky, *Chem. Commun.*, 2017, **53**, 10516–10519; (g) B. Li, S. Kundu, A. C. Stuckl, H. Zhu, H. Keil, R. Herbst-Irmer, D. Stalke, B. Schwederski, W. Kaim, D. M. Andrada, G. Frenking and H. W. Roesky, *Angew. Chem., Int. Ed.*, 2017, **56**, 397–400; (h) M. F. Silva Valverde, P. Schweyen, D. Gisinger, T. Bannenberg, M. Freytag, C. Kleeberg and

- M. Tamm, *Angew. Chem., Int. Ed.*, 2017, **56**, 1135–1140; (i) Y. Kim, K. Kim and E. Lee, *Angew. Chem., Int. Ed.*, 2018, **57**, 262–265.
- 11 (a) D. Martin, N. Lassauque, B. Donnadiou and G. Bertrand, *Angew. Chem., Int. Ed.*, 2012, **51**, 6172–6175; (b) J. P. Moerdyk, G. A. Blake, D. T. Chase and C. W. Bielawski, *J. Am. Chem. Soc.*, 2013, **135**, 18798–18801; (c) H. Song, Y. Kim, J. Park, K. Kim and E. Lee, *Synlett*, 2016, 477–485; (d) D. Martin, M. Soleilhavoup and G. Bertrand, *Chem. Sci.*, 2011, **2**, 389–399.
- 12 (a) S. Würtz and F. Glorius, *Acc. Chem. Res.*, 2008, **41**, 1523–1533; (b) C. Valente, S. Calimsiz, K. H. Hoi, D. Mallik, M. Sayah and M. G. Organ, *Angew. Chem., Int. Ed.*, 2012, **51**, 3314–3332; (c) Y. Kim, Y. Kim, M. Y. Hur and E. Lee, *J. Organomet. Chem.*, 2016, **820**, 1–7.
- 13 (a) M. Sajid, A. Stute, A. J. Cardenas, B. J. Culotta, J. A. Hepperle, T. H. Warren, B. Schirmer, S. Grimme, A. Studer, C. G. Daniliuc, R. Frohlich, J. L. Petersen, G. Kehr and G. Erker, *J. Am. Chem. Soc.*, 2012, **134**, 10156–10168; (b) J. C. Pereira, M. Sajid, G. Kehr, A. M. Wright, B. Schirmer, Z. W. Qu, S. Grimme, G. Erker and P. C. Ford, *J. Am. Chem. Soc.*, 2014, **136**, 513–519.
- 14 X. Tao, G. Kehr, X. Wang, C. G. Daniliuc, S. Grimme and G. Erker, *Chem. – Eur. J.*, 2016, **22**, 9504–9507.