




# INORGANIC CHEMISTRY

## FRONTIERS





## RESEARCH ARTICLE



Cite this: *Inorg. Chem. Front.*, 2016, **3**, 523

# Thioether-terminated nickel(II) coordination clusters with {Ni<sub>6</sub>} horseshoe- and {Ni<sub>8</sub>} rollercoaster-shaped cores†

Sebastian Schmitz,<sup>a</sup> Jan van Leusen,<sup>a</sup> Arkady Ellern,<sup>b</sup> Paul Kögerler<sup>\*a,c</sup> and Kirill Yu. Monakhov<sup>\*a</sup>

We report two polynuclear nickel(II) compounds whose supramolecular structures are controlled by small inorganic templating anions and  $\pi$ -conjugated Schiff-base ligands (L-SMe<sup>3-</sup> and HL-SMe<sup>2-</sup>) with peripheral, structurally exposed methylthioether groups. The central component of the compound [Ni<sub>6</sub>(L-SMe)<sub>3</sub>(CO<sub>3</sub>)(MeOH)<sub>6</sub>(THF)<sub>2</sub>]Cl·2MeOH (**1**, monoclinic space group *P2<sub>1</sub>/c*) displays a horseshoe-shaped {Ni<sub>6</sub>} core templated by  $\mu_6$ -CO<sub>3</sub><sup>2-</sup> moiety due to the fixation of atmospheric carbon dioxide. According to thermogravimetric analysis, compound **1** retains the structural integrity of its carbonate-bonded metal core [Ni<sub>6</sub>(L-SMe)<sub>3</sub>(CO<sub>3</sub>)]<sup>+</sup> up to ca. 390 °C. In the crystal lattice, compound **1** features an one-dimensional hydrogen-bonded chain structure [(CO<sub>3</sub> ⊂ Ni<sub>6</sub>)...Cl...{CO<sub>3</sub> ⊂ Ni<sub>6</sub>}]<sub>∞</sub>. The structure of the compound [Ni<sub>8</sub>(HL-SMe)<sub>2</sub>(L-SMe)<sub>2</sub>(OH)<sub>4</sub>(MeCN)<sub>4</sub>(H<sub>2</sub>O)<sub>4</sub>](NO<sub>3</sub>)<sub>2</sub>·11MeCN·2PhCN (**2**, triclinic space group *P1̄*) consists of a dicationic crown-like {Ni<sub>8</sub>} metallamacrocycle hosting two NO<sub>3</sub><sup>-</sup> anions in the upper and lower cavities. The {Ni<sub>6</sub>} complex is characterised by antiferromagnetic exchange interactions, whereas {Ni<sub>8</sub>} reveals predominantly ferromagnetic exchange coupling between the spin-1 Ni(II) centres. The presence of terminal thioether anchoring groups at the periphery of complex **1**, coupled with high thermal stability and solubility in several common organic solvents, renders this compound an interesting candidate for molecular surface-deposition from solution using electrospray ionisation mass spectrometry approach and, possibly, for supported-transition metal complex heterogeneous catalysis.

Received 5th December 2015,

Accepted 12th January 2016

DOI: 10.1039/c5qi00278h

rsc.li/frontiers-inorganic

## Introduction

Fabrication of nanomaterials<sup>1</sup> with well-defined morphology and properties derived from the contact of individual coordination complexes and clusters of magnetic, electrochemical, and catalytic interest with a metallic substrate (electrode) attracts great attention in modern surface chemistry<sup>2</sup> and physics<sup>3</sup> and is motivated by the continual search for multi-

functional, compact, and biocompatible structures for renewable and sustainable energy. The peripheral functionalisation of stimuli-responsive metalorganic compounds with the electron-rich “anchoring” groups<sup>4</sup> such as *e.g.* cyclic or acyclic thioether (–S–), 1,2-dithiolane (–S–S–) or thiol (–SH) constitutes one out of multiple chemical strategies<sup>5</sup> to anchor molecules to macroscopic solid-state electrodes<sup>6</sup> (*e.g.*, gold). The ligand environment of the metal centres as well as the structural, thermal and redox stability, solubility (for solution-driven molecular deposition), and volatility/sublimability (for deposition from the solid phase) of molecules are the key features in multilateral surface-oriented investigation toward hybrid catalytic systems and electronic devices.

The structural, spectroscopic, electrochemical, and catalytic properties of diamagnetic and paramagnetic S-terminated transition metal complexes and clusters were extensively studied during previous years: (i) electrochemical, electrocatalytic, and/or *in situ* infrared spectroscopic properties of some cobalt, iron, ruthenium, and osmium complexes adsorbed on gold electrodes *via* the sulfur-based binding groups (thiols, thiophenes, thiocyanates, *etc.*) were reported.<sup>7,8</sup> (ii) Reactivity of a few iron and rhodium complexes toward

<sup>a</sup>Institut für Anorganische Chemie, RWTH Aachen University, Landoltweg 1, 52074 Aachen, Germany. E-mail: paul.koegerler@ac.rwth-aachen.de, kirill.monakhov@ac.rwth-aachen.de

<sup>b</sup>Chemistry Department, Iowa State University, 1711 Gilman Hall, Ames, IA 50011, USA

<sup>c</sup>Jülich-Aachen Research Alliance (JARA-FIT) and Peter Grünberg Institute (PGI-6), Forschungszentrum Jülich, 52425 Jülich, Germany

†Electronic supplementary information (ESI) available: Synthesis of the H<sub>3</sub>L-SMe ligand and its proposed oxidation pathway, depiction of the non-functionalised H<sub>3</sub>L ligand, synthesis and solid-state molecular structures of compounds **A** and **B**, IR spectra and crystal data and structure refinement details of compounds **1**, **2**, **A** and **B**, and ESI-MS spectrum of compound **1**. CCDC 1440001–1440004. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c5qi00278h



thiolate-stabilised gold nanoparticles was discussed.<sup>9</sup> (iii) The mono- and dinuclear Cr(III)<sup>10</sup> and Ni(II)<sup>11</sup> complexes bearing thioether groups at the periphery of the PNP and PNP = S chelates were exploited for the catalytic polymerisation and oligomerisation of ethylene. (iv) In view of the high interest for the selective formation of self-assembly of highly ordered and oriented metal-organic monolayers on surfaces or the integration of metal complexes into mesoporous materials,<sup>12</sup> the deposition of S-terminated heterometallic complexes and clusters of Mo and Pd or Pt was studied on gold surfaces by scanning tunnelling microscopy (STM), ellipsometry, and X-ray photoelectron spectroscopy (XPS).<sup>13</sup> (v) The tetrahedral mixed-metal {RuCo<sub>3</sub>} carbonyl clusters functionalised with the phosphane-thiol ligands were studied on a gold surface by STM, XPS and X-ray absorption spectroscopy.<sup>14</sup> (vi) The chemical, structural, and thermodynamic characterisation of the mononuclear Pt(II) *cis*-complexes and the heterotrimeric {Co<sub>2</sub>Pt} clusters containing *N*-substituted bis(diphenylphosphanyl) amine ligands with the pendant thiobenzyl ester groups<sup>15</sup> as well as a trinuclear Ag(I) complex and a Ag(I) coordination polymer with a thioether function in their short-bite diphosphine ligands<sup>16</sup> were described. (vii) Sulfur-assisted phenyl migration from phosphorus to platinum in molecular {PtW<sub>2</sub>} and {PtMo<sub>2</sub>} clusters with thioether-functionalised short-bite ligands of the bis(diphenylphosphanyl)amine-type was demonstrated.<sup>17</sup> The synthesis and structural chemistry of several methylthio- or thiocyanate-furnished coordination complexes of Mn(II,III),<sup>18</sup> Fe(II),<sup>19</sup> and Fe(III)<sup>20</sup> which exhibit ferromagnetic and antiferromagnetic exchange interactions were also reported. To the best of our knowledge, the potential of the above-mentioned metal complexes in catalytic processes on substrate surfaces has not been explored so far.

By contrast, surface studies towards nano and quantum computation which used the thioether-functionalised molecules exhibiting slow-relaxation of magnetisation (single-molecule magnets, SMMs) were performed and dealt with the {Mn<sub>8</sub>Mn<sub>4</sub>IV},<sup>21</sup> {Fe<sub>4</sub>III}<sup>22</sup> (see also gold nanoparticles functionalised with a 1,2-dithiolane-decorated Fe<sub>4</sub>III SMM<sup>23</sup>), {Fe<sub>3</sub>III-Cr<sub>3</sub>III},<sup>24</sup> and {Ni<sub>4</sub>III}<sup>25</sup> complexes. The chemical and physical behaviour of the {Cr<sub>7</sub>III-Ni<sub>1</sub>III}-nuclearity rings<sup>26</sup> (with antiferromagnetic coupling in bulk) and the {Mn<sub>12</sub>III-Mn<sub>7</sub>III}-nuclearity compound<sup>27</sup> (with ferromagnetic coupling in bulk) upon deposition and anchoring to gold surfaces *via* sulfur bonds was assessed.

Our exploratory work is motivated by the potential use of isolated transition metal coordination complexes and clusters with the structurally exposed sulfur-based anchoring groups in industrially relevant catalytic processes on solid supports at low and high temperatures. Our strategy implies the synthesis and characterisation of polynuclear metal compounds with the functional nano-sized structures as well as their addressing from the perspectives of homogeneous catalysis, with the subsequent immobilisation of these molecules onto porous metallic substrates or noble metal-coated mesoporous silica nanoparticles<sup>28</sup> (*e.g.*, SBA-15) to eventually enable supported-transition metal complex heterogeneous (or heterogenised homogeneous<sup>29</sup>) catalysis. In particular, Ni(II) Schiff-base complexes,<sup>30</sup> used as catalysts in the hydrogenation (see, *e.g.* reduction of benzene<sup>31</sup>) and oxidation reactions as well as for the ethylene polymerisation,<sup>32</sup> constitute interesting target candidates for the above-mentioned functionalisation and application.

Herein, we report the preparation of two novel polynuclear nickel(II) complexes supported by a  $\pi$ -conjugated Schiff-base ligand with the chemically engineered -SMe functionalities



**Paul Kögerler**

at RWTH Aachen University and Group Leader for Molecular Magnetism at the Peter Grünberg Institute, Research Centre Jülich.

*Paul Kögerler graduated with a Dr rer. nat. degree with Prof. Achim Müller at the University of Bielefeld (Germany) in 2000, followed by a postdoctoral research stay at the Department of Physics and Astronomy at Iowa State University (USA). In 2003, he was appointed as a tenured Associate Scientist at the U.S. DOE Ames Laboratory, before returning to Germany in 2006 as Professor of Chemistry at the Institute of Inorganic Chemistry*




**Kirill Yu. Monakhov**

to join the group of Prof. Paul Kögerler at RWTH Aachen University. In 2015 he received a DFG Emmy Noether fellowship and now leads a junior research group at the Institute of Inorganic Chemistry at RWTH Aachen. His research interests include coordination and supramolecular chemistry, redox-active magnetic molecules, molecular electronics and spintronics, computational chemistry, catalysis and surface science.

*Kirill Monakhov received his Dr rer. nat. degree with Prof. Gerald Linti at Heidelberg University (Germany) in 2010. After two years as a postdoctoral fellow of the German Research Foundation (DFG) and Cercle Gutenberg with Prof. Pierre Braunstein (University of Strasbourg, France), and after being awarded the Academia Europaea Burgen Scholarship in 2011, he returned to Germany in 2013 with a DFG postdoctoral reintegration fellowship*





**Fig. 1** The hexadentate Schiff base  $\text{H}_3\text{L-SMe}$  ( $=\text{C}_{21}\text{H}_{18}\text{N}_2\text{O}_3\text{S}$ ) used in this work.

(see  $\text{H}_3\text{L-SMe}$  in Fig. 1), isolated as  $[\text{Ni}_6(\text{L-SMe})_3(\text{CO}_3)(\text{MeOH})_6(\text{THF})_2]\text{Cl}\cdot 2\text{MeOH}$  (**1**) and  $[\text{Ni}_8(\text{HL-SMe})_2(\text{L-SMe})_2(\text{OH})_4(\text{MeCN})_4(\text{H}_2\text{O})_4](\text{NO}_3)_2\cdot 11\text{MeCN}\cdot 2\text{PhCN}$  (**2**), which are of relevance to *in situ* ultra-high vacuum deposition on surfaces by electrospray ionisation<sup>33</sup> (ESI). The obtained ionic compounds were characterised by infrared (IR) spectroscopy, ESI mass spectrometry (ESI-MS), and thermogravimetric analysis (TGA) and their solid-state molecular structures were determined by single-crystal X-ray diffraction. The magnetic properties were assessed *via* dc and ac magnetic susceptibility measurements and our study thus expands knowledge on existing topologies and magnetism in nickel coordination chemistry.<sup>34</sup>

## Results and discussion


### Synthesis and thermal stability

Compounds **1** and **2** were synthesised *via* one-pot four-component *in situ* reactions of  $n$  equivalents of  $\text{NiX}_2\cdot 6\text{H}_2\text{O}$  [ $n = 2$  for  $\text{X} = \text{Cl}$  (**1**) and  $n = 1.4$  for  $\text{X} = \text{NO}_3$  (**2**)] with two equivalents of 2-aminophenol, one equivalent of freshly prepared 2-hydroxy-5-(methylthio)isophthalaldehyde and two equivalents of triethylamine under reflux conditions in a mixture of organic solvents (Scheme 1). Note that upon increasing  $n$  to 2 in the synthesis reaction for compound **2**, no solid product could be isolated. Likewise, changing or replacing the counteranions in the Ni(II) precursors also prevents precipitation of crystalline product. A methanol/tetrahydrofuran volume ratio of 4 : 1 was used for the formation of the moisture stable, crys-

talline compound  $[\text{Ni}_6(\text{L-SMe})_3(\text{CO}_3)(\text{MeOH})_6(\text{THF})_2]\text{Cl}\cdot 2\text{MeOH}$  (**1**) and a acetonitrile/benzonitrile volume ratio of 10 : 1 for the moisture sensitive, crystalline compound  $[\text{Ni}_8(\text{HL-SMe})_2(\text{L-SMe})_2(\text{OH})_4(\text{MeCN})_4(\text{H}_2\text{O})_4](\text{NO}_3)_2\cdot 11\text{MeCN}\cdot 2\text{PhCN}$  (**2**). The aerobic reaction affording compound **1** is furthermore characterised by the uptake of atmospheric  $\text{CO}_2$ . Dark-red compounds **1** and **2** with a potentially redox-active, fully-deprotonated  $\text{L-SMe}^{3-}$  ligand (see Fig. S1 and synthesis of  $\text{H}_3\text{L-SMe}$  in the ESI†) were obtained in *ca.* 39% and 55% yields, respectively. Compound **1** is well soluble in THF, DMF, and MeCN and modestly soluble in MeOH and  $\text{CH}_2\text{Cl}_2$ , while compound **2** is well soluble in THF, DMF, and MeOH and moderately soluble in MeCN and  $\text{CH}_2\text{Cl}_2$ .

In contrast to the gradual decomposition in TGA experiments of compound **2** occurring as an ill-defined mass loss step between 40 and 800 °C, the thermal stability of compound **1** in a  $\text{N}_2$  atmosphere is characterised by three distinctive steps: (i) the first step (40–210 °C) corresponds to a 13.8% mass loss of two uncoordinated MeOH solvent molecules (the compound was measured immediately following synthesis) as well as two THF and two MeOH ligands coordinated to the Ni1/Ni'1 and Ni2/Ni'2 atoms, respectively ( $\Delta m_{\text{calc}} = 13.8\%$ ). (ii) The second step shows a 6.6% mass loss of four remaining MeOH ligands coordinated to the Ni3/Ni'3 atoms ( $\Delta m_{\text{calc}} = 6.5\%$ ) between 210 °C and 390 °C. (iii) The third, small step (1.8% mass loss from 390 °C to 400 °C) indicates chloride counterion desorption ( $\Delta m_{\text{calc}} = 1.8\%$ ). Thus, the TGA suggests that the  $[\text{Ni}_6(\text{L-SMe})_3(\text{CO}_3)]^+$  core fragment is stable up to *ca.* 390 °C and this high thermal stability might enable the use of this magnetic complex in high-temperature catalytic reactions<sup>32</sup> on solid supports. It is noteworthy that the molecular ion peak of  $[\text{Ni}_6(\text{L-SMe})_3(\text{CO}_3)\text{H}_2]^+$  without coordinated solvent molecules could be detected by ESI mass spectrometry (Fig. S7†).

Interestingly, the use of 2-hydroxy-5-methylisophthalaldehyde (see Fig. S2† for the structure of the non-functionalised  $\text{H}_3\text{L}$  ligand) instead of its methylthio derivate in the synthesis under less basic conditions (in the absence of  $\text{Et}_3\text{N}$ ; Scheme S1†) has resulted in the formation of mononuclear nickel(II) Schiff-base complexes with compositions  $[\text{Ni}(\text{H}_3\text{L})_2]\text{Cl}_2\cdot 4\text{MeOH}$  (**A**) and  $[\text{Ni}(\text{H}_3\text{L})_2](\text{NO}_3)_2\cdot 4\text{MeOH}$  (**B**), see



**Scheme 1** Synthesis of the hexanuclear and octanuclear nickel(II) Schiff-base complex compounds **1** and **2**. Reaction conditions: (i)  $n = 2$  for  $\text{X} = \text{Cl}$  and  $\text{MeOH} : \text{THF} = 4 : 1$ ; (ii)  $n = 1.4$  for  $\text{X} = \text{NO}_3$  and  $\text{MeCN} : \text{PhCN} = 10 : 1$ . For the IR spectra of these compounds, see Fig. S3 and S4.†




Fig. S8 and Tables S5–S8† for solid-state structure information. The sulfur ligand constituents and the presence of triethylamine thus appear to be crucial for the synthesis of compounds **1** and **2**.

### X-ray crystal structure analysis


The solid-state molecular structures of compounds **1** and **2** were established by single-crystal X-ray diffraction analyses and are illustrated in Fig. 2–4. Compound **1** crystallises in the monoclinic space group  $P2_1/c$  (Table S1, the ESI†) and consists of the cationic complex  $[\text{Ni}_6(\text{L-SMe})_3(\text{CO}_3)(\text{MeOH})_6(\text{THF})_2]^+$  (Fig. 2a), counterbalanced by a  $\text{Cl}^-$  anion, and two crystal solvent (methanol) molecules. The nano-sized  $[\text{Ni}_6(\text{L-SMe})_3(\text{CO}_3)(\text{MeOH})_6(\text{THF})_2]^+$  complex contains a distorted horse-shoe-shaped  $\text{Ni}_6$  core structure. The Ni atoms are connected to a  $\mu_6\text{-CO}_3^{2-}$  group, a result of the fixation of atmospheric carbon dioxide (Fig. 2b). Molecular “horseshoe”-like structures have also been observed for  $\{\text{Cr}_6^{\text{III}}\}$ -nuclearity complexes.<sup>35</sup> Sorption of atmospheric  $\text{CO}_2$  under aerobic reaction conditions is well-known in coordination chemistry;<sup>36</sup> homonuclear and heteronuclear transition-metal and lanthanide complexes that incorporate the carbonate ion in a wide variety of bridging modes have been described.<sup>37</sup>

The  $\{\text{CO}_3 \subset \text{Ni}_6\}$  fragment in compound **1** is ligated by three *in situ*-formed Schiff bases (threefold-deprotonated  $\text{L-SMe}^{3-}$ ; see Fig. 1) acting as  $\text{N}_2\text{O}_3$ -chelates for the divalent nickel centres. Six MeOH and two THF solvent molecules complete the coordination spheres of metal ions (Fig. 3):  $\text{Ni}1$  or  $\text{Ni}'1$  site binds to one THF molecule,  $\text{Ni}2$  or  $\text{Ni}'2$  to one terminal MeOH molecule, and  $\text{Ni}3$  or  $\text{Ni}'3$  to two terminal MeOH molecules. Bond distances of the  $\text{NO}_5$ -coordinated nickel ions (in Å):  $\text{Ni}1\text{-O}_{\text{L-SMe}}$ : 2.0010(19)–2.158(2);  $\text{Ni}1\text{-N}_{\text{L-SMe}}$ : 1.976(2);  $\text{Ni}1\text{-O}_{\text{THF}}$ : 2.098(2);  $\text{Ni}1\text{-O}_{\text{CO}_3}$ : 2.0896(19);  $\text{Ni}2\text{-O}_{\text{L-SMe}}$ : 1.9726(18)–2.143(2);  $\text{Ni}2\text{-N}_{\text{L-SMe}}$ : 1.963(2);  $\text{Ni}2\text{-O}_{\text{MeOH}}$ : 2.119(2);  $\text{Ni}2\text{-O}_{\text{CO}_3}$ : 2.0306(18);  $\text{Ni}3\text{-O}_{\text{L-SMe}}$ : 2.000(2) and 2.0359(19);  $\text{Ni}3\text{-N}_{\text{L-SMe}}$ : 1.995(2);  $\text{Ni}3\text{-O}_{\text{MeOH}}$ : 2.095(2) and 2.112(2);  $\text{Ni}3\text{-O}_{\text{CO}_3}$ : 2.0856(19). Thus, each  $\text{Ni(II)}$  centre in compound **1** exhibits a pseudo-octahedral local geometry with axially distorted O positions. The  $[\text{Ni}_6(\text{L-SMe})_3(\text{CO}_3)(\text{MeOH})_6(\text{THF})_2]^+$  structure reveals the non-bonding  $\text{Ni}1\cdots\text{Ni}2$  (or  $\text{Ni}'1\cdots\text{Ni}'2$ ),  $\text{Ni}1\cdots\text{Ni}3$  (or  $\text{Ni}'1\cdots\text{Ni}'3$ ), and  $\text{Ni}2\cdots\text{Ni}'2$  separations of 2.9707(6), 3.1251(6), and 3.0323(7) Å, respectively (the sum of the covalent radii of two Ni atoms is 2.48 Å (ref. 38)). The magnetically relevant bond angles fall in the ranges  $90.313(74)\text{--}90.828(78)^\circ$  for  $\text{Ni}1\text{--}(\mu_3\text{-O})\text{--Ni}2$ ,  $96.916(79)\text{--}101.451(84)^\circ$  for  $\text{Ni}1\text{--}(\mu_3\text{-O})\text{--Ni}3$ , and  $96.608(17)\text{--}100.461(17)^\circ$  for  $\text{Ni}2\text{--}(\mu_3\text{-O})\text{--Ni}'2$ .




**Fig. 2** (a) Molecular structure of  $[\text{Ni}_6(\text{L-SMe})_3(\text{CO}_3)(\text{MeOH})_6(\text{THF})_2]^+$  in compound **1**. (b) Different views of the  $\text{Ni}_6$  core templated by  $\text{CO}_3^{2-}$  with a  $\mu_6\text{-}\eta^1\text{:}\eta^1\text{:}\eta^1\text{:}\eta^1\text{:}\eta^1\text{:}\eta^1$  binding mode. (c) A segment of the extended solid-state structure of compound **1**, highlighting the association of two  $\{\text{Ni}_6\}$  coordination clusters via  $\text{C-H}_{\text{L-SMe}}\cdots\text{Cl}$  hydrogen bonding interactions ( $\text{O-H}_{\text{MeOH}}\cdots\text{Cl}$  interactions are also shown). Hydrogen atoms not involved in hydrogen bonds are omitted for clarity. Colour code: H, black; C, grey; N, blue; O, red; S, yellow; Cl, light green; Ni, dark green. The nickel and sulfur atoms, carbonate ion, and co-crystallised MeOH molecules are represented as ball-and-stick models. The chloride ion is depicted as space-filling model. See Table S2† for bond lengths and angles.





**Fig. 3** A fragment of compound **1** representing pseudo-octahedral coordination geometries of the Ni(II) centres. For clarity, only one chelating Schiff-base ligand L-SMe<sup>3-</sup> is explicitly depicted. The nickel, nitrogen, oxygen, and sulfur atoms are represented as ball-and-stick models. The terminal-coordinated solvent molecules are highlighted in different colours: MeOH, purple; THF, sky blue. Elemental colour code: C, grey; N, blue; O, red; S, yellow; Ni, dark green.



**Fig. 4** (a) Molecular structure of  $[\text{Ni}_8(\text{HL-SMe})_2(\text{L-SMe})_2(\text{OH})_4(\text{MeCN})_4(\text{H}_2\text{O})_4]^{2+}$  in compound **2**. (b) Non-planar ring of eight nickel(II) atoms as metal core in compound **2**. (c) Side view of the solid-state structure of compound **2** showing two  $\text{NO}_3^-$  anions in the top and down cavities. Hydrogen atoms and solvent molecules are omitted for clarity. The nickel and sulfur atoms as well as the C and N atoms of MeCN in (c) are represented as ball-and-stick models and the nitrate ions as space-filling models. Colour code: C, grey; N, blue; O, red; S, yellow; Ni, dark green. See Table S4† for bond lengths and angles.

A further interesting feature of compound **1** is the infinite one-dimensional chain structure in the crystal lattice, made up from repeating  $[\text{Ni}_6(\text{L-SMe})_3(\text{CO}_3)(\text{MeOH})_6(\text{THF})_2]^+$  entites assembled *via* four  $\text{C-H}_{\text{L-SMe}} \cdots \text{Cl}$  ( $d_{\text{C} \cdots \text{Cl}} = 3.6385(4)$  Å and  $3.7100(4)$  Å) and two identical  $\text{O-H}_{\text{MeOH}} \cdots \text{Cl}$  ( $d_{\text{O} \cdots \text{Cl}} = 3.0958(3)$  Å) hydrogen bonds, as shown in Fig. 2c.

Compound **2** crystallises in the triclinic space group  $P\bar{1}$  (Table S3†) and shows a dicationic  $\{\text{Ni}_8\}$  metallamacrocycle  $[\text{Ni}_8(\text{HL-SMe})_2(\text{L-SMe})_2(\text{OH})_4(\text{MeCN})_4(\text{H}_2\text{O})_4]^{2+}$  (Fig. 4a), reminiscent of a rollercoaster, and counterbalanced by two  $\text{NO}_3^-$  anions. Eleven acetonitrile and two benzonitrile molecules co-crystallise in the unit cell. The central metal core in the nano-sized  $[\text{Ni}_8(\text{HL-SMe})_2(\text{L-SMe})_2(\text{OH})_4(\text{MeCN})_4(\text{H}_2\text{O})_4]^{2+}$  moiety displays a non-planar ring of eight nickel(II) atoms (Fig. 4b). This ring is maintained by two monoprotonated  $\text{HL-SMe}^{2-}$  and two fully-deprotonated  $\text{L-SMe}^{3-}$  Schiff-base ligands (see Fig. 1) as well as four bridging  $\text{OH}^-$  groups. The coordination spheres of four nickel (Ni2, Ni3, Ni6 and Ni8) atoms are completed by terminal MeCN molecules, those of the other four (Ni1, Ni4, Ni5 and Ni7) only by terminal  $\text{H}_2\text{O}$  molecules (one MeCN/ $\text{H}_2\text{O}$  per Ni). As a result, each of the Ni(II) ions in the ring resides in a pseudo-octahedral  $\text{N}_2\text{O}_4$  or  $\text{NO}_5$  coordination environment (ratio  $\text{N}_2\text{O}_4 : \text{NO}_5 = 4 : 4$ ). The  $\text{NiN}_2\text{O}_4$  bond distances (in Å) amount to Ni2- $\text{O}_{\text{L-SMe}}$ : 1.990(7)–2.151(7); Ni2- $\text{N}_{\text{L-SMe}}$ : 1.980(8); Ni2-OH: 2.045(7), Ni2- $\text{N}_{\text{MeCN}}$ : 2.091(8); Ni3- $\text{O}_{\text{L-SMe}}$ : 1.973(7)–2.171(7); Ni3- $\text{N}_{\text{L-SMe}}$ : 1.992(9), Ni3-OH: 2.061(7); Ni3- $\text{N}_{\text{MeCN}}$ : 2.075(10); Ni6- $\text{O}_{\text{L-SMe}}$ : 1.978(7)–2.162(7), Ni6- $\text{N}_{\text{L-SMe}}$ : 1.992(8); Ni6-OH: 2.029(7); Ni6- $\text{N}_{\text{MeCN}}$ : 2.112(9); Ni8- $\text{O}_{\text{L-SMe}}$ : 1.996(7)–2.141(7); Ni8- $\text{N}_{\text{L-SMe}}$ : 1.977(9), Ni8-OH: 2.056(7), Ni8- $\text{N}_{\text{MeCN}}$ : 2.080(10). For the  $\text{NiNO}_5$  environment: Ni1- $\text{O}_{\text{L-SMe}}$ : 2.027(7)–2.087(7); Ni1- $\text{N}_{\text{L-SMe}}$ : 2.013(8); Ni1-OH: 2.093(7), Ni1- $\text{O}_{\text{H}_2\text{O}}$ : 2.071(7); Ni4- $\text{O}_{\text{L-SMe}}$ : 2.039(7)–2.077(7); Ni4- $\text{N}_{\text{L-SMe}}$ : 2.018(9); Ni4-OH: 2.138(6); Ni4- $\text{O}_{\text{H}_2\text{O}}$ : 2.059(7); Ni5- $\text{O}_{\text{L-SMe}}$ : 2.038(7)–2.054(7); Ni5- $\text{N}_{\text{L-SMe}}$ : 2.010(8), Ni5-OH: 2.133(7); Ni5- $\text{O}_{\text{H}_2\text{O}}$ : 2.071(7); Ni7- $\text{O}_{\text{L-SMe}}$ : 2.043(7)–2.063(7); Ni7- $\text{N}_{\text{L-SMe}}$ : 1.991(8); Ni7-OH: 2.116(7); Ni7- $\text{O}_{\text{H}_2\text{O}}$ : 2.080(8). The non-equivalent, non-bonding Ni $\cdots$ Ni distances range from 3.0662(25) to 3.1001(25) Å. The magnetically relevant bond angles fall in the ranges 93.5(3)–100.7(3)° for Ni3-( $\mu_3$ - $\text{O}_{2\text{O}_{\text{L-SMe}}}$ )-Ni8 and Ni3-( $\mu_3$ - $\text{O}_{12\text{L-SMe}}$ )-Ni7, and 95.1(3)–96.6(3)° for Ni2-( $\mu_2$ - $\text{O}_{14\text{OH}}$ )-Ni5 and Ni1-( $\mu_2$ - $\text{O}_{15\text{OH}}$ )-Ni6.

In general, the  $[\text{Ni}_8(\text{HL-SMe})_2(\text{L-SMe})_2(\text{OH})_4(\text{MeCN})_4(\text{H}_2\text{O})_4]^{2+}$  architecture can be viewed as a metallocavitand<sup>39</sup> hosting two  $\text{NO}_3^-$  anions in a nesting fashion (Fig. 4c). These nitrate groups are situated 2.3973(80) Å above and 2.6004(83) Å below the nickel ring plane and connected to the internal oxygen atoms of the dicationic complex *via* four H-bonds: (i) O21 or O26 of  $\text{NO}_3^-$  is linked to two bridging  $\text{OH}^-$  groups (O21- $\text{O}_{\text{OH}}$ : 2.7958(123) Å–2.8692(117) Å; O26- $\text{O}_{\text{OH}}$ : 2.9280(101) Å–2.9428(123) Å) and one coordinated  $\text{H}_2\text{O}$  molecule (O21- $\text{O}_{\text{H}_2\text{O}}$ : 3.0352(115) Å; O26- $\text{O}_{\text{H}_2\text{O}}$ : 2.9195(145) Å). (ii) The fourth H-bond appears between O22 or O25 of  $\text{NO}_3^-$  and the  $\text{H}_2\text{O}$  ligands at the Ni1 and Ni4 centres (O22- $\text{O}_{\text{H}_2\text{O}}$ : 2.7542(125) Å; O25- $\text{O}_{\text{H}_2\text{O}}$ : 2.8609(123) Å).

We note that in the crystal structures of compounds **1** and **2**, the S atoms at the periphery of the Schiff-base ligands



L-SMe<sup>3-</sup> and HL-SMe<sup>2-</sup> are not involved in any intermolecular coordination bonds.

### Magnetism and magnetochemical modelling

Magnetic properties of compounds **1** and **2** were studied based on the temperature- and field-dependent dc (static) and ac (dynamic) magnetic susceptibility. Since the ac susceptibilities do not reveal any out-of-phase components for compounds **1** nor **2**, the experimental data of the magnetic dc measurements are presented as  $\chi_m T$  vs.  $T$  and  $M_m$  vs.  $B$  curves in Fig. 5. For compound **1**, a  $\chi_m T$  value of 7.68 cm<sup>3</sup> K mol<sup>-1</sup> at 290 K is within the interval of 5.88–9.19 cm<sup>3</sup> K mol<sup>-1</sup> expected for six non-interacting high-spin Ni(II) centres.<sup>40</sup> Lowering the temperature,  $\chi_m T$  slowly decreases for  $T > 100$  K and rapidly decreases below that point indicating predominantly antiferromagnetic exchange interactions between the Ni(II) centres. For compound **2**, the corresponding value at 290 K is 9.29 cm<sup>3</sup> K mol<sup>-1</sup> being in the range expected for eight non-interacting high-spin Ni(II) centres (7.84–12.25 cm<sup>3</sup> K mol<sup>-1</sup>).<sup>40</sup> In contrast to compound **1**,  $\chi_m T$  slowly increases by decreasing the temperature, reaching a maximum at  $T \approx 30$  K. The  $\chi_m T$  value rapidly decreases on further temperature decrease. Such a maximum results from predominantly ferromagnetic exchange interactions between the eight Ni(II) centres. Since both magnetisation curves of compounds **1** and **2** do not exhibit saturation, a reliable conclusion with respect to the upper limit of total effective spin ground state  $S_{\text{tot}}$  is impossible.

A model description of the magnetic susceptibilities was based on the following assumptions, in order to contain computational limitations and to restrict the number of independent fitting parameters: Since the ligand field of each Ni(II) centre is well approximated as octahedrally symmetric, the ground term is an orbital momentum singlet <sup>3</sup>A<sub>2</sub> originating from ground term <sup>3</sup>F of the free high-spin Ni(II) ion – according to the crystal structure, the distortion of the octahedral

symmetry is expected to be small. Therefore, the zero-field splitting is expected to be negligible and thus omitted. Hence, the magnetic features of compounds **1** and **2** are described in terms of an isotropic effective spin approach ( $S = 1$ ) including an effective Heisenberg-Hamiltonian to characterise the exchange interactions. The number of exchange parameters is restricted in a further approximation: exchange pathways exhibiting similar geometries are assumed as identical, see Fig. 6 for the resulting coupling schemes.

The exchange interaction Hamiltonian for compound **1**

$$\hat{H}_{\text{ex},1} = -2J_A(\hat{S}_1\hat{S}_2 + \hat{S}_5\hat{S}_6) - 2J_B(\hat{S}_2\hat{S}_3 + \hat{S}_4\hat{S}_5) - 2J_C\hat{S}_3\hat{S}_4$$

and accordingly for compound **2**

$$\begin{aligned} \hat{H}_{\text{ex},2} = & -2J_A(\hat{S}_1\hat{S}_2 + \hat{S}_5\hat{S}_6) - 2J_B(\hat{S}_2\hat{S}_3 + \hat{S}_4\hat{S}_5) \\ & - 2J_C(\hat{S}_6\hat{S}_7 + \hat{S}_1\hat{S}_8) - 2J_D\hat{S}_7\hat{S}_8 - 2J_E\hat{S}_3\hat{S}_4 \end{aligned}$$

in combination with the Zeeman Hamiltonian are employed to model the experimental data within the computational framework CONDON 2.0.<sup>41</sup> The least-squares fit to the magnetic data of compound **1** (quality-of-fit parameter SQ = 0.9%) results in the exchange parameters  $J_A = -7.7 \pm 0.1$  cm<sup>-1</sup>,  $J_B = +7.2 \pm 0.1$  cm<sup>-1</sup>,  $J_C = -2.6 \pm 0.1$  cm<sup>-1</sup>, and an isotropic  $g$ -value of  $g_{\text{iso}} = 2.25 \pm 0.01$ . These parameters confirm the predominantly antiferromagnetic exchange interaction between the six Ni(II) centres. The magnitudes of the exchange interaction energies reveal weak interactions within the compound that result in a ground state of a total effective spin  $S_{\text{tot}} = 0$ .

The least-squares fit to the magnetic susceptibility data of compound **2** is slightly worse (SQ = 1.5%) if a model including three independent exchange coupling parameters ( $J_A = J_D = -1.6 \pm 0.1$  cm<sup>-1</sup>,  $J_B = -1.2 \pm 0.1$  cm<sup>-1</sup>,  $J_C = J_E = +5.2 \pm 0.1$  cm<sup>-1</sup>,  $g_{\text{iso}} = 2.13 \pm 0.02$ ) is employed. A slightly better goodness-of-fit is achieved (SQ = 1.2%) in the approximation shown in Fig. 6 based on the less-symmetric structure of the compound, although the margin of error of the fit parameter  $J_E$  is larger. The corresponding least-squares parameters are  $J_A = -1.5 \pm 0.2$  cm<sup>-1</sup>,  $J_B = -1.1 \pm 0.1$  cm<sup>-1</sup>,  $J_C = +4.6 \pm 0.1$  cm<sup>-1</sup>,  $J_D = -1.6 \pm 0.1$  cm<sup>-1</sup>,  $J_E = +4.9 \pm 1.7$  cm<sup>-1</sup> and  $g_{\text{iso}} = 2.13 \pm 0.02$ . Even though the calculated values exhibit weaker net exchange interactions in compound **2** than in compound **1**, they confirm the predominantly ferromagnetic interaction between the eight Ni(II) centres. According to the calculated parameters,




Fig. 5 Temperature dependence of  $\chi_m T$  at 0.1 Tesla for compounds **1** and **2**; inset: field dependence of the molar magnetisation  $M_m$  at 2 K; circles: experimental data, solid graphs: least-squares fits.




Fig. 6 Coupling schemes used for the magnetochemical modelling for compounds **1** and **2**.



the ground state of compound **2** is characterised by a total effective spin  $S_{\text{tot}} = 2$ .

## Conclusion

We synthesised and characterised two polynuclear nickel(II) compounds **1** and **2** with structurally exposed thioether functional groups in the periphery of the cationic complexes  $[\text{Ni}_6(\text{L-SMe})_3(\text{CO}_3)(\text{MeOH})_6(\text{THF})_2]^+$  and  $[\text{Ni}_8(\text{HL-SMe})_2(\text{L-SMe})_2(\text{OH})_4(\text{MeCN})_4(\text{H}_2\text{O})_4]^{2+}$ , respectively. These coordination clusters with distinct nickel topologies are characterised by different total ground-state spins and exhibit different types of exchange interactions: predominantly antiferromagnetic for the horseshoe-like  $\{\text{Ni}_6\}$  with  $S_{\text{tot}} = 0$  and predominantly ferromagnetic for the rollercoaster-like  $\{\text{Ni}_8\}$  with  $S_{\text{tot}} = 2$ . Due to specific chemical properties (thermal stability, solubility and detectability following ESI-MS), the anion-templated self-assembly **1** appears to be a suitable candidate for molecular deposition and anchoring onto metallic surfaces *via* a thioether linkage and studies at solid–liquid interfaces. In forthcoming studies we will explore its potential as a heterogenised homogeneous catalyst, for example *via* the use of cation exchange<sup>42</sup> to incorporate the solvent-free  $[\text{Ni}_6(\text{L-SMe})_3(\text{CO}_3)]^+$  fragment into large-pore metal–organic frameworks. We also plan to assess compound **1** and its carbonate-bonded metal core  $[\text{Ni}_6(\text{L-SMe})_3(\text{CO}_3)]^+$  as catalysts for ethylene oligomerisation where we expect the catalytic activity and selectivity to also depend on the pendant thioether groups.<sup>11b,c,43</sup> Finally, we target the immobilisation of the horseshoe-like  $\{\text{Ni}_6\}$  complex with its structurally exposed thioether anchoring groups onto noble metal-coated mesoporous silica nanoparticles (such as SBA-15),<sup>44</sup> in order to probe industrially relevant catalytic oxidation reactions on solid supports at low and high temperatures.

## Experimental

### Materials and methods

The synthesis of compounds **1**, **2**, **A** and **B** was carried out under aerobic conditions. All starting materials were from commercial sources and used as received. Solvents were used without further purification. The IR spectra of compounds **1**, **2**, **A** and **B** were recorded on a Nicolet Avatar 360 FTIR spectrometer (KBr pellets,  $\nu = 4000\text{--}400\text{ cm}^{-1}$ ). TG/DT analyses of compounds **1** and **2** were performed under  $\text{N}_2$  flow and a heating rate of  $5\text{ K min}^{-1}$  in the temperature range  $25\text{--}800\text{ }^\circ\text{C}$  on a Mettler Toledo TGA/SDTA 851e instrument. The ESI-MS spectrum of compound **1** in the positive ion mode was recorded on a 4000 QTRAP mass spectrometer system, using the LC/LC-MS method with direct infusion.

**Synthesis of  $[\text{Ni}_6(\text{L-SMe})_3(\text{CO}_3)(\text{MeOH})_6(\text{THF})_2]\text{Cl}\cdot 2\text{MeOH}$  (**1**).**  $\text{NiCl}_2\cdot 6\text{H}_2\text{O}$  (0.095 g, 0.4 mmol), 2-hydroxy-5-(methylthio)isophthalaldehyde (0.040 g, 0.2 mmol) and 2-aminophenol (0.044 g, 0.4 mmol) were solved in a mixture of MeOH (8 mL),

THF (2 mL) and triethylamine (0.056 mL, 0.4 mmol). The resulting red coloured mixture was stirred for 90 minutes under reflux. The MeOH/THF/ $\text{Et}_3\text{N}$  solution was filtered off and the filtrate was kept in a capped vial at room temperature. Dark-red single crystals of compound **1** were obtained *via* slow evaporation after *ca.* two weeks and then washed with a small amount of ice-cold methanol. Yield of the air-dried crystals: 0.049 g (38.5% based on  $\text{H}_3\text{L}$ ).  $\text{C}_{78}\text{ClH}_{85}\text{N}_6\text{Ni}_6\text{O}_{20}\text{S}_3$  (1910.34 g  $\text{mol}^{-1}$ ; disregarding solvent). IR (KBr pellet),  $\nu_{\text{max}}/\text{cm}^{-1}$ : 3405 (br,m), 2919 (w), 1615 (m), 1586 (sh), 1524 (m), 1473 (s), 1456 (sh), 1431 (sh), 1389 (s), 1336 (sh), 1313 (m), 1256 (sh), 1187 (w), 1150 (w), 1109 (w), 1047 (m), 1035 (sh), 950 (w), 883 (w), 827 (w), 744 (m), 684 (w), 589 (w), 524 (m). MS (MeOH,  $^+\text{ESI}$ ):  $m/z$ : 1536.8 ( $\text{C}_{64}\text{H}_{47}\text{N}_6\text{Ni}_6\text{O}_{12}\text{S}_3^+$ , 100%).

**Synthesis of  $[\text{Ni}_8(\text{HL-SMe})_2(\text{L-SMe})_2(\text{OH})_4(\text{MeCN})_4(\text{H}_2\text{O})_4](\text{NO}_3)_2\cdot 11\text{MeCN}\cdot 2\text{PhCN}$  (**2**).**  $\text{Ni}(\text{NO}_3)_2\cdot 6\text{H}_2\text{O}$  (0.060 g, 0.2 mmol), 2-hydroxy-5-(methylthio)isophthalaldehyde (0.040 g, 0.2 mmol) and 2-aminophenol (0.044 g, 0.4 mmol) were solved in a mixture of acetonitrile (10 mL), benzonitrile (1 mL) and triethylamine (0.056 mL, 0.4 mmol). The resulting red coloured mixture was stirred for 70 minutes under reflux. The MeCN/PhCN/ $\text{Et}_3\text{N}$  solution was filtered off and the filtrate was kept in a capped vial at room temperature. Red single crystals of compound **2** were obtained *via* slow evaporation after *ca.* one week and then washed with a small amount of ice-cold acetonitrile. Yield of the air-dried crystals: 0.033 g (55% based on Ni).  $\text{C}_{92}\text{H}_{86}\text{N}_{14}\text{Ni}_8\text{O}_{26}\text{S}_4$  (2401.55 g  $\text{mol}^{-1}$ ; disregarding solvent). IR (KBr pellet),  $\nu_{\text{max}}/\text{cm}^{-1}$ : 3424 (br, m), 3056 (w), 2918 (w), 2224 (w), 1617 (s), 1587 (s), 1527 (s), 1478 (vs), 1457 (sh), 1434 (w), 1386 (vs), 1312 (s), 1292 (sh), 1278 (sh), 1252 (s), 1189 (w), 1150 (w), 1109 (w), 1051 (s), 953 (w), 882 (m), 827 (m), 748 (s), 687 (m), 589 (m), 549 (w), 517 (m).

### X-ray crystallography

Single-crystal X-ray diffraction data for all compounds were collected on a Bruker Apex II diffractometer with Mo  $\text{K}\alpha$  radiation ( $\lambda = 0.71073\text{ \AA}$ ) at 153 K. Crystals were mounted in a Hampton cryoloop with Paratone-N oil. Absorption corrections were applied numerically based on multifaceted crystal model using SADABS software.<sup>45</sup> A summary of the crystal data and structure refinements for compounds **1**, **2**, **A** and **B** are given in Tables S1, S3, S5 and S7 in the ESI,<sup>†</sup> respectively. Details are described in the CIF files. CCDC 1440001–1440004 contain the supplementary crystallographic data for this paper.

### Magnetic susceptibility measurements

Magnetic susceptibility data of compounds **1** and **2** were recorded using a Quantum Design MPMS-5XL SQUID magnetometer for direct current (dc) and alternating current (ac) measurements. The polycrystalline samples were compacted and immobilised into cylindrical PTFE capsules. The dc susceptibility data were acquired as a function of the field (0.1–5.0 T) and temperature (2.0–290 K). The ac susceptibility data were measured in the absence of a static bias field in the 10–1000 Hz frequency range ( $T = 2.0\text{--}50\text{ K}$ ,  $B_{\text{ac}} = 3\text{ G}$ ), but no out-of-phase signals were detected. Data were corrected for dia-



magnetic contributions from the sample holder and the compounds ( $\chi_{\text{dia}}(1) = -0.96 \times 10^{-3} \text{ cm}^3 \text{ mol}^{-1}$ ,  $\chi_{\text{dia}}(2): -1.20 \times 10^{-3} \text{ cm}^3 \text{ mol}^{-1}$ ).

## Acknowledgements

This work was supported by an ERC Starting Grant (MOLSPIN-TRON, 308051). K. Y. M. thanks the Excellence Initiative of the German federal and state governments for an RWTH Start-Up grant. We are grateful to Oliver Linnenberg for the synthesis and purification of the  $\text{H}_3\text{L-SMe}$  ligand.

## References

- 1 M. N. Rittner, *J. of Metals*, 2004, **56**, 22.
- 2 *Modern Surface Organometallic Chemistry*, ed. J.-M. Basset, R. Psaro, D. Roberto and R. Ugo, Wiley-VCH, Weinheim, 2009.
- 3 L. Bogani and W. Wernsdorfer, *Nat. Mater.*, 2008, **7**, 179.
- 4 See, e.g.: (a) I. S. Kristensen, D. J. Mowbray, K. S. Thygesen and K. W. Jacobsen, *J. Phys.: Condens. Matter*, 2008, **20**, 374101; (b) Z. Li, M. Smeu, M. A. Ratner and E. Borguet, *J. Phys. Chem. C*, 2013, **117**, 14890; (c) V. Kaliginedi, A. V. Rudnev, P. Moreno-García, M. Baghernejad, C. Huang, W. Hong and T. Wandlowski, *Phys. Chem. Chem. Phys.*, 2014, **16**, 23529. See also: (d) S. Schmitz, J. van Leusen, A. Ellern, P. Kögerler and K. Yu. Monakhov, *Inorg. Chem. Front.*, 2015, **2**, 1095.
- 5 A. Cornia, M. Mannini, P. Sainctavit and R. Sessoli, *Chem. Soc. Rev.*, 2011, **40**, 3076.
- 6 J. J. Parks, A. R. Champagne, T. A. Costi, W. W. Shum, A. N. Pasupathy, E. Neuscamman, S. Flores-Torres, P. S. Cornaglia, A. A. Aligia, C. A. Balseiro, G. K.-L. Chan, H. D. Abruña and D. C. Ralph, *Science*, 2010, **328**, 1370.
- 7 (a) T. T. T. Li, H. Y. Liu and M. J. Weaver, *J. Am. Chem. Soc.*, 1984, **106**, 1233; (b) K. I. Ozoemena, T. Nyokong and P. Westbroek, *Electroanalysis*, 2003, **15**, 1762; (c) W. Zhou, S. Ye, M. Abe, T. Nishida, K. Uosaki, M. Osawa and Y. Sasaki, *Chem. – Eur. J.*, 2005, **11**, 5040.
- 8 (a) H. O. Finklea and D. D. Hanshew, *J. Am. Chem. Soc.*, 1992, **114**, 3173; (b) M. S. Ravenscroft and H. O. Finklea, *J. Phys. Chem.*, 1994, **98**, 3843; (c) H. O. Finklea and M. S. Ravenscroft, *Isr. J. Chem.*, 1997, **37**, 179; (d) D. A. Brevnov and H. O. Finklea, *J. Electrochem. Soc.*, 2000, **147**, 3461; (e) R. M. Haddox and H. O. Finklea, *J. Phys. Chem. B*, 2004, **108**, 1694; (f) N. Tuccitto, V. Torrisi, M. Cavazzini, T. Morotti, F. Puntoriero, S. Quici, S. Campagna and A. Licciardello, *ChemPhysChem*, 2007, **8**, 227.
- 9 (a) M. Yamada and H. Nishihara, *Langmuir*, 2003, **19**, 8050; (b) M. Yamada, T. Tadera, K. Kubo and H. Nishihara, *J. Phys. Chem. B*, 2003, **107**, 3703; (c) T. Belser, M. Stoeckli and A. Pfaltz, *J. Am. Chem. Soc.*, 2005, **127**, 8720.
- 10 Z. Weng, S. Teo and T. S. A. Hor, *Dalton Trans.*, 2007, 3493.
- 11 (a) K. Song, H. Gao, F. Liu, J. Pan, L. Guo, S. Zai and Q. Wu, *Eur. J. Inorg. Chem.*, 2009, 3016; (b) A. Ghisolfi, C. Fliedel, V. Rosa, R. Pattacini, A. Thibon, K. Yu. Monakhov and P. Braunstein, *Chem. – Asian J.*, 2013, **8**, 1795; (c) A. Ghisolfi, C. Fliedel, V. Rosa, K. Yu. Monakhov and P. Braunstein, *Organometallics*, 2014, **33**, 2523.
- 12 P. Van Der Voort, C. Vercaemst, D. Schaubroeck and F. Verpoort, *Phys. Chem. Chem. Phys.*, 2008, **10**, 347.
- 13 P. Croizat, F. Müller, H. Mantz, A. Englisch, R. Welter, S. Hübner and P. Braunstein, *C. R. Chim.*, 2009, **12**, 1228.
- 14 A. Naitabdi, O. Toulemonde, J. P. Bucher, J. Rosé, P. Braunstein, R. Welter and M. Drillon, *Chem. – Eur. J.*, 2008, **14**, 2355.
- 15 V. Gallo, P. Mastroianni, C. F. Nobile, P. Braunstein and U. Englert, *Dalton Trans.*, 2006, 2342.
- 16 V. Rosa, C. Fliedel, A. Ghisolfi, R. Pattacini, T. Avilés and P. Braunstein, *Dalton Trans.*, 2013, **42**, 12109.
- 17 S. Todisco, P. Mastroianni, M. Latronico, V. Gallo, U. Englert, N. Re, F. Creati and P. Braunstein, *Inorg. Chem.*, 2015, **54**, 4777.
- 18 S. Mameri, A. M. Ako, F. Yesil, M. Hibert, Y. Lan, C. E. Anson and A. K. Powell, *Eur. J. Inorg. Chem.*, 2014, 4326.
- 19 C. Fliedel, V. Rosa, A. Falceto, P. Rosa, S. Alvarez and P. Braunstein, *Inorg. Chem.*, 2015, **54**, 6547.
- 20 A. Abhervé, J. M. Clemente-Juan, M. Clemente-León, E. Coronado, J. Boonmak and S. Youngme, *New J. Chem.*, 2014, **38**, 2105.
- 21 (a) A. Cornia, A. C. Fabretti, M. Pacchioni, L. Zobbi, D. Bonacchi, A. Caneschi, D. Gatteschi, R. Biagi, U. Del Pennino, V. De Renzi, L. Gurevich and H. S. J. Van der Zant, *Angew. Chem., Int. Ed.*, 2003, **42**, 1645; (b) L. Zobbi, M. Mannini, M. Pacchioni, G. Chastanet, D. Bonacchi, C. Zanardi, R. Biagi, U. Del Pennino, D. Gatteschi, A. Cornia and R. Sessoli, *Chem. Commun.*, 2005, 1640; (c) S. Voss, M. Burgert, M. Fonin, U. Groth and U. Rüdiger, *Dalton Trans.*, 2008, 499.
- 22 (a) M. Mannini, F. Pineider, P. Sainctavit, C. Danieli, E. Otero, C. Sciancalepore, A. M. Talarico, M.-A. Arrio, A. Cornia, D. Gatteschi and R. Sessoli, *Nat. Mater.*, 2009, **8**, 194; (b) F. Pineider, M. Mannini, C. Danieli, L. Armelao, F. M. Piras, A. Magnani, A. Cornia and R. Sessoli, *J. Mater. Chem.*, 2010, **20**, 187; (c) M. Mannini, F. Pineider, C. Danieli, F. Totti, L. Sorace, Ph. Sainctavit, M.-A. Arrio, E. Otero, L. Joly, J. Criginski Cesar, A. Cornia and R. Sessoli, *Nature*, 2010, **468**, 417; (d) M. Jesus Rodriguez-Douton, M. Mannini, L. Armelao, A.-L. Barra, E. Tancini, R. Sessoli and A. Cornia, *Chem. Commun.*, 2011, 47, 1467.
- 23 M. Perfetti, F. Pineider, L. Poggini, E. Otero, M. Mannini, L. Sorace, C. Sangregorio, A. Cornia and R. Sessoli, *Small*, 2014, **10**, 323.
- 24 M. Mannini, E. Tancini, L. Sorace, P. Sainctavit, M.-A. Arrio, Y. Qian, E. Otero, D. Chiappe, L. Margheriti, J. C. Cezar, R. Sessoli and A. Cornia, *Inorg. Chem.*, 2011, **50**, 2911.
- 25 (a) A. Ghisolfi, K. Yu. Monakhov, R. Pattacini, P. Braunstein, X. López, C. de Graaf, M. Speldrich, J. van Leusen, H. Schilder and P. Kögerler, *Dalton Trans.*, 2014, **43**, 7847;



- (b) V. Heß, F. Matthes, D. E. Bürgler, K. Yu. Monakhov, C. Besson, P. Kögerler, A. Ghisolfi, P. Braunstein and C. M. Schneider, *Surf. Sci.*, 2015, **641**, 210.
- 26 (a) V. Corradini, R. Biagi, U. del Pennino, V. De Renzi, A. Gambardella, M. Affronte, C. A. Muryn, G. A. Timco and R. E. P. Winpenny, *Inorg. Chem.*, 2007, **46**, 4937; (b) V. Corradini, F. Moro, R. Biagi, V. De Renzi, U. del Pennino, V. Bellini, S. Carretta, P. Santini, V. A. Milway, G. Timco, R. E. P. Winpenny and M. Affronte, *Phys. Rev. B: Condens. Matter*, 2009, **79**, 144419; (c) V. Corradini, A. Ghirri, U. del Pennino, R. Biagi, V. A. Milway, G. Timco, F. Tuna, R. E. P. Winpenny and M. Affronte, *Dalton Trans.*, 2010, **39**, 4928; (d) H. Rath, G. A. Timco, V. Corradini, A. Ghirri, U. del Pennino, A. Fernandez, R. G. Pritchard, C. A. Muryn, M. Affronte and R. E. P. Winpenny, *Chem. Commun.*, 2013, **49**, 3404.
- 27 J. Dreiser, A. M. Ako, C. Wäckerlin, J. Heidler, C. E. Anson, A. K. Powell, C. Piamonteze, F. Nolting, S. Rusponi and H. Brune, *J. Phys. Chem. C*, 2015, **119**, 3550.
- 28 N. Song and Y.-W. Yang, *Chem. Soc. Rev.*, 2015, **44**, 3474.
- 29 M. G. L. Petrucci and A. K. Kakkar, *Adv. Mater.*, 1996, **8**, 251.
- 30 A. R. Silva, C. Freire, B. de Castro, M. M. A. Freitas and J. L. Figueiredo, *Microporous Mesoporous Mater.*, 2001, **46**, 211 and references cited therein.
- 31 V. Arun, N. Sridevi, P. P. Robinson, S. Manju and K. K. M. Yusuff, *J. Mol. Catal. A: Chem.*, 2009, **304**, 191.
- 32 K. C. Gupta and A. K. Sutar, *Coord. Chem. Rev.*, 2008, **252**, 1420 and references cited therein.
- 33 C. Hamann, R. Woltmann, I.-Po Hong, N. Hauptmann, S. Karan and R. Berndt, *Rev. Sci. Instrum.*, 2011, **82**, 033903.
- 34 For recent studies, see e.g.: (a) J. Esteban, L. Alcázar, M. Torres-Molina, M. Monfort, M. Font-Bardia and A. Escuer, *Inorg. Chem.*, 2012, **51**, 5503; (b) J. Esteban, A. Escuer, M. Font-Bardia, O. Roubeau and S. J. Teat, *Polyhedron*, 2013, **52**, 339; (c) J. Esteban, M. Font-Bardia, J. S. Costa, S. J. Teat and A. Escuer, *Inorg. Chem.*, 2014, **53**, 3194; (d) A. Massard, G. Rogez and P. Braunstein, *Dalton Trans.*, 2014, **43**, 42; (e) A. Escuer, J. Esteban, J. Mayans and M. Font-Bardia, *Eur. J. Inorg. Chem.*, 2014, 5443; (f) M. Pait, A. Bauzá, A. Frontera, E. Colacio and D. Ray, *Inorg. Chem.*, 2015, **54**, 4709; (g) K. I. Alexopoulou, A. Terzis, C. P. Raptopoulou, V. Psycharis, A. Escuer and S. P. Perlepes, *Inorg. Chem.*, 2015, **54**, 5615.
- 35 (a) F. K. Larsen, J. Overgaard, S. Parsons, E. Rentschler, A. A. Smith, G. A. Timco and R. E. P. Winpenny, *Angew. Chem., Int. Ed.*, 2003, **42**, 5978; (b) M. Rancan, G. N. Newton, C. A. Muryn, R. G. Pritchard, G. A. Timco, L. Cronin and R. E. P. Winpenny, *Chem. Commun.*, 2008, 1560; (c) A. McRobbie, A. R. Sarwar, S. Yeninas, H. Nowell, M. L. Baker, D. Allan, M. Luban, C. A. Muryn, R. G. Pritchard, R. Prozorov, G. A. Timco, F. Tuna, G. F. S. Whitehead and R. E. P. Winpenny, *Chem. Commun.*, 2011, **47**, 6251.
- 36 (a) E. García-España, P. Gaviña, J. Latorre, C. Soriano and B. Verdejo, *J. Am. Chem. Soc.*, 2004, **126**, 5082; (b) L.-Y. Kong, Z.-H. Zhang, H.-F. Zhu, H. Kawaguchi, T.-A. Okamura, M. Doi, Q. Chu, W.-Y. Sun and N. Ueyama, *Angew. Chem., Int. Ed.*, 2005, **44**, 4352; (c) L. Natrajan, J. Pécaut and M. Mazzanti, *Dalton Trans.*, 2006, 1002; (d) P. C. Andrews, T. Beck, C. M. Forsyth, B. H. Fraser, P. C. Junk, M. Massi and P. W. Roesky, *Dalton Trans.*, 2007, 5651; (e) S.-D. Bian, J.-H. Jia and Q.-M. Wang, *J. Am. Chem. Soc.*, 2009, **131**, 3422; (f) X.-L. Tang, W.-H. Wang, W. Dou, J. Jiang, W.-S. Liu, W.-W. Qin, G.-L. Zhang, H.-R. Zhang, K.-B. Yu and L.-M. Zheng, *Angew. Chem., Int. Ed.*, 2009, **48**, 3499; (g) H. Ke, L. Zhao, G. F. Xu, Y. N. Guo, J. Tang, X. Y. Zhang and H. J. Zhang, *Dalton Trans.*, 2009, 10609; (h) A. M. García-Deibe, C. Portela-García, M. Fondo, A. J. Mota and J. Sanmartín-Matalobos, *Chem. Commun.*, 2012, **48**, 9915; (i) J.-B. Peng, Q.-C. Zhang, X.-J. Kong, Y.-Z. Zheng, Y.-P. Ren, L.-S. Long, R.-B. Huang, L.-S. Zheng and Z. Zheng, *J. Am. Chem. Soc.*, 2012, **134**, 3314; (j) T. D. Pasatoiu, A. Ghirri, A. M. Madalan, M. Affronte and M. Andruh, *Dalton Trans.*, 2014, **43**, 9136.
- 37 (a) T. N. Hooper, R. Inglis, M. A. Palacios, G. S. Nichol, M. B. Pitak, S. J. Coles, G. Lorusso, M. Evangelisti and E. K. Brechin, *Chem. Commun.*, 2014, **50**, 3498 and references cited therein; (b) A. K. Ghosh, M. Pait, M. Shatruck, V. Bertolasi and D. Ray, *Dalton Trans.*, 2014, **43**, 1970; (c) C. C. Wang, H. Y. Li, X. Q. Zhang, Y. M. Jiang and S. H. Zhang, *J. Cluster Sci.*, 2015, **26**, 1055; (d) V. Velasco, D. Aguilà, L. A. Barrios, I. Borilovic, O. Roubeau, J. Ribas-Ariño, M. Fumanal, S. J. Teat and G. Aromí, *Chem. Sci.*, 2015, **6**, 123; (e) W. Sethi, S. Sanz, K. S. Pedersen, M. A. Sørensen, G. S. Nichol, G. Lorusso, M. Evangelisti, E. K. Brechin and S. Piligkos, *Dalton Trans.*, 2015, **44**, 10315; (f) X.-M. Zhang, P. Li, W. Gao, J.-P. Liu and E.-Q. Gao, *Dalton Trans.*, 2015, **44**, 13581; (g) M. Holyńska, R. Clérac and M. Rouzières, *Chem. – Eur. J.*, 2015, **21**, 13321.
- 38 (a) R. D. Shannon, *Acta Crystallogr.*, 1976, **32**, 751; (b) B. Cordero, V. Gómez, A. E. Platero-Prats, M. Revés, J. Echeverría, E. Cremades, F. Barragán and S. Alvarez, *Dalton Trans.*, 2008, 2832.
- 39 See e.g.: (a) P. D. Frischmann and M. J. MacLachlan, *Chem. Soc. Rev.*, 2013, **42**, 871; (b) J.-J. Liu, Y.-F. Guan, Y. Chen, M.-J. Lin, C.-C. Huang and W.-X. Dai, *Dalton Trans.*, 2015, **44**, 9370.
- 40 H. Lueken, *Magnetochemie*, B.G. Teubner, Stuttgart, 1999.
- 41 (a) M. Speldrich, H. Schilder, H. Lueken and P. Kögerler, *Isr. J. Chem.*, 2011, **51**, 215; (b) J. van Leusen, M. Speldrich, H. Schilder and P. Kögerler, *Coord. Chem. Rev.*, 2015, **289–290**, 137–148.
- 42 D. T. Genna, A. G. Wong-Foy, A. J. Matzger and M. S. Sanford, *J. Am. Chem. Soc.*, 2013, **135**, 10586 and references cited therein.
- 43 W. Keim, *Angew. Chem., Int. Ed. Engl.*, 1990, **29**, 235–244.
- 44 (a) Y. T. Lai, T. C. Chen, Y. K. Lan, B. S. Chen, J. H. You, C. M. Yang, N. C. Lai, J. H. Wu and C. S. Chen, *ACS Catal.*, 2014, **4**, 3824–3836; (b) H. Guerba, B. Djellouli, C. Petit and V. Pitchon, *C. R. Chim.*, 2014, **17**, 775–784.
- 45 G. M. Sheldrick, *Acta Crystallogr., Sect. A: Found. Crystallogr.*, 2008, **64**, 112.

