


 Cite this: *RSC Adv.*, 2018, 8, 12785

A ratiometric fluorescence assay for acetylcholinesterase activity and inhibitor screening based on supramolecular assembly induced monomer–excimer emission transition of a perylene probe†

 Chunhua He,^{ab} Huipeng Zhou,^{*b} Ejaz Hussain,^{bc} Yunyi Zhang,^{bd} Niu Niu,^{bc} Yunhui Li,^a Yuqin Ma^{†a} and Cong Yu^{*bc}

A ratiometric fluorescence assay for acetylcholinesterase activity is established, which is based on controlled perylene probe assembly and monomer–excimer transition. In a buffer solution, a perylene probe with two negatively charged groups (PDI-DHA) mainly exists in monomeric form. In the presence of cationic lauroylcholine and lauric acid, PDI-DHA can form supramolecular assemblies and the perylene excimer emission can be observed. AChE can catalyze the hydrolysis of lauroylcholine to anionic lauric acid and choline. The hydrolysis process can trigger the breakdown of the supramolecular assemblies. The perylene excimer recovers to the monomeric form because of the de-aggregation of the probe. The excimer–monomer transition can be detected, and a ratiometric fluorescence assay for AChE activity and inhibitor screening is therefore established.

Received 9th February 2018

Accepted 19th March 2018

DOI: 10.1039/c8ra01274a

rsc.li/rsc-advances

Introduction

Acetylcholinesterase (AChE) is an important member of the carboxylesterase family that plays an essential role in the catalysis of hydrolysis of the neurotransmitter acetylcholine (ACh) localized in the central and peripheral nervous systems of animals.¹ Thus it maintains the level of ACh. The process is a pivotal step to control the neural response.^{2–4} The function of brain neurons can be damaged as a result of dysfunction of acetylcholine. Its deficiency or elevation can induce vital diseases such as decreased heart rate, dilation of the blood vessel, and Alzheimer's disease (AD).⁵ The decrease in the level of ACh from the standard levels in the cortex and hippocampus,⁶ could help the assembly of amyloid β peptides into amyloid fibrils, which results in Alzheimer's disease (AD).^{7–11} In fact, AChE plays an essential role in the development of AD. AChE inhibitors are usually used for the inhibition of ACh hydrolysis and increase the ACh concentration. The AChE

inhibitors are currently used as a principal drug for Alzheimer's disease and neuromuscular disorders.^{12–15} Therefore, development of highly sensitive methods for the detection of AChE activity and screening of its potential inhibitor has great practical significance.

In the past decades, various assays for AChE activity and screening for potential inhibitors of AChE have been established. For example, the electrochemical,^{16,17} colorimetric,¹⁸ chemiluminescent,¹⁹ and fluorometric assays, *etc.*^{20,21} Various fluorescent materials have been tested to improve the detection of AChE activity and screening of its inhibitors. Such as CdS quantum dots,²² gold nanoparticles,²³ and aromatic dyes.²⁴ Among them, organic fluorophores are highly attractive because of their high thermal and photochemical stability, longer lifetime and higher quantum yield.^{25,26}

Perylene diimide (PDI) derivatives are one of the outstanding large planar π -conjugated molecules possessing excellent optical and electronic properties.^{24,27,28} Furthermore, PDI derivatives as promising fluorescent probes have been applied to the biological analysis and sensing applications.^{29–31} PDI derivatives have higher aggregation tendency due to strong π – π , or π –C–H interactions and hydrophobic interactions. It results in the fluorescence quenching of monomers or turn-on excimer emission.³² Over last few years, our group has reported studies on the PDI derivatives based fluorescent sensors.^{33–37} In particular, PDI derivatives could be introduced into aggregation by polycation or polyanion, which result in excimer emission.^{38,39} However, small molecules like

^aChangchun University of Science and Technology, Weixing Road, No. 7989, Changchun 130022, P. R. China. E-mail: myq9393@sina.com

^bState Key Laboratory of Electroanalytical Chemistry, Changchun Institute of Applied Chemistry, Chinese Academy of Sciences, Changchun, 130022, P. R. China. E-mail: congyu@ciac.ac.cn; hpzhou@ciac.ac.cn

^cUniversity of the Chinese Academy of Sciences, Beijing, 100049, P. R. China

^dSchool of Materials Science and Engineering, Tianjin University, Tianjin, 300072, PR China

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c8ra01274a



This journal is © The Royal Society of Chemistry 2018

of lauroylcholine concentration (0–800 μM). PDI-DHA and lauroylcholine could form supramolecular assemblies due to electrostatic interactions, resulting in a CAC value of 310 μM for lauroylcholine. However, in this case, the concentration of enzyme substrate (lauroylcholine) remains high for the AChE activity detection and the sensitivity may be limited.

Herein, we selected lauric acid as the alternate counterion to reduce the concentration of lauroylcholine. As shown in Fig. 2, 130 μM of lauroylcholine which was lower than its CAC, 10 μM of PDI-DHA and varying amounts of lauric acid (100–500 μM) were added.

With the increase of the amount of lauric acid, the monomer emission decreased and excimer emission increased gradually. When 300 μM lauric acid was added, the value of intensity ratio (I_E/I_M , in which I_E and I_M refer to the emission intensities at 680 nm and 548 nm, respectively) reached its maximum. Fig. 2b illustrates the stable assemblies of 10 μM PDI-DHA and 300 μM lauric acid in the presence of 130 μM (much lower than 310 μM) of lauroylcholine.

For optimal AChE activity detection, the concentration of lauroylcholine should be as low as possible. The concentration of PDI-DHA was fixed at 10 μM . The fluorescence spectra of supramolecular assemblies were interrogated with the fixed ratio of [lauroylcholine]: [lauric acid] = 130 μM : 300 μM . As shown in Fig. S2,† the intensity ratio (the I_E/I_M value) increased gradually with the increase of lauroylcholine concentration. When the concentration of lauroylcholine was 130 μM , the I_E/I_M value reached its maximum. A further increase of the concentration of the mixture of lauroylcholine/lauric acid (130 μM /300 μM) caused no increase of the I_E/I_M value, indicating that most of the PDI-DHA molecules had been transformed to the aggregated form.

Meanwhile, both PDI-DHA and lauric acid contained carboxylic acid groups. Thus the supramolecular aggregation in mixed solutions can be controlled by varying the pH of the system. We investigated the fluorescence intensity ratio (the I_E/I_M value) in different pH buffer (see Fig. S3†). When the pH value is greater than 8.0, the PDI-DHA molecules may not aggregate. And when the pH value is 7.5, the I_E/I_M value of PDI-DHA reached the maximum in the assay system containing lauric acid (300 μM) and lauroylcholine (130 μM). The optimal pH value was chosen to be 7.5 throughout the whole experiment.

We also investigated the optimal concentration of PDI-DHA with the assay system containing lauroylcholine (130 μM) and

lauric acid (300 μM). As shown in Fig. S4,† different concentrations (5, 10, 20, 30 μM) of PDI-DHA were added, it was difficult to get the excimer emission when the concentrations were too low. When the concentration of PDI-DHA was 10 μM , the I_E/I_M value reached its maximum. Therefore the optimal concentration of PDI-DHA was chosen to be 10 μM .

Characterization of supramolecular assembly

The average size distribution of the supermolecular assembly was characterized in MOPS buffer by Dynamic Light Scattering at 25 $^{\circ}\text{C}$. As shown in Fig. S5,† the results confirmed that the supramolecular assemblies could not be formed in the absence of the PDI-DHA, lauric acid or lauroylcholine chloride. The supramolecular assemblies with sizes around 414 nm could be measured in the mixed solution when the three components present simultaneously.

Supramolecular assemblies triggering the development of AChE activity assay

AChE can hydrolyze lauroylcholine into lauric acid and choline, the amount of lauroylcholine will decrease with the addition of AChE. The hydrolysis process can trigger the breakdown of the supramolecular complex as a result of the coulombic repulsive interactions between negatively charged PDI-DHA and lauric acid. The monomeric emission of PDI-DHA is a turn-on because of the excimer-monomer transition due to de-aggregation. Thus, the ratio of excimer and monomer could be used to monitor the process of enzymatic hydrolysis. The time-dependence of the resulting sample was tested by recording the fluorescence spectra at regular time intervals (Fig. 3). In the presence of 150 mU mL^{-1} AChE, with the prolongation of time, the lauroylcholine was consumed, and the ratio of lauroylcholine and lauric acid continuously changed. Resulting in the transition of the PDI-DHA monomer state and excimer state. The excimer emission decreased gradually, meanwhile, the monomer emission increased gradually with increasing enzymatic reaction time. It is worth noting that there is no clear isoemissive point at around 640 nm because it is a more complex system than the simple two-component excimer-monomer equilibrium.⁴³ Maximum I_M/I_E value was observed after 180 min of enzymatic reaction (Fig. 3b). Therefore, 180 min was selected as the optimal enzyme reaction time.




Fig. 2 Changes in emission spectra (a) and corresponding I_E/I_M value (b) of PDI-DHA (10 μM) and lauroylcholine (130 μM) in the presence of the different concentrations (100–500 μM) of lauric acid.




Fig. 3 Emission spectra (a) and the corresponding I_M/I_E value (b) changes as a function of the enzymatic reaction time. Conditions: AChE = 120 mU mL^{-1} , lauric acid = 300 μM , lauroylcholine = 130 μM .



Our assay could be used to qualify the concentration of AChE. The changes in fluorescence emission of the supramolecular assemblies were monitored by varying the concentration of AChE. Fixed concentration of lauroylcholine (130 μM) and different concentrations of AChE (0, 5, 10, 20, 40, 60, 80, 100, 150, 200, 250 mU mL^{-1}) were mixed. The mixture was placed in 37 $^{\circ}\text{C}$ water bath for 180 min, after hydrolysis reaction, lauric acid (300 μM) and PDI-DHA (10 μM) were added. The fluorescence emission was detected at 20 $^{\circ}\text{C}$. As shown in Fig. 4, with an increase in the concentration of AChE, the monomer emission intensity of PDI-DHA increased gradually. The emission intensity ratios ($I_{\text{M}}/I_{\text{E}}$) of PDI-DHA plotted against the concentrations of AChE, which displayed a linear relationship at AChE concentration range of 5–150 mU mL^{-1} . The linear correlation equation is $IF = 0.0914 + 0.0204 [\text{AChE}]$, $R^2 = 0.999$. Where “IF” refers to the emission intensity ratio of the $I_{\text{M}}/I_{\text{E}}$, and “[AChE]” refers to the concentration of AChE. Our method had a better sensitivity and 5 mU mL^{-1} AChE could be easily detected.^{44–48}

Selectivity of the assay

To examine the specificity of the AChE activity assay, the fluorescence response of lauroylcholine towards various enzymes and proteins, including ALP, S1 nuclease, lysozyme, Exo I nuclease, trypsin, and BSA were recorded. The concentrations of ALP, S1 nuclease, lysozyme, Exo I nuclease and trypsin were kept at 1 U mL^{-1} , and the concentrations of BSA was 1 mg mL^{-1} . The samples were incubated for 180 min under the same conditions as for AChE. From Fig. 5, compared the value of $I_{\text{M}}/I_{\text{E}}$ for AChE (150 mU mL^{-1}) with that of other substances (1 U mL^{-1}). Significant changes were only seen for AChE. Other enzymes and proteins gave no noticeable monomer intensity recovery. The results suggest that our assay is highly selective for AChE, and could be applied to detect AChE activity in the presence of other enzymes.

Inhibitor screening of the assay

It is a fact that the activity of AChE to hydrolyze lauroylcholine could be restrained when an AChE inhibitor existed. The inhibitors of AChE were widely applied as drugs for the treatment of Alzheimer's disease (AD) *via* increasing the amounts of acetylcholine. Therefore, screening for AChE potential inhibitor is of great value. In order to confirm the potential application of




Fig. 5 Selectivity study. $I_{\text{M}}/I_{\text{E}}$ values of the PDI-DHA in the presence of AChE and the other proteins. Columns 0–6: AChE, ALP, S1 nuclease, lysozyme, Exo I nuclease, trypsin, and BSA. AChE: 150 mU mL^{-1} ; other enzymes: 1 U mL^{-1} , BSA: 1 mg mL^{-1} . Background is deducted.

the assay for AChE inhibitor screening, 3-hydroxycarbofuran and donepezil, two typical inhibitors of AChE were tested.⁴⁶ The fluorescence emission spectra of PDI-DHA with different concentration of both inhibitors were investigated (Fig. S6a and S7a†). The Fig. S6b and S7b† showed that the $I_{\text{M}}/I_{\text{E}}$ value reduced with the enhancement of the amounts of 3-hydroxycarbofuran and donepezil. The results indicate that the 3-hydroxycarbofuran and donepezil were more effective at higher concentrations. And the IC_{50} values of 3-hydroxycarbofuran and donepezil were estimated to be 149.2 nM and 57.5 nM, respectively. The results clearly suggest that our assay can be applied for inhibitor screening.

To further investigate the applicability of the supramolecular assemblies for screening AChE inhibitors. Lake water samples were used. Different concentrations of the AChE inhibitors 3-hydroxycarbofuran (50, 100, 200, 300, 400, 500 nM) and donepezil (25, 50, 100, 150, 200 nM) were added to the lake water samples. Their inhibition effects were compared with those in the buffer solution. As shown in Fig. 6, the inhibition effect of the samples prepared in the lake water is similar to those prepared in buffer solution. The results suggest that our assay can be applied for the screening of AChE inhibitors in real samples.




Fig. 4 (a) Fluorescence spectra changes of the assembly in the presence of different concentrations of AChE (5–250 mU mL^{-1}). (b) Corresponding $I_{\text{M}}/I_{\text{E}}$ value based calibration curve for AChE (5–250 mU mL^{-1}).




Fig. 6 Plot of the inhibition efficiency as a function of 3-hydroxycarbofuran (50, 100, 200, 300, 400, 500 nM) (a) and donepezil (25, 50, 100, 150, 200 nM) (b) concentration. Samples were prepared in deionized water (black line) or lake water (red line).



Conclusions

In summary, we report a ratiometric fluorescence method for the detection of AChE activity and its inhibitor screening based on the formation of a supramolecular assembly of a PDI derivative in the presence of small molecules (lauric acid and lauroylcholine) for the first time. It is based on the monomer-excimer emission transition and the controlled release of PDI-DHA from the supramolecular assemblies. AChE can catalyze the hydrolysis of lauroylcholine to lauric acid and choline. The hydrolysis process can trigger the breakdown of the supramolecular assemblies. The perylene excimer changes to the monomeric. The excimer-monomer transition can be detected, and a ratiometric fluorescence method for AChE activity is established. The assay is facile, efficient and inexpensive, which also exhibits high sensitivity and selectivity. Our assay can also be applied for the screening of AChE inhibitors in real samples.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (21561162004), and the Science and Technology Development Project of the Jilin Province (20170204038GX).

Notes and references

- M. Palkovit and D. M. Jacobowi, *J. Comp. Neurol.*, 1974, **157**, 29–41.
- H. Zejli, I. Naranjo-Rodriguez, B. Liu, K. R. Temsamani and J. L. Marty, *Talanta*, 2008, **77**, 217–221.
- J. W. Mason, C. L. Schmid, L. M. Bohn and W. R. Roush, *J. Am. Chem. Soc.*, 2017, **139**, 5865–5869.
- G. S. M. Hanan, Y. W. Barbas, E. K. Hussain and Y. Shoham, *Brain Struct. Funct.*, 2013, **218**, 59–72.
- A. V. Terryjr and J. J. Buccafusco, *J. Pharmacol. Exp. Ther.*, 2003, **306**, 821–827.
- A. M. S. Kar, D. Issa, D. S. Seto, B. Auld, R. Collier and J. Quirion, *Neurochem*, 1998, **70**, 2179–2187.
- M. Singh, M. Kaur, H. Kukreja, R. Chugh, O. Silakari and D. Singh, *Eur. J. Med. Chem.*, 2013, **70**, 165–188.
- L. A. Arribas, M. A. Lomillon, O. D. Renedo and M. J. Martínez, *Talanta*, 2013, **111**, 8–12.
- M. H. Pournaghi-Azar and A. Saadatirad, *J. Electroanal. Chem.*, 2008, **624**, 293–298.
- R. L. Zhang, S. S. Liang, M. Jin, T. He and Z. Q. Zhang, *Sens. Actuators, B*, 2017, **253**, 196–202.
- A. Makower, C. R. Lowe and F. W. Scheller, *J. Anal. Chem.*, 1999, **364**, 179–183.
- M. B. Colovic, D. Z. Krstic, T. D. Lazarevic-Pasti, A. M. Bondzic and V. M. Vasic, *Curr. Neuropharmacol.*, 2013, **1**, 3–6.
- J. W. Mason, C. L. Schmid, L. M. Bohn and W. R. Roush, *J. Am. Chem. Soc.*, 2017, **139**, 5865–5869.
- E. Kozma and M. Catellani, *Dyes Pigm.*, 2013, **98**, 160–179.
- L. P. Köse and I. Gulcin, *Rec. Nat. Prod.*, 2017, **11**, 558–561.
- H. Wang, J. Wang, C. Timchalk and Y. H. Lin, *Anal. Chem.*, 2008, **80**, 8477–8484.
- D. Du, Y. Tao, W. Y. Zhang, D. L. Liu and H. B. Li, *Biosens. Bioelectron.*, 2011, **26**, 4231–4235.
- M. Wang, X. G. Gu, G. X. Zhang, D. Q. Zhang and D. B. Zhu, *Langmuir*, 2009, **25**, 2504–2507.
- S. Sabelle, P. Y. Renard, K. Pecorella, S. S. Dezard, C. Creminon, J. Grassi and C. Mioskowski, *J. Am. Chem. Soc.*, 2002, **124**, 4874–4880.
- E. Golub, R. Freeman and I. Willner, *Anal. Chem.*, 2013, **85**, 12126–12133.
- K. Ma, L. Lu, Z. Qi, J. Feng, C. Zhuo and Y. Zhang, *Biosens. Bioelectron.*, 2015, **68**, 648–653.
- R. Gill, L. Bahshi, B. Freeman and I. Willner, *Angew. Chem., Int. Ed.*, 2008, **47**, 1676–1679.
- V. Pavlov, Y. Xiao and I. Willner, *Nano Lett.*, 2005, **5**, 649–653.
- E. Hussain, H. Zhou, N. Yang, S. A. Shahzad and C. Yu, *Dyes Pigm.*, 2017, **147**, 211–224.
- L. Z. Gai, H. C. Chen, B. Zou, H. Lu, G. Q. Lai, Z. F. Li and Z. Shen, *Chem. Commun.*, 2012, **48**, 10721–10723.
- Y. D. Zhang, Y. N. Cai, Z. L. Qi, L. Lu and Y. X. Qian, *Anal. Chem.*, 2013, **85**, 8455–8461.
- M. Wang, X. G. Gu, G. X. Zhang, D. Q. Zhang and D. B. Zhu, *Anal. Chem.*, 2009, **81**, 4444–4449.
- P. R. L. Malenfant, C. D. Dimitrakopoulos, J. D. Gelorme, L. L. Kosbar, T. O. Graham, A. Curioni and W. Andreoni, *Appl. Phys. Lett.*, 2002, **80**, 2517–2519.
- G. Grisci, E. Kozma, K. Pagano, L. Ragona and F. Galeotti, *RSC Adv.*, 2016, **6**, 64374–64382.
- Z. J. Chen, L. M. Wang, G. Zou, L. Zhang, G. J. Zhang, X. F. Cai and M. S. Teng, *Dyes Pigm.*, 2012, **94**, 410–415.
- S. S. Zhang, X. J. Wang, Y. W. Huang, H. Y. Zhai and Z. H. Liu, *Sens. Actuators, B*, 2018, **254**, 805–810.
- Q. F. Yan, Y. Zhou, Y. Q. Zheng, J. Pei and D. H. Zhao, *Chem. Sci.*, 2013, **4**, 4389–4395.
- D. L. Liao, J. Chen, H. P. Zhou, Y. Wang, Y. X. Li and C. Yu, *Anal. Chem.*, 2013, **85**, 2667–2672.
- J. Chen, D. L. Liao, Y. Wang, H. P. Zhou, W. Y. Li and C. Yu, *Org. Lett.*, 2013, **15**, 2133–2135.
- Y. Y. Zhang, C. Y. Zhang, J. Chen, Y. X. Li, M. D. Yang, H. P. Zhou, S. A. Shahzad, H. Qi, C. Yu and S. H. Jiang, *J. Mater. Chem. C*, 2017, **5**, 4691–4694.
- Y. Chen, W. Y. Li, Y. Wang, X. D. Yang, J. Chen, Y. G. Jiang, C. Yu and Q. Lin, *J. Mater. Chem. C*, 2014, **2**, 4080–4085.
- Y. Wang, J. Chen, Y. Chen, W. Y. Li and C. Yu, *Anal. Chem.*, 2014, **86**, 4371–4378.
- Y. J. Wu, P. H. Ni, M. Z. Zhang and X. L. Zhu, *Soft Matter*, 2010, **6**, 3751–3758.
- M. Wathier, B. A. Lakin, P. N. Bansal, S. S. Stoddart, B. D. Snyder and M. W. Grinstaf, *J. Am. Chem. Soc.*, 2013, **135**, 4930–4933.
- Y. Wang, J. Chen, H. Jiao, Y. Chen, W. Li, Q. Zhang and C. Yu, *Chem.–Eur. J.*, 2013, **19**, 12846–12852.



- 41 E. N. Savariar, S. Ghosh, D. C. Gonzalez and S. Thayumanavan, *J. Am. Chem. Soc.*, 2008, **130**, 5416–5417.
- 42 D. C. Gonzalez, E. N. Savariar and S. Thayumanavan, *J. Am. Chem. Soc.*, 2009, **131**, 7708–7716.
- 43 S. Nagatoishi, T. Nojima, B. Juskowiak and S. Takenaka, *Angew. Chem., Int. Ed.*, 2005, **117**, 5195–5198.
- 44 F. D. Feng, Y. L. Tang, S. Wang, Y. L. Li and D. B. Zhu, *Angew. Chem., Int. Ed.*, 2007, **46**, 7882–7886.
- 45 X. Shen, F. X. Liang, G. X. Zhang and D. Q. Zhang, *Analyst*, 2012, **137**, 2119–2123.
- 46 D. J. Selkoe, *Physiol. Rev.*, 2001, **81**, 741–766.
- 47 C. Ye, M. Q. Wang, X. Zhong, S. H. Chen, Y. Y. Chai and R. Yuan, *Biosens. Bioelectron.*, 2016, **79**, 34–40.
- 48 K. Ingkaninan, P. Temkitthawon, K. Chuenchom, T. Yuyaem and W. Thongnoi, *J. Ethnopharmacol.*, 2003, **89**, 261–264.

