


Chemical Science

rsc.li/chemical-science



ISSN 2041-6539



EDGE ARTICLE

Huaqiang Zeng *et al.*

A halogen bond-mediated highly active artificial chloride channel with high anticancer activity

Cite this: *Chem. Sci.*, 2018, 9, 4044

A halogen bond-mediated highly active artificial chloride channel with high anticancer activity†

Changliang Ren,^a Xin Ding,^a Arundhati Roy,^a Jie Shen,^a Shaoyuan Zhou,^b Feng Chen,^a Sam Fong Yau Li,^c Haisheng Ren,^b Yi Yan Yang^a and Huaqiang Zeng^{*,a}

Chloride-selective transmembrane carriers or channels might have possible uses in treating channelopathies or cancers. While chloride carriers have been extensively investigated, the corresponding chloride channels have remained limitedly studied. Moreover, all hitherto reported channel systems lack clearly definable and readily modifiable positions in their structures for the reliable construction and combinatorial optimization of their ion transport properties. As a result, the existing channels are limited by their large molecular weight, weak activity or low anion selectivity. In this report, we describe a readily accessible and robust monopeptide-based scaffold for the reliable construction of halogen bond-mediated artificial anion channels via directional assembly of electron-deficient iodine atoms, which create a transmembrane pathway for facilitating anion transport. The high intrinsic modularity of the backbone of the scaffold, which enables the rapid and combinatorial optimization of the transport activity and selectivity of channels, effectively delivers a highly active chloride channel **A10**. Such high activity in chloride transport subsequently leads to an excellent IC₅₀ value of 20 μM toward inhibiting the growth of human breast cancer cells (BT-474), an anticancer activity that is even higher than that of the well-known anticancer agent cisplatin.

Received 6th February 2018

Accepted 15th March 2018

DOI: 10.1039/c8sc00602d

rsc.li/chemical-science

Introduction

Tightly regulated by varying families of ion channels, precise maintenance of ion concentration gradients across biological membranes is crucial for many biological and cellular processes. Chloride is the most abundant anion in the human body due to its important roles in controlling the membrane potential, cell volume, cellular pH balance, secretion of trans-epithelial fluid and electrolytes, *etc.*¹ Misregulation of chloride ions could lead to a variety of life-shortening ‘channelopathies’ including cystic fibrosis, Bartter syndrome and Dent’s disease. Accordingly, significant research efforts have been made recently to coherently develop artificial chloride carriers² or channels³ as possible replacements for natural chloride channels for channelopathies.⁴ On the other hand, artificial anion transporters might also disrupt pH gradients^{4f,5a,b} or ion homeostasis^{5c} that may induce cell death, and therefore may have medical applications as anticancer agents.^{4f,5}

Recently, Matile *et al.* reported self-assembled chloride carriers^{2b,c} or channels,³ⁱ elegantly exploiting halogen bonding interaction as the driving force for anion transport. Halogen bonds offer good bond strength and also good anion-induced directionality.⁶ Therefore, artificial anion transporters comprising simple structural motifs based on halogen–anion interactions may indeed offer a new horizon of prospects to comprehend complex chloride transport phenomena, often manifested in protein channels. Nevertheless, most documented artificial chloride channels purposefully harness various non-covalent interactions such as electrostatic, H-bonding, anion-π and halogen bonding forces. Channels that primarily rely on halogen bonds for facilitating anion transport still remain limitedly investigated with just one precedent.³ⁱ Such a rare occurrence presumably results from the lack of suitable molecular scaffolds for aligning electron-deficient iodine atoms into a well-defined one-dimensional (1D) array. Moreover, the majority of existing chloride channels^{3a–m} suffer from relatively complex molecular structures and/or low selectivity in anion recognition, greatly restricting their application in drug discovery.

We recently described a novel class of readily accessible monopeptide-based scaffolds capable of self-assembling into 1D columnar structures through highly directional intermolecular H-bonding forces, subsequently enabling rapid room-temperature gelation of diverse types of crude oils (Fig. 1a).^{7a–c}

^aInstitute of Bioengineering and Nanotechnology, 31 Biopolis Way, The Nanos, Singapore 138669. E-mail: hqzeng@ibn.a-star.edu.sg

^bCollege of Chemical Engineering, Sichuan University, Chengdu, China 610065

^cNUS Environmental Research Institute, Department of Chemistry, National University of Singapore, 3 Science Drive 3, Singapore 117543

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c8sc00602d






Fig. 1 (a) Crystal structure of F-Phe-C4, illustrating the formation of the 1D columnar stack with directionally assembled side chains. (b) Molecular design of the chloride-transporting channel library with systematically tunable R_1 and R_2 groups. (c) Side and top views of the computationally optimized H-bonded 1D columnar ensemble formed from L8. (d) A possible multi-ion jumping mode responsible for chloride transport.

Revealed by the crystal structure of F-Phe-C4 (Fig. 1a), one unique feature of this monopeptide-based scaffold is that the same type of side chains such as Fmoc, R_1 and R_2 are always aligned to the same side as a result of the high directionality of H bonds (Fig. 1a).^{7a} On this basis, we envisioned that replacing the Fmoc group with a chloride-binding unit such as the tetrafluoroiodobenzyl group may lead to an interesting class of artificial anion channels with anion transport mediated by

halogen bonds (Fig. 1b–d).^{3i,7d} A high intrinsic modularity involving R_1 and R_2 groups in the backbone should allow for rapid and combinatorial optimization of transport activity and selectivity of channels. In this report, we demonstrate that this indeed is the case, with the identified chloride channel exhibiting a high activity in chloride transport with respect to many other types of anions. We further demonstrate that human breast cancer cells (BT-474) are sensitive to a sodium chloride concentration gradient across the membrane, and this feature can be utilized by artificial chloride channels to inhibit cancer cell growth for their potential uses in cancer chemotherapy.^{4f,5}

Results and discussion

Combinatorial identification of highly active anion-transporting artificial channels

We synthesized a total of 15 possible channel molecules (five amino acids \times three alkyl chains of different lengths, Fig. 1b) with the tetrafluoroiodobenzyl group as the chloride-binding and -transporting unit. Corroborated by the crystal structure of F-Phe-C4 (Fig. 1a) and the computationally optimized structure of L8 (Fig. 1c), the ability of these 15 molecules to pile up to form a H-bonded 1D structure can be convincingly demonstrated by the ability of A10, L8 and L10 to efficiently congeal in *n*-hexane *via* the formation of a 3D entangled fibrous network (ESI, Fig. S1†). The ion transport activities of the synthesized channels were then evaluated using the pH-sensitive HPTS (8-hydroxypyrene-1,3,6-trisulfonic acid) assay (Fig. 2 and S2†).⁸ In a typical experiment, large unilamellar vesicles (LUVs), encapsulating HPTS (100 μM) and NaCl (100 mM) at pH = 7.0, were diluted into the same buffer at pH 8.0 to generate a pH gradient across LUVs. After the addition of channel molecules, changes in the fluorescence intensity of HPTS were monitored over 5 min. Fig. 2b shows that, among all 15 channels tested at 10 μM , A10, L8 and L10 display the highest ion transport capacities with fractional activities (R_{Cl^-}) of 60%, 68% and 59%, respectively. The corresponding EC_{50} values, which are the




Fig. 2 (a) Schematic illustration of the HPTS assay for the Cl^- transport study conducted using a pH gradient of 7 to 8 and the pH-sensitive dye HPTS entrapped inside LUVs. (b) Normalized transport activities (R_{Cl^-}) obtained over 5 min at 10 μM for all channel molecules. $R_{\text{Cl}^-} = (I_{\text{Cl}^-} - I_0) / (I_{\text{Triton}} - I_0)$ wherein I_{Cl^-} and I_0 are the ratiometric values of I_{460}/I_{403} before the addition of triton at $t = 300$ s, and I_{Triton} is the ratiometric value of I_{460}/I_{403} at $t = 300$ s right after the addition of triton with internal/external buffers containing 100 mM NaCl. All the data were averaged over two measurements.




Fig. 3 (a) Transport selectivity of alkali metal ions by L8 (3.6 μM) obtained from the HPTS assay by varying the extravesicular MCl (M⁺ = Li⁺, Na⁺, K⁺, Rb⁺ and Cs⁺). (b) Concentration-dependent changes in the fluorescence intensity of the chloride-sensitive SPQ dye (λ_{ex} = 360 nm, λ_{em} = 430 nm) after the addition of L8 at different concentrations. Inside LUV: 200 mM NaNO₃ and 0.5 mM SPQ. Outside LUV: 200 mM NaCl. (c) Ion transport activities of L8 (3.6 μM) determined in the absence or presence of proton transporter FCCP (1 μM), indicating a preferential transport of Cl⁻ over H⁺. (d) Ion transport activities of L8 (3.6 μM) determined in the absence or presence of a potassium carrier, valinomycin (VA, 25 pM), indicating a preferential transport of Cl⁻ over OH⁻. (e) Ion transport activities of four control compounds 1–4 at 3.6 μM, suggesting a critical role of halogen bonds in mediating the transport of Cl⁻ and OH⁻. (f) Anion selectivity assayed for A10 (For L8 and L10, see Fig. S10†). (g) Single current trace of L8 recorded at -100 mV with symmetric baths (cis chamber = trans chamber = 1 M KCl). (h) I–V curve for obtaining the Cl⁻ conductance (γ_{Cl⁻}) of L8. All the data were averaged over two measurements.

in a bilayer of 128 POPC molecules (17 152 atoms) solvated on two sides by 2 × 2397 water molecules (Fig. 4a). This leads to a simulation system of 32 062 atoms with a dimension of 70 Å

(w) × 70 Å (w) × 74 Å (h). After equilibration steps, the production run was carried out for 30 ns. The last 20 ns trajectories with 1000 structural snapshots were used for analysing




Fig. 4 (a) Embedding computationally optimized H-bonded L8 inside a simulation box of 70 Å (w) × 70 Å (w) × 74 Å (h), comprising 128 POPC molecules and 4794 water molecules. (b) Probability distribution patterns of θ_{234} , θ_{345} , θ_{456} and θ_{567} obtained after analysing 1000 structures. (c) A representative highly populated structure with the four θ angles approaching those having the highest probabilities shown in (b). POPC = 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine.

both the hydrophobic thickness of the POPC membrane and the distribution probability of angle θ (e.g., the angle formed by the three immediately adjacent iodine atoms such as iodine atoms 2–4, Fig. 4a and b) using a probability density function.^{10f}

To estimate the hydrophobic membrane thickness, the Z-coordinates along the membrane normal of all 254 ester O-atoms from 64 POPC molecules located either at the top or on the bottom layers were averaged. The separation distance between the two averaged Z-coordinates of O-atoms from the top and bottom layers was calculated with deduction of a van der Waals diameter of 3.1 Å for the O-atom. The same calculation was performed for all 1000 structural snapshots to derive the probability distribution of the hydrophobic membrane thickness (Fig. S12†). From these computations, 633 structures have a hydrophobic thickness of 27 to 29 Å, and the average thickness over 1000 structures is 28.1 Å, a value that could be corroborated by the experimentally and computationally determined values of 27.1 and 27.8 Å, respectively, for the POPC membrane.¹¹

Embedding eight molecules of L8 in the lipid membrane shows that the first and eighth molecules are not aligned well with the central six molecules (Fig. 4a and c). The most highly populated angles for θ_{234} , θ_{345} , θ_{456} and θ_{567} are 159°, 162°, 163° and 157°, respectively, with a representative structure, closely capturing these angles, shown in Fig. 4c. In this highly populated structure, the largest intermolecular separation distance along the Z axis among the eight iodine atoms occurs between the 1st and 7th iodine atoms (25 Å). Consistent with the intermolecular separation of 5.0 Å in **Fmoc-Phe-C4** in the solid state (Fig. 1a)^{7a} and of 4.9 Å in the computationally optimized H-bonded L8 (Fig. 1c), this separation distance of 25 Å suggests that six or seven molecules might be sufficient to span the hydrophobic core distance of 28.1 Å.

Through these MD simulations, the most important point to note is that, regardless of angle θ of varying magnitudes, all eight molecules of L8 remain H-bonded to each other in all 1000 structures surveyed (Fig. 4c). This demonstrates the reliability of intermolecular H-bonds in linking six or seven anion-binding molecules together to form a self-assembled 1D pathway, which spans the hydrophobic membrane region to facilitate highly efficient anion transport across the membrane.

High activity in chloride transport

The use of halogen bonds to mediate anion transport was first explored by Matile,^{2b,c,3i} with compounds **5** (EC_{50} = 3.1 μM)^{2c} and **8** (EC_{50} = 0.88 μM in terms of chloride binding units)³ⁱ as the most active carrier and channel molecules among their own

Table 1 EC_{50} values of channels A10, L8 and L10 as well as other halogen bond-mediated chloride carriers (5–7) or channels (8)^a


	A10	L8	L10	5 ^b	6 ^c	7 ^d	8 ^e
EC_{50} (μM)	2.37	0.39	0.93	3.6	260	68	0.88
n value ^f	3.5	3.6	3.1	3.3	3.6	1.9	0.8

^a Determined using Matile's conditions (100 mM NaCl, 10 mM HEPES, pH gradient from 7 to 8, and the total lipid concentration without cholesterol = 31.25 μM; see ref. 2b, c and 3i). ^b EC_{50} value from ref. 2c is 3.1 μM with a Hill coefficient of 3.3. ^c From ref. 2c. ^d From ref. 2b. ^e From ref. 3i with the value normalized by the authors on the basis of eight halogen-binding units contained in channel 8. ^f Hill coefficient.



categories, respectively. These EC_{50} values, including those of **6** (260 μ M) and **7** (68 μ M) presented in Table 1, were determined using cholesterol-free LUVs with a total lipid concentration of 31.3 μ M, while our assay conditions contain lipid and cholesterol at concentrations of 73.9 and 36.9 μ M, respectively. For a fair comparison, we have therefore re-determined the EC_{50} values of **A10**, **L8** and **L10** using Matile's conditions (100 mM NaCl, 10 mM HEPES, pH gradient from 7 to 8, and the total lipid concentration without cholesterol = 31.3 μ M).^{2b,c,3i}

To confirm that we are able to produce LUVs with properties similar to those prepared by Matile and his co-workers under the same assay conditions, we have performed Hill analysis on commercially available compound **5**, and obtained an EC_{50} value of 3.6 μ M with a Hill coefficient of 3.3 (Fig. S13†). These values are similar or identical to the reported values of EC_{50} (3.1 μ M) and the Hill coefficient (3.3).^{2c} Under these assay conditions, the EC_{50} values of **A10**, **L8** and **L10** were



determined to be 2.37, 0.39 and 0.93 μ M (Table 1 and Fig. S13†), respectively. The most active **L8** is 8.2 and 1.2 times more active than carrier **5** (3.6 μ M) and unimolecular channel molecule **8** (0.88 μ M in terms of effective chloride-binding units). When normalized based on the molecular weight, the ion transport activity of **L8** is about 6.5 and 1.6 times those of **5** and **8**, respectively. That **L8** is more potent than unimolecular channel **8** is a clear indication of the excellent ability of the non-covalently associated peptidic scaffold in inducing linearly arrayed iodine atoms into a conformation that is more conducive to chloride transport than the rod-like scaffold in **8**.

In addition, chloride carrier **5**, when evaluated under our assay conditions containing 33 mol% cholesterol that makes membrane less fluid,¹² shows a moderate transport activity of 38% at 40 μ M, and such moderate activity remains unchanged even when the concentration increases to 100 μ M (Fig. S14†). These values suggest the ion transport activity of **5** to be at least 10 times lower than that of **L8** having an EC_{50} value of 3.6 μ M in the cholesterol-containing environment.




Fig. 5 (a) Sensitivity of human breast cancer cells BT-474 toward five different concentrations of NaCl from 6.4 to 19.2 g mL⁻¹ after 24 h. (b) Viabilities of BT-474 cells in the presence of various concentrations of **L8** and **A10** at [NaCl] = 6.4 g L⁻¹.

High anticancer activity

The above verified highly efficient chloride transport through a channel mechanism displayed by these artificial anion channels prompted us to examine the possibility of their uses in cancer chemotherapy.^{4f,5} Human breast cancer cells (BT-474, obtained from the American Type Culture Collection, USA) were cultured in Dulbecco's Modified Eagle Medium in the presence of up to 19.2 g L⁻¹ NaCl (6.4 g L⁻¹ is the typical concentration for cell growth) with concentrations of **L8** or **A10** varying from 0 to 100 μ M (Fig. 5 and S15†). The viabilities of cells, determined after culturing the cells for 1 day at 37 °C with 5% CO₂ in the absence of channel molecules, show that BT-474 cells are sensitive to the sodium chloride concentration gradient (Fig. 5a). That is, compared to cell viability at [NaCl] = 6.4 g L⁻¹, 12, 79 and 99% fewer cells survive at [NaCl] of 12.8, 16.0 and 19.2 g L⁻¹, respectively. Continued testing shows that **A10** is considerably more potent than **L8** in inhibiting cell growth across all three different concentrations of NaCl (e.g., 6.4, 9.6 and 12.8 g L⁻¹) with an IC₅₀ value of 20 μ M for **A10** at [NaCl] = 6.4 g L⁻¹ (>100 μ M for **L8**, Fig. 5b and S15†). For comparison, highly effective anticancer agent cisplatin has an IC₅₀ value of 37 μ M against the same BT-474 cells.¹³ Assuming all **A10** molecules associate to form channels with each channel comprising six or more such molecules, the IC₅₀ value in terms of effective channel concentration is lower than 3.3 μ M.

Conclusions

In conclusion, we have developed a class of artificial chloride channels with high activity and good selectivity *via* directional



self-assembly of polarized electron-deficient iodine atoms. While these linearly arrayed acidic iodine atoms, which form a series of chloride-binding and -transporting sites, are responsible for facilitating chloride transport likely *via* a multi-ion jumping mode, the high modularity of the mono-peptide backbone enables rapid optimization of the transport activity and selectivity of channels in a combinatorial format. The identified channels **A10**, **L8** and **L10** all turn out to be very active with the best EC₅₀ values for chloride transport reaching 0.39 μ M (1.2 mol% relative to lipid) and 3.6 μ M (3.2 mol% relative to lipid/cholesterol) in cholesterol-free and -containing LUVs. In particular, the highly active **A10** exhibits not only a fractional transport activity for chloride much better than other monovalent anions including bromide, but also an excellent inhibitory activity toward human breast cancer cells with an IC₅₀ value of 20 μ M. Further refinement to enhance the anion selectivity and cytotoxicity of channels toward cancer cells is possible and expected.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by the Institute of Bioengineering and Nanotechnology (Biomedical Research Council, Agency for Science, Technology and Research, Singapore) and the Singapore National Research Foundation under its Environment and Water Research Programme administered by PUB.

Notes and references

- (a) R. Benz and R. E. W. Hancock, *J. Gen. Physiol.*, 1987, **89**, 275; (b) C. Duran, C. H. Thompson, Q. Xiao and C. Hartzell, *Annu. Rev. Physiol.*, 2010, **72**, 95; (c) N. Busschaert, C. Caltagirone, W. Van Rossom and P. A. Gale, *Chem. Rev.*, 2015, **115**, 8038.
- (a) P. V. Santacrose, J. T. Davis, M. E. Light, P. A. Gale, J. C. Iglesias-Sánchez, P. Prados and R. Quesada, *J. Am. Chem. Soc.*, 2007, **129**, 1886; (b) A. Vargas Jentzsch, D. Emery, J. Mareda, P. Metrangolo, G. Resnati and S. Matile, *Angew. Chem., Int. Ed.*, 2011, **50**, 11675; (c) A. V. Jentzsch, D. Emery, J. Mareda, S. K. Nayak, P. Metrangolo, G. Resnati, N. Sakai and S. Matile, *Nat. Commun.*, 2012, **3**, 905; (d) H. Valkenier, L. W. Judd, H. Li, S. Hussain, D. N. Sheppard and A. P. Davis, *J. Am. Chem. Soc.*, 2014, **136**, 12507; (e) A. Roy, D. Saha, A. Mukherjee and P. Talukdar, *Org. Lett.*, 2016, **18**, 5864; (f) J. Gravel, J. Kempf and A. Schmitzer, *Chem.-Eur. J.*, 2015, **21**, 18642; (g) E. B. Park and K.-S. Jeong, *Chem. Commun.*, 2015, **51**, 9197; (h) M. Lisbjerg, H. Valkenier, B. M. Jessen, H. Al-Kerdi, A. P. Davis and M. Pittelkow, *J. Am. Chem. Soc.*, 2015, **137**, 4948.
- (a) V. Sidorov, F. W. Kotch, G. Abdrakhmanova, R. Mizani, J. C. Fetting and J. T. Davis, *J. Am. Chem. Soc.*, 2002, **124**, 2267; (b) P. H. Schlesinger, R. Ferdani, J. Liu, J. Pajewska, R. Pajewski, M. Saito, H. Shabany and G. W. Gokel, *J. Am. Chem. Soc.*, 2002, **124**, 1848; (c) N. Madhavan, E. C. Robert and M. S. Gin, *Angew. Chem., Int. Ed.*, 2005, **44**, 7584; (d) V. Gorteau, G. Bollot, J. Mareda, A. Perez-Velasco and S. Matile, *J. Am. Chem. Soc.*, 2006, **128**, 14788; (e) J. D. Sadowsky, W. D. Fairlie, E. B. Hadley, H. S. Lee, N. Umezawa, Z. Nikolovska-Coleska, S. M. Wang, D. C. S. Huang, Y. Tomita and S. H. Gellman, *J. Am. Chem. Soc.*, 2007, **129**, 139; (f) J. Mareda and S. Matile, *Chem.-Eur. J.*, 2009, **15**, 28; (g) X. Li, B. Shen, X.-Q. Yao and D. Yang, *J. Am. Chem. Soc.*, 2009, **131**, 13676; (h) C. R. Yamnitz, S. Negin, I. A. Carasel, R. K. Winter and G. W. Gokel, *Chem. Commun.*, 2010, **46**, 2838; (i) A. Vargas Jentzsch and S. Matile, *J. Am. Chem. Soc.*, 2013, **135**, 5302; (j) T. Saha, S. Dasari, D. Tewari, A. Prathap, K. M. Sureshan, A. K. Bera, A. Mukherjee and P. Talukdar, *J. Am. Chem. Soc.*, 2014, **136**, 14128; (k) T. Saha, A. Gautam, A. Mukherjee, M. Lahiri and P. Talukdar, *J. Am. Chem. Soc.*, 2016, **138**, 16443; (l) X. Wei, G. Zhang, Y. Shen, Y. Zhong, R. Liu, N. Yang, F. Y. Al-mkhaizim, M. A. Kline, L. He, M. Li, Z.-L. Lu, Z. Shao and B. Gong, *J. Am. Chem. Soc.*, 2016, **138**, 2749; (m) H. Behera and N. Madhavan, *J. Am. Chem. Soc.*, 2017, **139**, 12919; (n) C. J. E. Haynes, J. Zhu, C. Chimere, S. Hernández-Ainsa, I. A. Riddell, T. K. Ronson, U. F. Keyser and J. R. Nitschke, *Angew. Chem., Int. Ed.*, 2017, **56**, 15388.
- (a) G. W. Gokel and N. Barkey, *New J. Chem.*, 2009, **33**, 947; (b) J. T. Davis, O. Okunola and R. Quesada, *Chem. Soc. Rev.*, 2010, **39**, 3843; (c) P. R. Brotherhood and A. P. Davis, *Chem. Soc. Rev.*, 2010, **39**, 3633; (d) S. Matile, A. Vargas Jentzsch, J. Montenegro and A. Fin, *Chem. Soc. Rev.*, 2011, **40**, 2453; (e) D. S. Kim and J. L. Sessler, *Chem. Soc. Rev.*, 2015, **44**, 532; (f) X. Wu, L. W. Judd, E. N. W. Howe, A. M. Withecombe, V. Soto-Cerrato, H. Li, N. Busschaert, H. Valkenier, R. Pérez-Tomás, D. N. Sheppard, Y.-B. Jiang, A. P. Davis and P. A. Gale, *Chem.*, 2016, **1**, 127.
- (a) N. Busschaert, M. Wenzel, M. E. Light, P. Iglesias-Hernández, R. Pérez-Tomás and P. A. Gale, *J. Am. Chem. Soc.*, 2011, **133**, 14136; (b) V. Soto-Cerrato, P. Manuel-Manresa, E. Hernando, S. Calabuig-Fariñas, A. Martínez-Romero, V. Fernández-Dueñas, K. Sahlholm, T. Knöpfel, M. García-Valverde, A. M. Rodilla, E. Jantus-Lewintre, R. Farràs, F. Ciruela, R. Pérez-Tomás and R. Quesada, *J. Am. Chem. Soc.*, 2015, **137**, 15892; (c) S.-K. Ko, S. K. Kim, A. Share, V. M. Lynch, J. Park, W. Namkung, W. Van Rossom, N. Busschaert, P. A. Gale, J. L. Sessler and I. Shin, *Nat. Chem.*, 2014, **6**, 885; (d) J. L. Sessler, L. R. Eller, W.-S. Cho, S. Nicolaou, A. Aguilar, J. T. Lee, V. M. Lynch and D. J. Magda, *Angew. Chem., Int. Ed.*, 2005, **44**, 5989; (e) B. A. Smith, M. M. Daschbach, S. T. Gammon, S. Xiao, S. E. Chapman, C. Hudson, M. Suckow, D. Piwnica-Worms, G. W. Gokel and W. M. Leevy, *Chem. Commun.*, 2011, **47**, 7977; (f) S. N. Berry, V. Soto-Cerrato, E. N. W. Howe, H. J. Clarke, I. Mistry, A. Tavassoli, Y.-T. Chang, R. Perez-Tomas and P. A. Gale, *Chem. Sci.*, 2016, **7**, 5069; (g) T. Saha, M. S. Hossain, D. Saha, M. Lahiri and P. Talukdar, *J. Am. Chem. Soc.*, 2016, **138**, 7558.



- 6 L. C. Gilday, S. W. Robinson, T. A. Barendt, M. J. Langton, B. R. Mullaney and P. D. Beer, *Chem. Rev.*, 2015, **115**, 7118.
- 7 (a) C. Ren, G. H. B. Ng, H. Wu, K.-H. Chan, J. Shen, C. Teh, J. Y. Ying and H. Zeng, *Chem. Mater.*, 2016, **28**, 4001; (b) C. Ren, F. Chen, F. Zhou, J. Shen, H. Su and H. Zeng, *Langmuir*, 2016, **32**, 13510; (c) C. L. Ren, J. Shen, F. Chen and H. Q. Zeng, *Angew. Chem., Int. Ed.*, 2017, **56**, 3847; (d) C. L. Ren, J. Shen and H. Q. Zeng, *J. Am. Chem. Soc.*, 2017, **139**, 12338.
- 8 H. Zhao, S. Sheng, Y. Hong and H. Zeng, *J. Am. Chem. Soc.*, 2014, **136**, 14270.
- 9 A. S. Verkman, R. Takla, B. Sefton, C. Basbaum and J. H. Widdicombe, *Biochemistry*, 1989, **28**, 4240.
- 10 (a) S. Jo, T. Kim, V. G. Iyer and W. Im, *J. Comput. Chem.*, 2008, **29**, 1859; (b) E. L. Wu, X. Cheng, S. Jo, H. Rui, K. C. Song, E. M. Davila-Contreras, Y. Qi, J. Lee, V. Monje-Galvan, R. M. Venable, J. B. Klauda and W. Im, *J. Comput. Chem.*, 2014, **35**, 1997; (c) S. Jo, J. B. Lim, J. B. Klauda and W. Im, *Biophys. J.*, 2009, **97**, 50; (d) K. Vanommeslaeghe, E. Hatcher, C. Acharya, S. Kundu, S. Zhong, J. Shim, E. Darian, O. Guvench, P. Lopes, I. Vorobyov and A. D. MacKerell, *J. Comput. Chem.*, 2010, **31**, 671; (e) W. L. Jorgensen, J. Chandrasekhar, J. D. Madura, R. W. Impey and M. L. Klein, *J. Chem. Phys.*, 1983, **79**, 926; (f) U. Essmann, L. Perera, M. L. Berkowitz, T. Darden, H. Lee and L. G. Pedersen, *J. Chem. Phys.*, 1995, **103**, 8577; (g) G. J. Martyna, D. J. Tobias and M. L. Klein, *J. Phys. Chem.*, 1994, **101**, 4177; (h) H. Sun, *J. Phys. Chem. B*, 1998, **102**, 7338–7364; (i) D. N. Bopege, M. Petrowsky, A. M. Fleshman, R. Frech and M. B. Johnson, *J. Phys. Chem. B*, 2012, **116**, 71.
- 11 Y. Guo, S. Pogodin and V. A. Baulin, *J. Chem. Phys.*, 2014, **140**, 174903.
- 12 (a) G. M. Cooper, *The Cell: A Molecular Approach*, Sinauer Associates, Sunderland, MA, 2nd edn, 2000; (b) D. Marsh, *Biochim. Biophys. Acta*, 2009, **1788**, 2114.
- 13 M. Buccioni, D. Dal Ben, C. Lambertucci, F. Maggi, F. Papa, A. Thomas, C. Santinelli and G. Marucci, *Sci. World J.*, 2014, **2014**, 264829.

