

REVIEW

View Article Online
View Journal | View Issue



Cite this: *Nat. Prod. Rep.*, 2023, 40, 1521

S-Adenosylmethionine: more than just a methyl donor

Yu-Hsuan Lee, ^{†a} Daan Ren, ^{†a} Byungsun Jeon^{‡a} and Hung-wen Liu ^{*ab}

Covering: from 2000 up to the very early part of 2023

S-Adenosyl-L-methionine (SAM) is a naturally occurring trialkyl sulfonium molecule that is typically associated with biological methyltransfer reactions. However, SAM is also known to donate methylene, aminocarboxypropyl, adenosyl and amino moieties during natural product biosynthetic reactions. The reaction scope is further expanded as SAM itself can be modified prior to the group transfer such that a SAM-derived carboxymethyl or aminopropyl moiety can also be transferred. Moreover, the sulfonium cation in SAM has itself been found to be critical for several other enzymatic transformations. Thus, while many SAM-dependent enzymes are characterized by a methyltransferase fold, not all of them are necessarily methyltransferases. Furthermore, other SAM-dependent enzymes do not possess such a structural feature suggesting diversification along different evolutionary lineages. Despite the biological versatility of SAM, it nevertheless parallels the chemistry of sulfonium compounds used in organic synthesis. The question thus becomes how enzymes catalyze distinct transformations *via* subtle differences in their active sites. This review summarizes recent advances in the discovery of novel SAM utilizing enzymes that rely on Lewis acid/base chemistry as opposed to radical mechanisms of catalysis. The examples are categorized based on the presence of a methyltransferase fold and the role played by SAM within the context of known sulfonium chemistry.

Received 1st December 2022

DOI: 10.1039/d2np00086e

rsc.li/npr

- | | | | |
|-------|--|-------|--|
| 1 | Introduction | 2.4.2 | SpnF and SpnL in spinosyn biosynthesis |
| 2 | Enzymes with a methyltransferase (MT) fold | 2.4.3 | SlnM in salinomycin biosynthesis |
| 2.1 | SAM-dependent alkylation reactions | 2.4.4 | SAM-dependent hydroxylase |
| 2.1.1 | 3-Amino-3-carboxypropyl transfer (small molecules) | 2.4.5 | SAM-dependent decarboxylase |
| 2.1.2 | ACP transfer (RNA) | 2.5 | SAM as a nucleophile |
| 2.1.3 | Carboxymethyl transfer | 2.5.1 | Carboxy-SAM |
| 2.1.4 | Adenosyl transfer | 2.5.2 | Queuosine |
| 2.2 | Intramolecular cyclization of SAM | 2.6 | SAM as an amino donor |
| 2.2.1 | Azetidine 2-carboxylic acid | 3 | Enzymes without a typical MT-fold |
| 2.2.2 | Cyclopropane | 3.1 | Pyridoxal 5'-phosphate-dependent enzymes |
| 2.3 | Methylation induced cascade reaction | 3.1.1 | Cyclopropylation |
| 2.3.1 | Terpene cyclization | 3.1.2 | Elimination |
| 2.3.2 | Pictet–Spengler rearrangement | 3.2 | Polyketide synthase |
| 2.3.3 | Stevens rearrangement | 3.2.1 | Cyclopropylation |
| 2.3.4 | Alkaloid cyclization | 4 | Conclusions |
| 2.4 | SAM as a prosthetic group | 5 | Conflicts of interest |
| 2.4.1 | LepI in leporin biosynthesis | 6 | Acknowledgements |
| | | 7 | References |

^aDepartment of Chemistry, University of Texas at Austin, Austin, TX 78712, USA.
E-mail: H.w.liu@mail.utexas.edu

^bDivision of Chemical Biology & Medicinal Chemistry, College of Pharmacy, University of Texas at Austin, Austin, TX 78712, USA

[†] These authors contributed equally.

[‡] Current affiliation: Center for Neuro-Medicine, Korea Institute of Science & Technology, Seoul, Korea 02792.

1 Introduction

S-Adenosyl-L-methionine (SAM or AdoMet, **1**) is a sulfonium-containing primary metabolite found in both prokaryotic and eukaryotic cells.¹ SAM is best known for the role it plays in one-



carbon metabolism where it serves as a methyl donor in many biological reactions,²⁻⁴ and a search in InterPro reveals that approximately 3 million protein sequences have been annotated as SAM-dependent methyltransferases.⁵ The molecular mechanism of SAM-dependent methylation generally involves a nucleophilic substitution at the sulfonium methyl carbon of SAM to generate a variety of C-, O-, N- and S-methylated products.⁶

Not all enzymes, however, utilize SAM to catalyze methylation reactions. In particular, a significant number of enzymes also bind an [Fe₄S₄] cluster typically *via* a highly conserved CX₃CX₂C motif in the active site that serves to reduce SAM leading to the homolytic cleavage of the C5'-S linkage.^{7,8} The resulting 5'-deoxyadenosyl radical equivalent (2) can then initiate a broad range of radical-mediated transformations. Radical SAM enzymes thus constitute a protein superfamily with more than 700 thousand protein sequences⁹ arguably becoming one of the focal points of modern enzymological study as summarized in many recent reviews.¹⁰⁻¹⁷

A number of other enzymes are also dependent on SAM for their activity; however, they are not easily classified as either SAM-dependent transferases or radical SAM enzymes.¹⁸⁻²¹ Among these enzymes, SAM may serve as an alkyl donor as opposed to a strict methyl donor or stabilize reaction intermediate(s) *via* the positive charge of its cationic sulfonium moiety. Alternatively, SAM may only play a structural role helping to maintain the integrity of the enzyme and its active site and thus remain intact throughout the catalytic cycle. While many of these enzymes are structurally related to the SAM-dependent methyltransferases,²²⁻²⁴ this is not universally the case.

This review emphasizes the varied enzymology of SAM in biological transformations that are not expected to involve radical intermediates (Fig. 1). The enzymes will therefore be discussed based on whether they are structurally related to the SAM-dependent methyltransferases (Section 2) or appear to represent different classes of enzymes altogether (Section 3). As will become evident, however, even structurally related enzymes can exhibit unexpected behavior thereby complicating bioinformatics efforts to infer their activities and placing an



Yu-Hsuan Lee was born in Taipei (Taiwan) and received her bachelor's degree from National Taiwan University in 2016, where she spent the last year as an exchange student at the Technical University of Munich. She then joined the graduate program in chemistry at the University of Texas at Austin. In the research group led by Prof. Hung-wen (Ben) Liu, she has studied the mechanisms of

radical SAM enzymes involved in the biosynthesis of nucleoside natural products.



Byungsun Jeon received his PhD from the University of Texas at Austin, where he studied the enzymes and biosynthetic pathway of spinosyn A using kinetic analysis and organic synthesis under the guidance of Prof. Hung-wen (Ben) Liu. Upon completing his studies in 2017, he worked with Prof. Christopher J. Chang at University of California, Berkeley as a post-doctoral researcher. In 2019, he

started his independent career at Korea Institute of Science and Technology (KIST). His current research focuses on organic synthesis of natural products and biased-ligand development for G-protein coupled receptors.



Daan Ren was born in Nanjing, China. He received his bachelor's degree from Nanjing University in 2016. He then joined Prof. Hung-wen Liu's group at the University of Texas at Austin as a graduate student. His graduate research has focused on the biosynthesis of C-nucleoside natural products and elucidating the mechanisms of enzymatic reactions.



Hung-wen (Ben) Liu was born in Taipei (Taiwan). He completed his PhD degree with Prof. Koji Nakanishi from Columbia University in natural product chemistry and continued his postdoctoral research with Prof. Christopher Walsh at MIT. He started his independent career in the chemistry department at the University of Minnesota in 1984 and moved to University of Texas at Austin in 2000. His

most recent research focuses on atypical sugar biosynthesis, radical SAM enzymology and mechanistic investigations of enzyme catalyzed transformations during natural product biosynthesis.






Fig. 1 Distinct uses of SAM in enzymatic reactions.

emphasis on experimental investigation. Research has nevertheless made steady progress, and since 2010 there has been a considerable expansion in the recognized scope of SAM

enzymology observed in nature. Furthermore, continued study in the future is certain to fill in many of the gaps in the current understanding of these enzymes as well as identify new and potentially unforeseen biological reactions involving SAM.

2 Enzymes with a methyltransferase (MT) fold

The electrophilicity of the SAM sulfonium moiety is illustrated by its involvement in a number of different biological alkylation reactions (Fig. 2). However, the regioselectivity of nonenzymatic alkyl transfer from sulfonium ions bearing three different primary alkyl groups cannot be precisely controlled by tuning either the steric bulkiness of the substituents or other reaction conditions such as the solvent, counter ion or polarizability.^{25–28} Consequently, the regioselectivity of enzyme catalyzed alkyl transfer from SAM appears to originate in variations in the enzyme active sites that tune the relative positioning between the alkyl donor and the nominal substrate, which serves as the alkyl acceptor. Moreover, a number of enzymes have also evolved active site arrangements that preferentially recognize derivatives of SAM thereby further increasing the scope of biological alkylation reactions. Protein crystallography has thus become a critical tool in understanding the details of SAM-dependent alkylation reactions. The following section is




Fig. 2 Summary of SAM-dependent alkylation reactions, showing the by-product generated following each group transfer. (A) Methyl transfer yields *S*-adenosyl homocysteine (SAH, 3), (B) ACP (3-amino-3-carboxypropyl) transfer yields 5'-methylthioladenosine (MTA, 4) and (C) adenosyl transfer yields L-methionine (5).



focused mainly on more recent findings regarding transfer of the 3-amino-3-carboxypropyl group and the modified methyl group from SAM, each catalyzed by an enzyme adopting a methyltransferase fold. Finally, a brief discussion is included of the adenosyl transfer enzymes, which lack a methyltransferase fold yet utilize SAM as the adenosyl donor.

2.1 SAM-dependent alkylation reactions

A number of enzymes have been discovered to selectively catalyze transfer of the 3-amino-3-carboxypropyl (ACP) group from the SAM sulfonium to an acceptor nucleophile expelling methylthioadenosine (MTA) as the leaving group. This type of chemistry has been described in the biosynthesis of several natural products as well as modification of tRNA and rRNA.^{29–31} Moreover, analogous reactions have also been reported for derivatives of SAM and in particular decarboxy-SAM (dc-SAM, **6**) leading to aminopropyl transfer reactions observed during the biosynthesis of spermidine (**7**) (Fig. 3).^{32–34} Catalysis of both ACP transfer and aminopropyl transfer involves an inversion of the stereocenter consistent with a direct nucleophilic displacement

reaction.^{35–37} Substrate binding is of particular importance for regioselective alkyl transfer, and thus the emphasis of the following discussion will be placed on how SAM binding allows for selective C–S bond cleavage.

2.1.1 3-Amino-3-carboxypropyl transfer (small molecules). Nicotianamine (**8**) is a precursor to phytosiderophores, which facilitate metal acquisition by plants.^{38,39} Its biosynthesis involves nicotianamine synthase (NAS),⁴⁰ which catalyzes the condensation of three ACP moieties each from one molecule of SAM and azetidine ring cyclization, though the timing of cyclization remains elusive (Fig. 4A).^{41–44} Although NAS had previously only been found in plants and fungi, recent studies have indicated that NAS-like enzymes exist in both archaea and bacteria as well.^{45,46}

Archaeal NAS from *Methanothermobacter thermautotrophicus* (MtNAS) was found to copurify with thermonicotianamine (tNA, **9**), which is structurally distinct from nicotianamine in that the first condensation partner is a glutamate (**10**) rather than an ACP moiety (Fig. 4B).⁴⁵ Various crystallographic snapshots of MtNAS in complex with its substrate, a proposed intermediate and its product have led to a mechanistic proposal involving two rounds of ACP transfer to glutamate leading to tNA.^{45,47} In particular, the active site of MtNAS extends deep into the protein interior from a solvent exposed entrance allowing for sequential binding of glutamate and SAM.⁴⁵ As SAM binds, glutamate translocates from the donor site at the entrance of the reaction chamber to the buried acceptor site. Transfer of the ACP moiety from SAM to glutamate or the intermediate AP₂-Glu₁ (**11**) is facilitated by Glu81 and Tyr107, which reside in the middle of the active site and deprotonate the nucleophilic amine on the acceptor substrate (*i.e.*, Glu or AP₂-Glu₁) (Fig. 4C).⁴⁵




Fig. 3 Aminopropyl group transfer during the biosynthesis of spermidine. SAM is first decarboxylated to form decarboxy-SAM (dc-SAM, **6**) before the 3-amino-3-carboxypropyl group is transferred to putrescine to yield spermidine (**7**).




Fig. 4 (A) Nicotianamine synthase (NAS) catalyzes the formation of nicotianamine (**8**) from three molecules of SAM. (B) Molecular mechanism proposed for MtNAS during the biosynthesis of thermonicotianamine (tNA, **9**). (C) Crystallographic snapshot of MtNAS (PDB ID: 3FPF) complexed with MTA (pink) and tNA (tan). Nucleophilic displacement was facilitated by deprotonation of the substrate nitrogen by Tyr107 and Glu81 (cyan). (D) The NAS-like enzyme CntL catalyzes transfer of one ACP to D-His (**13**) providing x-NA (**14**) as the precursor of bacterial metallopeptide, staphylopine (**12**).



CntL is a bacterial NAS-like enzyme that participates in the biosynthesis of staphylopine (12),⁴⁶ a metallophore in the pathogenic bacterium *Staphylococcus aureus*.^{48–50} Mutant strains that do not produce CntL show reduced virulence and decreased intracellular metal content.⁴⁶ CntL catalyzes the transfer of ACP from SAM to D-histidine (13) forming xNA (14) (Fig. 4D).⁴⁶ The structure of CntL was solved at 1.81 Å and found to be similar to that of MtNAS.⁵¹ Both MtNAS and CntL possess a Rossmann fold MT C-terminal domain with a characteristic glycine-rich motif that binds SAM as the ACP donor.^{45,51} An open to closed conformational change upon substrate binding was noted for CntL and involves the partially disordered inter-domain linker in the CntL–SAM binary complex becoming an ordered α -helix in the CntL–MTA–xNA ternary complex.⁵¹ The mechanism of CntL catalyzed ACP transfer is otherwise analogous to that of MtNAS. In addition to NAS and CntL, many other NAS-like enzymes have been identified or proposed in the biosynthesis of structurally diverse metallophores with an amino group serving as the acceptor nucleophile, which underscores the importance of SAM-dependent ACP modification of metallophores in bacteria.^{52–54}

Microcin C (15) produced from *E. coli* is a potent inhibitor of aspartyl-tRNA synthetase.⁵⁵ The biosynthesis of microcin C involves aminopropylation of the phosphate which enhances the inhibitory effect by approximately 10-fold.⁵⁶ Two enzymes have been proposed to catalyze aminopropylation including a SAM-dependent methyltransferase MccD (PDB ID: 5FCD) and the N-terminal pyridoxal 5'-phosphate (PLP)-dependent decarboxylase domain of MccE (Fig. 5, route a).⁵⁶ *In vitro* characterization of the two enzymes initially suggested that aminopropylation may proceed with ACP transfer to Mcc¹¹²⁰ (16) catalyzed by MccD followed by decarboxylation of the alkylated intermediate Mcc¹²²¹ (17) catalyzed by MccE.⁵⁷ However, an alternative route similar to polyamine biosynthesis, where decarboxy-SAM (6) is the aminopropyl donor such that MccE and MccD operate in the reversed order (Fig. 5, route b), has not been completely ruled out. Moreover, the activity of




Fig. 6 Other natural products with ACP groups derived from SAM.

MccD can be significantly enhanced by the nucleosidase Mtn, which can hydrolyze the MTA by-product and presumably alleviate product inhibition.⁵⁷

Additional ACP transferases with small molecule substrates are also involved in the biosynthesis of isonocardicin C (18),^{58,59} betaine lipid (19),⁶⁰ discadenine (20)^{61,62} and 2-(3-amino-3-carboxypropyl)-isoxazolin-5-one (ACI) (21) (Fig. 6),^{63,64} however, none of these enzymes appear to be homologous to the aforementioned ACP transferases NAS, MtNAS, CntL and MccD. With the exception of ACI, ACP transfer from SAM is respectively catalyzed in these cases by NAT, BtaA and discadenine synthase. However, while the *in vitro* formation of ACI is known to require a partially purified enzyme fraction from sweet pea seedlings as well as the presence of ATP and magnesium,^{63,64} the responsible enzyme as well as the role of ATP remains elusive.

2.1.2 ACP transfer (RNA). Another class of ACP transferases is involved in the maturation of eukaryotic and archaeal 18S rRNA, where a uridine base (U1191 in yeast, U1248 in human) in the loop capping helix 31 undergoes a 3-step modification consisting of pseudouridylation, methylation and ACP transfer to yield 1-methyl-3-(3-amino-3-carboxypropyl)-pseudouridine (m¹acp³Ψ, 22, see Fig. 7A).⁶⁵ The participation of Tsr3 in ACP transfer was revealed by differences in the nucleoside profile of 18S rRNA isolated from a collection of mutant strains with random gene deletions.⁶⁶ Crystal structures of archaeal Tsr3 from *Vulcanisaeta distributa* (VdTsr3) and *Sulfolobus solfataricus* (SsTsr3) have been solved with and without SAM-bound, respectively.⁶⁶ The C-terminus of Tsr3 is closely packed with the N-terminal domain and adopts a SPOUT fold, which is characterized by a deep trefoil knot and often found in the structures of RNA methyltransferases.^{67,68} In contrast, its functional counterpart Tyw2 has a Rossmann fold and also catalyzes ACP transfer from SAM during the biosynthesis of wybutosine (23).⁶⁹ Archaeal Tsr3 and its closest structural homolog Trm10,^{70,71} which is a tRNA guanidine N-methyltransferase (G, 24, to m¹G, 25) found in archaeans and eukaryotes (Fig. 7B),⁷² employ a monomeric quaternary structure while the majority of enzymes in this superfamily are homodimers.⁷³

Comparison of the crystal structures of VdTsr3 and Trm10 with SAM bound reveals overall similarity at the interface of the N- and C-terminal domains.⁶⁶ This is where SAM binds with the adenine moiety buried in a hydrophobic pocket and the ribose




Fig. 5 Aminopropylation during maturation of microcin C (15). MccD can catalyze transfer of an ACP group onto Mcc¹¹²⁰ (16), which can further be decarboxylated in a reaction catalyzed by MccE (route a). An alternative pathway starts with the formation of dc-SAM catalyzed by MccE followed by MccD mediated aminopropyl transfer (route b).






Fig. 7 Enzymatic functions of (A) Tsr3 and (B) its closest homolog Trm10. Tsr3 catalyzes ACP modification of a pseudouridine residue whereas Trm10 catalyzes methylation of a guanosine or adenosine residue. (C) Structure of wybutosine highlighting the ACP moiety transferred under the action of Tyw2. Solvent accessible area of (D) Tsr3 (PDB ID: 5APG) and (E) Trm10 (PDB ID: 4JWF) is shown colored with the most hydrophilic residues in cyan to the most hydrophobic in gold. In Tsr3, the ACP side chain of Se-SAM (pink) is exposed to solvent and stabilized by hydrophilic interactions. Asp70, Ser72 and Trp73 in the DTW domain are in purple. In contrast, the ACP side chain of SAH (pink) in Trm10 is buried, and the entrance to the binding cavity is lined with hydrophobic residues. (F) Enzymatic function of TapT. (G) Sequence alignment of Tsr3 and TapT at the DTW domain (gray) (Ec: *Escherichia coli*; Vp: *Variovorax paradoxus*; Cs: *Clostridium saccharoperbutylacetonicum*; Vd: *Vulcanisaeta distributa*; Ss: *Sulfolobus solfataricus*; Hs: *Homo sapiens*; Sc: *Saccharomyces cerevisiae*).

moiety hydrogen-bonded to the protein backbone. However, closer inspection of the VdTsr3 active site shows the carboxyl moiety of ACP positioned towards the entrance of the substrate binding pocket surrounded by hydrophilic residues, while the methyl group is directed into a hydrophobic pocket consisting of an unconserved aliphatic residue and a conserved tryptophan (Trp73 in VdTsr3, see Fig. 7D).⁶⁶ Mutation of the Trp to Ala results in both diminished SAM affinity and ACP transferase activity in SsTsr3.⁶⁶ Furthermore, Tsr3 employs a conserved residue (Asp70 in VdTsr3) that serves as a general base, whereas the equivalent Asp210 in Trm10 is not essential for methyl transfer.⁷⁴ The above catalytically important residues in Tsr3 constitute the DTW motif with the conserved primary sequence (D/E)X(T/S)W (Fig. 7G),⁷⁵ in which Asp is the possible catalytic base and Trp assists in methyl group positioning.⁷⁶ Proteins with this domain are classified into the TDD superfamily, which is named after the three representative protein members Tsr3, DTWD1 and DTWD2.⁷⁵

Although ACP transfer catalyzed by Tsr3 is the final step in $m^1\text{acp}^3\Psi$ modification during 18S rRNA maturation, studies of yeast mutants lacking enzymes for pseudouridylation and methylation have suggested that *in vivo* ACP transfer activity is independent of either modification such that the corresponding mutants can produce acp^3U (27) or $\text{acp}^3\Psi$, respectively.^{77,78} These results suggest that Tsr3 and its homologs can catalyze modification of unmodified uridine residues in RNA. Two recent studies involving comparative genomic and ribonucleome analysis have supported this hypothesis by showing that the bacterial Tsr3 homolog TapT from *E. coli* catalyzes acp^3U (27) modification at U47 (26) in the V-loop of several bacterial tRNAs (Fig. 7F).^{76,79} While a structure of TapT is not yet available, sequence analyses and mutational studies imply that TapT

harbors the catalytically important DTW motif similar to that of archaeal Tsr3 despite overall low sequence identity (Fig. 7G).⁷⁶

2.1.3 Carboxymethyl transfer. In Section 2.1.2, several SAM-dependent enzymes were described that catalyze transfer of the 3-amino-3-carboxypropyl (ACP) group to a nucleobase on tRNA thereby improving the thermostability and maintaining its normal function.^{76,77,79,80} 5'-Carboxymethoxyuridine (cmo^5U , 28), also known as 5'-oxyacetyluridine, is a modification on the wobble uridine of tRNA often observed in *E. coli* and many other Gram-negative bacteria that broadens recognition of degenerate codons during protein synthesis.^{81–84} Gene manipulation studies have suggested that formation of this unusual nucleobase involves the activity of two putative SAM-dependent methyltransferases CmoA and CmoB (Fig. 8A).⁸⁵ CmoA catalyzes an unprecedented carboxylation of SAM yielding carboxy-SAM (Cx-SAM, 29),^{86,87} the details of which will be discussed in Section 2.5.1. Subsequently, CmoB transfers the carboxymethyl group from 29 to 5'-hydroxyl uridine (30) and generates 5'-carboxymethoxyuridine (28).⁸⁶

While CmoB can catalyze alkyl transfer from both SAM and Cx-SAM, it exhibits a strong preference for Cx-SAM.⁸⁸ In order to identify the structural determinants of this selectivity, the crystal structure of CmoB was solved by Kim and co-workers (Fig. 8B).^{88,89} It was found that the C-terminal domain of CmoB adopts a Rossmann fold shared among typical methyltransferases, whereas the structure of the N-terminal domain, which spans around 100 amino acid residues, is unique to CmoB.⁸⁸ Lys91 was found adjacent to the anionic carboxymethyl moiety and mutation of Lys to Ala completely abolished the production of cmo^5U (28) *in vivo*, the effect of which is attributed to a 19-fold increase in affinity towards SAM and a 275-fold reduction in affinity towards Cx-SAM.⁸⁸ Consequently, Lys91 likely serves to stabilize the carboxymethyl moiety in the active






Fig. 8 (A) Enzymatic functions of CmoA and CmoB. Two CmoA homologs TglE and MccS^{Bam} also catalyze the formation of 29. (B) Crystal structure of Cx-SAM bound to CmoB (PDB ID: 4QNV). The carboxymethyl moiety of Cx-SAM is coordinated by Tyr200, Lys91 and Arg315. The unique 100 residues at the N-terminus of CmoB are shown in green.

site *via* electrostatic interactions thereby enhancing the specificity of CmoB towards Cx-SAM (29) *versus* SAM. Furthermore, unlike typical SAM methyltransferases, CmoB does not appear to employ general base catalysis to increase the nucleophilicity of the acceptor oxygen,⁸⁸ because the substrate hydroxyl has an apparent pK_a of 7.68 which is sufficient for specific acid/base catalysis.⁹⁰

Following discovery of the CmoA/CmoB carboxymethylation cascade, a homologous enzyme pair was found in the biosynthetic pathway of 3-thiaglutarate (31), a nonproteinogenic amino acid with unknown function.⁹¹ TglE/F show high sequence identity with CmoA/B, and TglF has been shown to catalyze the *trans*-carboxymethylation of a norcysteine thiol in a ribosomally synthesized and post-translationally modified peptide (32 → 33) prior to amide hydrolysis by the protease TglG to release 3-thiaglutarate (31) (Fig. 9A).⁹¹ Primary sequence alignment between CmoB and TglF showed that the essential residues including Lys91 for selective binding of Cx-SAM in CmoB are all conserved in TglF. Furthermore, the biosynthetic gene cluster (Mcc^{Bam}) of the Trojan horse peptide-cytidylate antibiotic microcin C (34) encodes MccB^{Bam} which has been shown to catalyze transfer of the carboxymethyl moiety from Cx-SAM synthesized by MccS^{Bam} to its tRNA substrate 35 (Fig. 9B).⁹² The C-terminal domain of MccB^{Bam} (MccB^{Bam}_{CTD}) adopts a methyltransferase fold that does not show

significant similarity with CmoB and is strictly specific for Cx-SAM (29).⁹² Although the site of carboxymethylation in 34 has yet to be determined, this finding implies that the two-enzyme cascade for this modification is operative in both Gram-negative and Gram-positive bacteria.⁹²

The discovery of Cx-SAM demonstrates that SAM can be naturally modified leading to additional diversity among the available types of biological SAM-dependent alkyl transfer reactions. Given the high intracellular concentration of SAM *versus* its modified counterparts such as Cx-SAM,^{88,93} enzymes must employ recognition mechanisms in order to differentiate between the SAM congeners and thereby avoid inhibition or nonproductive shunt reactions. The structures of CmoB demonstrate that interactions between specific residues and the bound SAM analog at least play a part in this specificity. Consequently, it may be possible to engineer new alkyl transferases *via* optimized point mutations to facilitate the selective binding of other non-natural SAM analogs and thereby further expand the range of biocatalytic alkylation reactions for biotechnological purposes.^{94,95}

2.1.4 Adenosyl transfer. Of the three possible alkyl transfer reactions involving SAM (Fig. 2), adenosyl transfer reactions are only known in the hydrolysis of SAM to methionine (5) and adenosine as well as the biosynthesis of 5'-deoxy-5'-haloadenosine.^{96,97} FIA from *Streptomyces cattleya* and SaIL from *Salinispora tropica* are homologous enzymes sharing 35% sequence identity and catalyze adenosyl transfer from SAM to fluoride and chloride ions, respectively (Fig. 10A).^{98–101} The utilization of SAM in these biosynthetic pathways is not immediately obvious, because the resulting 5'-fluoro-5'-deoxyadenosine (5'-FDA, 36) and 5'-chloro-5'-deoxyadenosine (5'-CIDA, 37) are each metabolized by downstream enzymes before yielding the final halogenated natural products (39–42) with only part of the ribose skeleton from SAM remaining (Fig. 10A).^{101,102}

Crystallographic studies of both FIA and SaIL revealed an overall structure distinct from SAM-dependent methyltransferases. The structures of FIA and SaIL instead resemble those of proteins belonging to the DUF62 superfamily despite none of the members having been functionally characterized at




Fig. 9 SAM-dependent carboxymethylation during the biosynthesis of (A) 3-thiaglutarate (31) and (B) McC-like peptidyl cytidylate (34).




Fig. 13 Examples of natural products that contain an azetidine 2-carboxylate (AZE) moiety.

methyltransferase (MT) fold despite each utilizing SAM as the substrate. Therefore, only MT-like cyclases are included in the following section, and the rest will be discussed in Section 3.

2.2.1 Azetidine 2-carboxylic acid. Azetidine 2-carboxylic acid (AZE, 45) is a non-proteinogenic amino acid first isolated from *Convallaria majalis*.¹¹⁴ While it has been suggested that the four-membered ring was biosynthesized *via* intramolecular cyclization of aminocarboxypropyl of SAM, the details remained unelucidated.^{115–117} Since the discovery of AZE, several natural products carrying the AZE moiety have been isolated and their biosyntheses have been proposed to utilize SAM as the precursor to AZE, including nicotianamine (8) and the mugineic acid family of phytosiderophores (47).^{118–120} Recently, AZE

formation during the biosynthesis of both vioprolide (48) and azetidomonamide (49) was independently reported by the Li and Müller groups.^{121–123} The responsible enzymes (*i.e.*, VioH and AzeJ) share only 28% sequence identity and catalyze the conversion of SAM to AZE (45). Similar chemistry has also been proposed for the homologous enzyme BnVI during the biosynthesis of bonnevillamides (50) (Fig. 13).¹²⁴ Both AzeJ and VioH have been reported to display relatively low turnover numbers with the k_{cat} of AzeJ being 17 min^{−1} and that of VioH being too low to readily measure.^{122,123} Furthermore, VioH is most active under alkaline conditions indicative of general base catalysis to facilitate nucleophilic attack of the amino group at C-γ.¹²² However, none of these enzymes have been structurally characterized, and homology models have been inconclusive due to the lack of a suitable template.¹²² Consequently, it remains uncertain how the ACP side chain of SAM is arranged in the active site to facilitate the 4-*exo*-tet cyclization and avoid lactonization.

2.2.2 Cyclopropane. SAM-dependent two-enzyme cascades have recently been reported to install a spirocyclopropyl moiety in CC-1065 (51)¹²⁵ and possibly gilvusmycin (52)¹²⁶ and yatakemycin (53)¹²⁷ as well (Fig. 14). CC-1065 biosynthesis involves the HemN-like radical SAM enzyme C10P and a SAM-dependent methyltransferase C10Q.¹²⁵ In a coupled assay, the two enzymes together catalyze the cyclopropylation of 55 yielding CC-1065 (51), the mechanism of which was proposed to involve intermediate 56 generated from the addition of a SAM methylene radical (54) to the pyrrole ring of the nominal substrate (55) followed by reduction under the action of C10P. The subsequent cyclopropane ring formation is hypothesized to be catalyzed by C10Q and proposed to involve base catalyzed intramolecular elimination of SAH.¹²⁵ The reaction proposed to be catalyzed by C10Q is analogous to other intramolecular elimination reactions involving SAM; however, in this case the methyl group has





Fig. 14 The proposed two enzyme cascade for C10P and C10Q catalyzed formation of the cyclopropane ring during biosynthesis of CC-1065 (51) and related compounds (52 & 53).



type of chemistry has also been found in natural product biosynthesis where SAM serves as the Lewis acid catalyst by essentially donating a methyl cation to the electron rich π -system of the terpenoid precursor. Cyclization of the resulting carbocation thus proceeds in tandem with alkylation before deprotonation, which in the case of cyclopropyl fatty acid synthases is performed by an active site bicarbonate ion (**60**) (Fig. 15B).^{135,136}

2.3.1 Terpene cyclization. Sodorifen (**61**) is a 16-carbon terpene natural product isolated from *Serratia plymuthica*,¹³⁷ which possesses the four-gene *sod* cluster responsible for sodorifen biosynthesis.^{138,139} Biosynthesis involves SodC catalyzed cyclization of farnesyl pyrophosphate (FPP, **59**) to yield the biosynthetic intermediate pre-sodorifen pyrophosphate (PSPP, **62**), which is subsequently converted to sodorifen under the action of SodD (Fig. 16).^{139,140} The tertiary structure of SodC was predicted by *in silico* homology modeling, which allowed for further computation of the reaction mechanism.¹⁴⁰ It has thus been proposed that PSPP formation proceeds through a cyclopropane intermediate (**65**) that is generated upon the carbocationic rearrangement of FPP following SodC catalyzed π -methylation by SAM (**59** \rightarrow **63**).¹⁴⁰ The initially generated cyclohexane carbocation first undergoes ring contraction in a stepwise process involving a neutral bicyclic intermediate (**64** \rightarrow **65** \rightarrow **66**).¹⁴⁰ The resulting carbocation **66** can then undergo additional hydride and methyl shifts directed by an active site Tyr-His dyad leading to PSPP formation with an overall release of energy.¹⁴⁰ Bioinformatics analysis of clustered *sodC* and *sodD* homologs has identified 28 *sod*-like biosynthetic gene clusters that may also encode additional methyltransferases, Rieske proteins and flavodoxin proteins.¹⁴¹ This suggests a number of additional exotic terpenoid ring systems remain to be discovered.

Fig. 15 Cation-induced cyclizations catalyzed by (A) canonical terpene cyclases and by (B) cyclopropane fatty acid synthases.



This journal is © The Royal Society of Chemistry 2023




Fig. 17 (A) Methylation induced cyclization catalyzed by TleD in teleocidin biosynthesis. Crystal structures of (B) TleD hexamer (PDB ID: 5GM2) and (C) TleD active site formed at the interface of two monomers (green & blue) with bound teleocidin A1 (72, tan) and product SAH (pink).

→ 74), which is consistent with deuterium incorporation experiments, followed by intramolecular electrophilic aromatic substitution to yield an intermediate containing an unstable spiro-ring system (75).¹⁴⁴ Subsequent Pictet-Spengler rearrangement^{145,146} and deprotonation would then yield 69–71.

Crystal structures of TleD have been solved with SAH bound (2.5 Å) as well as both SAH and substrate bound (2.8 Å) in the active site.¹⁴⁷ These structures place the sulfur of SAH approximately 4.4 Å away from C25 of the teleocidin A1 (72), which is consistent with the proposed methyltransfer mechanism.¹⁴⁷ Unlike typical SAM methyltransferases, TleD appears to be active as a hexamer with each active site formed at the interface of two TleD monomers in a domain-swapped fashion (Fig. 17B and C).¹⁴⁷ The active site of TleD is tightly lined with hydrophobic residues and thus water is excluded and prevented from quenching the putative intermediate carbocations.¹⁴⁷ The geranyl moiety of the substrate appears to be flexible and assumes two different conformations in the crystal structure; however, molecular dynamics analysis suggests that only one of them is likely to account for the observed product distribution.¹⁴⁷

2.3.3 Stevens rearrangement. Alkylation induced carbocationic rearrangements can also be initiated *via* the methylation of moieties other than alkenes and do not necessarily need to involve cyclization. For example, this type of chemistry appears to be relevant to the biosynthesis of echinomycin (76), which is also known as quinomycin A and is a nonribosomal peptide antiproliferative and antibiotic agent.^{148–150} The structure of echinomycin features a methyl thioacetal linkage, which is introduced *via* a redox-neutral rearrangement of the disulfide bridged precursor triostatatin (78)¹⁵¹ and is believed to lend




Fig. 18 (A) Mechanism of Ecm18 catalyzed rearrangement of a disulfide bridge to a thioacetal. (B) Hydrophobic surface of the Ecm18 substrate binding pocket (PDB ID: 4NEC) colored according to most hydrophobic (gold) and most hydrophilic (cyan) residues. (C) Ecm18/76/SAH ternary complex. The distance between the sulfur of SAH and the methyl group in 76 is consistent with the proposed methyltransfer reaction. The basic nitrogen on His115 (cyan) is near the disulfide bridge and could facilitate ylide 80 formation *via* deprotonation of the methylated sulfonium intermediate 79.

additional stability to the dimer compared to its disulfide counterpart (Fig. 18A).^{152,153} The enzyme responsible for this transformation is the SAM-dependent methyltransferase Ecm18 (Qui8)^{154–156} or Swb8 in the case of the related compound SW-163D (77) and its derivatives.^{157,158}

The catalytic cycle of Ecm18 is hypothesized to proceed through an intermediary sulfonium ion generated *via* SAM-dependent methylation of one of the two sulfur atoms of the disulfide bridge in triostatatin (78 → 79).¹⁵⁴ The sulfonium ion can then undergo α -deprotonation in the presence of a general base to yield an ylide (80), which is capable of a Stevens rearrangement to yield the methylthioacetal linkage (80 → 76). To probe the mechanistic details of Ecm18 catalysis, Hotta and co-workers solved the 1.5 Å crystal structure of Ecm18 with the reaction products SAH and echinomycin (76) bound in the active site.¹⁵⁹ Ecm18 was thus found to adopt a characteristic SAM binding Rossmann fold featuring a compact hydrophobic substrate/product binding site (Fig. 18B).¹⁵⁹ The proposed methylation reaction is consistent with the linear arrangement of the sulfur center of SAH, the transferred methyl group and



the sulfur center on the product. A polar histidine residue resides sufficiently close to the thioacetal and could serve as a general base for α -deprotonation of the substrate sulfonium to yield the putative sulfur ylide **80** (Fig. 18C).¹⁵⁹ Rearrangement of the ylide may then proceed *via* a 3-*endo*-tet cyclization and ring-opening to yield the methyl thioacetal in **76** (Fig. 18A, mechanism a).¹⁵⁹ However, two other mechanistic hypotheses for the rearrangement have also been proposed that do not require intermediary cyclization *via* **81** to effect the Stevens rearrangement as shown in Fig. 18A.^{107,160,161} One involves heterolytic cleavage of the S–S bond followed by thiolate addition to the resulting thionium **82** (mechanism b). The other starts with homolytic cleavage of the S–S linkage and a subsequent radical–radical recombination to construct the C–S bond in **76** (mechanism c).

2.3.4 Alkaloid cyclization. Biosynthesis of a subclass of pyrroloindoline alkaloids has also been shown to involve an alkylation–annulation cascade.¹⁶² Natural products in this family such as physostigmine (**84**) are characterized by complex, fused ring systems derived from tryptophan.¹⁶³ Liu and co-workers discovered that the SAM-dependent methyltransferase PsmD is an early participant in the biosynthesis of physostigmine where it catalyzes cyclization of **85**.¹⁶⁴ PsmD catalysis begins with methylation at C3 of the indole moiety resulting in an iminium ion **86** that undergoes intramolecular nucleophilic addition by a nearby amide nitrogen to afford the heterocycle in **84** (Fig. 19A).¹⁶⁴ A similar methylation induced cyclization has been reported in the biosynthesis of nocardioazine B (**87**) and proposed in indimicin (**88**).^{165,166} Evidence for such chemistry comes from model systems, which have demonstrated that C3 alkylation of the indole moiety always results in intramolecular addition of the adjacent amide.^{167,168} Thus, only the alkylation step appears to require enzyme catalysis with subsequent cyclization following nonenzymatically. Such alkylation–annulation tandem reactions have also been utilized in the total synthesis of an expanded scope of alkylated pyrroloindoline scaffolds in which alkyl bromides are used as the alkyl donor (Fig. 19B).¹⁶⁸

2.4 SAM as a prosthetic group

Apart from direct participation in the group transfer reactions, SAM can also serve as a prosthetic group for many enzymes that

are annotated as SAM-dependent methyltransferases. Among these examples, SAM does not undergo any covalent changes in its structure during the reaction.

2.4.1 LepI in leporin biosynthesis. Leporins (**89**) are pyridone alkaloids exhibiting mild antibacterial activity.¹⁶⁹ Within the *lep* biosynthetic gene cluster, LepI is annotated as a SAM-dependent *O*-methyltransferase and contains a C-terminal Rossmann fold for SAM binding.¹⁷⁰ However, *in vitro* analysis suggests that LepI accelerates a stereoselective dehydration reaction of its nominal substrate (**91**, the stereochemistry of the carbon with the departing hydroxyl group is not yet known) to (*E*)-**92**, and facilitates a subsequent regioselective hetero-Diels–Alder (HDA) reaction (**92** → **89**) (Fig. 20A). Meanwhile, LepI can also convert a shunt intramolecular-Diels–Alder (IMDA) product **93** to **89** *via* a retro-Claisen rearrangement.¹⁷⁰ Consistent with its annotation, however, catalytically active LepI copurifies with SAM at almost 90% occupancy. Furthermore, sinefungin (SFG, **94**), which is an isosteric analog of SAM, also supports LepI catalysis, while SAH (**3**) is a strong inhibitor of both the dehydration and retro-Claisen rearrangement. Therefore, the positive charge on SAM or sinefungin appears to play a key role in LepI catalysis even if SAM itself remains intact.¹⁷⁰

The positive charge on SAM and sinefungin was originally proposed to play an electrostatic role in catalysis, since coordination of the substrate to a sulfonium moiety is suggested by computational simulations to accelerate the cyclization and drive the periselectivity towards the favored product.¹⁷⁰ However, the crystal structure of LepI complexed with SAM and an unreactive substrate analog (**90**) revealed that the distance between the two ligands is too remote for coordination (Fig. 20A and 21),¹⁷¹ whereas a product complex between LepI/SAM/**89** indicated that the guanidinium group of Arg295 forms a hydrogen bond with the amide carbonyl of the LepI product (**89**).¹⁷¹ In addition, the His133 residue essential for LepI catalysis is near the 4-O of **89**, implying its role in deprotonation of **91** (Fig. 20B). The resulting imidazolium of His133, along with the guanidinium of Arg295, is thus hypothesized to stabilize the transition-state during cycloaddition *via* **92**.¹⁷¹ This conclusion has also been corroborated by three other independent crystallography studies.^{172–174} Sequence and structural comparison between LepI and its closest homolog OxaC, which is an *O*-methyltransferase catalyzing the methyltransfer of meleagrins (**95**) to yield oxaline (**96**) (Fig. 20D),¹⁷⁵ revealed that Arg295 in LepI aligns with an essential histidine (His313) in OxaC (Fig. 20G).¹⁷¹ His313, along with Glu369, is conserved among functional *O*-methyltransferases for substrate deprotonation, and the substitution of the former in LepI may provide an evolutionary perspective on the mechanistic shift from performing methylation to dehydration and cyclization.

Such repurposed methyltransferases catalyzing cyclization reactions have also been found to participate in other biosynthetic pathways. For example, fischerin (**97**) features a *cis*-decalin moiety, the biosynthesis of which is not immediately obvious. The dedicated cyclase FinI shows 25% sequence identity and structural homology with LepI and also co-purifies with SAM as does LepI.^{170,176} Moreover, FinI catalyzed cyclization of **98** to **99** is believed to involve an *exo*-selective DA reaction




Fig. 19 (A) Methylation induced cyclization in pyrroloindoline alkaloid biosynthesis. (B) Synthesis of pyrroloindolines *via* alkylation–cyclization cascade.






Fig. 20 (A) LepI catalyzed dehydration–cyclization cascade during leporin C biosynthesis. Although the stereochemistry at C7 in **91** is currently unclear, it has been shown that the diastereomer of **91** is not accepted by LepI. Comparison of crystal structures of (B) LepI/SAM/**89** complex (PDB ID: 6IX9) and (C) OxaC/SFG/**95** complex (PDB ID: 5W7S). In LepI, SAM is too far from the product (**89**, in tan) for methyl transfer; however, in the methyltransferase OxaC, the substrate mealegrin (**95**) (tan) is located in between SFG and the His313–Glu369 dyad. In contrast, the catalytic histidine is substituted with Arg295 in LepI such that deprotonation cannot occur. Green ribbons in LepI are residues from the second monomeric peptide. (D) OxaC catalyzed methylation during formation of oxaline (**96**). (E) Structure of sinefungin (**94**). (F) Cyclization of **98** catalyzed by FinI. (G) Sequence alignment of LepI, FinI and OxaC highlighting the conserved SAM binding arginine residue (Arg310) that interacts with SAM, the catalytic base (His313) in OxaC (in gray boxes) and the essential arginine (Arg295/Arg313) in the pericyclases in red.




Fig. 21 Crystal structures of (A) LepI (PDB ID: 6IX5), (B) SpnF (PDB ID: 4PNE) and (C) SpnL (PDB ID: 7V6H). Each enzyme contains a C-terminal Rossmann fold catalytic domain containing seven β -strands (cyan) surrounded by five α -helices (blue), while the N-terminal domains differ between each protein. The cofactor, SAM (**1**) for LepI and SAH (**3**) for SpnF and SpnL (pink with a gray surface), binds similarly at the C-terminal domain of all three enzymes. The substrate analog (**90**) bound in the LepI active site is shown with a green surface.

(Fig. 20F), which is rare among known pericyclases.¹⁷⁶ Unlike the case of LepI, however, SAM and the substrate **98** bind the FinI active site so as to place the sulfonium methyl of SAM close to the amide nitrogen of **98** at a distance of 3.4 Å suggesting an electrostatic interaction and a possible role for SAM in substrate

recognition. Whereas in the case of LepI Arg295 can act as a Lewis acid to activate the diene for cyclization,¹⁷¹ the homologous residue Arg313 in FinI is unable to do so given the structure of **98** (Fig. 20F). Instead, Arg310 may play a similar role serving to activate the dienophile in the FinI catalyzed reaction by forming a hydrogen bond with the carbonyl group at C7 of **98**. Although Arg310 in FinI is conserved in homologous methyltransferases such as OxaC, where it interacts with SAM, this residue appears to have been evolutionarily repurposed in the FinI active site to interact and likely activate the substrate for cyclization (Fig. 20G).

2.4.2 SpnF and SpnL in spinosyn biosynthesis. SpnF and SpnL participate in the biosynthesis of the tetracyclic polyketide spinosyn A (**100**) and are both annotated as SAM-dependent methyltransferases with a SAM binding Rossmann fold at the C-terminal;¹⁷⁷ however, neither enzyme catalyzes a methyl transfer reaction. SpnF instead catalyzes an intramolecular [4 + 2]-cycloaddition (**101** → **102**), whereas SpnL catalyzes the cyclization of **103**, yielding the first biosynthetic intermediate **108** with the same ring skeleton as spinosyn A.¹⁷⁷ The catalytic properties of SpnF were identified when an approximately 500-fold increase in reaction rate was observed for the cyclization reaction in the presence of the enzyme *versus* its absence (Fig. 22).¹⁷⁷ Furthermore, measurement of secondary kinetic






Fig. 22 Biosynthesis of spinosyn A highlighting the [4 + 2] cycloaddition catalyzed by SpnF and two proposed mechanisms for the trans-annulation catalyzed by SpnL.

isotope effects (KIE) suggested that both non-enzymatic and SpnF-catalyzed cyclization likely proceed *via* a stepwise rather than a concerted mechanism,¹⁷⁸ in agreement with computational studies.^{179,180} In contrast to LepI, however, active SpnF copurifies with SAH rather than SAM.¹⁸¹ The crystal structure of the SpnF–SAH binary complex was solved at 1.50 Å (Fig. 21B), and computational docking of substrate or product into the hydrophobic cavity of SpnF implied that neither substrate nor product exhibits direct contact with the SAH moiety.¹⁸¹ Therefore, SAH may serve as a component necessary for the integrity of the SpnF structure, and the positive charge of the sulfonium ion of SAM is less likely to be essential for catalysis.¹⁸¹

Following SpnF catalyzed [4 + 2] cycloaddition and SpnG catalyzed glycosylation,¹⁸² SpnL catalyzes the transannular carbon–carbon bond formation between C3 and C14 of **103** to generate **108** (Fig. 22).¹⁷⁷ Two possible mechanisms have been postulated, the first involving deprotonation at C12 to form an enolate (**104**) poised for Michael addition at C3 giving **105**. Alternatively, a nucleophilic residue in the active site may first

add to C13 to yield intermediate **106**, which may then cyclize into **107** prior to elimination of the assisting nucleophile. Involvement of a nucleophilic residue during modification of the β -carbon of an enone moiety is indicative of a Rauhut–Currier (RC) reaction^{146,183} and is reminiscent of thymidylate synthase chemistry.¹⁸⁴ When the SpnL reaction was conducted in buffered D₂O, only a single solvent deuterium was incorporated into the product; however, no significant primary KIE was observed with the C12-deuterated substrate.¹⁸⁵ Moreover, a C13-fluorinated substrate analog **109** led to inactivation of SpnL *via* the formation of a covalent enzyme–substrate adduct (**112**) demonstrated by mass spectrometry (Fig. 23A).¹⁸⁵ Thus, all the biochemical evidence is consistent with an RC mechanism. SpnL shares 35% sequence identity with SpnF and also copurifies with bound SAH.¹⁸⁵ The binary complex of SpnL with SAH bound was solved at 3.05 Å (Fig. 21C) and demonstrated that Cys60 resides nearby the bound SAH with its thiol moiety buried and apparently inaccessible from the active site cavity (Fig. 23B).¹⁸⁵ Consequently, a conformational change upon




Fig. 23 (A) C13-fluorinated substrate (**109**) leads to covalent modification of SpnL. (B) Crystal structure of SpnL/SAH binary complex (PDB ID: 7V6H) highlighting the SAH (**3**) binding site and the catalytic Cys60 (carbons as cyan and sulfur as yellow spheres), which is buried in the protein interior.






Fig. 24 (A) Cyclization catalyzed by IccD during the biosynthesis of ilicicolin H. (B) Dehydration–cyclization reactions catalyzed by SAM-independent pericyclases that each adopt a SAM-binding fold.




Fig. 25 SlnM catalyzes spirocyclization during salinomycin biosynthesis. SAM is proposed to stabilize an anionic aspartate residue via electrostatic interactions to facilitate substrate protonation.

substrate binding was proposed to allow for Cys60 exposure to the substrate thereby facilitating catalysis.

During biosynthesis of the fungal secondary metabolite ilicicolin H (**113**), the cyclohexene moiety is believed to be constructed *via* an inverse-electron demand Diels–Alder (DA) reaction catalyzed by IccD, which harbors an intact SAM-binding motif yet does not copurify with SAM (Fig. 24A).¹⁸⁶ Moreover, two additional groups of methyltransferase-like proteins were also found to catalyze pericyclic reactions. Epil, Upil and Hpil are proposed to catalyze a hetero-DA reaction yielding **119** on the quinone methide substrate (**117**) generated upon dehydration of **116**, while PdxI, AdxI and ModxI catalyze conversion of the same substrate to **121** *via* a proposed Alder-ene reaction (Fig. 24B).¹⁸⁷ The catalytic cycles of these pericyclases, however, do not depend on either SAM or SAH indicating they are mechanistically distinct from those of LepI or SpnF and SpnL.^{186,187} Pericyclases characterized by a SAM-binding fold appear both evolutionarily and catalytically distinct.^{188–190}


2.4.3 SlnM in salinomycin biosynthesis. Salinomycin (**122**) is a broadly utilized coccidiostat featuring a bis-spiro tricyclic structure with a polyether linkage.¹⁹¹ Its derivatives have recently drawn attention in the field of oncology due to their antitumor potential.^{192,193} Within the biosynthetic gene cluster, SlnM is annotated as a SAM-dependent *O*-methyltransferase containing a glycine-rich motif canonical for SAM binding.^{194,195} Gene inactivation of SlnM results in the accumulation of a ring-opened intermediate (**123**) suggesting that SlnM catalyzes the dehydration and subsequent spirocyclization of the putative precursor species **124** (Fig. 25).¹⁹⁶ The presence of SAM is essential for SlnM activity, though it is not consumed over the course of the reaction, and SAM can be replaced by its isosteric analog sinefungin (**94**) but not SAH.¹⁹⁶ Consequently, SAM likely

serves to both help maintain the structural integrity of SlnM and also contribute directly to catalysis in some manner that requires the positively charged sulfonium moiety. On the basis of sequence alignment, mutational studies and computational modelling, Asp58 and Asp186 were identified as catalytically important residues, with one presumably serving as the general acid that facilitates the dehydration reaction.¹⁹⁶ It has therefore been hypothesized that the sulfonium of SAM facilitates elimination of water by stabilizing the negative charge buildup on the aspartate residues as they protonate the leaving group (Fig. 25).¹⁹⁶ However, the exact positioning between SAM, substrate and these Asp residues has yet to be fully established due to the lack of a crystal structure.


2.4.4 SAM-dependent hydroxylase. The first example of a SAM-dependent hydroxylase is RdmB, which catalyzes decarboxylative hydroxylation of 15-demethoxy- ϵ -rhodomycin (**125**) to form the anthracycline precursor β -rhodomycin (**126**).¹⁹⁷ While RdmB requires SAM for activity, it operates as an O_2 -dependent oxidase that recognizes several different substrates such as **127** in addition to **125** (Fig. 26A and B).^{197–201} These substrates all feature a carboxylate moiety at C10, which upon decarboxylation forms the anion **128** that is stabilized by extended conjugation. Moreover, sinefungin (**94**) but not SAH (**3**) is able to support hydroxylase activity, which led to the hypothesis that electrostatic stabilization of the anionic intermediate **128** by SAM is essential for catalysis.²⁰⁰ The intermediate **129** can then activate molecular oxygen to yield the peroxide **130**. In the presence of reduced glutathione (GSH), **130** is converted to the hydroxylation product **131**.

There is reason to believe that RdmB may have diverged from an evolutionary lineage of SAM-dependent *O*-methyltransferases upon losing methyltransferase activity. DnrK is a homolog of RdmB (52% sequence identity) that catalyzes methylation at O-4 of carminomycin (**133**) yielding





Recent structural comparisons of RdmB and DnrK coupled with extensive mutational studies by Grocholski and co-workers have provided even more insight into this question.²⁰⁵ The Michaelis complex of RdmB is solvent inaccessible,²⁰⁰ which prevents protonation of the anionic intermediate **129** by water. Closure of the active site is facilitated by a loop that includes Phe300 and blocks the channel connecting the substrate binding site to the external media (Fig. 27C).²⁰⁵ In contrast, the corresponding Phe296 in DnrK faces away from the channel leaving the active site of DnrK solvent accessible (Fig. 27D). Notably, DnrK catalyzes the conversion of **127** to **132** *in vitro* (Fig. 26B).²⁰⁵ Although the timing of methylation at C4-OH is unclear, this observation is consistent with the hypothesis that upon decarboxylation at C15, the resulting anion (*i.e.*, **129**) is quenched by water in the DnrK active site.²⁰⁵ Moreover, the different orientation of Phe300 in RdmB and Phe296 in DnrK is due to Ser301 in RdmB, which is missing in the primary sequence of DnrK²⁰⁵ (Fig. 27B); however, Phe296 in DnrK can be rotated towards the channel thus blocking solvent access by insertion of a serine residue between Phe296 and Thr297, which mimics the Ser301 in RdmB (Fig. 27B).²⁰⁵ As expected, this DnrK insertion mutant (DnrK-Ser) did exhibit hydroxylase activity, implying the importance of solvent extrusion to support hydroxylation activity.²⁰⁵






Fig. 28 NDUFAF5 catalyzes modification of the substrate protein NDUFS7 at Arg73. The precise site of hydroxylation is unknown.

Human complex I is a respiratory chain protein that drives proton pumping for ATP synthesis by mediating electron transfer between NADH and ubiquinone.^{206,207} NDUFAF5 is an essential protein in the biogenesis of human complex I and is characterized by a conserved SAM-binding motif at the C-terminus. Suppression of NDUFAF5 expression results in reduced hydroxylation of an arginine residue in the protein NDUFS7, which is a subunit in human complex I.²⁰⁸ Thus, NDUFAF5 has been hypothesized to be an arginine hydroxylase targeting NDUFS7 (137 → 138) (Fig. 28). The secondary structure of NDUFAF5 is predicted to be similar to that of RdmB in both the substrate and SAM binding regions indicating that NDUFAF5 might adopt an active site structure similar to that of RdmB.²⁰⁸ While the *in vivo* activity of NDUFAF5 has been correlated with arginine hydroxylation, the *in vitro* activity and mechanistic details regarding whether the hydroxylase activity is SAM or SAH dependent and how the arginine is activated for oxidation remains unclear.

2.4.5 SAM-dependent decarboxylase. Brevione E (139) is a hexacyclic meroditerpenoid demonstrating allelopathic activity.²⁰⁹ The biosynthesis of brevione E involves BrvO, which is annotated as a SAM-dependent methyltransferase and proposed to catalyze the decarboxylation of 140/141 to 145 (Fig. 29).²¹⁰ Decarboxylation of 140/141 can be accelerated in the




Fig. 29 Proposed mechanism for BrvO catalyzed decarboxylation.

presence of BrvO and SAM. The mechanism is hypothesized to involve initial rearrangement of the 2',3'-alkene in 140/141 to the 3',4'-alkene in 143 prior to decarboxylation and generation of the oxyanionic intermediate 144. It is possible that SAM participates by helping to stabilize the negatively charged intermediate (144) *via* electrostatic interactions similar to the role it has been proposed to play during the catalytic cycles of SlnM and RdmB.^{196,200} However, the characterization of BrvO is currently incomplete including whether SAH and SFG can also support BrvO activity. Consequently, further studies will be required to understand the mechanism of this enzyme.

2.5 SAM as a nucleophile

2.5.1 Carboxy-SAM. In Section 2.1.3, the formation of 5'-carboxymethoxyuridine (28) residues at the wobble position of tRNA catalyzed by CmoB was discussed as an example of an alkylation reaction involving carboxy-SAM (Cx-SAM, 29) as the alkyl donor.⁸⁶ However, the biosynthesis of Cx-SAM itself requires carboxylation of the SAM methyl group catalyzed by CmoA with prephenate (146) being the carboxylate donor.⁸⁶ CmoA homologs have also been found in a broad range of organisms and natural product biosynthetic pathways including 3-thioglutamate (31) in *Pseudomonas* (TglE)⁹¹ as well as microcin C (34) in *Bacillus* (MccS)⁹² (Fig. 8, Section 2.1.3). The CmoA catalyzed reaction was proposed to involve the nucleophilic addition of a sulfonium methylide (148), generated from α -deprotonation of SAM, to the carbon dioxide liberated upon decarboxylative dehydration of prephenate (Fig. 30A).⁸⁶ Therefore, the CmoA catalyzed reaction differs from other SAM-dependent alkylation reactions in that the methyl substituent on the sulfonium moiety is acting as a nucleophile rather than an electrophile.

The first half of the CmoA catalyzed reaction is hypothesized to proceed analogous to that of prephenate hydratase, where decarboxylation of prephenate facilitates elimination of a hydroxide ion thereby generating CO₂ and phenylpyruvate (147).²¹¹ In the catalytic cycle of prephenate hydratase, the eliminated hydroxide is believed to abstract a proton from an active site threonine residue thereby completing the reaction.²¹¹ As the pK_a of the threonine hydroxyl is close to that measured for the trimethylsulfonium ion,²¹² the same hydroxide is expected to be competent for deprotonating the methyl of SAM. The resulting sulfonium methylide (148) can then add to the eliminated molecule of CO₂ to yield Cx-SAM (29). Moreover, the X-ray crystal structure of Cx-SAM-bound CmoA from *E. coli* revealed an overall Rossmann fold^{86,87} and a cavity adjacent to the Cx-SAM binding site that was predicted to be the prephenate (146) binding site by molecular docking.⁸⁶ Notably, the region of the active site adjacent to the eliminated prephenate hydroxide is lined primarily with hydrophobic residues leaving the methyl group of SAM the only available proton source nearby (Fig. 30B).⁸⁶

Computational substrate docking has also suggested an arrangement of the prephenate hydroxyl and SAM methyl that disallows a typical methylation reaction.⁸⁶ This is consistent with the observation that CmoA does not catalyze O-methylation








Fig. 32 (A) Transamination of SAM catalyzed by BioA. (B) Sequence alignment of RquA with related functional methyltransferases. RquA has two essential aspartate residues (Asp118 and Asp143, numbers in blue) for SAM binding whereas the DXXXGXG (red box) motif typically found in Rossmann fold MTs is missing in RquA. (C and D) Two proposed mechanisms for RquA catalyzed transamination during the biosynthesis of ridoquinone (159).

SAM to its α -keto acid congener while reductively aminating 7-keto-8-aminopelargonic acid (KAPA, **156**) to 7,8-diaminopelargonic acid (DAPA, **157**) during the biosynthesis of biotin (**155**) (Fig. 32A).^{223,224} The mechanism of BioA is thus no different from other PLP transaminases,²²⁵ despite SAM serving as the amine donor rather than a free amino acid. In contrast, the second example appears to be unrelated to PLP chemistry and contributes to the biosynthesis of ridoquinone in *Rhodospirillum rubrum*.²²⁶

Ridoquinone (**159**) differs structurally from ubiquinone (**160**) by replacing an *O*-methoxy group with an amine in a reaction that requires the enzyme RquA.^{227,228} RquA was recently characterized as a manganese-dependent enzyme that catalyzes transfer of the α -amino group from SAM to ubiquinone (**160**) to yield ridoquinone (**159**).²²⁹ While RquA resembles SAM-dependent methyltransferases, only a subset of the conserved residues in the predicted Rossmann fold generally associated with SAM binding are required for activity (Fig. 32B).^{227,229} In addition to a divalent metal ion, RquA catalysis also depends on aerobic conditions as well and is expected to involve a redox component. Indeed, isotope labelling experiments have suggested the oxidative decomposition of SAM to CO₂, MTA and an aldehyde fragment during the transformation.²²⁹ Mechanistic hypotheses for this transformation are shown in Fig. 32C and D; however, much work is still left to be done before the chemistry of this unique enzyme is fully understood.

3 Enzymes without a typical MT-fold

As discussed in the previous sections, evolutionary diversification of SAM-dependent methyltransferases has led to the

emergence of several different types of chemistry. However, there also exist groups of SAM utilizing enzymes that lack the MT-fold making their evolutionary relationship with the SAM-dependent methyltransferases less clear. These enzymes also show a broad range of different activities again emphasizing the mechanistic versatility and intrinsic reactivity of SAM as a biological substrate.

3.1 Pyridoxal 5'-phosphate-dependent enzymes

3.1.1 Cyclopropylation. The sulfonium moiety of SAM represents a good leaving group for substitution reactions thereby explaining its prevalence in various biological alkylation reactions. However, this feature of SAM also suggests that it may be susceptible to intramolecular cyclization reactions. Indeed, under alkaline conditions, SAM undergoes nonenzymatic decomposition to yield methylthioadenosine (MTA, **4**) and homoserine lactone (**44**) as one of its main decomposition products.^{108,109} In principle, SAM should also be capable of a favorable 3-*exo*-tet cyclization to form 1-aminocyclopropane-1-carboxylic acid (ACC, **46**) again with the elimination of MTA if it were not for the nonacidic C- α proton, which has a pK_a of approximately 29.²³⁰ ACC is indeed found as a metabolite from SAM in plants and is the precursor to the important plant hormone ethylene.^{231–233} This difficulty with deprotonation has been overcome in the case of plant ACC synthase by acidifying C1 *via* formation of an external aldimine with PLP (**167**).²³⁴ Deprotonation at C1 thus creates a highly conjugated quinonoid (**168**) and is favorable (Fig. 33). The reaction mechanism of plant ACC synthase has been extensively studied and reviewed;²³⁵ however, the existence of a bacterial counterpart had been previously unknown despite a number of






Fig. 33 Proposed mechanism for the transformation of SAM to ACC (46) and Me-ACC (170) catalyzed by GnmY and Orf29/Orf30, respectively.

cyclopropane containing natural products having been isolated from bacteria.^{111,112,236–238} Therefore, it is of interest to know whether similar chemistry is found in bacteria.

Recently, guanganmycin (GNM, 172), which is a structural homolog of the antitumor antibiotic leinamycin (LNM), was discovered through genome mining.²³⁹ Apart from a hybrid peptide–polyketide backbone, GNM also features an unusual ACC moiety (Fig. 34).²³⁹ Comparative analysis of *lnm*-type biosynthetic gene clusters between guanganmycin producers and non-producers revealed that the gene product of *gnmY* may catalyze the formation of ACC from SAM. This functional assignment was then validated by *in vivo* inactivation of *gnmY*, which rendered the mutant strain unable to produce guanganmycin unless exogenous ACC is supplemented.²⁴⁰ Sequence analysis revealed that GnmY belongs to the aspartate aminotransferase superfamily, and subsequent *in vitro* characterizations confirmed that GnmY catalyzes the conversion of SAM to ACC in a PLP-dependent manner.²⁴⁰ GnmY also exhibits steady-

state kinetic parameters comparable to those of plant ACC synthases and utilizes a catalytic lysine residue (Lys251) to form an internal aldimine (166) with PLP during catalysis.²⁴⁰ Nevertheless, bioinformatics analyses have suggested that GnmY is phylogenetically distinct from the family of plant ACC synthases despite sharing a similar mechanism of catalysis.²⁴⁰

Homologs of GnmY have also been found in the putative biosynthetic gene clusters for two other peptide antibiotics, SW-163C (173) and Q6402A (174),^{157,241} which are each decorated with an unusual 2-methyl-ACC (MeACC, 170) moiety (Fig. 34).^{242,243} In the biosynthetic gene cluster of Q6402A, the gene product of *orf30* shows homology with GnmY suggesting that its activity may involve the formation of MeACC (170).²⁴¹ Initial characterization of purified Orf30 revealed its ability to catalyze the conversion of SAM to ACC.²⁴¹ However, Orf30 exhibits a 1900-fold lower value of k_{cat} compared to GnmY such that SAM may not be the native substrate of Orf30.²⁴¹ The B₁₂-dependent radical SAM methylase Orf29 encoded by a gene fragment upstream of Orf30 was later demonstrated to catalyze methylation of SAM yielding (4'*R*)-4'-methyl-SAM (171).²⁴⁴ This SAM derivative (171) can be efficiently cyclized to give MeACC (170) under the action of Orf30 (Fig. 33).²⁴⁴ It remains to be established how Orf30 distinguishes between SAM and methylated-SAM.

3.1.2 Elimination. The muraymycins (175), sphaerimycin A (177) and caprazamycins/liposidomycins (178) are peptidyl nucleoside natural products that inhibit bacterial cell wall assembly.^{245,246} These antibiotics are structurally characterized by a unique disaccharide core consisting of 5'-amino-5'-deoxyribose (ADR) and 5'-C-glycyridine (GlyU) sugar residues (Fig. 35).²⁴⁷ These nucleoside antibiotic families are distinguished, however, by highly diverse appendages at the amino acid end of GlyU. Feeding experiments with a variety of amino acids enriched with stable isotopes revealed that the aminopropyl linker in muraymycin D1 (176) is derived from *L*-Met (Fig. 35).²⁴⁸ Although such a feature could be explained by an ACP transferase catalyzed reaction in the biosynthetic pathway as described in Section 2.1.1 followed by decarboxylation, no ACP transferase or SAM methyltransferase homolog is




Fig. 34 Natural products that contain SAM-derived cyclopropane moieties.




Fig. 35 Natural products that contain the ADR-GlyU disaccharide. The bold alkyl chain indicates the putative methionine-derived aminopropyl moiety.





Purification and *in vitro* characterization of Mur24 and its homologs LipJ²⁵⁰ and SphL²⁵² indicated that all three enzymes do indeed catalyze ACP transfer from SAM to the substrate (182) amine coupled with the elimination of MTA (Fig. 36).²⁴⁸ To investigate the role of PLP in this reaction, the conserved lysine residue necessary for internal aldimine formation with PLP was mutated leading to a complete loss of ACP transfer activity.²⁴⁸ The catalytic mechanism of Mur24 was further studied using SAM isotopologs, the results of which provided evidence that

More recently, the PLP-dependent enzyme SbzP was found to catalyze formation of the 6,5-bicyclic core of 6-azatetrahydroindane containing anticancer natural products (**184–186**).^{253–255} Feeding experiments established that the



The enzymes described above employ a SAM–PLP quinonoid intermediate (**168**) that can react *via* several different pathways eliminating MTA to either yield an unsaturated iminium or induce an intramolecular cyclization. In the former case, the iminium is susceptible to both electrophilic and nucleophilic addition. Moreover, electrophilic addition can occur at either C-

3.2 Polyketide synthase

3.2.1 Cyclopropylation.

3.2.1 Cyclopropylation. Colibactin (199) is a genotoxic peptide–polyketide metabolite isolated from *E. coli* featuring a cyclopropane ring, which is believed to be responsible for its DNA alkylation activity.^{262–264} Unlike plant ACC synthases and GnmY, which catalyze the conversion of SAM to free ACC (46),^{235,240} the cyclopropane ring in colibactin is constructed by




Fig. 39 Proposed mechanism for the Clbl catalyzed cyclopropylation during the formation of colibactin biosynthetic intermediate **203**.

cyclizing SAM as a component residue during peptide–polyketide chain assembly.²⁶⁵ SAM is first loaded onto a peptidyl carrier protein domain in the NRPS protein ClbH followed by a condensation step, which transfers the acyl chain from the upstream PKS (ClbC, **200**) to the amino group of ClbH–SAM (**201**) to yield intermediate **202** (Fig. 39).²⁶⁵ ClbI, which is annotated as a PKS enzyme, then catalyzes cyclization of the tethered SAM–acyl chain. Sequence alignment reveals that an otherwise conserved cysteine residue in the ketosynthase (KS) domain of ClbI is replaced by a serine (Ser178), and mutation of this residue to alanine abolishes the activity of ClbI.²⁶⁵ It was therefore hypothesized that the serine could act as a general base to deprotonate the α -proton (Fig. 39, mechanism a).²⁶⁵ The generated carbanion can then undergo 3-*exo*-tet cyclization and eliminate MTA (**202** \rightarrow **203**). Alternatively, serine can also serve as a nucleophile to displace MTA thereby forming an enzyme–substrate adduct **204** (Fig. 39, mechanism b) prior to deprotonation induced cyclization.²⁶⁵ As the crystal structure of ClbI or ClbI with SAM-loaded ClbH is currently unavailable, the precise role of the Ser178 and how the non-acidic proton at C- α in **202** is deprotonated is presently uncertain.

4 Conclusions

SAM is ranked among the most frequently observed substrates in enzymatic transformations.¹⁸ The majority of known SAM dependent biological reactions involve methylation; however, regioselective transfer of all three alkyl substituents of the SAM sulfonium have been reported (Sections 2.1 and 2.2) and may represent components of more complex biosynthetic cascades (Section 2.3). In addition to alkyl transfer, several other types of SAM-dependent reactions catalyzed by enzymes have been established including α -deprotonation with the potential for ylide formation (Section 2.5) as well as β -elimination concomitant with elimination of a neutral disulfide (Section 3).

In all cases, these enzyme catalyzed reactions conform to the well-known chemistry of the sulfonium cation, such that the role of the enzyme is to selectively stabilize those transition states representing one mode of reaction over another. Structural investigations of these enzymes have been critical in defining these constraints thereby explaining both the modes of catalysis as well as the observed regiochemistry underlying these various enzyme catalyzed reactions. Moreover, the principles underlying sulfonium enzymology are not limited to SAM-dependent enzymes alone and have also been appreciated among the dimethylsulfoniopropionate (DMSP) lyases found in marine bacteria²⁶⁶ as well as BurG during the biosynthesis of malleicypols.^{267,268}

Many aspects of SAM enzymology, however, remain to be explored. In particular, several enzymes as described in Section 2.4 require SAM even though the catalytic cycles of these enzymes do not appear to involve any sort of covalent rearrangement or fragmentation of SAM itself. In these instances, SAM may be required for proper folding and structural integrity of the enzyme,²⁶⁹ or it may play a more direct role in catalysis by modifying the structural and electrostatic features of the active site.^{270,271}

Many of the SAM-dependent enzymes also exhibit structural features that imply homology with the SAM methyltransferases, whereas others do not, suggesting different evolutionary lineages. Consequently, how these enzymes diversified in some cases from a common evolutionary origin to exhibit the broad range of biochemical activities and unique catalytic roles of SAM seen today is yet another question that remains open to investigation. In any case, SAM has become a focal point of modern enzymology that will likely bring many new discoveries and surprises in the years to come.

5 Conflicts of interest

The authors declare no conflicts of interest.

6 Acknowledgements

We thank Dr Mark W. Ruszczycky for his help with the English and his comments on the accuracy of the content. This work was supported by grants from the National Institutes of Health (GM035906 and GM040541) and the Welch Foundation (F-1511).

7 References

- 1 S. C. Lu, *Int. J. Biochem. Cell Biol.*, 2000, **32**, 391–395.
- 2 P. K. Chiang, R. K. Gordon, J. Tal, G. C. Zeng, B. P. Doctor, K. Pardhasaradhi and P. P. McCann, *FASEB J.*, 1996, **10**, 471–480.
- 3 W. A. M. Loenen, *Biochem. Soc. Trans.*, 2006, **34**, 330–333.
- 4 D. K. Liscombe, G. V. Louie and J. P. Noel, *Nat. Prod. Rep.*, 2012, **29**, 1238–1250.
- 5 M. Blum, H.-Y. Chang, S. Chuguransky, T. Grego, S. Kandasamy, A. Mitchell, G. Nuka, T. Paysan-Lafosse, M. Qureshi, S. Raj, L. Richardson, G. A. Salazar, L. Williams, P. Bork, A. Bridge, J. Gough, D. H. Haft, I. Letunic, A. Marchler-Bauer, H. Mi, D. A. Natale, M. Necci, C. A. Orengo, A. P. Pandurangan, C. Rivoire, C. J. A. Sigrist, I. Sillitoe, N. Thanki, P. D. Thomas, S. C. E. Tosatto, C. H. Wu, A. Bateman and R. D. Finn, *Nucleic Acids Res.*, 2021, **49**, D344–D354.
- 6 A.-W. Struck, M. L. Thompson, L. S. Wong and J. Micklefield, *ChemBioChem*, 2012, **13**, 2642–2655.
- 7 H. J. Sofia, G. Chen, B. G. Hetzler, J. F. Reyes-Spindola and N. E. Miller, *Nucleic Acids Res.*, 2001, **29**, 1097–1106.
- 8 P. A. Frey and O. T. Magnusson, *Chem. Rev.*, 2003, **103**, 2129–2148.
- 9 N. Oberg, T. W. Precord, D. A. Mitchell and J. A. Gerlt, *ACS Bio Med Chem Au*, 2022, **2**, 22–35.
- 10 J. B. Broderick, B. R. Duffus, K. S. Duschene and E. M. Shepard, *Chem. Rev.*, 2014, **114**, 4229–4317.
- 11 W. E. Broderick and J. B. Broderick, *J. Biol. Inorg. Chem.*, 2019, **24**, 769–776.
- 12 M. W. Ruszczycky, Y. Ogasawara and H.-w. Liu, *Biochim. Biophys. Acta*, 2012, **1824**, 1231–1244.
- 13 J. L. Vey and C. L. Drennan, *Chem. Rev.*, 2011, **111**, 2487–2506.



- 14 M. R. Challand, R. C. Driesener and P. L. Roach, *Nat. Prod. Rep.*, 2011, **28**, 1696–1721.
- 15 K. Boswinkle, J. McKinney and K. D. Allen, *J. Bacteriol.*, 2022, **204**, e0019722.
- 16 Y. Nicolet, *Nat. Catal.*, 2020, **3**, 337–350.
- 17 M. W. Ruszczycky, A. Zhong and H.-w. Liu, *Nat. Prod. Rep.*, 2018, **35**, 615–621.
- 18 M. Fontecave, M. Atta and E. Mulliez, *Trends Biochem. Sci.*, 2004, **29**, 243–249.
- 19 S. Roje, *Phytochemistry*, 2006, **67**, 1686–1698.
- 20 Q. Sun, M. Huang and Y. Wei, *Acta Pharm. Sin. B*, 2021, **11**, 632–650.
- 21 L. Barra, T. Awakawa and I. Abe, *JACS Au*, 2022, **2**, 1950–1963.
- 22 H. L. Schubert, R. M. Blumenthal and X. Cheng, *Trends Biochem. Sci.*, 2003, **28**, 329–335.
- 23 P. Z. Kozbial and A. R. Mushegian, *BMC Struct. Biol.*, 2005, **5**, 19.
- 24 J. L. Martin and F. M. McMillan, *Curr. Opin. Struct. Biol.*, 2002, **12**, 783–793.
- 25 K. Umemura, H. Matsuyama and N. Kamigata, *Bull. Chem. Soc. Jpn.*, 1990, **63**, 2593–2600.
- 26 M. Kobayashi, K. Umemura, N. Watanabe and H. Matsuyama, *Chem. Lett.*, 1985, **14**, 1067–1070.
- 27 H. M. R. Hoffmann, *J. Chem. Soc.*, 1965, 823.
- 28 H. Matsuyama, H. Minato and M. Kobayashi, *Bull. Chem. Soc. Jpn.*, 1975, **48**, 3287–3292.
- 29 H. Lin, *Bioorg. Chem.*, 2011, **39**, 161–170.
- 30 A. Noma, Y. Kirino, Y. Ikeuchi and T. Suzuki, *EMBO J.*, 2006, **25**, 2142–2154.
- 31 K. E. Sloan, A. S. Warda, S. Sharma, K. D. Entian, D. L. J. Lafontaine and M. T. Bohnsack, *RNA Biol.*, 2017, **14**, 1138–1152.
- 32 H. Tabor and C. W. Tabor, *Adv. Enzymol. Relat. Areas Mol. Biol.*, 1972, **36**, 203–268.
- 33 Y. Ikeguchi, M. C. Bewley and A. E. Pegg, *J. Biochem.*, 2006, **139**, 1–9.
- 34 A. E. Pegg and A. J. Michael, *Cell. Mol. Life Sci.*, 2010, **67**, 113–121.
- 35 B. T. Golding and I. K. Nassereddin, *J. Am. Chem. Soc.*, 1982, **104**, 5815–5817.
- 36 B. T. Golding and I. K. Nassereddin, *J. Chem. Soc., Perkin Trans. 1*, 1985, 2017–2024.
- 37 G. R. Orr, D. W. Danz, G. Pontoni, P. C. Prabhakaran, S. J. Gould and J. K. Coward, *J. Am. Chem. Soc.*, 1988, **110**, 5791–5799.
- 38 N. von Wiren, S. Klair, S. Bansal, J. F. Briat, H. Khodr, T. Shioiri, R. A. Leigh and R. C. Hider, *Plant Physiol.*, 1999, **119**, 1107–1114.
- 39 C. Curie, G. Cassin, D. Couch, F. Divol, K. Higuchi, M. Le Jean, J. Misson, A. Schikora, P. Czernic and S. Mari, *Ann. Bot.*, 2009, **103**, 1–11.
- 40 J. S. McFarlane and A. L. Lamb, in *Comprehensive Natural Products III: Chemistry and Biology*, ed. H.-w. Liu and T. Begley, Elsevier, Amsterdam, 3rd edn, 2020, ch. 16, vol. 5, pp. 395–414.
- 41 K. Higuchi, K. Kanazawa, N.-K. Nishizawa, M. Chino and S. Mori, *Plant Soil*, 1994, **165**, 173–179.
- 42 A. Herbig, G. Koch, H. P. Mock, D. Dushkov, A. Czihal, J. Thielmann, U. W. Stephan and H. Bäumlein, *Eur. J. Biochem.*, 1999, **265**, 231–239.
- 43 K. Higuchi, K. Suzuki, H. Nakanishi, H. Yamaguchi, N. K. Nishizawa and S. Mori, *Plant Physiol.*, 1999, **119**, 471–480.
- 44 H. Q. Ling, G. Koch, H. Bäumlein and M. W. Ganai, *Proc. Natl. Acad. Sci. U. S. A.*, 1999, **96**, 7098–7103.
- 45 C. Dreyfus, D. Lemaire, S. Mari, D. Pignol and P. Arnoux, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**, 16180–16184.
- 46 G. Ghssein, C. Brutesco, L. Ouerdane, C. Fojcik, A. Izaute, S. Wang, C. Hajjar, R. Lobinski, D. Lemaire, P. Richaud, R. Voulhoux, A. Espaillet, F. Cava, D. Pignol, E. Borezée-Durant and P. Arnoux, *Science*, 2016, **352**, 1105–1109.
- 47 C. Dreyfus, M. Larrouy, F. Cavelier, J. Martinez, D. Pignol and P. Arnoux, *Chem. Commun.*, 2011, **47**, 5825–5827.
- 48 L. Remy, M. Carrière, A. Derré-Bobillot, C. Martini, M. Sanguinetti and E. Borezée-Durant, *Mol. Microbiol.*, 2013, **87**, 730–743.
- 49 K. P. Grim, B. San Francisco, J. N. Radin, E. B. Brazel, J. L. Kelliher, P. K. Párraga Solórzano, P. C. Kim, C. A. McDevitt and T. E. Kehl-Fie, *mBio*, 2017, **8**, DOI: [10.1128/mbio.01281-17](https://doi.org/10.1128/mbio.01281-17).
- 50 L. Song, Y. Zhang, W. Chen, T. Gu, S.-Y. Zhang and Q. Ji, *Proc. Natl. Acad. Sci. U. S. A.*, 2018, **115**, 3942–3947.
- 51 Z. Luo, S. Luo, Y. Ju, P. Ding, J. Xu, Q. Gu and H. Zhou, *FASEB J.*, 2021, **35**, e21575.
- 52 S. Lhospice, N. O. Gomez, L. Ouerdane, C. Brutesco, G. Ghssein, C. Hajjar, A. Liratni, S. Wang, P. Richaud, S. Bleves, G. Ball, E. Borezée-Durant, R. Lobinski, D. Pignol, P. Arnoux and R. Voulhoux, *Sci. Rep.*, 2017, **7**, 17132.
- 53 J. S. McFarlane and A. L. Lamb, *Biochemistry*, 2017, **56**, 5967–5971.
- 54 M. C. Mastropasqua, M. D'Orazio, M. Cerasi, F. Pacello, A. Gismondi, A. Canini, L. Canuti, A. Consalvo, D. Ciavardelli, B. Chirullo, P. Pasquali and A. Battistoni, *Mol. Microbiol.*, 2017, **106**, 543–561.
- 55 K. Severinov, E. Semenova, A. Kazakov, T. Kazakov and M. S. Gelfand, *Mol. Microbiol.*, 2007, **65**, 1380–1394.
- 56 A. Metlitskaya, T. Kazakov, G. H. Vondenhoff, M. Novikova, A. Shashkov, T. Zatsepin, E. Semenova, N. Zaitseva, V. Ramensky, A. Van Aerschot and K. Severinov, *J. Bacteriol.*, 2009, **191**, 2380–2387.
- 57 A. Kulikovskiy, M. Serebryakova, O. Bantysh, A. Metlitskaya, S. Borukhov, K. Severinov and S. Dubiley, *J. Am. Chem. Soc.*, 2014, **136**, 11168–11175.
- 58 B. A. Wilson, S. Bantia, G. M. Salituro, A. M. Reeve and C. A. Townsend, *J. Am. Chem. Soc.*, 1988, **110**, 8238–8239.
- 59 A. M. Reeve, S. D. Breazeale and C. A. Townsend, *J. Biol. Chem.*, 1998, **273**, 30695–30703.
- 60 R. M. Klug and C. Benning, *Proc. Natl. Acad. Sci. U. S. A.*, 2001, **98**, 5910–5915.
- 61 M. Ihara, Y. Tanaka, K. Yanagisawa, Y. Taya and S. Nishimura, *Biochim. Biophys. Acta*, 1986, **881**, 135–140.



- 62 Y. Taya, Y. Tanaka and S. Nishimura, *FEBS Lett.*, 1978, **89**, 326–328.
- 63 F. Ikegami, R. Sakai, T. Ishikawa, Y.-H. Kuo, F. Lambein and I. Murakoshi, *Biol. Pharm. Bull.*, 1993, **16**, 732–734.
- 64 A. Callebaut, F. Lambein and R. Van Parijs, *Arch. Int. Physiol. Biochim.*, 1977, **85**, 157–158.
- 65 A. G. Saponara and M. D. Enger, *Biochim. Biophys. Acta*, 1974, **349**, 61–77.
- 66 B. Meyer, J. P. Wurm, S. Sharma, C. Immer, D. Pogoryelov, P. Kötter, D. L. J. Lafontaine, J. Wöhnert and K. D. Entian, *Nucleic Acids Res.*, 2016, **44**, 4304–4316.
- 67 K. L. Tkaczuk, S. Dunin-Horkawicz, E. Purta and J. M. Bujnicki, *BMC Bioinf.*, 2007, **8**, 73.
- 68 S. E. Strassler, I. E. Bowles, D. Dey, J. E. Jackman and G. L. Conn, *J. Biol. Chem.*, 2022, **298**, 102393.
- 69 M. Umitsu, H. Nishimasu, A. Noma, T. Suzuki, R. Ishitani and O. Nureki, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**, 15616–15621.
- 70 Z. Shao, W. Yan, J. Peng, X. Zuo, Y. Zou, F. Li, D. Gong, R. Ma, J. Wu, Y. Shi, Z. Zhang, M. Teng, X. Li and Q. Gong, *Nucleic Acids Res.*, 2014, **42**, 509–525.
- 71 B. Van Laer, M. Roovers, L. Wauters, J. M. Kasprzak, M. Dyzma, E. Deyaert, R. Kumar Singh, A. Feller, J. M. Bujnicki, L. Droogmans and W. Versées, *Nucleic Acids Res.*, 2016, **44**, 940–953.
- 72 J. E. Jackman, R. K. Montange, H. S. Malik and E. M. Phizicky, *RNA*, 2003, **9**, 574–585.
- 73 K. L. Tkaczuk, S. Dunin-Horkawicz, E. Purta and J. M. Bujnicki, *BMC Bioinf.*, 2007, **8**, 73.
- 74 A. Krishnamohan and J. E. Jackman, *Nucleic Acids Res.*, 2017, **45**, 9019–9029.
- 75 A. M. Burroughs and L. Aravind, *Front. Genet.*, 2014, **5**, 424.
- 76 B. Meyer, C. Immer, S. Kaiser, S. Sharma, J. Yang, P. Watzinger, L. Weiß, A. Kotter, M. Helm, H.-M. Seitz, P. Kötter, S. Kellner, K.-D. Entian and J. Wöhnert, *Nucleic Acids Res.*, 2020, **48**, 1435–1450.
- 77 X.-H. Liang, Q. Liu and M. J. Fournier, *RNA*, 2009, **15**, 1716–1728.
- 78 B. Meyer, J. P. Wurm, P. Kötter, M. S. Leisegang, V. Schilling, M. Buchhaupt, M. Held, U. Bahr, M. Karas, A. Heckel, M. T. Bohnsack, J. Wöhnert and K.-D. Entian, *Nucleic Acids Res.*, 2011, **39**, 1526–1537.
- 79 M. Takakura, K. Ishiguro, S. Akichika, K. Miyauchi and T. Suzuki, *Nat. Commun.*, 2019, **10**, 5542.
- 80 A. Babaian, K. Rothe, D. Girodat, I. Minia, S. Djondovic, M. Milek, S. E. Spencer Miko, H.-J. Wieden, M. Landthaler, G. B. Morin and D. L. Mager, *Cell Rep.*, 2020, **31**, 107611.
- 81 P. F. Agris, F. A. P. Vendeix and W. D. Graham, *J. Mol. Biol.*, 2007, **366**, 1–13.
- 82 A. Weixlbaumer, F. V. Murphy IV, A. Dziergowska, A. Malkiewicz, F. A. Vendeix, P. F. Agris and V. Ramakrishnan, *Nat. Struct. Mol. Biol.*, 2007, **14**, 498–502.
- 83 A. M. Edwards, M. A. Addo and P. C. Dos Santos, *Genes*, 2020, **11**, 907.
- 84 P. Boccaletto, F. Stefaniak, A. Ray, A. Cappannini, S. Mukherjee, E. Purta, M. Kurkowska, N. Shirvanizadeh, E. Destefanis, P. Groza, G. Avşar, A. Romitelli, P. Pir, E. Dassi, S. G. Conticello, F. Aguilo and J. M. Bujnicki, *Nucleic Acids Res.*, 2022, **50**, D231–D235.
- 85 S. J. Nasvall, P. Chen and G. R. Bjork, *RNA*, 2004, **10**, 1662–1673.
- 86 J. Kim, H. Xiao, J. B. Bonanno, C. Kalyanaraman, S. Brown, X. Tang, N. F. Al-Obaidi, Y. Patskovsky, P. C. Babbitt, M. P. Jacobson, Y.-S. Lee and S. C. Almo, *Nature*, 2013, **498**, 123–126.
- 87 R. T. Byrne, F. Whelan, P. Aller, L. E. Bird, A. Dowle, C. M. Lobley, Y. Reddivari, J. E. Nettleship, R. J. Owens, A. A. Antson and D. G. Waterman, *Acta Crystallogr., Sect. D: Biol. Crystallogr.*, 2013, **69**, 1090–1098.
- 88 J. Kim, H. Xiao, J. Koh, Y. Wang, J. B. Bonanno, K. Thomas, P. C. Babbitt, S. Brown, Y. S. Lee and S. C. Almo, *Nucleic Acids Res.*, 2015, **43**, 4602–4613.
- 89 S. Jeong and J. Kim, *Biochem. Biophys. Res. Commun.*, 2021, **534**, 604–609.
- 90 C. J. La Francois, Y. H. Jang, T. Cagin, W. A. Goddard III and L. C. Sowers, *Chem. Res. Toxicol.*, 2000, **13**, 462–470.
- 91 C. P. Ting, M. A. Funk, S. L. Halaby, Z. Zhang, T. Gonen and W. A. van der Donk, *Science*, 2019, **365**, 280–284.
- 92 M. Serebryakova, D. Tsibulskaya, O. Mokina, A. Kulikovskiy, M. Nautiyal, A. Van Aerschot, K. Severinov and S. Dubiley, *J. Am. Chem. Soc.*, 2016, **138**, 15690–15698.
- 93 B. D. Bennett, E. H. Kimball, M. Gao, R. Osterhout, S. J. Van Dien and J. D. Rabinowitz, *Chem. Biol.*, 2009, **5**, 593–599.
- 94 R. Wang and M. Luo, *Curr. Opin. Chem. Biol.*, 2013, **17**, 729–737.
- 95 A. Y. Rudenko, S. S. Mariasina, P. V. Sergiev and V. I. Polshakov, *Mol. Biol.*, 2022, **56**, 229–250.
- 96 H. Deng and D. O'Hagan, *Curr. Opin. Chem. Biol.*, 2008, **12**, 582–592.
- 97 D. O'Hagan and H. Deng, *Chem. Rev.*, 2015, **115**, 634–649.
- 98 D. O'Hagan, C. Schaffrath, S. L. Cobb, J. T. G. Hamilton and C. D. Murphy, *Nature*, 2002, **416**, 279.
- 99 C. Schaffrath, H. Deng and D. O'Hagan, *FEBS Lett.*, 2003, **547**, 111–114.
- 100 C. Dong, F. Huang, H. Deng, C. Schaffrath, J. B. Spencer, D. O'Hagan and J. H. Naismith, *Nature*, 2004, **427**, 561–565.
- 101 A. S. Eustáquio, F. Pojer, J. P. Noel and B. S. Moore, *Nat. Chem. Biol.*, 2008, **4**, 69–74.
- 102 D. B. Harper and D. O'Hagan, *Nat. Prod. Rep.*, 1994, **11**, 123–133.
- 103 H. Deng, C. H. Botting, J. T. Hamilton, R. J. Russell and D. O'Hagan, *Angew. Chem., Int. Ed.*, 2008, **47**, 5357–5361.
- 104 A. S. Eustáquio, J. Härle, J. P. Noel and B. S. Moore, *ChemBioChem*, 2008, **9**, 2215–2219.
- 105 H. Deng, S. A. McMahon, A. S. Eustáquio, B. S. Moore, J. H. Naismith and D. O'Hagan, *ChemBioChem*, 2009, **10**, 2455–2459.
- 106 T. Kornfuehrer, S. Romanowski, V. de Crécy-Lagard, A. D. Hanson and A. S. Eustáquio, *ChemBioChem*, 2020, **21**, 3495–3499.
- 107 D. Kaiser, I. Klose, R. Oost, J. Neuhaus and N. Maulide, *Chem. Rev.*, 2019, **119**, 8701–8780.
- 108 J. L. Hoffman, *Biochemistry*, 1986, **25**, 4444–4449.



- 109 D. F. Iwig and S. J. Booker, *Biochemistry*, 2004, **43**, 13496–13509.
- 110 J. Baldwin, *J. Chem. Soc., Chem. Commun.*, 1976, 734–736.
- 111 S. Ma, D. Mandalapu, S. Wang and Q. Zhang, *Nat. Prod. Rep.*, 2022, **39**, 926–945.
- 112 P. Diana and G. Cirrincione, *Biosynthesis of Heterocycles: From Isolation to Gene Cluster*, Wiley, Hoboken, 2015.
- 113 M. E. Churchill and L. Chen, *Chem. Rev.*, 2011, **111**, 68–85.
- 114 L. Fowden, *Nature*, 1955, **176**, 347–348.
- 115 E. Leete, *J. Am. Chem. Soc.*, 1964, **86**, 3162.
- 116 E. Leete, G. E. Davis, C. R. Hutchinson, K. W. Woo and M. R. Chedekel, *Phytochemistry*, 1974, **13**, 427–433.
- 117 E. Leete, L. L. Louters and H. S. Prakash Rao, *Phytochemistry*, 1986, **25**, 2753–2758.
- 118 T. Takemoto, K. Nomoto, S. Fushiya, R. Ouchi, G. Kusano, H. Hikino, S. Takagi, Y. Matsuura and M. Kakudo, *Proc. Jpn. Acad., Ser. B*, 1978, **54**, 469.
- 119 S. Mori and N. Nishizawa, *Plant Cell Physiol.*, 1987, **28**, 1081–1092.
- 120 S. Shojima, N. K. Nishizawa, S. Fushiya, S. Nozoe, T. Irifune and S. Mori, *Plant Physiol.*, 1990, **93**, 1497–1503.
- 121 F. Yan, D. Auerbach, Y. Chai, L. Keller, Q. Tu, S. Hüttel, A. Glemser, H. A. Grab, T. Bach, Y. Zhang and R. Müller, *Angew. Chem., Int. Ed.*, 2018, **57**, 8754–8759.
- 122 F. Yan and R. Müller, *ACS Chem. Biol.*, 2019, **14**, 99–105.
- 123 Z. Hong, A. Bolard, C. Giraud, S. Prévost, G. Genta-Jouve, C. Deregnacourt, S. Häussler, K. Jeannot and Y. Li, *Angew. Chem., Int. Ed.*, 2019, **58**, 3178–3182.
- 124 Y.-H. Shin, Y. H. Ban, J. Shin, I. W. Park, S. Yoon, K. Ko, J. Shin, S.-J. Nam, J. M. Winter, Y. Kim, Y. J. Yoon and D.-C. Oh, *J. Org. Chem.*, 2021, **86**, 11149–11159.
- 125 W. B. Jin, S. Wu, X.-H. Jian, H. Yuan and G.-L. Tang, *Nat. Commun.*, 2018, **9**, 2771.
- 126 X. Wang, S. Wu, W. Jin, B. Xu, G. Tang and H. Yuan, *Acta Biochim. Biophys. Sin.*, 2018, **50**, 516–518.
- 127 W. Huang, H. Xu, Y. Li, F. Zhang, X.-Y. Chen, Q.-L. He, Y. Igarashi and G.-L. Tang, *J. Am. Chem. Soc.*, 2012, **134**, 8831–8840.
- 128 I. Abe, *J. Antibiot.*, 2018, **71**, 763–768.
- 129 J. D. Rudolf and C. Y. Chang, *Nat. Prod. Rep.*, 2020, **37**, 425–463.
- 130 J. S. Dickschat, *Nat. Prod. Rep.*, 2016, **33**, 87–110.
- 131 C. Schmidt-Dannert, *Adv. Biochem. Eng./Biotechnol.*, 2015, **148**, 19–61.
- 132 D. J. Tantillo, *Chem. Soc. Rev.*, 2010, **39**, 2847–2854.
- 133 D. W. Christianson, *Chem. Rev.*, 2017, **117**, 11570–11648.
- 134 J. F. Quílez del Moral, Á. Pérez and A. F. Barrero, *Phytochem. Rev.*, 2020, **19**, 559–576.
- 135 D. F. Iwig, A. Uchida, J. A. Stromberg and S. J. Booker, *J. Am. Chem. Soc.*, 2005, **127**, 11612–11613.
- 136 J. E. Cronan and T. Luk, *Microbiol. Mol. Biol. Rev.*, 2022, **86**, e0001322.
- 137 M. Kai, E. Crespo, S. M. Cristescu, F. J. M. Harren, W. Francke and B. Piechulla, *Appl. Microbiol. Biotechnol.*, 2010, **88**, 965–976.
- 138 R. Schmidt, V. de Jager, D. Zühlke, C. Wolff, J. Bernhardt, K. Cankar, J. Beekwilder, W. van Ijcken, F. Sleutels, W. de Boer, K. Riedel and P. Garbeva, *Sci. Rep.*, 2017, **7**, 862.
- 139 S. von Reuss, D. Domik, M. C. Lemfack, N. Magnus, M. Kai, T. Weise and B. Piechulla, *J. Am. Chem. Soc.*, 2018, **140**, 11855–11862.
- 140 M. C. Lemfack, W. Brandt, K. Krüger, A. Gurowietz, J. Djifack, J.-P. Jung, M. Hopf, H. Noack, B. Junker, S. von Reuß and B. Piechulla, *Sci. Rep.*, 2021, **11**, 3182.
- 141 E. R. Duell, P. M. D'Agostino, N. Shapiro, T. Woyke, T. M. Fuchs and T. A. M. Gulder, *Microb. Cell Fact.*, 2019, **18**, 32.
- 142 T. Awakawa and I. Abe, *Org. Biomol. Chem.*, 2018, **16**, 4746–4752.
- 143 T. Awakawa, *J. Nat. Med.*, 2021, **75**, 467–474.
- 144 T. Awakawa, L. Zhang, T. Wakimoto, S. Hoshino, T. Mori, T. Ito, J. Ishikawa, M. E. Tanner and I. Abe, *J. Am. Chem. Soc.*, 2014, **136**, 9910–9913.
- 145 J. Stöckigt, A. P. Antonchick, F. Wu and H. Waldmann, *Angew. Chem., Int. Ed.*, 2011, **50**, 8538–8564.
- 146 C.-I. Lin, R. M. McCarty and H.-w. Liu, *Angew. Chem., Int. Ed.*, 2017, **56**, 3446–3489.
- 147 F. Yu, M. Li, C. Xu, B. Sun, H. Zhou, Z. Wang, Q. Xu, M. Xie, G. Zuo, P. Huang, H. Guo, Q. Wang and J. He, *Biochem. J.*, 2016, **473**, 4385–4397.
- 148 K. Katagiri, J. Shoji and T. Yoshida, *J. Antibiot.*, 1962, **15**, 273.
- 149 M. J. Waring and L. P. Wakelin, *Nature*, 1974, **252**, 653–657.
- 150 D. Kong, E. J. Park, A. G. Stephen, M. Calvani, J. H. Cardellina, A. Monks, R. J. Fisher, R. H. Shoemaker and G. Melillo, *Cancer Res.*, 2005, **65**, 9047–9055.
- 151 A. Cornish, M. J. Waring and R. D. Nolan, *J. Antibiot.*, 1983, **36**, 1664–1670.
- 152 J. S. Lee and M. J. Waring, *Biochem. J.*, 1978, **173**, 115–128.
- 153 K. R. Fox, N. L. Harrison and M. J. Waring, *FEBS Lett.*, 1981, **133**, 305–310.
- 154 K. Watanabe, K. Hotta, A. P. Praseuth, K. Koketsu, A. Migita, C. N. Boddy, C. C. C. Wang, H. Oguri and H. Oikawa, *Nat. Chem. Biol.*, 2006, **2**, 423–428.
- 155 C. Zhang, L. Kong, Q. Liu, X. Lei, T. Zhu, J. Yin, B. Lin, Z. Deng and D. You, *PLoS One*, 2013, **8**, e56772.
- 156 M. Sato, T. Nakazawa, Y. Tsunematsu, K. Hotta and K. Watanabe, *Curr. Opin. Chem. Biol.*, 2013, **17**, 537–545.
- 157 K. Watanabe, K. Hotta, M. Nakaya, A. P. Praseuth, C. C. C. Wang, D. Inada, K. Takahashi, E. Fukushima, H. Oguri and H. Oikawa, *J. Am. Chem. Soc.*, 2009, **131**, 9347–9353.
- 158 C. L. Lim, T. Nogawa, M. Uramoto, A. Okano, Y. Hongo, T. Nakamura, H. Koshino, S. Takahashi, D. Ibrahim and H. Osada, *J. Antibiot.*, 2014, **67**, 323–329.
- 159 K. Hotta, R. M. Keegan, S. Ranganathan, M. Fang, J. Bibby, M. D. Winn, M. Sato, M. Lian, K. Watanabe, D. J. Rigden and C. Y. Kim, *Angew. Chem., Int. Ed.*, 2014, **53**, 824–828.
- 160 S. H. Pine, B. A. Catto and F. G. Yamagishi, *J. Org. Chem.*, 1970, **35**, 3663–3666.
- 161 J. B. Sweeney, *Chem. Soc. Rev.*, 2009, **38**, 1027–1038.



- 162 C. Sun, W. Tian, Z. Lin and X. Qu, *Nat. Prod. Rep.*, 2022, **39**, 1721–1765.
- 163 S. Funayama and G. A. Cordell, in *Alkaloids*, ed. S. Funayama and G. A. Cordell, Academic Press, Massachusetts, 1st edn, 2015, ch. 2, pp. 63–102.
- 164 J. Liu, T. Ng, Z. Rui, O. Ad and W. Zhang, *Angew. Chem., Int. Ed.*, 2014, **53**, 136–139.
- 165 H. Li, Y. Qiu, C. Guo, M. Han, Y. Zhou, Y. Feng, S. Luo, Y. Tong, G. Zheng and S. Zhu, *Chem. Commun.*, 2019, **55**, 8390–8393.
- 166 L. Ma, W. Zhang, Y. Zhu, G. Zhang, H. Zhang, Q. Zhang, L. Zhang, C. Yuan and C. Zhang, *Appl. Microbiol. Biotechnol.*, 2017, **101**, 6123–6136.
- 167 N. Alqahtani, S. K. Porwal, E. D. James, D. M. Bis, J. A. Karty, A. L. Lane and R. Viswanathan, *Org. Biomol. Chem.*, 2015, **13**, 7177–7192.
- 168 T. M. Khopade, K. Ajayan, S. S. Joshi, A. L. Lane and R. Viswanathan, *ACS Omega*, 2021, **6**, 10840–10858.
- 169 M. R. TePaske, J. B. Gloer, D. T. Wicklow and P. F. Dowd, *Tetrahedron Lett.*, 1991, **32**, 5687–5690.
- 170 M. Ohashi, F. Liu, Y. Hai, M. Chen, M. C. Tang, Z. Yang, M. Sato, K. Watanabe, K. N. Houk and Y. Tang, *Nature*, 2017, **549**, 502–506.
- 171 Y. Cai, Y. Hai, M. Ohashi, C. S. Jamieson, M. Garcia-Borras, K. N. Houk, J. Zhou and Y. Tang, *Nat. Chem.*, 2019, **11**, 812–820.
- 172 Z. Chang, T. Ansbacher, L. Zhang, Y. Yang, T.-P. Ko, G. Zhang, W. Liu, J.-W. Huang, L. Dai, R.-T. Guo, D. T. Major and C.-C. Chen, *Org. Biomol. Chem.*, 2019, **17**, 2070–2076.
- 173 M. Chang, Y. Zhou, H. Wang, Z. Liu, Y. Zhang and Y. Feng, *Biochem. Biophys. Res. Commun.*, 2019, **515**, 255–260.
- 174 Q. Sun, Y. Hu, Y. Gu, J. Huang, J. He, L. Luo, Y. Yang, S. Yin, C. Dou, T. Wang, X. Fu, L. He, S. Qi, X. Zhu, S. Yang, X. Wei and W. Cheng, *Signal Transduction Targeted Ther.*, 2019, **4**, 17.
- 175 S. A. Newmister, S. Romminger, J. J. Schmidt, R. M. Williams, J. L. Smith, R. G. S. Berlinck and D. H. Sherman, *Org. Biomol. Chem.*, 2018, **16**, 6450–6459.
- 176 M. Ohashi, D. Tan, J. Lu, C. S. Jamieson, D. Kanayama, J. Zhou, K. N. Houk and Y. Tang, *J. Am. Chem. Soc.*, 2023, **145**, 3301–3305.
- 177 H. J. Kim, M. W. Ruszczycky, S.-h. Choi, Y.-n. Liu and H.-w. Liu, *Nature*, 2011, **473**, 109–112.
- 178 B.-S. Jeon, M. W. Ruszczycky, W. K. Russell, G.-M. Lin, N. Kim, S.-h. Choi, S.-A. Wang, Y.-n. Liu, J. W. Patrick, D. H. Russell and H.-w. Liu, *Proc. Natl. Acad. Sci. U. S. A.*, 2017, **114**, 10408–10413.
- 179 A. Patel, Z. Chen, Z. Yang, O. Gutiérrez, H.-w. Liu, K. N. Houk and D. A. Singleton, *J. Am. Chem. Soc.*, 2016, **138**, 3631–3634.
- 180 Z. Yang, S. Yang, P. Yu, Y. Li, C. Doubleday, J. Park, A. Patel, B.-s. Jeon, W. K. Russell, H.-w. Liu, D. H. Russell and K. N. Houk, *Proc. Natl. Acad. Sci. U. S. A.*, 2017, **115**, E848–E855.
- 181 C. D. Fage, E. A. Isiorho, Y.-n. Liu, D. T. Wagner, H.-w. Liu and A. T. Keatinge-Clay, *Nat. Chem. Biol.*, 2015, **11**, 256–258.
- 182 E. A. Isiorho, H.-w. Liu and A. T. Keatinge-Clay, *Biochemistry*, 2012, **51**, 1213–1222.
- 183 C. E. Aroyan, A. Dermenci and S. J. Miller, *Tetrahedron*, 2009, **65**, 4069–4084.
- 184 C. W. Carreras and D. V. Santi, *Annu. Rev. Biochem.*, 1995, **64**, 721–762.
- 185 S.-h. Choi, B. Jeon, N. Kim, H.-H. Wu, T.-P. Ko, M. W. Ruszczycky, E. A. Isiorho, Y.-n. Liu, A. T. Keatinge-Clay, M.-D. Tsai and H.-w. Liu, *J. Am. Chem. Soc.*, 2021, **143**, 20291–20295.
- 186 Z. Zhang, C. S. Jamieson, Y.-L. Zhao, D. Li, M. Ohashi, K. N. Houk and Y. Tang, *J. Am. Chem. Soc.*, 2019, **141**, 5659–5663.
- 187 M. Ohashi, C. S. Jamieson, Y. Cai, D. Tan, D. Kanayama, M.-C. Tang, S. M. Anthony, J. V. Chari, J. S. Barber, E. Picazo, T. B. Kakule, S. Cao, N. K. Garg, J. Zhou, K. N. Houk and Y. Tang, *Nature*, 2020, **586**, 64–69.
- 188 B.-S. Jeon, S.-A. Wang, M. W. Ruszczycky and H.-w. Liu, *Chem. Rev.*, 2017, **117**, 5367–5388.
- 189 C. S. Jamieson, M. Ohashi, F. Liu, Y. Tang and K. N. Houk, *Nat. Prod. Rep.*, 2019, **36**, 698–713.
- 190 K. Watanabe, *J. Nat. Med.*, 2021, **75**, 434–447.
- 191 Y. Miyazaki, M. Shibuya, H. Sugawara, O. Kawaguchi and C. Hirsoe, *J. Antibiot.*, 1974, **27**, 814–821.
- 192 S. Zhou, F. Wang, E. T. Wong, E. Fonkem, T.-C. Hsieh, J. M. Wu and E. Wu, *Curr. Med. Chem.*, 2013, **20**, 4095–4101.
- 193 J. Dewangan, S. Srivastava and S. K. Rath, *Tumor Biol.*, 2017, **39**, DOI: [10.1177/1010428317695035](https://doi.org/10.1177/1010428317695035).
- 194 C. Jiang, H. Wang, Q. Kang, J. Liu and L. Bai, *Appl. Environ. Microbiol.*, 2012, **78**, 994–1003.
- 195 M. E. Yurkovich, P. A. Tyrakis, H. Hong, Y. Sun, M. Samborsky, K. Kamiya and P. F. Leadlay, *ChemBioChem*, 2012, **13**, 66–71.
- 196 C. Jiang, Z. Qi, Q. Kang, J. Liu, M. Jiang and L. Bai, *Angew. Chem., Int. Ed.*, 2015, **54**, 9097–9100.
- 197 A. Jansson, J. Niemi, Y. Lindqvist, P. Mäntsälä and G. Schneider, *J. Mol. Biol.*, 2003, **334**, 269–280.
- 198 Y. Wang, J. Niemi, K. Airas, K. Ylihönko, J. Hakala and P. Mäntsälä, *Biochim. Biophys. Acta*, 2000, **1480**, 191–200.
- 199 Y. Wang, J. Niemi and P. Mäntsälä, *FEMS Microbiol. Lett.*, 2002, **208**, 117–122.
- 200 A. Jansson, H. Koskiniemi, A. Erola, J. Wang, P. Mäntsälä, G. Schneider and J. Niemi, *J. Biol. Chem.*, 2005, **280**, 3636–3644.
- 201 C. Gui, X. Mo, Y.-C. Gu and J. Ju, *Org. Lett.*, 2017, **19**, 5617–5620.
- 202 K. Madduri, F. Torti, A. L. Colombo and C. R. Hutchinson, *J. Bacteriol.*, 1993, **175**, 3900–3904.
- 203 H. A. Blair, *Drugs*, 2018, **78**, 1903–1910.
- 204 A. Jansson, H. Koskiniemi, P. Mäntsälä, J. Niemi and G. Schneider, *J. Biol. Chem.*, 2004, **279**, 41149–41156.
- 205 T. Grocholski, P. Dinis, L. Niiranen, J. Niemi and M. Metsä-Ketelä, *Proc. Natl. Acad. Sci. U. S. A.*, 2015, **112**, 9866–9871.
- 206 J. Hirst, *Annu. Rev. Biochem.*, 2013, **82**, 551–575.
- 207 L. A. Sazanov, *Nat. Rev. Mol. Cell Biol.*, 2015, **16**, 375–388.
- 208 V. F. Rhein, J. Carroll, S. Ding, I. M. Fearnley and J. E. Walker, *J. Biol. Chem.*, 2016, **291**, 14851–14860.



- 209 F. A. Macías, R. M. Varela, A. M. Simonet, H. G. Cutler, S. J. Cutler, F. M. Dugan and R. A. Hill, *J. Org. Chem.*, 2000, **65**, 9039–9046.
- 210 D. Yan and Y. Matsuda, *Angew. Chem., Int. Ed.*, 2022, **61**, e202210938.
- 211 J. Van Vleet, A. Kleeb, P. Kast, D. Hilvert and W. W. Cleland, *Biochim. Biophys. Acta*, 2010, **1804**, 752–754.
- 212 J. Crosby and C. J. M. Stirling, *J. Chem. Soc. B*, 1970, 671–679.
- 213 W. E. Stites, A. G. Gittis, E. E. Lattman and D. Shortle, *J. Mol. Biol.*, 1991, **221**, 7–14.
- 214 F. Jordan, H. Li and A. Brown, *Biochemistry*, 1999, **38**, 6369–6373.
- 215 J. J. Dwyer, A. G. Gittis, D. A. Karp, E. E. Lattman, D. S. Spencer, W. E. Stites and B. García-Moreno, *Biophys. J.*, 2000, **79**, 1610–1620.
- 216 D. G. Isom, C. A. Castañeda, B. R. Cannon and B. García-Moreno, *Proc. Natl. Acad. Sci. U. S. A.*, 2011, **108**, 5260–5265.
- 217 F. G. Bordwell, *Acc. Chem. Res.*, 1988, **21**, 456–463.
- 218 S. D. Kinzie, B. Thern and D. Iwata-Reuyl, *Org. Lett.*, 2000, **2**, 1307–1310.
- 219 R. M. McCarty and V. Bandarian, *Bioorg. Chem.*, 2012, **43**, 15–25.
- 220 S. G. Van Lanen and D. Iwata-Reuyl, *Biochemistry*, 2003, **42**, 5312–5320.
- 221 I. Mathews, R. Schwarzenbacher, D. McMullan, P. Abdubek, E. Ambing, H. Axelrod, T. Biorac, J. M. Canaves, H.-J. Chiu, A. M. Deacon, M. DiDonato, M.-A. Elsliger, A. Godzik, C. Grittini, S. K. Grzechnik, J. Hale, E. Hampton, G. W. Han, J. Haugen, M. Hornsby, L. Jaroszewski, H. E. Klock, E. Koesema, A. Kreusch, P. Kuhn, S. A. Lesley, I. Levin, M. D. Miller, K. Moy, E. Nigoghossian, J. Ouyang, J. Paulsen, K. Quijano, R. Reyes, G. Spraggon, R. C. Stevens, H. van den Bedem, J. Velasquez, J. Vincent, A. White, G. Wolf, Q. Xu, K. O. Hodgson, J. Wooley and I. A. Wilson, *Proteins*, 2005, **59**, 869–874.
- 222 C. Grimm, R. Ficner, T. Sgraja, P. Haebel, G. Klebe and K. Reuter, *Biochem. Biophys. Res. Commun.*, 2006, **351**, 695–701.
- 223 G. L. Stoner and M. A. Eisenberg, *J. Biol. Chem.*, 1975, **250**, 4029–4036.
- 224 S. Mann and O. Ploux, *Biochim. Biophys. Acta*, 2011, **1814**, 1459–1466.
- 225 R. S. Breen, D. J. Campopiano, S. Webster, M. Brunton, R. Watt and R. L. Baxter, *Org. Biomol. Chem.*, 2003, **1**, 3498–3499.
- 226 B. C. Brajcich, A. L. Iarocci, L. A. G. Johnstone, R. K. Morgan, Z. T. Lonjers, M. J. Hotchko, J. D. Muhs, A. Kieffer, B. J. Reynolds, S. M. Mandel, B. N. Marbois, C. F. Clarke and J. N. Shepherd, *J. Bacteriol.*, 2010, **192**, 436–445.
- 227 Z. T. Lonjers, E. L. Dickson, T.-P. Chu, J. E. Kreutz, F. A. Neacsu, K. R. Anders and J. N. Shepherd, *J. Bacteriol.*, 2012, **194**, 965–971.
- 228 A. C. Bernert, E. J. Jacobs, S. R. Reinl, C. C. Y. Choi, P. M. Roberts Buceta, J. C. Culver, C. R. Goodspeed, M. C. Bradley, C. F. Clarke, G. J. Basset and J. N. Shepherd, *Biochim. Biophys. Acta, Mol. Cell Biol. Lipids*, 2019, **1864**, 1226–1234.
- 229 T. Neupane, L. R. Chambers, A. J. Godfrey, M. M. Monlux, E. J. Jacobs, S. Whitworth, J. E. Spawn, S. H. K. Clingman, K. L. Vergunst, F. M. Niven, J. J. Townley, I. W. Orion, C. R. Goodspeed, K. A. Cooper, J. D. Cronk, J. N. Shepherd and D. N. Langelaan, *Commun. Chem.*, 2022, **5**, 89.
- 230 T. L. Amyes and J. P. Richard, *Synlett*, 2017, **28**, 2407–2421.
- 231 G. D. Peiser, T.-T. Wang, N. E. Hoffman, S. F. Yang, H.-w. Liu and C. T. Walsh, *Proc. Natl. Acad. Sci. U. S. A.*, 1984, **81**, 3059–3063.
- 232 S. F. Yang and N. E. Hoffmann, *Annu. Rev. Plant Physiol.*, 1984, **35**, 155–189.
- 233 F. X. Nascimento, M. J. Rossi and B. R. Glick, *Front. Plant Sci.*, 2018, **9**, 114.
- 234 Y. Li, L. Feng and J. F. Kirsch, *Biochemistry*, 1997, **36**, 15477–15488.
- 235 M. Jakubowicz, *Acta Biochim. Pol.*, 2002, **49**, 757–774.
- 236 H.-P. Zhang, H. Kakeya and H. Osada, *Tetrahedron Lett.*, 1998, **39**, 6947–6948.
- 237 C. J. Thibodeaux, W.-c. Chang and H.-w. Liu, *Chem. Rev.*, 2012, **112**, 1681–1709.
- 238 Y. Y. Fan, X. H. Gao and J. M. Yue, *Sci. China: Chem.*, 2016, **59**, 1126–1141.
- 239 G. Pan, Z. Xu, Z. Guo, Hindra, M. Ma, D. Yang, H. Zhou, Y. Gansemans, X. Zhu, Y. Huang, L.-X. Zhao, Y. Jiang, J. Cheng, F. Van Nieuwerburgh, J.-W. Suh, Y. Duan and B. Shen, *Proc. Natl. Acad. Sci. U. S. A.*, 2017, **114**, E11131–E11140.
- 240 Z. Xu, G. Pan, H. Zhou and B. Shen, *J. Am. Chem. Soc.*, 2018, **140**, 16957–16961.
- 241 C. Maruyama, Y. Chinone, S. Sato, F. Kudo, K. Ohsawa, J. Kubota, J. Hashimoto, I. Kozono, T. Doi, K. Shin-Ya, T. Eguchi and Y. Hamano, *Biomolecules*, 2020, **10**, 775.
- 242 M. Hiramoto, A. Niwa, A. Miyake, H. Yamamoto, Y. Takebayashi, T. Nishikawa, M. Shibazaki and K. Nagai, *Tennen Yuki Kagobutsu Toronkai Koen Yoshishu*, 1993, **35**, 678–684.
- 243 K. Kurosawa, K. Takahashi and E. Tsuda, *J. Antibiot.*, 2001, **54**, 615–621.
- 244 F. Kudo, A. Minato, S. Sato, N. Nagano, C. Maruyama, Y. Hamano, J. Hashimoto, I. Kozono, K. Shin-Ya and T. Eguchi, *Org. Lett.*, 2022, **24**, 8975–8979.
- 245 C. T. Walsh and W. Zhang, *ACS Chem. Biol.*, 2011, **6**, 1000–1007.
- 246 M. McErlean, X. Liu, Z. Cui, B. Gust and S. G. Van Lanen, *Nat. Prod. Rep.*, 2021, **38**, 1362–1407.
- 247 X. Chi, P. Pahari, K. Nonaka and S. G. Van Lanen, *J. Am. Chem. Soc.*, 2011, **133**, 14452–14459.
- 248 Z. Cui, J. Overbay, X. Wang, X. Liu, Y. Zhang, M. Bhardwaj, A. Lemke, D. Wiegmann, G. Niro, J. S. Thorson, C. Ducho and S. G. Van Lanen, *Nat. Chem. Biol.*, 2020, **16**, 904–911.
- 249 L. Kaysser, L. Lutsch, S. Siebenberg, E. Wemakor, B. Kammerer and B. Gust, *J. Biol. Chem.*, 2009, **284**, 14987–14996.



- 250 M. Funabashi, S. Baba, K. Nonaka, M. Hosobuchi, Y. Fujita, T. Shibata and S. G. Van Lanen, *Chembiochem*, 2010, **11**, 184–190.
- 251 L. Cheng, W. Chen, L. Zhai, D. Xu, T. Huang, S. Lin, X. Zhou and Z. Deng, *Mol. BioSyst.*, 2011, **7**, 920–927.
- 252 M. Funabashi, S. Baba, T. Takatsu, M. Kizuka, Y. Ohata, M. Tanaka, K. Nonaka, A. P. Spork, C. Ducho, W.-C. L. Chen and S. G. Van Lanen, *Angew. Chem., Int. Ed.*, 2013, **52**, 11607–11611.
- 253 A. Takahashi, S. Kurasawa, D. Ikeda, Y. Okami and T. Takeuchi, *J. Antibiot.*, 1989, **42**, 1556–1561.
- 254 A. L. Stefanska, R. Cassels, S. J. Ready and S. R. Warr, *J. Antibiot.*, 2000, **53**, 357–363.
- 255 C. S. Houge-Frydrych, M. L. Gilpin, P. W. Skett and J. W. Tyler, *J. Antibiot.*, 2000, **53**, 364–372.
- 256 L. Barra, T. Awakawa, K. Shirai, Z. Hu, G. Bashiri and I. Abe, *Nature*, 2021, **600**, 754–758.
- 257 R. C. Kelly, M. E. Bolitho, D. A. Higgins, W. Lu, W.-L. Ng, P. D. Jeffrey, J. D. Rabinowitz, M. F. Semmelhack, F. M. Hughson and B. L. Bassler, *Nat. Chem. Biol.*, 2009, **5**, 891–895.
- 258 N. Jahan, J. A. Potter, M. A. Sheikh, C. H. Botting, S. L. Shirran, N. J. Westwood and G. L. Taylor, *J. Mol. Biol.*, 2009, **392**, 763–773.
- 259 D. A. Higgins, M. E. Pomianek, C. M. Kraml, R. K. Taylor, M. F. Semmelhack and B. L. Bassler, *Nature*, 2007, **450**, 883–886.
- 260 Y. Wei, L. J. Perez, W.-L. Ng, M. F. Semmelhack and B. L. Bassler, *ACS Chem. Biol.*, 2011, **6**, 356–365.
- 261 Y. L. Du and K. S. Ryan, *Nat. Prod. Rep.*, 2019, **36**, 430–457.
- 262 A. R. Healy, H. Nikolayevskiy, J. R. Patel, J. M. Crawford and S. B. Herzon, *J. Am. Chem. Soc.*, 2016, **138**, 15563–15570.
- 263 M. Xue, C. S. Kim, A. R. Healy, K. M. Wernke, Z. Wang, M. C. Frischling, E. E. Shine, W. Wang, S. B. Herzon and J. M. Crawford, *Science*, 2019, **365**, eaax2685.
- 264 M. R. Wilson, Y. Jiang, P. W. Villalta, A. Stornetta, P. D. Boudreau, A. Carrá, C. A. Brennan, E. Chun, L. Ngo, L. D. Samson, B. P. Engelward, W. S. Garrett, S. Balbo and E. P. Balskus, *Science*, 2019, **363**, eaar7785.
- 265 L. Zha, Y. Jiang, M. T. Henke, M. R. Wilson, J. X. Wang, N. L. Kelleher and E. P. Balskus, *Nat. Chem. Biol.*, 2017, **13**, 1063–1065.
- 266 J. S. Dickschat, P. Rabe and C. A. Citron, *Org. Biomol. Chem.*, 2015, **13**, 1954–1968.
- 267 F. Trottman, K. Ishida, J. Franke, A. Stanišić, M. Ishida-Ito, H. Kries, G. Pohnert and C. Hertweck, *Angew. Chem., Int. Ed.*, 2020, **59**, 13511–13515.
- 268 F. Trottman, K. Ishida, M. Ishida-Ito, H. Kries, M. Groll and C. Hertweck, *Nat. Chem.*, 2022, **14**, 884–890.
- 269 P. Wittung-Stafshede, *Acc. Chem. Res.*, 2002, **35**, 201–208.
- 270 A. Warshel, P. K. Sharma, M. Kato, Y. Xiang, H. Liu and M. H. M. Olsson, *Chem. Rev.*, 2006, **106**, 3210–3235.
- 271 S. D. Fried and S. G. Boxer, *Annu. Rev. Biochem.*, 2017, **86**, 387–415.

