

COMMUNICATION

[View Article Online](#)
[View Journal](#) | [View Issue](#)

Cite this: *Dalton Trans.*, 2025, **54**, 8035

Received 26th February 2025,
Accepted 27th February 2025

DOI: 10.1039/d5dt00480b

rsc.li/dalton

Isolation of the parent triplet titanocene *via* NHC stabilisation†

Felix Meyer,^a Serhiy Demeshko,^a Christopher Golz,^b Oliver P. E. Townrow[✉]*^c and Malte Fischer[✉]*^a

We present the synthesis and characterization of the parent isolable monomeric triplet titanocene complex, stabilized by the N-heterocyclic carbene (NHC) IMe₄. Investigated by SQUID magnetometry and quantum-chemical calculations, reactivity studies of the titanium precursor [Cp₂Ti(btmsa)] (2) with the NHC I^tPr₂Me₂ and the zirconocene complex [Cp₂Zr(py)(btmsa)] (1) revealed divergent reactivity patterns, highlighting the role of steric and electronic effects in stabilization.

Although the Group 4 elements titanium, zirconium, and hafnium show a strong thermodynamic preference for the +IV oxidation state, their complexes in the +II state are becoming increasingly significant in synthetic applications, ranging from catalysis to materials development.¹ Combining Lewis acid reactivity in the +IV state and reductive transformations in the +II state, systems capable of reversible redox cycling between these have the potential to unlock exciting new possibilities for utilizing these Earth-abundant elements in efficient redox catalysis.^{1,2} Additionally, paramagnetic titanium(III) organometallics have also found recent attention as molecular spintronic components based on Earth-abundant metals, adding further relevance to finding novel examples.³

However, due to the thermodynamic stability of the +IV oxidation state, accessing low-valent species typically requires ‘masking’ through the incorporation of redox non-innocent ligands or ligands with pronounced backbonding capabilities (e.g. alkenes and alkynes). These ligands provide access to con-

trolled sources of the low-valent form, serving as synthons for its generation.^{1,2,4} A key factor in advancing this field is a deep understanding of the electronic structure and nature of truly isolable low-valent Group 4 compounds and intermediates. Unlike their later transition metal homologues, titanocene and zirconocene [Cp₂M] (Cp = η⁵-cyclopentadienyl; M = Ti, Zr) are not isolable and decompose through C–H activation processes.⁵ It must be noted that even the permethylated derivative [Cp*₂Ti] (Cp* = η⁵-pentamethylcyclopentadienyl) is unstable and, for example, readily coordinates dinitrogen.⁶ However, its magnetic moment could be determined and was found to be in between 2.48 (129 K) and 2.60 μ_B (298 K), slightly lower than the spin-only value of 2.83 μ_B, and consistent with the presence of two unpaired electrons.⁶ Through the incorporation of bulky silyl groups at the cyclopentadienyl ligands, several triplet titanocene derivatives have been isolated and structurally characterized, such as [C₅Me₄(SiMe₂^tBu)₂Ti], which has a magnetic moment of 2.40 μ_B.⁷ The reactive parent Group 4 metallocene fragment [Cp₂M] is typically generated *in situ* by use of soluble, closed-shell-singlet synthons through the release of CO, PMe₃, alkenes, alkynes, or by mixing [Cp₂MCl₂] with strong reducing agents.^{1c,4,8,9} Notable examples include ([Cp₂ZrCl₂]/ⁿBuLi),¹⁰ [Cp₂Ti(PMe₃)₂],¹¹ [Cp₂Ti(CO)₂] (Scheme 1),¹² and btmsa (bis(trimethylsilyl)acetylene) complexes [Cp₂Zr(py)(btmsa)] (1) (py = pyridine), and [Cp₂Ti(btmsa)] (2).^{1c,4,8,13} Their chemistry

^aInstitut für Anorganische Chemie, Georg-August-Universität Göttingen, Tammannstraße 4, D-37077 Göttingen, Germany.


E-mail: malte.fischer@uni-goettingen.de

^bInstitut für Organische und Biomolekulare Chemie, Georg-August-Universität Göttingen, Tammannstraße 2, D-37077 Göttingen, Germany

^cInstitute of Nanotechnology, Karlsruhe Institut für Technologie, Hermann-von-Helmholtz-Platz 1, D-76344 Eggenstein-Leopoldshafen, Germany.

E-mail: oliver.townrow@kit.edu


†Electronic supplementary information (ESI) available: Experimental procedures, characterization of the reported compounds, crystallographic and computational details. CCDC 2419251–2419254. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d5dt00480b>



Scheme 1 Examples of bis(cyclopentadienyl)titanium(II) complexes.

has been extensively reviewed on many occasions.^{1c,4,8,14} In contrast to these commonly employed diamagnetic Group 4 metallocenes, we present the synthesis and isolation of a crystalline *paramagnetic* parent titanocene, stabilised by an NHC which better mimics the electronic state of free [Cp₂Ti].⁵ We further demonstrate how the choice of the metal precursor and NHC is crucial to metallocene formation, as varying these caused either ligand exchange, imidazole tautomerisation, or C–H activation. The reaction of **1** with IMe₄ (1,3,4,5-tetramethylimidazol-2-ylidene) in C₆D₆ at room temperature resulted in an expected clean ligand exchange reaction in which the NHC replaces pyridine to yield [Cp₂Zr(IMe₄)(btmsa)] (**3**) (Scheme 2). The ligand exchange is visibly evident by an immediate colour change of the reaction mixture from intense purple to yellow–brown upon NHC addition. Inspection of the ¹H NMR spectrum revealed the chemical shifts of free pyridine, in addition to divergence of the Me₃Si signals of the btmsa ligand, indicating that free rotation is now sterically prohibited (Fig. S2†). The ¹³C chemical shift of the NHC carbene centre is located at δ¹³C{¹H} = 197.8 ppm, confirmed by ¹H/¹³C HMBC NMR (Fig. S4†), shifted when compared to that of the free carbene (δ¹³C{¹H} = 213.3 ppm).¹⁵ It is worth noting that neither the addition of a second equivalent of IMe₄ to **3** nor the direct addition of two equivalents of IMe₄ to **1** led to any exchange of btmsa for IMe₄, even at elevated temperatures (Fig. S5†). Yellow crystals of **3** suitable for single-crystal X-ray diffraction (SCXRD) were obtained from a saturated *n*-hexane solution at –30 °C, which allowed for structural confirmation. The coordination environment at zirconium is best described as distorted square pyramidal according to the τ₅ structural parameter (τ₅ = 0.11) (Fig. S6†).¹⁶ The C_{NHC}–Zr bond length of 2.466(3) Å is in good agreement to other IMe₄ ligated zirconium complexes, *e.g.* [(η⁷-C₇H₇)(η⁵-C₅H₅)Zr(IMe₄)] (2.4452(17) Å).^{17,18} Interestingly, no reaction readily occurs between **1** and the slightly larger NHC ⁱPr₂Me₂ (1,3-diisopropyl-4,5-dimethylimidazol-2-ylidene). However, upon prolonged heating of this mixture up to 90 °C, **1** and ⁱPr₂Me₂ are completely consumed, and the characteristic signals of free pyridine

are observed by ¹H NMR spectroscopy. In addition, three signals in the typical range of cyclopentadienyl ligands and two new signals typical of hydride species are found, indicative of the formation of multiple products (Fig. S7†). After several attempts, a few dark yellow crystals were obtained from a saturated *n*-hexane solution at –30 °C, and the structure of the κ¹-imidazole complex **4** was confirmed by SCXRD (Fig. S8†), with structural parameters in good agreement to those of **3**.¹⁹ The formation of **4** suggests that propene is formed as a by-product. However, the expected ABX coupling pattern in the ¹H NMR spectrum could not be unambiguously assigned due to the formation of additional byproducts and overlapping signals. Therefore, the headspace of the reaction, performed in toluene as the solvent, was analysed *via* GC-MS, confirming the presence of propene (Fig. S19 and S20†). Furthermore, trimethylsilane, ethyltrimethylsilane, vinyltrimethylsilane, pyridine and the employed solvents were detected, with the presence of the various silanes underscoring the complexity of the reaction's outcome.¹⁹ It is worth noting that although the NHC-to-imidazole transformation is unusual, the reverse imidazole to NHC reactivity pattern has precedence in the literature.²⁰ We then explored the reactions of these NHCs with the lighter, titanium, analogue [Cp₂Ti(btmsa)] (**2**). Contrary to the formation of **4**, the reaction of **2** with one equivalent of ⁱPr₂Me₂ in aromatic solvents occurs at room temperature, resulting in the immediate release of btmsa, as confirmed by its ¹H NMR chemical shift at 0.16 ppm (Fig. S9†). Although the reaction onset occurred rapidly, full conversion of **2** proceeds slowly and plateaus at around 50% after 72 hours. Heating the mixture to 90 °C for an extended period eventually drives the reaction toward near completion; however, approximately half an equivalent of the NHC remains unreacted, and signs of a paramagnetic product begin to emerge.¹⁹ Upon work-up at this stage, dark violet crystals suitable for SCXRD are consistently obtained, revealing the formation of the hydride-bridged dititanium complex **5** (Fig. 1). Additionally, formal loss of dihydrogen and C–H activation of one cyclopentadienyl ligand led to the formation of bridging fulvalene and cyclopentadienyl



Scheme 2 Reactions of [Cp₂Ti(btmsa)] (**1**) and [Cp₂Zr(py)(btmsa)] (**2**) towards IMe₄ and ⁱPr₂Me₂ to give complexes **3**, **4**, **5**, and **6**.






Fig. 1 Molecular structure of **5** in the crystal. Anisotropic displacement parameters are drawn at the 50% probability level, with data collected at 100 K (hydrogen atoms attached to carbon omitted for clarity). Selected bond lengths (Å) and angles (deg): Ti1...Ti2 3.1183(6), Ti1–C1 2.318(2), Ti1–H1 1.92(4), Ti2–C13 2.166(2), Ti2–H1 1.84(4), Ti1–H1–Ti2 112.6, Ct1–Ti1–Ct2 136.0, Ct3–Ti2–Ct4 133.5 (Ct1 = C12–C15, Ct2 = C22–C26, Ct3 = C17–C20, Ct4 = C27–C31).

ligands to both titanium atoms and only one NHC ligation, which explains why approximately half an equivalent of the NHC remains unreacted. The Ti1–C1 bond length is 2.318(2) Å and clearly exceeds the respective single bond covalent radii of the respective elements ($\Sigma_{\text{cov}}(\text{Ti}–\text{C}) = 2.11$ Å),²¹ indicating expected donor–acceptor interactions. This bond length aligns well with other NHC-ligated titanium hydrides, such as $[(\text{C}_8\text{H}_4(1,4\text{-Si}^i\text{Pr}_3)_2)_2\{\text{IME}_4\}\text{Ti}(\mu\text{-H})\text{TiH}]$ (**I**) reported by Tsoureas *et al.*, in which also only one of the titanium(III) centres is coordinated by an NHC (2.300(2) Å).²² The Ti–H bond lengths (Ti1–H1 1.92(4) Å; Ti2–H1 1.84(4) Å) also compare favourably to **I**, as well as to $[(\eta^5\text{-C}_5\text{H}_3\text{-1,2-Me}_2)_2\text{Ti}(\mu\text{-H})\text{Ti}(\eta^5\text{-C}_5\text{H}_3\text{-1,2-Me}_2)(\eta^5, \eta^1\text{-C}_5\text{H}_2\text{-1,2-Me}_2)]$ (**II**) (1.79(2) Å and 1.91(2) Å).^{22,23} Both titanium atoms exhibit a distorted tetrahedral coordination environment according to the τ_4 and τ_4' structural parameters (Ti1: 0.83, 0.73; Ti2: 0.79, 0.73).²⁴ The Ti2–C13 bond length of 2.166(2) Å is typical of a single bond, falling between the single-bond covalent radii of titanium and carbon (*vide supra*) and matching the analogous bond length in **II** (2.204(2) Å).^{21,23} The folding of the fulvalene ligand is evident from a dihedral angle of 17.6(2)° between the two planes defined by both five-membered rings, as well as the inter-ring carbon–carbon bond length of 1.452(4) Å, both of which are diagnostic of fulvalene dititanium complexes (*cf.* $[\{\text{Cp}_2\text{Ti}(\mu\text{-F})\}_2(\mu\text{-}\eta^5\text{-C}_{10}\text{H}_8)]$ (1.464(8) Å and 15.3(3)°)).²⁵ Inspection of the ^1H NMR spectrum from crystalline **5** in C_6D_6 at room temperature revealed several broad signals between $\delta^1\text{H} = -25$ and +35 ppm, indicative of a paramagnetic compound (Fig. S14†).

Finally, we turned our attention to the remaining combination: $[\text{Cp}_2\text{Ti}(\text{btmsa})]$ (**2**) and IME_4 . Upon adding one equivalent of IME_4 to a C_6D_6 solution of **2**, three new broad signals at –29.48, 9.92, and 93.46 ppm were detected alongside the characteristic signal of free btmsa in the ^1H NMR spectrum, indicating a rapid ligand exchange reaction that occurs within

a few minutes, yielding $[\text{Cp}_2\text{Ti}(\text{IME}_4)]$ (**6**) (Scheme 2 and Fig. S16, S17). Slow diffusion of *n*-pentane into a benzene solution of **6** produced dark yellow crystals, allowing for the unambiguous elucidation of the molecular structure by SCXRD (Fig. 2, top). Notably, complex **6** crystallises as dark yellow plates in the triclinic space group $P\bar{1}$, with three independent molecules in the unit cell, mainly due to different orientations of the Cp ligands (eclipsed and staggered conformers observed); the other structural parameters of all three molecules are in very good agreement. The average titanium– C_{NHC} bond length is 2.27 Å, which is slightly shorter compared to the herein reported complex **5**. In relation to a related formal three-coordinate and NHC-ligated titanium complex, such as $[(\text{C}_6\text{H}_6)_2\text{Ti}(\text{IME}_2\text{H}_2)]$ (2.323(3) Å), this bond length is reduced and corresponds to the different titanium oxidation states in the two complexes.²⁶ The titanium centre is in a trigonal planar coordination environment with a pronounced widening of the Ct1–Ti1–Ct2 angle (144.5°). The IME_4 ligand is aligned almost perpendicular to the Ct1–Ti1–Ct2 axis, exhibiting a dihedral angle of 81.8°. The UV-vis spectrum of $\text{Cp}_2\text{Ti}(\text{IME}_4)$ reveals an absorption maximum at 389 nm ($\epsilon = 1 \text{ g L}^{-1}$), in accordance to its (dark) yellow colour (Fig. S18†).

To gain further insight into the electronic ground state of complex **6**, it was investigated using SQUID magnetometry (see




Fig. 2 Top: Molecular structure of $[\text{Cp}_2\text{Ti}(\text{IME}_4)]$ (**6**) in the crystal. Anisotropic displacement parameters are drawn at the 50% probability level, with data collected at 100 K (hydrogen atoms omitted for clarity). Selected bond lengths (Å) and angles (deg): Ti1–C1 2.2712(9), C1–N1 1.3624(10), C1–N2 1.3603(10), C4–C5 1.3543(13), Ct1–Ti1–Ct2 144.5, Ct1–Ti1–C1 106.8, Ct2–Ti1–C1 108.6; Bottom: variable-temperature $\chi_{\text{M}}T$ product for **6** (open circles) and best fit curve (red line); Insert: calculated spin density (Triplet; UPBE0-D3BJ/Def2-TZVP; isovalue = 0.15 Å^{–3}).



sition numbers 2419251–2419254 for (3–6)[†] and can be obtained from <https://www.ccdc.cam.ac.uk/structures/>.

There are no conflicts to declare.

Financial support (VCI Liebig fellowships for O.P.E.T. and M. F.; YIG Prep Pro fellowship for O.P.E.T.) is gratefully acknowledged. The NMR, MS, and EA services at the Faculty of Chemistry of the Georg-August-Universität Göttingen are kindly acknowledged for measurements and technical assistance. Support from the DFG (INST 186/1237-1 and INST 186/1324-1) is also gratefully acknowledged. The authors acknowledge support by the state of Baden-Württemberg through bwHPC and the German Research Foundation (DFG) through grant no INST 40/575-1 FUGG for access to the JUSTUS 2 cluster.

- 1 Selected recent review articles: (a) M. Manßen and L. L. Schafer, *Chem. Soc. Rev.*, 2020, **49**, 6947–6994; (b) S. Fortier and A. Gomez-Torres, *Chem. Commun.*, 2021, **57**, 10292–10316; (c) U. Rosenthal, *ChemistryOpen*, 2021, **10**, 1234–1243.
- 2 E. P. Beaumier, A. J. Pearce, X. Y. See and I. A. Tonks, *Nat. Rev. Chem.*, 2019, **3**, 15–34.
- 3 (a) L. C. de Camargo, M. Briganti, F. S. Santana, D. Stinghen, R. R. Ribeiro, G. G. Nunes, J. F. Soares, E. Salvadori, M. Chiesa, S. Benci, R. Torre, L. Sorace, F. Totti and R. Sessoli, *Angew. Chem., Int. Ed.*, 2020, **60**, 2588–2593; (b) M. Briganti, G. Serrano, L. Poggini, A. L. Sorrentino, B. Cortigiani, L. C. de Camargo, J. F. Soares, A. Motta, A. Caneschi, M. Mannini, F. Totti and R. Sessoli, *Nano Lett.*, 2022, **22**, 8626–8632; (c) S. Wisbeck, A. L. Sorrentino, F. S. Santana, L. C. de Camargo, R. R. Ribeiro, E. Salvadori, M. Chiesa, N. Giaconi, A. Caneschi, M. Mannini, L. Poggini, M. Briganti, G. Serrano, J. F. Soares and R. Sessoli, *Chem. Sci.*, 2024, **15**, 14390–14398.
- 4 U. Rosenthal, *Chem. Soc. Rev.*, 2020, **49**, 2119–2139.
- 5 (a) M. Polasek and J. Kubista, *J. Organomet. Chem.*, 2007, **692**, 4073–4083; (b) P. J. Chirik, *Organometallics*, 2010, **29**, 1500–1517.
- 6 J. E. Bercaw, *J. Am. Chem. Soc.*, 1974, **96**, 5087–5095.
- 7 (a) P. B. Hitchcock, F. M. Kerton and G. A. Lawless, *J. Am. Chem. Soc.*, 1998, **120**, 10264–10265; (b) L. Lukesova, M. Horacek, P. Stepnicka, K. Fejfarova, R. Gyepes, I. Cisarova, J. Kubista and K. Mach, *J. Organomet. Chem.*, 2002, **663**, 134–144; (c) L. Lukesova, J. Pinkas, M. Horacek,

The data supporting this article have been included as part of the ESI.† Additional references have been cited within the ESI.†^{27–47} Crystallographic data have been deposited at the Cambridge Crystallographic Data Centre (CCDC) under depo-

- R. Gyepes, J. Kubista and K. Mach, *J. Organomet. Chem.*, 2006, **691**, 748–758.
- 8 (a) U. Rosenthal, *Organometallics*, 2020, **39**, 4403–4414; (b) U. Rosenthal, *ChemistryOpen*, 2019, **8**, 1036–1047.
- 9 See for example: (a) D. W. Stephan, *Organometallics*, 1992, **11**, 996–999; (b) G. Bousrez, I. Dechamps, J.-L. Vasse and F. Jaroschik, *Dalton Trans.*, 2015, **44**, 9359–9362; (c) G. Bousrez, D. Harakat, S. Chevreux, I. Déchamps-Olivier and F. Jaroschik, *Dalton Trans.*, 2024, **53**, 15595–15601.
- 10 (a) E. Negishi, F. E. Cederbaum and T. Takahashi, *Tetrahedron Lett.*, 1986, **27**, 2829–2832; (b) E. Negishi and S. Huo, in *Titanium and Zirconium in Organic Synthesis*, ed. I. Marek, Wiley-VCH, Weinheim, 2002, pp. 1–49.
- 11 (a) L. B. Kool, M. D. Rausch, H. G. Alt, M. Herberhold, U. Thewalt and B. Wolf, *Angew. Chem., Int. Ed. Engl.*, 1985, **24**, 394–401; (b) L. B. Kool, M. D. Rausch, H. G. Alt, M. Herberhold, B. Honold and U. Thewalt, *J. Organomet. Chem.*, 1987, **320**, 37–45.
- 12 (a) J. G. Murray, *J. Am. Chem. Soc.*, 1959, **81**, 752–753; (b) J. L. Atwood, K. E. Stone, H. G. Alt, D. C. Hrnčir and M. D. Rausch, *J. Organomet. Chem.*, 1975, **96**, C4–C6.
- 13 (a) V. V. Burlakov, A. V. Polyakov, A. I. Yanovsky, Y. T. Struchkov, V. B. Shur, M. E. Vol'pin, Z. Rosenthal and H. Görls, *J. Organomet. Chem.*, 1994, **476**, 197–206; (b) U. Rosenthal, A. Ohff, W. Baumann, A. Z. Tillack, H. Görls, V. V. Burlakov and V. B. Shur, *Z. Anorg. Allg. Chem.*, 1995, **621**, 77–83; (c) J. R. Nitschke, S. Zürcher and T. D. Tilley, *J. Am. Chem. Soc.*, 2000, **122**, 10345–10352; (d) U. Rosenthal, V. V. Burlakov, P. Arndt, W. Baumann and A. Spannenberg, *Organometallics*, 2003, **22**, 884–900; (e) M. Fischer, L. Vincent-Heldt, M. Hillje, M. Schmidtman and R. Beckhaus, *Dalton Trans.*, 2020, **49**, 2068–2072.
- 14 See for example: (a) P. Chirik, *Organometallics*, 2010, **29**, 1500–1517; (b) Z. Xi and Z. Li, *Top. Organomet. Chem.*, 2004, **8**, 27–56.
- 15 J. Klauke, N. Sinthathurai, C. Golz, O. P. E. Townrow and M. Fischer, *ChemRxiv*, 2024, preprint, DOI:DOI: [10.26434/chemrxiv-2024-4lzc5](https://doi.org/10.26434/chemrxiv-2024-4lzc5).
- 16 A. W. Addison, T. N. Rao, J. Reedijk, J. van Rijn and G. C. Verschoor, *J. Chem. Soc., Dalton Trans.*, 1984, 1349–1356.
- 17 R. J. Baker, T. Bannenberg, A. Kunst, S. Randoll and M. Tamm, *Inorg. Chim. Acta*, 2006, **359**, 4797–4801.
- 18 For a summary of Group 14 NHC complexes and structural parameters see: C. D. Zhang and G. Zi, *Chem. Soc. Rev.*, 2015, **44**, 1898–1921.
- 19 For details see the ESI.†
- 20 See for example: (a) J. Ruiz and B. F. Perandones, *J. Am. Chem. Soc.*, 2007, **129**, 9298–9299; (b) M. Brill, J. Diaz, M. A. Huertos, R. Lopez, J. Perez and L. Riera, *Chem. – Eur. J.*, 2011, **17**, 8584–8595.
- 21 (a) P. Pyykkö and M. Atsumi, *Chem. – Eur. J.*, 2009, **15**, 186–197; (b) P. Pyykkö and M. Atsumi, *Chem. – Eur. J.*, 2009, **15**, 12770–12779.
- 22 N. Tsoureas, J. C. Green and F. G. N. Cloke, *Chem. Commun.*, 2017, **53**, 13117–13120.
- 23 S. S. Semproni, C. Milsman and P. J. Chirik, *Organometallics*, 2012, **31**, 3672–3682.
- 24 (a) L. Yang, D. R. Powell and R. P. Houser, *Dalton Trans.*, 2007, 955–964; (b) A. Okuniewski, D. Rosiak, J. Chojnacki and B. Becker, *Polyhedron*, 2015, **90**, 47–57.
- 25 P. Yu, E. F. Murphy, H. W. Roesky, P. Lubini, H.-G. Schmidt and M. Noltemeyer, *Organometallics*, 1997, **16**, 313–316.
- 26 H. Braunschweig, C. Brückner, M. A. Celik, K. Dück, F. Hupp, T. Kramer, J. Krebs and I. Krummenacher, *Chem. – Eur. J.*, 2015, **21**, 11056–11064.
- 27 C. Barnett, M. L. Cole and J. B. Harper, *Eur. J. Inorg. Chem.*, 2021, 4954–4958.
- 28 N. Kuhn and T. Kratz, *Synthesis*, 1993, 561–562.
- 29 E. Bill, *Simulation of the experimental magnetic data was performed with the julX_2S program*, Max-Planck Institute for Chemical Energy Conversion, Mülheim/Ruhr, Germany.
- 30 Bruker, *SAINT, V8.40B*, Bruker AXS Inc., Madison, Wisconsin, USA.
- 31 L. Krause, R. Herbst-Irmer, G. M. Sheldrick and D. Stalke, *J. Appl. Crystallogr.*, 2015, **48**, 3–10.
- 32 G. M. Sheldrick, *Acta Crystallogr., Sect. A:Found. Adv.*, 2015, **71**, 3–8.
- 33 G. M. Sheldrick, *Acta Crystallogr., Sect. C:Struct. Chem.*, 2015, **71**, 3–8.
- 34 O. V. Dolomanov, L. J. Bourhis, R. J. Gildea, J. A. K. Howard and H. Puschmann, *J. Appl. Crystallogr.*, 2009, **42**, 339–341.
- 35 C. R. Groom, I. J. Bruno, M. P. Lightfoot and S. C. Ward, *Acta Crystallogr.*, 2016, **B72**, 171–179.
- 36 D. Kratzert, *FinalCif, V132*, <https://dkratzert.de/finalcif.html>.
- 37 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. V. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery Jr, J. E. Peralta, F. Ogliaro, M. J. Bearpark, J. J. Heyd, E. N. Brothers, K. N. Kudin, V. N. Staroverov, T. A. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. P. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman and D. J. Fox, *Gaussian 16, Revision A.03*, Gaussian, Inc., Wallingford CT, 2016.
- 38 A. D. Becke, *Phys. Rev. A:At., Mol., Opt. Phys.*, 1998, **38**, 3098–3100.
- 39 J. P. Perdew, *Phys. Rev. B:Condens. Matter Mater. Phys.*, 1986, **33**, 8822–8824.
- 40 J. P. Perdew, K. Burke and M. Ernzerhof, *Phys. Rev. Lett.*, 1996, **77**, 3865–3868.



- 41 C. Adamo and V. Barone, *J. Chem. Phys.*, 1999, **110**, 6158–6170.
- 42 T. Yanai, D. P. Tew and N. C. Handy, *Chem. Phys. Lett.*, 2004, **393**, 51–57.
- 43 F. Weigend and R. Ahlrichs, *Phys. Chem. Chem. Phys.*, 2005, **7**, 3297–3305.
- 44 S. Grimme, J. Antony, S. Ehrlich and H. Krieg, *J. Chem. Phys.*, 2010, **132**, 154104.
- 45 S. Grimme, S. Ehrlich and L. Goerigk, *J. Comput. Chem.*, 2011, **32**, 1456–1465.
- 46 *NBO 6.0*, E. D. Glendening, J. K. Badenhoop, A. E. Reed, J. E. Carpenter, J. A. Bohmann, C. M. Morales, C. R. Landis and F. Weinhold, Theoretical Chemistry Institute, University of Wisconsin, Madison, WI, 2013.
- 47 A. Gilbert, *IQMol 3.1.4*, 2024.

