

RESEARCH ARTICLE

View Article Online
View Journal

Cite this: DOI: 10.1039/d5qi01092f

Design of porous self-assembled homoleptic and heteroleptic Pd²⁺ cages incorporating silicon-based fluoride acceptors: the way towards nuclear imaging applications†Marika Drexler,^a Ioannis Kanavos,^b Daniela Krauss,^a Guillermo Moreno-Alcántar,^a Lydia M. Smith,^c Madeleine E. George,^c Timothy H. Witney,^c Ryszard Lobinski,^{b,d} Luisa Ronga^{ID} *^b and Angela Casini^{ID} *^a

Supramolecular self-assembly is a promising tool to develop multifunctional theranostic agents and has recently entered the field of radiochemistry. In this work, lantern-shaped [Pd₂L₄]⁴⁺ metallacages featuring a blood–brain barrier-penetrating peptide were designed for dual-modality imaging. Two silicon-based fluoride acceptors were incorporated in the cage ligand for radiolabeling with fluorine-18, an isotope used in positron emission tomography (PET). Ligands (**L1**, **L2**) and the respective homoleptic cages (**C1_{hom}** [Pd₂(**L1**)₄]⁴⁺, **C2_{hom}** [Pd₂(**L2**)₄]⁴⁺) were fully characterized by NMR spectroscopy, liquid chromatography and high-resolution electrospray mass spectrometry (HR-ESI-MS). Cage stability was assessed in different solvents and concentrations by HPLC. ¹⁸F-labelled cages were obtained by radiolabeling the ligands pre-assembly under mild conditions within four hours via [¹⁹F]-to-[¹⁸F] isotopic exchange. The high lipophilicity of the ligands was also assessed *in vitro* (log *D*_{pH7.4}) and *in ovo*, using the chick chorioallantoic membrane (CAM) model. Furthermore, formation of host–guest complexes between the metallacages and perrhenate (ReO₄[−]), the ‘cold’ surrogate of radioactive ^{99m}TcO₄[−] (used for single photon computed tomography, SPECT), could be detected by mass spectrometry, although the adduct did not sustain chromatographic separation. To improve stability of the supramolecular system, heteroleptic cages of general formula [Pd₂(L)_{*m*}(**L0**)_{*n*}]⁴⁺ (*m* = 1, 2, *n* = 4 − *m*) were synthesized by statistical self-assembly and separated by liquid chromatography in good yield. Overall, this study demonstrates the feasible adaptation of supramolecular principles to achieve innovative theranostic agents.

Received 8th May 2025,

Accepted 31st May 2025

DOI: 10.1039/d5qi01092f

rsc.li/frontiers-inorganic

Introduction

In the last few decades, supramolecular coordination complexes (SCCs) have been explored in different fields of chemistry due to the broad span of possible applications, including in medicine.^{1–3} SCCs are discrete molecular arrangements composed by metal nodes bound to organic ligands that come

together *via* coordination-driven self-assembly (CDSA).^{4,5} A plethora of two-dimensional and three-dimensional shapes can be created upon thoughtful combination of the aforementioned building blocks, which morphologically are classified broadly into three main types, namely metallacycles, metallacages and helicates.^{6,7} The coordination geometry of the metal ion, ligand curvature, size and number of coordination sites encode the self-assembly of the proposed architecture.^{3,8} In this context, the case of the metallacages (MCgs) is particularly appealing since their porous nature favors their use to encapsulate and transport small molecules.⁹ Moreover, *exo*- and *endo*-functionalization of the ligands pre- and post-CDSA allows for further tuning of the supramolecular system. The host–guest chemistry and versatility of MCgs towards functionalization led to their investigation as catalysts,^{10,11} sensors^{12,13} and gas storage systems¹⁴ amongst other possible uses.¹⁵ Concerning biomedical applications, MCgs have been initially explored as potential carriers of cargos of pharmaceutical interest^{16–20} either for therapy or imaging (Fig. 1A), or

^aChair of Medicinal and Bioinorganic Chemistry, School of Natural Sciences, Technical University of Munich, Garching b., München, Germany.

E-mail: angela.casini@tum.de

^bUniversité de Pau et des Pays de l'Adour, E2S UPPA, CNRS, IPREM, 2 Av. du Président Pierre Angot, Pau, France. E-mail: luisa.ronga@univ-pau.fr

^cSchool of Biomedical Engineering and Imaging Sciences, King's College London, St Thomas' Hospital, London, SE1 7EH, UK

^dChair of Analytical Chemistry, Warsaw University of Technology, ul. Noakowskiego 3, 00-664 Warszawa, Poland

† Electronic supplementary information (ESI) available: NMR spectra, chromatograms, MS data, gamma counter data, *in ovo* imaging data. See DOI: <https://doi.org/10.1039/d5qi01092f>








Fig. 2 (A) Design and synthesis of BBB-targeted ligands L1 and L2 by Fmoc-SPPS. The exo-functionalization consists of a BBB shuttle peptide α PepH3, a silicon-based fluoride acceptor (SiFA) for radiolabeling with the PET nuclide fluoride-18 and a PEG linker (4-((2-(2-(2-aminoethoxy)ethoxy)ethyl)amino)-4-oxobutanoic acid) that acts as a spacer. (B) Coordination-driven self-assembly of homoleptic and heteroleptic MCs, C_{hom} and C_{het} . Reaction conditions: 4 eq. ligand (in total), 2 eq. $[Pd(MeCN)_4](BF_4)_2$, DMSO, RT, 1 h. (C) 1H NMR (top) and 1H -DOSY NMR (bottom) spectra in DMSO- d_6 (400 MHz) of $C2_{hom}$ (purple) compared to L2 (blue). The α -pyridinyl protons a and b shift upon metal coordination as explained by the scheme. (D) Mass spectra of $C2_{hom}$ (top) and the host-guest complex $ReO_4^- \cdot C2_{hom}$ (bottom) obtained by DI-HR-ESI-MS. Corresponding theoretical and experimental isotopic patterns are shown in the graph insets.



bioconjugated to MCgs structures.⁴⁸ Interestingly, the peptide has been reported to cross the BBB *via* an active transport mechanism, namely adsorptive-mediated transcytosis (AMT), due to the presence of positively charged amino acid residues in its sequence.⁵² Here, the physiologically more stable *D*-enantiomer of PepH3 (*D*-PepH3) was applied.⁵³ The suitability of this design strategy was challenged by exploring the fundamentals of homoleptic and heteroleptic CDSA (Fig. 2B). The lipophilic character of the ligands ($\log D_{\text{pH}7.4}$) was determined, and in one case, also investigated *in ovo* in glioblastoma U87 tumors engrafted on the chicken chorioallantoic membrane (CAM) model by PET. Further, the obtained homoleptic MCgs were characterized by different methods, including ¹H NMR spectroscopy and high-resolution electrospray ionization mass spectrometry (HR-ESI-MS). In addition, the encapsulation properties of the homoleptic systems were investigated by using the guest molecule perrhenate as a cold surrogate for ^{99m}TcO₄[−]. Further, a protocol for the synthesis of ¹⁸F-labeled MCgs *via* pre-assembly labeling was optimized, which held radiochemically pure MCgs. In parallel, the synthesis of heteroleptic MCgs, of general formula [Pd₂(L)_{*m*}(LO)_{*n*}]⁴⁺ (*m* = 1, 2, *n* = 4 − *m*) was achieved by statistical self-assembly of two different types of ligands,⁵⁴ followed by chromatographic separation.

Experimental

General information

All reagents and solvents were purchased commercially and were used without further purification. The synthesis of SiFAs building block was achieved as previously reported,^{55,56} and their purity >95% were determined by reverse-phase high performance liquid chromatography (RP-HPLC). Ligand **L0** (2-(3,5-bis(pyridin-3-ylethynyl)phenyl)acetic acid) was synthesized according to literature-known procedure.⁵⁷ The product was obtained as a white solid in a total yield of 66% and 97% purity by RP-HPLC.

Fluoride-18 in target water ([¹⁸O]H₂O) was obtained from Klinikum rechts der Isar (Munich, Germany) and Guys and St Thomas' Hospital (London, UK).

¹H NMR and ¹H-DOSY NMR spectra were recorded on a Bruker Avance III HD 400 NMR instrument by Bruker Cooperations (Billerica, USA). Chemical shifts are given in parts per million (ppm) and are referenced to the residual proton signal of the solvent. The coupling constants *J* are reported in hertz (Hz) and the signal splitting patterns are expressed as singlet (s), doublet (d), doublet of doublets (dd), triplet (t), doublet of triplets (dt), doublet of quartets (dq) and multiplet (m).

Analytical and semi-preparative RP-HPLC measurements were performed using gradient systems by Shimadzu Corp. (Kyoto, Japan), consisting of a SPD-20A dual wavelength UV/Vis detector (*l* = 220 nm, *l* = 254 nm), two LC-20AD gradient pumps, a CBM-20A communication unit and a CTO-20A column oven. For analytical RP-HPLC measurements a flow

rate of 1 mL min^{−1} and for semi-preparative RP-HPLC measurements a flow rate of 10 mL min^{−1} was used. Quality controls of the ligands and analysis of cage formation reactions were performed using a MultoKrom® 100-5 C-18 column (150 × 4.6 mm, 5 μm particle size) from CS-Chromatographie Service GmbH (Langerwehe, Germany). For analytical separation of ligands and cages, different gradients of H₂O (0.1% trifluoroacetic acid (TFA)) (A) and MeCN (2% H₂O, 0.1% TFA) (B) were used during RP-HPLC measurements. For semi-preparative purification of ligands, different gradients of H₂O (0.1% TFA) (A) and MeCN (5% H₂O, 0.1% TFA) (B) were used.

Quality controls of radiolabeled products was conducted using a gradient system by Shimadzu Corp. (Kyoto, Japan), consisting of an SPD-20A dual wavelength UV/Vis detector (*l* = 220 nm, *l* = 254 nm), two LC-20AD gradient pumps, a HERM LB500 (NaI scintillation crystal) as well as a FlowStar² LB514 radio detector from Berthold Technologies GmbH (Bad Wilbad, Germany) and a CBM-20A communication unit. Radiolabeled ligands were analysed using a MultoKrom® 100-5 C-18 column (125 × 4 mm, 5 μm particle size) with a constant flow rate of 1 mL min^{−1}. Analysis of the radiolabeled cages was performed using a MultoHigh® Bio 300-5 C4 column (150 × 4.6 mm, 5 μm particle size) at a flow rate of 1 mL min^{−1}. Radio thin-layer chromatography (TLC) was performed on a ScanRAM radio TLC detector by LabLogic (Sheffield, UK) and analysed with the Laura software.

Direct-infusion (DI) and liquid chromatography (LC) high-resolution electrospray-ionization mass spectrometry (HR-ESI-MS) was applied for the analysis of ligands and MCgs using an Orbitrap Q-Exactive Plus Mass Spectrometer from Thermo Fisher Scientific. For LC-MS, the MS system was coupled to Dionex ultimate 3000 series UHPLC from Thermo Fisher Scientific for LC-MS. The column used for the C2_{hom} analysis was a SUPELCO BIOShell A400 Protein C4 (3.4 μm, 400 Å, 15 cm × 1 mm) from Sigma-Aldrich. The mobile phases were A: H₂O and B: MeCN, both with 0.1% formic acid (FA). The flow rate was 0.05 mL min^{−1} and sample elution was performed by using a gradient from 20% to 40% of B over 3 min, followed by a gradient from 40% to 80% of B over 15 min. Sample injection volume was 10 μL. For the C1_{het} analysis the column used was an AcclaimTM 120: C18 (5 μm, 120 Å, 4.6 mm × 100 mm), Dionex Bonded Silica Products from Thermo Fisher Scientific. The mobile phases were A: H₂O and B: ACN, both with 0.1% formic acid (FA). The flow rate was 0.5 mL min^{−1} and sample elution was performed by using a gradient from 20% to 40% of B over 3 min, followed by a gradient from 40% to 80% of B over 15 min. Ionization was performed using an ESI source operating in positive ion mode with a capillary voltage of 3.50 kV and capillary temperature 320 °C. Sheath gas, auxiliary gas and sweep gas flow rate were set at 20, 12 and 1 (arbitrary units), respectively. Auxiliary gas temperature was set at 100 °C.

For DI, the sample was infused at 0.01 mL min^{−1} and ionization was performed using an ESI source operating in positive ion mode with a capillary voltage of 3.50 kV and capillary temperature of 300 °C. Sheath gas, auxiliary gas and sweep



gas flow rate were set at 20, 12 and 0 (arbitrary units), respectively. Auxiliary gas temperature was set at 50 °C.

Ligand synthesis

Synthesis of the BBB-shuttle peptide $_D\text{PepH3}$. $_D\text{PepH3}$ (AGILKRW) was synthesized *via* microwave-assisted solid phase peptide synthesis on a Liberty Blue™ instrument manufactured by CEM (Matthews, USA). ProTide Rink Amide resin (loading capacity = 0.65 mmol g⁻¹) in combination with Fmoc-protected amino acids were used for the synthesis. No final Fmoc-deprotection was performed and the peptide was stored on the resin until further use.

Synthesis of $_D\text{PepH3-}_D\text{Dap}((\text{SiFA})\text{BA})\text{-Ebes-L0 (L1)}$

Fmoc-deprotection. The N-terminal Fmoc-protected $_D\text{PepH3}$, immobilized on resin, (60.0 μmol, 1.0 eq.) was deprotected by addition of 20% piperidine in dimethyl formamide (DMF, 5 mL) at RT. The deprotection step was carried out twice (1 × 15 min, 1 × 5 min) and afterwards the resin was washed with DMF (6 × 5 mL).

Fmoc- $_D\text{Dap}(\text{Dde})\text{-OH}$ (*N*-alpha-(9-fluorenylmethyl-oxycarbonyl)-*N*-beta-[(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene) ethyl]-D-2,3-diaminopropionic acid) (90.0 μmol, 1.5 eq.), TBTU (*N,N,N',N'*-tetramethyluronium-tetrafluoroborate) (90.0 μmol, 1.5 eq.), HOAt (1-hydroxy-7-azabenzotriazole) (90.0 μmol, 1.5 eq.) and 2,4,6-trimethylpyridine (300 μmol, 5.0 eq.) were dissolved in 5 mL DMF. After 2 min preincubation, the solution was added to the deprotected resin-bound peptide and shaken for 2 h at RT. Afterwards the resin was washed with DMF (6 × 5 mL) and *N*-methyl-2-pyrrolidone (NMP, 6 × 5 mL). For a coupling in branched position the Dde-protection group was cleaved by adding a solution of NH₂OH·HCl (6.0 mmol, 100 eq.) and imidazole (4.5 mmol, 75 eq.) in 7 mL NMP/DCM (5/2). After shaking for 2 h at RT, the resin was washed with NMP (6 × 5 mL) and DMF (6 × 5 mL).

Amide coupling. Fmoc-(SiFA)BA (72.0 μmol, 1.2 eq.), TBTU (90.0 μmol, 1.5 eq.) and HOAt (90.0 μmol, 1.5 eq.) were dissolved in 5 mL DMF. After the addition of DIPEA (*N,N*-diisopropylethylamine) (210 μmol, 3.5 eq.) the solution was preincubated for 30 min. The coupling solution was added to the deprotected resin-bound compound and the reaction mixture was shaken at RT for 2.5 h. Afterwards the resin was washed with DMF (6 × 5 mL).

Fmoc-Ebes (*N*-[8-(9-fluorenylmethyl-oxycarbonyl)amino-3,6-dioxaoctyl]succinamic acid) (90.0 μmol, 1.5 eq.) and **L0** (90.0 μmol, 1.5 eq.) were coupled consecutively to the resin-bound compound by first performing Fmoc-deprotection and then amide coupling.

Resin cleavage. The ligand was cleaved off the resin by shaking with 2 × 5 mL of a mixture of TFA/triisopropyl silane (TIPS)/H₂O (95/2.5/2.5) for 2 × 1 h at RT. The ligand solution was collected in a round-bottom flask and stirred for 1 h before being concentrated in an N₂ stream. Then, 5 mL of TFA was added and the mixture was stirred at RT for another 2 h. Afterwards, the TFA was removed again in an N₂ stream and the ligand was purified *via* semi-preparative RP-HPLC and lyophilized.

$_D\text{PepH3-}_D\text{Dap}((\text{SiFA})\text{BA})\text{-Ebes-L0}$ was obtained as a white solid (11.5 mg, 11% yield, 98% purity).

RP-HPLC (10–90% B): t_R = 9.7 min.

¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm) = 10.78 (s), 8.78 (d, J = 3.1 Hz, 2H), 8.61 (d, J = 6.5 Hz, 2H), 8.49 (d, J = 4.4 Hz), 8.26 (d, J = 6.6 Hz), 8.21 (d, J = 5.9 Hz), 8.14–8.08 (m), 8.04–7.97 (m), 7.98–7.92 (m), 7.89–7.79 (m), 7.68 (s), 7.67–7.61 (m), 7.61–7.53 (m), 7.52–7.43 (m), 7.39 (s), 7.32 (d, J = 7.1 Hz), 7.11 (d, J = 2.6 Hz), 7.08–7.01 (m), 7.00–6.92 (m), 4.43 (dq, J = 13.6, 6.7 Hz), 4.33–3.92 (m), 3.81–3.58 (m), 3.53–3.46 (m), 3.43 (t, J = 5.8 Hz), 3.36 (t, J = 5.9 Hz), 3.20 (dt, J = 30.4, 5.7 Hz), 3.12–2.92 (m), 2.72 (d, J = 7.5 Hz), 2.36 (s), 1.85–1.36 (m), 1.25 (d, J = 7.4 Hz), 1.11 (s), 1.01 (d, J = 1.1 Hz), 0.85 (d, J = 6.4 Hz), 0.83–0.73 (m).

MS (ESI positive): calculated monoisotopic mass for C₉₀H₁₂₄FN₁₉O₁₄Si: 1741.93; found by ESI-MS: m/z = 582 [$M + 3H$]³⁺, 872 [$M + 2H$]²⁺.

Synthesis of $_D\text{PepH3}(\text{SiFA})\text{SeFe-Ebes-L0 (L2)}$. Fmoc-SiFA (SeFe) (72.0 μmol, 1.2 eq.), Fmoc-Ebes (90.0 μmol, 1.5 eq.) and **L0** (90.0 μmol, 1.5 eq.) were coupled consecutively to the resin-bound peptide by first performing Fmoc-deprotection and then amide coupling as described above. A resin cleavage was carried out to obtain $_D\text{PepH3}(\text{SiFA})\text{SeFe-Ebes-L0}$ as a white solid (17.6 mg, 17% yield, 98% purity) as described for the synthesis of **L1**.

RP-HPLC (10–90% B): t_R = 9.5 min.

¹H NMR (400 MHz, DMSO-*d*₆) δ (ppm) = 10.79 (s), 8.79 (s, 2H), 8.73 (d, J = 6.5 Hz), 8.61 (dd, J = 4.9, 1.7 Hz, 2H), 8.52–8.43 (m), 8.39–8.30 (m), 8.28–8.17 (m), 8.00 (dt, J = 7.9, 1.8 Hz), 7.98–7.78 (m), 7.75–7.62 (m), 7.61–7.53 (m), 7.53–7.45 (m), 7.39 (s), 7.32 (d, J = 7.9 Hz), 7.12 (d, J = 2.4 Hz), 7.09–7.01 (m), 7.00–6.92 (m), 4.44 (dt, J = 19.4, 6.9 Hz), 4.33 (d, J = 5.9 Hz), 4.30–3.93 (m), 3.74 (d, J = 5.8 Hz), 3.50 (dd, J = 4.3, 2.6 Hz), 3.43 (t, J = 5.7 Hz), 3.37 (t, J = 5.9 Hz), 3.29–3.14 (m), 3.14–2.94 (m), 2.73 (d, J = 7.5 Hz), 2.46–2.29 (m), 2.24 (d, J = 0.6 Hz), 2.08 (s), 1.97–1.90 (m), 1.71–1.58 (m), 1.56–1.42 (m), 1.37 (d, J = 7.1 Hz), 1.23 (s), 1.11 (s), 1.03–0.97 (m), 0.88–0.70 (m).

MS (ESI positive): calculated monoisotopic mass for C₈₈H₁₂₁FN₁₈O₁₃Si: 1684.91; found by ESI-MS: m/z = 562 [$M + 3H$]³⁺, 844 [$M + 2H$]²⁺, 1686 [$M + H$]⁺.

Synthesis of metallacages by CDSA and RP-HPLC analysis

To achieve homoleptic MCgs assembly, ligand **L1** or **L2** (2.0 eq., 5 mM in DMSO) and [Pd(MeCN)₄](BF₄)₂ (1.0 eq., 20 mM in DMSO) were mixed in DMSO to reach a theoretical cage concentration of 1 mM. Instead, CDSA of the heteroleptic cages was achieved by either mixing 1.0 eq. of **L1** (5 mM in DMSO) and 3.0 eq. of **L0** (5 mM in DMSO) for **C1_{het}**, or 2.7 eq. of **L2** (5 mM in DMSO) and 1.3 eq. of **L0** (5 mM in DMSO) for **C2_{het}**, with 2.0 eq. of [Pd(MeCN)₄](BF₄)₂ (20 mM in DMSO) and adding DMSO to reach a theoretical cage concentration of 1 mM.

In all cases, the reaction was complete after 1 h at RT and analysed by analytical RP-HPLC. For the homoleptic cages a gradient of 40–70% B in 15 min was applied. In the case of the



RP-HPLC (40–70% B in 15 min): t_R = 7.3 min. Yield 82.1%.

num TLC plated F₂₅₄, MeCN/H₂O, 8/2, +10% sodium acetate (2 M), +1% TFA).

¹⁸F-*D*PepH3-*D*Dap((SiFA)BA)-Ebes-L0 (¹⁸F-L1)

RCY: 44%, RCP: 98%.

Radio RP-HPLC (10–80% B in 15 min): *t*_R = 12.4 min.

¹⁸F-*D*PepH3-(SiFA)SeFe-Ebes-L0 (¹⁸F-L2)

RCY: 47%, RCP: 99%.

Radio RP-HPLC (30–60% B in 15 min): *t*_R = 10.4 min.

Self-assembly of ¹⁸F-[Pd₂(L1)₄](BF₄)₄ (¹⁸F-C1_{hom}) and ¹⁸F-[Pd₂(L2)₄](BF₄)₄ (¹⁸F-C2_{hom}). 30 nmol of L1 or L2 (2.0 eq., 5 mM in DMSO) were labeled according to the previously described procedure. The resulting solution of ¹⁸F-L1 or ¹⁸F-L2 was evaporated at 80 °C with an applied N₂ flow. 3 μL of [Pd(MeCN)₄](BF₄)₂ (4.0 eq., 20 mM in DMSO) and 3 μL of DMSO were added to the labeled ligand. After 4 h, the reaction was complete and analysed *via* radio RP-HPLC (40–70% B in 15 min, C4 column).

¹⁸F-[Pd₂(L1)₄](BF₄)₄ (¹⁸F-C1_{hom})

Radio RP-HPLC (40–70% B in 15 min): *t*_R = 9.6 min.

¹⁸F-[Pd₂(L2)₄](BF₄)₄ (¹⁸F-C2_{hom})

Radio RP-HPLC (40–70% B in 15 min): *t*_R = 7.7 min.

Lipophilicity determination (log *D*_{pH7.4})

The octanol-PBS partition coefficient (log *D*_{pH7.4} values) were determined by addition of 0.5 MBq of the labeled ligand to a 1.5 mL LoBind Eppendorf tube, filled with 500 μL octanol and 500 μL PBS (*n* = 8). The tubes were vortexed for 3 min at RT and centrifuged (9000 rpm, 5 min, RT) using a Heraeus Pico 17 centrifuge by Thermo Scientific (Waltham, USA) (24 × 1.5/2.0 mL rotor with ClickSeal-Lid). Out of each tube 200 μL of each solvent layer were taken out and the activity was quantified separately using a γ-counter by PerkinElmer Inc. (Langerwehe, Germany). Lipophilicity of the cages could not be determined due to stability issues upon vortexing of their solution.

$$\log D_{\text{pH}7.4} (^{18}\text{F-L1}) = 2.53 \pm 0.05.$$

$$\log D_{\text{pH}7.4} (^{18}\text{F-L2}) = 2.07 \pm 0.07.$$

Cell culture

U87 cells were cultivated at 37 °C (+5% CO₂) in MEME (Sigma Life Science) containing 10% fetal bovine serum (Thermo Fisher), 1% L-glutamine (Sigma Life Science) and 1% penicillin–streptomycin (Sigma Aldrich). The cells were split 1/6 twice a week using trypsin–EDTA solution (Thermo Fisher) for detachment. The cells were tested Mycoplasma-free every month.

In ovo studies

All *in ovo* experiments were carried out on fertilized Dekalb white or brown eggs (Henry Stewart & co. Ltd, UK). Experiments took place before embryonic day 14 (E14). The eggs were incubated for up to 14 days at 12–14 °C in a wine cooler (Haller) with humidified atmosphere until use. To grow tumors on the chick chorioallantoic membrane (CAM), eggs were cleaned with Brinsea disinfectant (100×) and moved to an incubator (Brinsea) with a temperature of 38.7 °C and 48%

humidity. The first day of incubation at this temperature was embryonic day 0 (E0). To loosen the CAM from the eggshell the incubator trays were slowly tilted from one side to the other until E3. On E3, eggs were taken from the incubator for window cutting. The eggs were rolled to prevent the CAM sticking to the shell and placed on a cushioned holder. They were pierced at the wide base where the air cell is located and approximately 5 mL of albumin was removed through the hole using a syringe with a 19G needle. The hole was then resealed with Scotch Magic tape. Next, a square of tape was placed onto one side of the egg and four rectangularly arranged holes were punched into the shell through the tape. A rectangular window (1 × 2 cm) was made with sharp dissection scissors by carefully cutting three sides of a rectangle into the shell using the holes for orientation and without damaging the inner shell membrane. The window was closed with tape and the eggs placed in the incubator again until E7. U87 cells were maintained as described previously and cell culture media was renewed 24 h prior to harvesting. On the day of inoculations (E7), cells were detached from the flask, resuspended in media and aliquots with 3 × 10⁶ cells were prepared. The aliquots were centrifuged for 3 min at 500g, 4 °C in a Centrifuge 5424 R (Eppendorf) and stored on ice. Meanwhile, Matrigel® Matrix Basement Membrane (Corning) was defrosted on ice. Eggs were taken from the incubator, placed on an egg holder and the windows were opened to locate the CAM. After dabbing the CAM dry with a sterile lens tissue, a suspension of the U87 cell pellet in 20 μL of Matrigel was pipetted onto the CAM. The eggs were closed with tape, labelled accordingly and placed in the incubator for a further 7 days. On E14 the eggs were removed from the incubator and placed on a cushioned holder. For tumor imaging the shell window was enlarged to allow for direct injection of the radiotracer into a CAM blood vessel.

Preparation of the radiotracer for *in ovo* studies. The radio-labeled ligand ¹⁸F-L1 was prepared as described before. The obtained solution in EtOH was reduced to 100 μL in an N₂ stream at 60 °C and then diluted with PBS to obtain until EtOH <25%. A sample was taken for quality control *via* radio RP-HPLC on a 1260 Infinity II LC-instrument (Agilent Technologies, MeCN (+5% H₂O + 0.1% TFA)/H₂O (+0.1% TFA), 10% → 80% MeCN gradient over 15 min, 1.5 mL min⁻¹ flow rate) with a GABI Nova detector (Elysia-Raytest) on a ZORBAX Eclipse XDB-C18 Analytical 4.6 × 250 mm 5 Micron column (Agilent Technologies). For *in ovo* injection, 150 μL an activity of 5 MBq were prepared by further dilution with saline (0.9 wt% NaCl).

In ovo PET imaging. On E14 a CAM vein was cannulated and 90 μL of a 1 mg mL⁻¹ solution of the anaesthetic medetomidine (Virbac) was pipetted onto the surface of the CAM. Eggs were left for 15 min at RT before receiving an intravenous bolus injection of ~3 MBq of the labelled radiotracer on the imaging bed (<150 μL), followed by 50 μL PBS (Sigma Life Science). A 60 min dynamic PET scan was performed using a Mediso NanoScan PET/CT system (1–5 coincidence mode; 3D reconstruction; computed tomography (CT) attenuation cor-



Table 1 Calculated and experimental m/z values (most abundant isotope) for MCg species and host–guest adducts with perhenate

Compound	Formula	m/z		Observed adducts
		Theoretical	Measured	
C1_{hom}	C ₃₆₀ H ₄₉₆ F ₄ N ₇₆ O ₅₆ Pd ₂ Si ₄	661.8758 ($z = 11$)	661.8747 664.2367	[M ⁴⁺ + FA + ACN + 7H ⁺] ¹¹⁺ [M ⁴⁺ + 2FA + Na + 7H ⁺] ¹¹⁺
[Pd ₂ (L1) ₃]	C ₂₇₀ H ₃₇₂ F ₃ N ₅₇ O ₄₂ Pd ₂ Si ₃	686.4560 ($z = 8$)	686.4570 692.3329 695.2059 700.5811 704.9545 709.2046	[M ⁴⁺ + FA + 4H ⁺] ⁸⁺ [M ⁴⁺ + 2FA + 4H ⁺] ⁸⁺ [M ⁴⁺ + 2FA + Na ⁺ + 4H ⁺] ⁸⁺ [M ⁴⁺ + 2DMSO + 4H ⁺] ⁸⁺ [M ⁴⁺ + Cl ⁻ + 2DMSO + 5H ⁺] ⁸⁺ [M ⁴⁺ + 2Cl ⁻ + 2DMSO + 6H ⁺] ⁸⁺
C2_{hom}	C ₃₅₂ H ₄₈₄ F ₄ N ₇₂ O ₅₂ Si ₄ Pd ₂	637.3200 ($z = 11$)	637.3203 641.0483 643.5020	[M ⁴⁺ + FA + 7H ⁺] ¹¹⁺ [M ⁴⁺ + FA + ACN + 7H ⁺] ¹¹⁺ [M ⁴⁺ + 2FA + Na + 7H ⁺] ¹¹⁺
ReO ₄ C C2 _{hom}	C ₃₅₂ H ₄₈₄ F ₄ N ₇₂ O ₅₆ Si ₄ Pd ₂ Re	705.5519 ($z = 10$) 655.9506 ($z = 11$)	705.5536 655.9501	[M ⁴⁺ + 2FA + 6H ⁺] ¹⁰⁺ [M ₃ ⁺ + 8H ⁺] ¹¹⁺
[Pd ₂ (L2) ₃]	C ₂₆₄ H ₃₆₃ F ₃ N ₅₄ O ₃₉ Si ₃ Pd ₂	670.8237 ($z = 8$)	670.8237 683.4458 687.8202	[M ⁴⁺ + 2FA + 4H ⁺] ⁸⁺ [M ⁴⁺ + 2DMSO + Cl ⁻ + 5H ⁺] ⁸⁺ [M ⁴⁺ + 2DMSO + 2Cl ⁻ + 6H ⁺] ⁸⁺
ReO ₄ C [Pd ₂ (L2) ₃]	C ₂₆₄ H ₃₆₃ F ₃ N ₅₄ O ₄₃ Si ₃ Pd ₂ Re	696.4408 ($z = 8$)	696.4399	[M ³⁺ + FA + 5H ⁺] ⁸⁺
C1_{het}	C ₁₅₆ H ₁₆₆ FN ₂₅ O ₂₀ Pd ₂ Si	594.2134 ($z = 5$) 742.7652 ($z = 4$)	594.2142 742.7651	[M ⁴⁺ + H ⁺] ⁵⁺ [M ⁴⁺] ⁴⁺
C2_{het}	C ₁₅₄ H ₁₆₃ FN ₂₄ O ₁₉ SiPd ₂	582.8092 ($z = 5$)	582.8102 605.8088	[M ⁴⁺ + H ⁺] ⁵⁺ [M ⁴⁺ + 2FA + Na] ⁵⁺
[Pd ₂ (L2) ₂ (L0) ₂]	C ₂₂₀ H ₂₇₀ F ₂ N ₄₀ O ₃₀ Si ₂ Pd ₂	609.2671 ($z = 7$)	609.2677 625.4096	[M ⁴⁺ + 3H ⁺] ⁷⁺ [M ⁴⁺ + 2FA + Na ⁺ + 2H ⁺] ⁷⁺




Fig. 3 Characterization of the homoleptic cage **C2_{hom}** by LC-HR-ESI-MS. Left: TIC (mass range between 600–820 m/z) and extracted ion chromatograms (XIC) of ions m/z 637.32 ($z = 11$, **C2_{hom}** at $t_R = 13.94$ min) and 670.83 ($z = 8$, [Pd₂(L2)₃] + 2 FA at $t_R = 14.37$ min). Right: mass spectrum of **C2_{hom}** + FA at $t_R = 13.94$ min (comparison of theoretical and isotopic pattern is provided as insert) and [Pd₂(L2)₃] + 2 FA at $t_R = 14.37$ min. The mobile phases were A: H₂O and B: MeCN, both with 0.1% formic acid (FA). The flow rate was 0.05 mL min⁻¹ and sample elution was performed by using a gradient from 20% to 40% of B over 3 min, followed by a gradient from 40% to 80% of B over 15 min.



Heteroleptic metallacages. Heterolepticity in MCs design is a promising strategy to increase multi-functionalization as ligands with different properties are combined in one unique structure. In recent work, the CDSA of a heteroleptic Pd₂L(**L0**)₃-type cage was a viable solution to introduce a peptide-functionalized ligand featuring DOTA as chelator for lutetium-177.⁵⁰ In order to favor the self-assembly of the heteroleptic cage **C1_{het}** featuring a **L1/L0** ratio of 1 : 3 – [Pd₂(**L1**)(**L0**)₃]⁴⁺ – 1 eq. of **L1** and 3 eq. of **L0** were reacted with [Pd(MeCN)₄](BF₄)₄ in DMSO for 1 hour at RT. ¹H NMR spectroscopy revealed a deshielding of the α-pyridinyl protons to 9.76 and 9.40 ppm, respectively, compared to the reference spectra of **L1** and **L0** (Fig. S25[†]). Moreover, the ¹H-DOSY NMR spectrum showed more than one newly formed species, upon addition of the Pd²⁺ precursor, suggesting a mixture of supramolecular assemblies (Fig. S26[†]). The self-assembly process was further investigated by RP-HPLC, where the species could be separated and analysed by MS (Fig. S27A and B[†]). By comparing the chroma-

A) Bar chart showing the percentage of intact cages for C1_{hom}, C2_{hom}, and C1_{hom} + C2_{hom} in water, DMSO, MeCN, and HPLC solvent at 1 h, 4 h, and 24 h. The y-axis represents % intact cage (0 to 100). The x-axis shows the solvents. The legend indicates: C1_{hom} (white), C2_{hom} (hatched), 1 h (light blue), 4 h (medium blue), and 24 h (dark blue).

Solvent	Condition	1 h	4 h	24 h
water	C1 _{hom}	~98	~95	~95
	C2 _{hom}	~88	~75	~62
	C1 _{hom} + C2 _{hom}	~92	~81	~74
DMSO	C1 _{hom}	~92	~81	~74
	C2 _{hom}	~48	~34	~27
	C1 _{hom} + C2 _{hom}	~81	~63	~52
MeCN	C1 _{hom}	~81	~63	~52
	C2 _{hom}	~72	~52	~51
	C1 _{hom} + C2 _{hom}	~72	~52	~51
HPLC solvent *	C1 _{hom}	~92	~81	~49
	C2 _{hom}	~90	~86	~74
	C1 _{hom} + C2 _{hom}	~90	~86	~74


B) HPLC chromatograms showing the elution of C1_{hom} and L1 at different time points (0 min, 1 h, 4 h, 24 h) in HPLC solvent. The x-axis represents retention time t_R [min] (0 to 15). The y-axis represents intensity. The legend indicates: DMSO (black), 1 h (light blue), 4 h (medium blue), and 24 h (dark blue). The peaks are labeled C1_{hom} and L1.

This journal is © the Partner Organisations 2025

For the assembly of the heteroleptic $[\text{Pd}_2(\text{L2})(\text{L0})_3]^{4+}$ (C2_{het}) cage, tuning of the ratio **L2/L0** was required. Eventually, a mixture of heteroleptic species was obtained by combining 2.7 eq. of **L2** and 1.3 eq. of **L0** and reacting with $[\text{Pd}(\text{MeCN})_4]$


Encapsulation studies

One of the strategies to radiolabel metallacages is *via* encapsulation of the radioactive guest molecule. The encapsulation of $^{99m}\text{TcO}_4^-$ and stability of different host-guest systems *in vivo* have already been demonstrated successfully by the groups of Lusby and Archibald as well as by Correia and Casini.⁴⁸ In both studies perrhenate was used as a ‘cold’ surrogate for pertechnetate and the host-guest interaction was investigated either by NMR spectroscopy or *via in silico* methods. Following these promising results, we investigated the encapsulation of perrhenate in our MCgs. Thus, 2 eq. of NaReO_4 were added to solutions of **C1_{hom}** and **C2_{hom}** in $\text{D}_2\text{O}/\text{DMSO-}d_6$ (1/1) and ^1H NMR spectra were recorded after 30 min. Deuterated water was




Inorg. Chem. Front.

Therefore, the ability of the cages to encapsulate perhenate in predominantly aqueous media was additionally evaluated by DI-HR-ESI-MS in the case of **C2_{hom}**. The comparison of the mass spectra of **C2_{hom}** and the host-guest system (ReO₄⁻)**C2_{hom}** in Fig. 2D, unambiguously revealed the formation of the host-guest adduct through the appearance of signals at *m/z* 655.9501 and 721.3443, which can be attributed to the 11⁺ and 10⁺ charge state of (ReO₄⁻)**C2_{hom}**, respectively (Table 1). While the encapsulation is not quantitative, this result confirms some affinity of ReO₄⁻ for the cage cavity in aqueous solution, at concentrations more relevant to the clini-



Since the *D*PepH3-conjugated ligands **L1** and **L2** are modified with a SiFA moiety, labeling of the metallacage structure with

In the case of **L1**, a $\log D_{\text{pH}7.4}$ of 2.53 ± 0.05 of the radio-labeled ligand was determined. To assess the effects of such high lipophilicity *in vivo*, the organ uptake of **¹⁸F-L1** was evaluated in the CAM model (Fig. 7A). This preclinical tool can be used for high-throughput tumor imaging in a time- and cost-effective way and is in line with the ethical principles of reduction, refinement and replacement (3Rs).⁶⁵ Glioblastoma



F-18 is possible *via exo*-functionalization. Here, the radiolabeling can either proceed before CDSA (pre-assembly), during CDSA (one-pot) or after CDSA (post-assembly).⁴³ In our case, to avoid cage disassembly in the relatively ‘harsh’ radiofluorination conditions, the ligands were radiolabeled pre-assembly and then CDSA was performed to form the homoleptic cages. Thus, after labeling 30 nmol of ligand (1 eq.) with ~200 MBq of fluoride-18 and elution in EtOH, the solvent was fully evaporated and CDSA was conducted by adding 2 eq. of [Pd(MeCN)₄](BF₄)₂ in 6 μL of DMSO (Fig. 6A). These conditions enabled the optimal concentration of each component (mM range) for successful CDSA. The self-assembly was monitored by radio RP-HPLC at 15 min, 60 min and 4 h (Fig. 6C). As depicted in Fig. 6B, 73% conversion to ¹⁸F-C1_{hom} (*t_R* = 9.6 min) was reached after 4 h with 18% remaining ¹⁸F-L1 (*t_R* = 6.7 min), while full conversion to ¹⁸F-C2_{hom} (*t_R* = 7.7 min) was achieved. In both cases slight defluorination (¹⁸F-C1_{hom}: 9%, ¹⁸F-C2_{hom}: 3%) occurred as evidenced by the F-18 peak at *t_R* = 2.4 min. Further purification of the reaction mixtures by analytical radio RP-HPLC or *via* cartridges was not carried out due to the short half-life of F-18 of 109.7 min, as well as possible instability of the radiolabeled cages.

Conclusions

In this work, we aimed at further substantiating the applicability of supramolecular coordination complexes, namely porous MCgs, for radiopharmaceutical applications. These nano-structures can benefit from the self-assembly multicomponent design for functionalization and pharmacokinetics modulation, but at the same time their size range (1-few nm) enables their precise characterization by classical separation and analytical methods; thus, offering an ideal versatile chemical space to achieve next generation nuclear imaging agents.

Here, two bispyridinyl ligands featuring a BBB shuttle peptide- and different silicon-based fluoride acceptor synthons (L1 and L2) were successfully synthesized by solid-phase peptide synthesis and radiolabeled with fluorine-18. The lipophilicity of the ligands was determined *in vitro*, *via* standard log *D*_{PH7.4} determination, and for L1 also *ex vivo* using the CAM model. Afterwards, formation of the respective homoleptic metallacages C1_{hom} and C2_{hom} was achieved by CDSA and confirmed by NMR spectroscopy, RP-HPLC and HR-ESI-MS. Importantly, ¹⁸F-radiolabeled homoleptic cages were achieved by labeling the ligands pre-assembly and *via* formation of the cages *in situ*. The optimization of the isotopic exchange protocol was an important step towards the future applicability of MCgs for PET imaging in a clinical setting.

Additionally, partial encapsulation of ReO₄[−] into the cages following mixing of the components was evidenced by HR-ESI-MS; however, the extent and stability of the (ReO₄[−])CMCg host-guest adducts appeared less pronounced with respect to previously reported Pd₂L₄ cage structures.⁴⁸ Thus, further ligand optimization is necessary to validate the suitability of these porous systems for encapsulation of

anionic radioactive counterparts, such as ^{99m}TcO₄[−] for SPECT imaging or the β[−]-emitting ¹⁸⁸ReO₄[−].

In order to improve the stability of the homoleptic Pd-based cages in solution, heteroleptic MCgs were also synthesized by statistical self-assembly and characterized by ESI-MS. The obtained results showed that [Pd₂L(L0)₃]⁴⁺ (C1/2_{het}) and [Pd₂L₂(L0)₂]⁴⁺ were the main assembled species.

In the future, an option to increase MCg's stability and to enable optimized radiofluorination of the cages post-assembly, could be replacing Pd²⁺ with the more kinetically inert Pt²⁺ to achieve [Pt₂L₄]⁴⁺ cages. The latter have already shown to be more stable with respect to ligand exchange than their Pd-based counterparts,^{67,68} which would also improve their stability in physiological conditions. Alternatively, organometallic building blocks endowed with higher metal-carbon bond stability could also be integrated in the cage scaffold.¹

Overall, our results highlight the current challenges in the construction of targeted theranostics based on SCCs for future clinical applications in nuclear medicine. While much effort still needs to be spent on the development of metallacages for biomedical applications, their design flexibility and easiness of functionalization offer a unique toolbox for the obtainment of multimodal theranostics.

Author contributions

A. C. and L. R. were responsible for the project conceptualization and supervision. M. D. and A. C. wrote the first draft of the manuscript and all co-authors contributed to the Results and discussion sections. M. D., I. K. and A. C. designed the graphics. M. D. and D. K. performed the synthesis and radiolabeling of ligands and cages. I. K., L. R. and R. L. carried out high-resolution mass spectrometry studies and performed the data analysis. G. M.-A. and A. C. provided expertise in MS and NMR data analysis. M. D., L. M. S., M. E. G. and T. H. W. contributed to, and supervised, the *in ovo* studies.

Data availability

The data supporting this article have been included as part of the ESI.† Raw data for this article, including chromatograms, NMR spectra, MS data, gamma counter data for log *D* determination, *in ovo* imaging data are available at Zenodo at [<https://doi.org/10.5281/zenodo.15358484>].

Conflicts of interest

The authors declare no conflict of interest.

Acknowledgements

We acknowledge funding from EU Horizon Europe Research and Innovation program, EIC Pathfinder Open network



“SMARTdrugs” (grant agreement ID: 101129886). Funding for T. H. W. was provided by UK Research and Innovation (UKRI) under the UK Government’s Horizon Europe funding guarantee (grant no. 10091247) to T. H. W., arising from the above mentioned EIC Pathfinder grant. The TUM International Graduate School of Science and Engineering (IGSSE) is kindly acknowledged for financial support (International Project Team: Supramolecular Organometallic and Coordination Complexes for Medical Applications, SOCoMED). This research has also received funding from the European Union’s Horizon H2020 Research and Innovation under the Marie Skłodowska-Curie grant agreement no. 945416 through the PhD fellowship of IK. The authors thank Dr Simon Godin for training and technical assistance with LC-MS/MS instrument.

References

- 1 A. Pöthig and A. Casini, Recent Developments of Supramolecular Metal-based Structures for Applications in Cancer Therapy and Imaging, *Theranostics*, 2019, **9**, 3150.
- 2 A. Casini, B. Woods and M. Wenzel, The Promise of Self-Assembled 3D Supramolecular Coordination Complexes for Biomedical Applications, *Inorg. Chem.*, 2017, **56**, 14715.
- 3 H. Sepehrpour, W. Fu, Y. Sun and P. J. Stang, Biomedically Relevant Self-Assembled Metallacycles and Metallacages, *J. Am. Chem. Soc.*, 2019, **141**, 14005.
- 4 D. L. Caulder and K. N. Raymond, Supermolecules by Design, *Acc. Chem. Res.*, 1999, **32**, 975–982.
- 5 L. Chen, Q. Chen, M. Wu, F. Jiang and M. Hong, Controllable Coordination-Driven Self-Assembly: From Discrete Metallocages to Infinite Cage-Based Frameworks, *Acc. Chem. Res.*, 2015, **48**, 201–210.
- 6 J. Malina, P. Scott and V. Brabec, Recognition of DNA/RNA bulges by antimicrobial and antitumor metallohelices, *Dalton Trans.*, 2015, **44**, 14656–14665.
- 7 J. Malina, H. Kosthrunova, P. Scott and V. Brabec, FeII Metallohelices Stabilize DNA G-Quadruplexes and Downregulate the Expression of G-Quadruplex-Regulated Oncogenes, *Chem. – Eur. J.*, 2021, **27**, 11682–11692.
- 8 S. Pullen, J. Tessarolo and G. H. Clever, Increasing structural and functional complexity in self-assembled coordination cages, *Chem. Sci.*, 2021, **12**, 7269–7293.
- 9 G. Moreno-Alcántar and A. Casini, Bioinorganic supramolecular coordination complexes and their biomedical applications, *FEBS Lett.*, 2023, **597**, 191.
- 10 W.-X. Gao, H.-N. Zhang and G.-X. Jin, Supramolecular catalysis based on discrete heterometallic coordination-driven metallacycles and metallacages, *Coord. Chem. Rev.*, 2019, **386**, 69–84.
- 11 R. Ham, C. J. Nielsen, S. Pullen and J. N. H. Reek, Supramolecular Coordination Cages for Artificial Photosynthesis and Synthetic Photocatalysis, *Chem. Rev.*, 2023, **123**, 5225–5261.
- 12 N. Dey and C. J. E. Haynes, Supramolecular Coordination Complexes as Optical Biosensors, *ChemPlusChem*, 2021, **86**, 418–433.
- 13 S. Ganta and D. K. Chand, Multi-Stimuli-Responsive Metallogel Molded from a Pd2L4-Type Coordination Cage: Selective Removal of Anionic Dyes, *Inorg. Chem.*, 2018, **57**, 3634–3645.
- 14 M. M. Deegan, M. R. Dworzak, A. J. Gosselin, K. J. Korman and E. D. Bloch, Gas Storage in Porous Molecular Materials, *Chem. – Eur. J.*, 2021, **27**, 4531–4547.
- 15 E. G. Percástegui, T. K. Ronson and J. R. Nitschke, Design and Applications of Water-Soluble Coordination Cages, *Chem. Rev.*, 2020, **120**, 13480–13544.
- 16 J. E. M. Lewis, E. L. Gavey, S. A. Cameron and J. D. Crowley, Stimuli-responsive Pd2L4 metallocupramolecular cages: towards targeted cisplatin drug delivery, *Chem. Sci.*, 2012, **3**, 778.
- 17 B. Therrien, G. Süß-Fink, P. Govindaswamy, A. K. Renfrew and P. J. Dyson, The “Complex-in-a-Complex” Cations [(acac)2MCRu6(p-IPrC6H4Me)6(tpt)2(dhbcq)3]6+: A Trojan Horse for Cancer Cells, *Angew. Chem., Int. Ed.*, 2008, **47**, 3773.
- 18 J. W. Yi, N. P. E. Barry, M. A. Furrer, O. Zava, P. J. Dyson, B. Therrien, *et al.*, Delivery of Floxuridine Derivatives to Cancer Cells by Water-Soluble Organometallic Cages, *Bioconjugate Chem.*, 2012, **23**, 461–471.
- 19 A. Schmidt, V. Molano, M. Hollering, A. Pöthig, A. Casini and F. E. Kühn, Evaluation of New Palladium Cages as Potential Delivery Systems for the Anticancer Drug Cisplatin, *Chem. – Eur. J.*, 2016, **22**, 2253.
- 20 J. Han, A. Schmidt, T. Zhang, H. Permentier, G. M. M. Groothuis, R. Bischoff, *et al.*, Bioconjugation strategies to couple supramolecular exo-functionalized palladium cages to peptides for biomedical applications, *Chem. Commun.*, 2017, **53**, 1405.
- 21 X. Wang, Q. Su, Z. Zhang, J. Yang, Y. Zhang and M. Zhang, Biotinylated platinum(II) metallacage towards targeted cancer theranostics, *Chem. Commun.*, 2020, **56**, 8460–8463.
- 22 G. Yu, B. Zhu, L. Shao, J. Zhou, M. L. Saha, B. Shi, *et al.*, Host–guest complexation-mediated codelivery of anti-cancer drug and photosensitizer for cancer phototherapy, *Proc. Natl. Acad. Sci. U. S. A.*, 2019, **116**, 6618–6623.
- 23 Y. Li, F. Huang, P. J. Stang and S. Yin, Supramolecular Coordination Complexes for Synergistic Cancer Therapy, *Acc. Chem. Res.*, 2024, **57**, 1174–1187.
- 24 J. Zhang, W. Ma, B. Yang, T. Shi, S. Liao, Y. Li, *et al.*, Biomimetic Metallacage Nanoparticles with Aggregation-Induced Emission for NIR-II Fluorescence Imaging-Guided Synergistic Immuno-Phototherapy of Tumors, *ACS Appl. Mater. Interfaces*, 2024, **16**, 69028–69044.
- 25 D. Xu, Y. Li, S. Yin and F. Huang, Strategies to address key challenges of metallacycle/metallacage-based supramolecular coordination complexes in biomedical applications, *Chem. Soc. Rev.*, 2024, **53**, 3167–3204.



- 26 R. D. Mukhopadhyay, Y. Kim, J. Koo and K. Kim, Porphyrin Boxes, *Acc. Chem. Res.*, 2018, **51**, 2730–2738.
- 27 G. Yu, S. Yu, M. L. Saha, J. Zhou, T. R. Cook, B. C. Yung, *et al.*, A discrete organoplatinum(II) metallacage as a multimodality theranostic platform for cancer photochemotherapy, *Nat. Commun.*, 2018, **9**, 4335.
- 28 X. Lin, F. Chen, X. Yu, H. Wang, H. Qiu, Y. Li, *et al.*, Phenylthiol-BODIPY-based supramolecular metallacycles for synergistic tumor chemo-photodynamic therapy, *Proc. Natl. Acad. Sci. U. S. A.*, 2022, **119**, e2203994119.
- 29 T. Rodríguez-Prieto, D. Wragg, N. Heiduk, M. Park, N. Strittmatter, R. A. Fischer, *et al.*, A Golden Touch in the Design of Multifunctional Porphyrin Metallacages: Host-Guest Chemistry for Drug-Target Interactions, *CCS Chem.*, 2024, **6**, 1662–1671.
- 30 J. Han, A. F. B. Räder, F. Reichart, B. Aikman, M. N. Wenzel, B. Woods, *et al.*, Bioconjugation of Supramolecular Metallacages to Integrin Ligands for Targeted Delivery of Cisplatin, *Bioconjugate Chem.*, 2018, **29**, 3856.
- 31 M. L. Saha, X. Yan and P. J. Stang, Photophysical Properties of Organoplatinum(II) Compounds and Derived Self-Assembled Metallacycles and Metallacages: Fluorescence and its Applications, *Acc. Chem. Res.*, 2016, **49**, 2527–2539.
- 32 B. Woods, D. Döllerer, B. Aikman, M. N. Wenzel, E. J. Sayers, F. E. Kühn, *et al.*, Highly luminescent metallacages featuring bispyridyl ligands functionalised with BODIPY for imaging in cancer cells, *J. Inorg. Biochem.*, 2019, **199**, 110781.
- 33 B. Aikman, R. Bonsignore, B. Woods, D. Doellerer, R. Scotti, C. Schmidt, *et al.*, Highly-fluorescent BODIPY-functionalised metallacages as drug delivery systems: synthesis, characterisation and cellular accumulation studies, *Dalton Trans.*, 2022, **51**, 7476–7490.
- 34 E. O. Bobylev, R. A. Knol, S. Mathew, D. A. Poole, I. Kotsogianni, N. I. Martin, *et al.*, In vivo biodistribution of kinetically stable Pt2L4 nanospheres that show anti-cancer activity, *Chem. Sci.*, 2023, **14**, 6943–6952.
- 35 C. J. Brown, F. D. Toste, R. G. Bergman and K. N. Raymond, Supramolecular Catalysis in Metal-Ligand Cluster Hosts, *Chem. Rev.*, 2015, **115**, 3012–3035.
- 36 A. Dissanayake, J. A. Sperry and J. R. Morrow, An octahedral coordination cage with six Fe(III) centers as a T1 MRI probe, *Chem. Commun.*, 2024, **60**, 12249–12252.
- 37 P. R. Sahoo, J. A. Sperry, S. G. Turowski and J. R. Morrow, Self-Assembled Iron(III) Coordination Cage as an MRI-Active Carrier for a Gold(I) Drug, *Bioconjugate Chem.*, 2024, **35**, 1618–1626.
- 38 R. Wang, L. An, J. He, M. Li, J. Jiao and S. Yang, A class of water-soluble Fe(III) coordination complexes as T1-weighted MRI contrast agents, *J. Mater. Chem. B*, 2021, **9**, 1787–1791.
- 39 G. E. Sokolow, M. R. Crawley, D. R. Morphet, D. Asik, J. A. Sperry, A. J. R. McGray, *et al.*, Metal–Organic Polyhedron with Four Fe(III) Centers Producing Enhanced T1 Magnetic Resonance Contrast in Tumors, *Inorg. Chem.*, 2022, **61**, 2603–2611.
- 40 B. P. Burke, W. Grantham, M. J. Burke, G. S. Nichol, D. Roberts, I. Renard, *et al.*, Visualizing Kinetically Robust CoIII4L6 Assemblies in Vivo: SPECT Imaging of the Encapsulated [99mTc]TcO4, Anion, *J. Am. Chem. Soc.*, 2018, **140**, 16877.
- 41 A. Ahmedova, Biomedical Applications of Metallosupramolecular Assemblies—Structural Aspects of the Anticancer Activity, *Front. Chem.*, 2018, **6**, 620.
- 42 J. Xu, J. Wang, J. Ye, J. Jiao, Z. Liu, C. Zhao, *et al.*, Metal-Coordinated Supramolecular Self-Assemblies for Cancer Theranostics, *Adv. Sci.*, 2021, **8**, 2101101.
- 43 G. Moreno-Alcántar, M. Drexler and A. Casini, Assembling a new generation of radiopharmaceuticals with supramolecular theranostics, *Nat. Rev. Chem.*, 2024, **8**, 893–914.
- 44 G. Yu, T. R. Cook, Y. Li, X. Yan, D. Wu, L. Shao, *et al.*, Tetraphenylethene-based highly emissive metallacage as a component of theranostic supramolecular nanoparticles, *Proc. Natl. Acad. Sci. U. S. A.*, 2016, **113**, 13720–13725.
- 45 A. B. S. Elliott, J. E. M. Lewis, H. van der Salm, C. J. McAdam, J. D. Crowley and K. C. Gordon, Luminescent Cages: Pendant Emissive Units on [Pd2L4]4+ “Click” Cages, *Inorg. Chem.*, 2016, **55**, 3440.
- 46 A. Schmidt, M. Hollering, M. Drees, A. Casini and F. E. Kühn, Supramolecular exo-functionalized palladium cages: fluorescent properties and biological activity, *Dalton Trans.*, 2016, **45**, 8556.
- 47 A. Schmidt, M. Hollering, J. Han, A. Casini and F. E. Kühn, Self-assembly of highly luminescent heteronuclear coordination cages, *Dalton Trans.*, 2016, **45**, 12297.
- 48 B. Woods, R. D. M. Silva, C. Schmidt, D. Wragg, M. Cavaco, V. Neves, *et al.*, Bioconjugate Supramolecular Pd2+ Metallacages Penetrate the Blood Brain Barrier In Vitro and In Vivo, *Bioconjugate Chem.*, 2021, **32**, 1399.
- 49 R. Cosiáls, C. Simó, S. Borrós, V. Gómez-Vallejo, C. Schmidt, J. Llop, *et al.*, PET Imaging of Self-Assembled 18F-Labelled Pd2L4 Metallacages for Anticancer Drug Delivery, *Chem. – Eur. J.*, 2023, **29**, e202202604.
- 50 S. Deiser, M. Drexler, G. Moreno-Alcántar, M. Irl, C. Schmidt, T. Günther, *et al.*, Synthesis of 177Lu-Labeled, Somatostatin-2 Receptor-Targeted Metalla-Assemblies: Challenges in the Design of Supramolecular Radiotherapeutics, *Inorg. Chem.*, 2023, **62**, 20710–20720.
- 51 V. Neves, F. Aires-da-Silva, M. Morais, L. Gano, E. Ribeiro, A. Pinto, *et al.*, Novel Peptides Derived from Dengue Virus Capsid Protein Translocate Reversibly the Blood-Brain Barrier through a Receptor-Free Mechanism, *ACS Chem. Biol.*, 2017, **12**, 1257.
- 52 D. Wu, Q. Chen, X. Chen, F. Han, Z. Chen and Y. Wang, The blood-brain barrier: Structure, regulation and drug delivery, *Signal Transduction Targeted Ther.*, 2023, **8**, 217.
- 53 M. Cavaco, J. Valle, R. da Silva, J. D. G. Correia, M. A. R. B. Castanho, D. Andreu, *et al.*, DPepH3, an Improved Peptide Shuttle for Receptor-independent

- Transport Across the Blood-Brain Barrier, *Curr. Pharm. Des.*, 2020, **26**, 1495.
- 54 J. E. M. Lewis, Molecular engineering of confined space in metal-organic cages, *Chem. Commun.*, 2022, **58**, 13873–13886.
 - 55 L. Iovkova, B. Wängler, E. Schirmmacher, R. Schirmmacher, G. Quandt, G. Boening, *et al.*, para-Functionalized Aryl-di-tert-butylfluorosilanes as Potential Labeling Synthons for ¹⁸F Radiopharmaceuticals, *Chem. – Eur. J.*, 2009, **15**, 2140–2147.
 - 56 S. Deiser, S. Fenzl, V. König, M. Drexler, L. M. Smith, M. E. George, *et al.*, (SiFA)SeFe: A Hydrophilic Silicon-Based Fluoride Acceptor Enabling Versatile Peptidic Radiohybrid Tracers, *J. Med. Chem.*, 2024, **67**, 14077–14094.
 - 57 B. Woods, M. N. Wenzel, T. Williams, S. R. Thomas, R. L. Jenkins and A. Casini, Exo-Functionalized Metallacages as Host-Guest Systems for the Anticancer Drug Cisplatin, *Front. Chem.*, 2019, **7**, 68.
 - 58 T. Günther, N. Holzleitner, D. Di Carlo, N. Urtz-Urban, C. Lapa and H.-J. Wester, Development of the First ¹⁸F-Labeled Radiohybrid-Based Minigastrin Derivative with High Target Affinity and Tumor Accumulation by Substitution of the Chelating Moiety, *Pharmaceutics*, 2023, **15**, 826.
 - 59 A. Höhne, L. Mu, M. Honer, P. A. Schubiger, S. M. Ametamey, K. Graham, *et al.*, Synthesis, ¹⁸F-Labeling, and in Vitro and in Vivo Studies of Bombesin Peptides Modified with Silicon-Based Building Blocks, *Bioconjugate Chem.*, 2008, **19**, 1871–1879.
 - 60 C. Wängler, B. Waser, A. Alke, L. Iovkova, H.-G. Buchholz, S. Niedermoser, *et al.*, One-Step ¹⁸F-Labeling of Carbohydrate-Conjugated Octreotate-Derivatives Containing a Silicon-Fluoride-Acceptor (SiFA): In Vitro and in Vivo Evaluation as Tumor Imaging Agents for Positron Emission Tomography (PET), *Bioconjugate Chem.*, 2010, **21**, 2289–2296.
 - 61 F. Kaiser, A. Schmidt, W. Heydenreuter, P. J. Altmann, A. Casini, S. A. Sieber, *et al.*, Self-Assembled Palladium and Platinum Coordination Cages: Photophysical Studies and Anticancer Activity, *Eur. J. Inorg. Chem.*, 2016, **2016**, 5189.
 - 62 R. Evans, G. Dal Poggetto, M. Nilsson and G. A. Morris, Improving the Interpretation of Small Molecule Diffusion Coefficients, *Anal. Chem.*, 2018, **90**, 3987–3994.
 - 63 E. Puig, C. Desmarests, G. Gontard, M. N. Rager, A. L. Cooksy and H. Amouri, Capturing a Square Planar Gold(III) Complex Inside a Platinum Nanocage: A Combined Experimental and Theoretical Study, *Inorg. Chem.*, 2019, **58**, 3189.
 - 64 A. K. Ghose, T. Herbertz, R. L. Hudkins, B. D. Dorsey and J. P. Mallamo, Knowledge-Based, Central Nervous System (CNS) Lead Selection and Lead Optimization for CNS Drug Discovery, *ACS Chem. Neurosci.*, 2012, **3**, 50–68.
 - 65 L. M. Smith, H. E. Greenwood, W. E. Tyrrell, R. S. Edwards, V. de Santis, F. Baark, *et al.*, The chicken chorioallantoic membrane as a low-cost, high-throughput model for cancer imaging, *npj Imaging*, 2023, **1**, 1.
 - 66 Y. Li, C. Schiepers, R. Lake, S. Dadparvar and G. R. Berenji, Clinical utility of ¹⁸F-fluoride PET/CT in benign and malignant bone diseases, *Bone*, 2012, **50**, 128–139.
 - 67 Z. T. Avery, M. G. Gardiner and D. Preston, Using a New Pt(II) Source to Make Pt(II) Lantern-Shaped Cages, Including Low-Symmetry, Heteroleptic, and Multicavity Examples, *Angew. Chem., Int. Ed.*, 2024, e202418079.
 - 68 Ž. D. Bugarčić, J. Bogojeski and R. van Eldik, Kinetics, mechanism and equilibrium studies on the substitution reactions of Pd(II) in reference to Pt(II) complexes with biomolecules, *Coord. Chem. Rev.*, 2015, **292**, 91–106.

