

# ChemComm

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



Journal Name


## Dynamic Foldamer Chemistry

Bryden A. F. Le Bailly<sup>a</sup> and Jonathan Clayden\*<sup>a</sup>Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

[www.rsc.org/](http://www.rsc.org/)

Foldamers can be made more than pieces of static, conformationally uniform molecular architecture by designing into their structure the conformational dynamism characteristic of functional molecular machines. We show that these dynamic foldamers display biomimetic properties reminiscent of allosteric proteins and receptor molecules. They can translate chemical signals into conformational changes, and hence into chemical outputs such as control of reactivity and selectivity. Future developments could see dynamic foldamers operating in the membrane phase providing artificial mechanisms for communication and control that integrate synthetic chemistry into synthetic biology.



<sup>a</sup> School of Chemistry, University of Bristol, Cantock's Close, Bristol BS8 1TS, UK.



## Journal Name

## Synthetic structures and synthetic function

Nature extracts vast diversity of function from a limited collection of molecular structures. The requirement for the structures of functional molecules to be encoded in genetic information in a parsimonious and evolvable way means that the core features of both structure and reactivity in living systems emerge from simple linear sequences of proteins and peptides. The ribosome is the most marvellous molecular structure in the known universe,<sup>1</sup> but it can do only one thing: make polymers of a small collection of  $\alpha$ -amino acids. The control of biosynthetic pathways by enzymes means that even the wider chemistry of other primary and secondary metabolites is but an 'extended phenotype'<sup>2</sup> of protein function (Fig. 1).




Fig 1: Origin of chemical function in (a) nature and (b) designed systems.

Biological function is honed by Darwinian evolution. Chemical ingenuity is not constrained by evolutionary limitations in such a way.<sup>3</sup> In the imagination of a chemist, alternative extended functional molecules can be envisaged built from all kinds of building blocks. But within such huge variety of potential, guiding principles are needed: what molecules to make, and how? For much of the 20<sup>th</sup> century, chemists took the structures of nature as their lead. The chemical synthesis of proteins and peptides is now routine, and the synthetic construction of the molecules of secondary metabolism without direct recourse to the chemistry of life (the 'total synthesis of natural products') reached its zenith in the last decade of the millennium.<sup>4</sup> But in parallel a new approach was developing: the direct synthesis of biomimetic function.<sup>5</sup> Functional mimicry need not imply structural homology, and molecules were designed and synthesised having some of the key conformational characteristics of their natural counterparts, but built from non-biological precursors.

These molecules are *foldamers*<sup>6-10</sup> – extended molecular structures with (in Gellman's definition) 'a strong tendency to adopt a specific compact conformation.'<sup>6</sup>

Foldamer chemistry has been immensely successful in mimicking structural motifs such as helices, sheets, columns, and cavities in the solid state, and in many cases also in solution. Well defined foldamer architectures can imitate biomolecular structures by binding guests<sup>11-14</sup> or biomolecules,<sup>15</sup> or catalysing reactions selectively.<sup>16-18</sup>

Although a classically ideal foldamer adopts a single conformation, we shall contend in this article that an over-emphasis on conformational uniformity may limit chemists' ability to design and synthesise foldamers that mimic some of the wider functions of biomolecules. Most classes of proteins are not fully rigid,<sup>19</sup> and indeed many proteins have segments fully devoid of tertiary structure, or have tertiary structures that become well defined only on binding a ligand.<sup>20,21</sup> In several classic examples,<sup>22</sup> binding of a ligand induces conformational changes that lend a degree of environmental control to the function of a protein. In allosteric proteins such as haemoglobin, the conformational change induced by binding of one oxygen ligand enhances the protein's affinity for others.<sup>22</sup> In membrane bound receptors of the G protein-coupled receptor (GPCR) class, a change in conformation induced by binding to an appropriate ligand leads to modulation of the affinity of the receptor for its partner G protein.<sup>23</sup> In rhodopsin, a switch from *E* to *Z* in the geometry of one double bond of a bound ligand (retinal) stimulates switching between alternative alignments of membrane-spanning helices, also inducing binding of the G protein transducin, and leading ultimately to induction of a nerve signal.<sup>24</sup>

In this review, we describe our response to these aspects of biomolecular function in the design, synthesis and exploitation of foldamers that display dynamic features of conformational switchability and environmental responsiveness. We show how the conceptual range of foldamers may be expanded to include extended molecular structures with a 'strong tendency'<sup>6</sup> to adopt not just one, but multiple favoured conformations, while still retaining structural definition. We shall also review some simple spectroscopic methods that we developed for quantifying that term *strong*. Finally we outline some structural features that have been introduced into dynamic foldamer molecules that allow controlled switching between these alternative structures.

### Dynamic helical structures

One of the most common structural motifs adopted by foldamers (and indeed by natural biomolecules) is the helix.<sup>25-28</sup> This is for good reason: a helix results when a sequence of identical three-dimensional shapes are stacked in such a way

that adjacent members of the stack are all related by the same translational vector (Fig 2).<sup>29</sup> Much of the work that has been carried out in the field of foldamer chemistry has been essentially directed towards ensuring the conformational uniformity implied by this repeated translational vector takes on a physical reality. A helix is a chiral structure, and a pair of left and right-handed (*M* and *P*) conformations are enantiomeric (and therefore of equal energy) if the helix is built from achiral monomers (Fig 2a), but diastereoisomeric (and therefore of unequal energy) if the helix is built from chiral monomers (Fig 2b). Natural biopolymers, and many synthetic foldamers, are built from enantiomerically pure chiral monomers, and hence adopt preferentially one of these two (right-handed, in the case of protein  $\alpha$ -helices and a typical DNA double helix; left-handed, in the case of Z-DNA and collagen)<sup>30-32</sup> diastereoisomeric conformational alternatives.




Fig. 2 (a) A helix built from achiral monomers can adopt either of two enantiomeric screw senses; (b) a helix built from chiral monomers adopts a single screw sense

Although they are far less abundant than helical oligomers of chiral monomers (for reasons that we plan to discuss in detail elsewhere), a number of helical polymers and oligomers built from *achiral* monomers have been described.<sup>7,28</sup> Some of

these (polyisocyanides,<sup>33</sup> and longer polyamides of aromatic amino acids<sup>34</sup> for example) exist as stable, separable left- and right-handed screw-sense enantiomers and are chiral structures, even in solution. Others (aromatic polyureas for example<sup>35</sup>) are helical in the solid state but tend to collapse into random coils in solution.

Importantly, though, a few classes of achiral polymers retain helicity in solution while interconverting rapidly (on a timescale of seconds or less) between their enantiomeric conformers.<sup>36</sup> We call these inverting helices 'configurationally achiral', because even though their global conformation is chiral, they contain no configurationally stable elements of stereochemistry (Fig 3). Configurationally achiral helices include polyisocyanates (Fig 4a),<sup>37</sup> shorter polyamides of aromatic amino acids (Fig 4b),<sup>38</sup> polyamides of the achiral aliphatic amino acids 2-aminoisobutyric acid (Aib) (Fig 4c)<sup>39</sup> and related achiral quaternary  $\alpha$ -amino acids,<sup>40</sup> and polyamides containing alternating Aib and dihydrophenylalanine ( $\Delta$ Phe) residues (Fig 4d).<sup>41</sup> Oligomers of Aib typically adopt a  $3_{10}$  helical geometry<sup>42-44</sup> consisting of repeated  $\beta$ -turns, and Fig 4e and 4f illustrate the three-and-a-half helical turns of the screw-sense labile but reliably helical conformation adopted by an achiral Aib undecamer  $N_3Aib_{11}OCH_2CH_2SiMe_3$ .<sup>45</sup> Related configurationally achiral *supramolecular* helical structures are also known.<sup>27,46-48</sup>




Fig. 3: The pair of rapidly interconverting enantiomeric screw sense conformers of a configurationally achiral helix. Local environment within each conformer is unchanged with respect to displacement along the axis of the helix

## Journal Name




Fig 4: Some oligomers forming configurationally achiral helices: (a) polyisocyanates; (b) oligoamides of aromatic amino acids; (c) oligomers  $Aib_n$ ; (d) oligomers  $(Aib-\Delta Phe)_n$ ; (e) a conformational representation and X-ray crystal structure of the  $3_{10}$  helix formed by  $N_3Aib_{11}OCH_2CH_2SiMe_3$ .<sup>45</sup>

These dynamic helical structures have a valuable property that we may loosely term 'line chirality' (by analogy with the term 'point chirality'). This is to say, within each conformer every point in the helix experiences the same local chiral environment, irrespective of its linear coordinate along the axis of the helix (Fig 3). And, importantly, inversion of helical screw sense in a structure of complete helical fidelity (that is, one in which only uninterrupted helical conformations are populated) likewise inverts the local chiral environment of every point within the helix. A dynamic, screw-sense invertible helix may thus in principle be used as a communication device: conformational uniformity ensures that the adoption of a particular helical screw sense by any point in the helix must be communicated to all other points in the helix.

Circular dichroism (CD) spectroscopy has generally been the technique of choice for the study of helical oligomers in solution: the repeated geometric relationship between chromophores in adjacent monomers gives rise to an identifiable Cotton effect associated with a specific detailed helical geometry.<sup>49</sup> However, for configurationally achiral helices, in which left and right-handed screw senses are equally populated, the Cotton effects cancel and no CD spectrum can be observed. This has limited chemists' ability to

observe directly the solution-state conformations of helices built purely from achiral monomers. The barriers to inversion of screw sense in achiral polyisocyanates<sup>37</sup> ( $\Delta G^\ddagger \approx 90 \text{ kJmol}^{-1}$  at  $100^\circ\text{C}$ , Fig 4a) and oligo-Aib amides<sup>50</sup> ( $\Delta G^\ddagger \approx 46 \text{ kJmol}^{-1}$  at  $-8^\circ\text{C}$ , Fig 4b) were established by dynamic NMR studies.

These and other helical but achiral oligomers (such as oligomers of aromatic achiral amino acids) trivially fail to conform to Gellman's definition<sup>6</sup> of a foldamer: they populate two (enantiomeric) conformations rather than one. This may seem a quibble, but it is a conceptually fecund one, because the equal populations of screw-sense conformers of a configurationally achiral parent oligomer may be usefully perturbed by the introduction of a relatively small chiral conformational influence. In the polyisocyanates, even asymmetric isotopic substitution in a fraction of the monomers (the 'sergeants and soldiers' principle)<sup>51</sup> or attachment of a (photochemically switchable) chiral terminator unit<sup>52</sup> is sufficient to bias the helical conformation towards one screw sense.<sup>53,54</sup> In oligo-Aib structures (Fig 4c),<sup>55-57</sup> and their close relatives,<sup>58,59</sup> including the didehydrophenylalanine-containing oligomers studied by Inai and coworkers (Fig 4d),<sup>60-63</sup> a single chiral terminal or medial residue is sufficient to induce some

degree of preferred screw sense,<sup>57</sup> as is a hydrogen-bonded interaction with a chiral partner.<sup>41,64-67</sup>

Because these dynamic helices can exhibit *helicity* with or without a preferred *screw sense*, we find it productive to distinguish clearly between these two terms. A compound displays 'helicity' if it adopts a helical secondary structure (rather than a linear, zig-zag, or random coil alternative); we use the term 'screw sense' to refer to each enantiomeric conformation of that overall helical secondary structure. Thus 'helicity preference' (helical vs. random coil for example) may be distinguished from 'screw-sense preference' (*M* vs. *P* screw sense): the achiral helices of Fig 4 have a powerful helicity preference but exhibit no screw-sense preference.

### NMR quantifies conformational preference in dynamic foldamers

Detailed NMR studies of the structures of achiral Aib oligomers bearing terminal chiral substituents allowed us to quantify accurately the level of screw-sense control induced in the helix by different chiral terminal residues at either the N<sup>39,68</sup> or the C terminus.<sup>69</sup> Values of screw-sense preference had previously been derived in other oligomers by comparing CD spectra with compounds supposed to have a single enantiomeric preference,<sup>70,71</sup> or by using NMR to measure ratios of diastereoisomeric conformers interconverting slowly on the NMR timescale.<sup>38</sup> Our quantitative NMR approach was made possible by a spectroscopic feature displayed by two 'reporter' groups, A and B, embedded in a helical but configurationally achiral oligomer (Fig 5a). In the chiral environment of the helix, A and B become diastereotopic, experiencing different chemical environments and giving distinct signals in an NMR spectrum (<sup>13</sup>C in the case of A = B = CH<sub>3</sub>; <sup>1</sup>H in the case of A = B = H; <sup>19</sup>F in the case of A = B = CH<sub>2</sub>F or FC<sub>6</sub>H<sub>4</sub>). If the screw sense of the helix inverts slowly on the NMR timescale, the two signals will be resolvable, with chemical shift difference  $\Delta\delta_{\text{slow}}$ . If the helix inverts rapidly on the NMR timescale (typically rates of 0.1-1 × 10<sup>3</sup> Hz) the two signals will coalesce and give rise to a single signal. Fig 5b

shows this effect in the <sup>13</sup>C NMR spectrum of an achiral Aib<sub>9</sub> helix in which the middle residue (Aib\*\*) is <sup>13</sup>C labelled in both of its methyl groups on raising the temperature from 233 K through the coalescence temperature to 273 K.

The situation is subtly different if the two screw-sense conformers of the helical oligomer are in equilibrium but are not equally populated – for example by virtue of a covalently or non-covalently bound chiral influence that affects the screw-sense conformer distribution but does not directly influence the chemical shift of A and B (represented schematically in Fig 5c, and with an example in Fig 5d). In such an event, the slow exchange spectrum will be identical with the equally-populated system, but at fast exchange weighted averaging leads not to a single signal but to two new signals, separated by a new chemical shift difference  $\Delta\delta_{\text{fast}}$ . Fig 5d illustrates the change in line shape of the <sup>13</sup>C NMR signals in oligomers Cbz-L-ValAib<sub>9</sub>Ot-Bu <sup>13</sup>C-labelled in its middle Aib residue, on raising the temperature from 233 K to 273 K.

Because the two screw-sense conformers *P* and *M* contributing to the fast exchange spectrum are present in relative populations of 1 :  $K_{\text{eq}}$ , the ratio

$$\Delta\delta_{\text{fast}} / \Delta\delta_{\text{slow}} = (K_{\text{eq}} - 1) / (K_{\text{eq}} + 1)$$

or

$$\Delta\delta_{\text{fast}} / \Delta\delta_{\text{slow}} = ([M]-[P]) / ([M]+[P])$$

Given its similarity with the formula for enantiomeric excess (e.e.), we call this value  $([M]-[P]) / ([M]+[P])$  'helical excess' (h.e.). We have used the method with <sup>1</sup>H-containing,<sup>68,72</sup> <sup>13</sup>C-containing<sup>30,39,69,73</sup> and <sup>19</sup>F-containing<sup>74</sup> reporters, and the quantitative interpretation of  $\Delta\delta_{\text{fast}} / \Delta\delta_{\text{slow}}$  as a measure of screw-sense preference has been validated by comparison with data obtained by line shape analysis,<sup>39</sup> from the linear relationship between  $\Delta\delta_{\text{fast}} / \Delta\delta_{\text{slow}}$  using different NMR reporters<sup>74</sup> and also between  $\Delta\delta_{\text{fast}} / \Delta\delta_{\text{slow}}$  and molar ellipticity in CD spectra.<sup>72</sup>




Figure 5: Detection of screw-sense preference by NMR in the fast exchange regime. (a) and (b) Coalescence between signals in an equally populated pair of enantiomeric conformers; (c) and (d) Coalescence between signals in an unequally populated pair of diastereoisomeric conformers.

Table 1: Screw-sense preference induced in a helical chain of achiral Aib residues bearing a chiral residue at the (a) N terminus<sup>68</sup> or (b) the C terminus,<sup>69,75</sup> detected at ambient temperatures in methanol by NMR reporters A and B (A, B =  $^1\text{H}$  or  $^{13}\text{C}$ ). Helical excess (h.e.) is negative in a left-handed helix and positive in a right-handed helix. Values have been corrected for decay of conformational preference between the chiral inducer and the NMR spectroscopic reporter, as outlined in references<sup>68</sup> and <sup>76</sup>.


| (a) Varying the N terminus of $\text{Cbz-Xxx-Aib}_n\text{-R(AB)}$ . |      | (b) Varying the C terminus of $\text{Cbz-R(AB)-Aib}_n\text{-Yyy-NHt-Bu}$ . |      |
|---|------|--|------|
| Xxx =   | h.e. | Yyy =  | h.e. |
|   | 4    |  | +98  |
|   | -20  |  | +94  |
|   | -33  |  | +92  |
|   | -35  |  | +85  |
|   | -42  |  | +26  |
|   | -44  |  | +21  |

|  |     |  |      |
|--|-----|--|------|
| Cbz-L-Pro                                    | -45 | L- $\alpha$ -methylvaline-NHt-Bu                 | +92  |
| Cbz-L-Phe                                    | -52 | (L- $\alpha$ -methylvaline) <sub>2</sub> -NHt-Bu | +100 |
| Cbz-L-isovaline                              | +25 | L-Ala-Ot-Bu                                      | -70  |
| Cbz-L- $\alpha$ -methylphenylalanine         | +56 | L-Phe-Ot-Bu                                      | -44  |
| Cbz-L- $\alpha$ -methylvaline                | +68 | L-Val-Ot-Bu                                      | -60  |
| Cbz-(L- $\alpha$ -methylvaline) <sub>2</sub> | +95 | L- <i>tert</i> -Leu-Ot-Bu                        | -46  |
|  |     | L- $\alpha$ -methylvaline-Ot-Bu                  | -56  |

A drawback of this NMR method using a geminal pair of reporter groups in an achiral residue is that it cannot report on whether an *M* or a *P* helix is preferred—in other words, the sign of the screw-sense preference remains undetermined. However, CD studies correlated with data from the enantiomerically enriched, isotopically labelled <sup>13</sup>C NMR probe discussed below<sup>30</sup> have allowed the major screw sense to be assigned with confidence in most cases. Some values for the screw-sense preference induced by covalently linked residues at the N or C terminus are summarised in Table 1.

### Conformational persistence and signal decay

The values for helical excess (h.e.) obtained from these studies can be considered comparable in meaning to the CD-derived screw-sense excess (s.e.) values reported by Sugimoto.<sup>70,71</sup> However, unlike CD, NMR reports the conformational preference at a single site in the oligomer chain.<sup>77</sup> For this reason, it can be used not only to quantify global conformational preferences, but also to assess the variation of conformational preference within a foldamer chain. By synthesising a series of oligomers carrying <sup>13</sup>C labels at each position in a chain of Aib residues,<sup>78</sup> we were thus able to observe the way that an initial screw-sense preference induced locally by a chiral residue decays exponentially (but remarkably slowly) as the chiral influence becomes more distant (Fig. 6).<sup>76</sup>



(b)




Fig 6. The screw-sense preference exhibited by an Aib oligomer (as reported by chemical shift difference  $\Delta\delta$ ) decays as a terminal chiral influence becomes more distant. (a) Chemical shift differences were measured in a series of three 'frame-shifted' oligomers each containing three pairs of abundance-labelled (100%, 50% or 25%) <sup>13</sup>C labels. (b) Decay is more rapid in polar solvents and at higher temperature (dashed lines 0 °C; solid lines 23 °C; dotted lines 40 °C).

The spatial decay constant is at a minimum in non-polar solvents, but is increased by solvents with more powerful dipoles or hydrogen-bonding properties. Quantitative comparisons can be made: in THF, at 23 °C, there is only a 0.5% drop in helical excess at the chain terminus each time the chain is extended by a further Aib residue, while the equivalent value in MeOH is 6.1%. A separate study<sup>79</sup> measured the ability of a range of alternative achiral amino acids to 'communicate' a screw-sense preference between two helical domains and found that in MeOH quaternary amino acids communicated a conformational preference highly effectively (>90% fidelity), while a single Gly or  $\beta$ -alanine led to loss of about 50% of the conformational signal. More bulky achiral quaternary amino acids (e.g. 1,1-diphenylglycine) disrupted the helical conformation and prevented communication of helical screw sense. Similar NMR measurements should be possible in other achiral helical foldamers containing paired enantiotopic signals, provided NMR spectra can be measured in the fast exchange régime.

This decay in screw-sense preference must arise from the intrusion of occasional non-helical conformational features into the overall helical structure of the molecule – an indication of just how 'strong' its 'strong preference'<sup>6</sup> for

helicity is. These conformational features are presumably localised, rare helix reversals that occur transiently. Their detailed structure is of interest because they offer an insight into the mechanisms of screw-sense reversal in helical foldamers, and are consequently the subject of further studies currently under way.<sup>80</sup>

Nature apparently also makes use of the conformational properties of dynamic foldamers which are reliably helical but in which screw sense is poorly controlled. Part of the structural inspiration for our work in this area was the observation that the fungal metabolites known as peptaibiotics<sup>81</sup> apparently adopt well defined screw senses in the solid state, despite the fact that parts of their structure contain relatively few chiral residues embedded in a largely achiral chain. The solid state structure of cephaibol C (Fig 7) and cephaibol A contain at their N terminus a stretch of right-handed  $3_{10}$  helix in which only the N-terminal Phe residue and the Leu at residue 7 are chiral.<sup>82</sup> During our work on conformational preferences in structures such as those illustrated in Table 1, we noted that an N-terminal L amino acid adjacent to a sequence of Aib residues does not typically induce a right-handed screw sense,<sup>30,83</sup> and in a sequence Ac-L-Phe-Aib<sub>4</sub>-L-Leu... the L-Phe and L-Leu residues are expected to be mismatched in their conformational properties, inducing opposite screw senses and disfavoured the adoption of a uniform conformation. To test this hypothesis, we made four 'cephaibologues'—diastereoisomers of the N-terminal nonapeptide of the cephaibols (Fig 7). We found that indeed the diastereoisomers A and D that correspond most closely to the relative configuration of the natural structures are less conformationally uniform than the unnatural diastereoisomers.<sup>84</sup> The *Trichoderma* fungi producing the cephaibols seem to have evolved to produce compounds which have dynamic conformational properties, possibly because these properties enhance the membrane activity required for them to function as antibiotics.<sup>45</sup>




Fig 7: (a) The structure of cephaibol C, and (b) the CD spectra of four diastereoisomeric 'cephaibologues' corresponding to its N-terminal nonapeptide sequence. The most conformationally well defined (B and C) are diastereoisomers of the less well defined structures A and D whose structures match most closely the relative configuration of cephaibol C itself.

### Dynamic screw sense switching detected by <sup>13</sup>C NMR

In the case of <sup>13</sup>C labelled Aib foldamers, an additional piece of conformational information may be gained by the incorporation of an isotopic <sup>13</sup>C label enantioselectively into one of the two geminal methyl groups of an Aib residue.<sup>85</sup> The pro-*R* methyl group of an Aib residue embedded in a right-

handed helix appears reliably at higher field than the pro-*S* residue<sup>86</sup> (though the locations are inverted in an esterified C-terminal Aib residue). Thus, the major <sup>13</sup>C signal from a single <sup>13</sup>C-labelled residue *R*-Aib\* in a right-handed helix will appear as the upfield member of the diastereotopic pair (so, memorably, *R* @ *P* = 'on the right'). While the chemical shift separation of the two signals reports the magnitude of the local screw-sense preference of the oligomer, the location of the major <sup>13</sup>C NMR signal in the upfield or downfield signal therefore indicates the left or right-handed chirality of the screw-sense preference at that point in the oligomer.<sup>30</sup>

Thus the starting chiral oligomer shown in Fig 8, which is preferentially labelled with <sup>13</sup>C in the pro-*R* methyl group of its C-terminal Aib\* residue, displays in its <sup>13</sup>C NMR spectrum a pair of signals separated by 0.75 ppm. This indicates that it adopts a screw-sense preference of about 33% h.e. In addition the major peak is upfield of the minor, confirming that the helix is an interconverting mixture of which the predominant conformer is left handed.<sup>87</sup> The ester-bearing stereogenic centre is however also invertible: methanolysis to the alcohol and treatment under Mitsunobu conditions gives its enantiomer. The <sup>13</sup>C NMR spectrum clearly indicates that a local inversion of conformational preference has taken place at the C terminus of the oligomer. Inversion of the configuration of the single controlling centre in the oligomer leads necessarily to inversion of the screw-sense preference of the equilibrating mixture, and information about this stereochemical inversion is transmitted through the induced conformational preference in the helix.




Fig 8. The stereochemical consequence of inversion of configuration at an ester-bearing centre is detected by a remote <sup>13</sup>C NMR reporter of conformational preference. The two diastereotopic methyl groups of the C-terminal labelled Aib residue are differentially enriched in <sup>13</sup>C (pro-*R* in red 75% <sup>13</sup>C; pro-*S* in green 25% <sup>13</sup>C). Inversion of screw-sense conformation that results from a remote inversion of N-terminal configuration exchanges their positions in the <sup>13</sup>C NMR spectrum. Note that in this case, the major signal is

(unusually) upfield in an *M* helix because the *R*-Aib\* residue is a C-terminal ester.

A similar dynamic inversion of screw-sense preference can be achieved by exploiting the fact that the functionality at the C terminus of an Aib oligomer affects local secondary structure in such a way that the presence or absence of a single hydrogen bond can lead to a reversal of screw-sense preference. This feature, first observed in the tendency of C-terminal esters to exhibit reversed screw-sense preference relative to C-terminal secondary amides<sup>69</sup> (See Table 1), was exploited in a pH-dependent switch<sup>88</sup> in which the removal of a hydrogen bond by base-promoted deprotonation led to inversion of screw-sense preference. The C-terminal NH group in an Aib oligomer was selectively acidified by amidation with a tren ligand and complexation with a Lewis acidic Zn<sup>2+</sup> cation (Figure 9a, b).<sup>89</sup> Sequential addition of OH<sup>-</sup> (Fig 9c, e) and H<sup>+</sup> (Fig 9 d, f) ions caused repeated switching of the foldamer between a predominantly right-handed and a predominantly left-handed conformation.




Fig 9 (a) A C terminal tren-Zn moiety acidifies an adjacent NH group (orange) in a dynamic foldamer allowing it to be selectively deprotonated and reprotonated by base and acid. The two diastereotopic methyl groups of the N-terminal labelled Aib residue are differentially enriched in  $^{13}\text{C}$  (pro-*R* in red 70%  $^{13}\text{C}$ ; pro-*S* in green 30%  $^{13}\text{C}$ ). (b)-(e)  $^{13}\text{C}$  NMR spectra illustrating (b) the right handed screw sense of the original foldamer; (c) the left handed screw sense induced on deprotonation by base; (d) the right handed screw sense restored by addition of acid; (e) and (f) further screw sense inversions during a second cycle of base/acid treatment.

## A dynamic foldamer as a biomimetic artificial receptor

In a typical biological receptor, a conformational change results from the reversible binding of a ligand with a binding site.<sup>90</sup> An artificial mimic of this function would provide an important link between synthetic chemistry and synthetic biology, and offer the possibility of designing complex artificial molecular communication systems. In our early investigations, we chose to ensure an intimate connection between ligand and receptor by using reversible (dynamic) covalent chemistry. Boronic acids are well known to form, reversibly, cyclic boronate esters by condensation with diols.<sup>91</sup> A series of Aib\*-labelled oligomers were made carrying an amino-boronate function at their N terminus (Figure 10a).<sup>92</sup> In the absence of any diol ligand,  $^{11}\text{B}$  NMR indicated that in deuterated methanol the receptor mimic adopted a cyclic azaborolidine structure at the N terminus, and  $^{13}\text{C}$  NMR reported, as expected, no screw-sense preference in the achiral foldamer. Adding a chiral diol ligand led to spontaneous formation of the cyclic boronate ester, whose combined steric bulk and Lewis acidity promoted the formation of the methanol-bridged boronate structure. Additionally, with a chiral ligand such as (–)-diisopropyl tartrate (DIPT) a preferred left-handed screw sense was induced in the Aib oligomer that was detectable in the  $^{13}\text{C}$  NMR signals of the remote enantioselectively labelled Aib\* residue (Fig 10 c). By contrast, an achiral ligand (the meso diastereoisomer of a diol for example) was unable to induce a screw-sense preference.

Just as ligands might vary in their binding strength at a receptor binding site, different diols displayed different equilibrium constants for formation of boronate esters under the conditions of the experiments. The characteristic binding strength of different ligands could hence be exploited in a series of experiments in which sequential ligands compete for the same binding site. After adding 2 equiv. of (–)-DIPT ( $K = 300 \text{ M}^{-1}$ ) to the receptor (which induced a predominantly left-handed screw sense) the addition of 2 equiv. of the more strongly binding (+)-hydrobenzoin ( $K = 1500 \text{ M}^{-1}$ ) led to the displacement of DIPT from the binding site and the consequent inversion of the screw sense in the receptor to a right-handed preference, indicated by an exchange in position of the major and minor peaks in the  $^{13}\text{C}$  NMR spectrum (Fig 10d). (+)-Hydrobenzoin thus acts as a *competitive antagonist*

of (–)-DIPT. Even stronger binding was exhibited by (–)-pinanediol ( $K \gg 2000 \text{ M}^{-1}$ ). Although chiral, pinanediol induced no screw-sense preference, probably because its greater steric hindrance prevents any interaction of the aminoboronate N atom with B. Pinanediol is an irreversible inhibitor of the artificial receptor. The further biomimetic credentials of the boronate-based receptor were evident in its ability to respond even to biological messengers such as adenosine ( $K \approx 625 \text{ M}^{-1}$ ) and guanosine ( $K \approx 600 \text{ M}^{-1}$ ), which both induced a left-handed screw sense, and uridine and cytidine ( $K > 2000 \text{ M}^{-1}$ ) the last of which induced a right-handed screw sense.





Fig 10. (a) A dynamic foldamer carrying an N-terminal boronate ester bonding site (purple) and a C-terminal labelled Aib residue are differentially enriched in  $^{13}\text{C}$  (pro-*R* in red 75%  $^{13}\text{C}$ ; pro-*S* in green 25%  $^{13}\text{C}$ ). (b) A cartoon of the receptor in its resting state. (c) Reversible esterification with a weakly binding ligand (–)-diisopropyl tartrate (DIPT, blue),  $K = 300 \text{ M}^{-1}$ , induces a left-handed screw-sense preference. (d) A more strongly binding ligand (+)-hydrobenzoin (red),  $K = 1500 \text{ M}^{-1}$ , acts a competitive antagonist, displacing the (blue) DIPT from

the binding site and inducing a right-handed screw-sense preference. (e) An even more strongly binding ligand, pinanediol (orange) with  $K \gg 2000 \text{ M}^{-1}$  acts as an irreversible inhibitor, displacing all other ligands, and turning the screw-sense preference off.

### Dynamic modulation of foldamer conformation by competitive noncovalent binding in a multicomponent system


Despite the biomimetic character of the reversible and selective ligand binding at this 'purinergic' receptor, it nonetheless employs non-biomimetic reversible covalent bonding. Typical biological receptors bind ligands through non-covalent interactions. Inai has shown that achiral oligomers carrying basic binding sites exhibited a conformational response to binding whose strength depended on the nature of the hydrogen bonding between the ligand and the receptor.<sup>41,64-66</sup> Advancing this concept, we sought a basic binding site that displayed selective 1:1 binding interactions with acidic ligands, and found that the 2-pyridylacetamide-terminated oligomers illustrated in Fig 11 showed this property in the presence of either carboxylic or phosphoric acid ligands (Fig 11a).<sup>93</sup> Adding a chiral phosphoric acid to the foldamer induced a switch to a preferred screw-sense preference, evident in the induced  $^{13}\text{C}$  NMR chemical shift difference in the C-terminal Aib\*\* labels (Fig 11b). Adding ammonia to the mixture disrupted the interaction with the chiral acid, turning off the screw-sense preference. Adding hydrochloric acid restored the effect of the chiral ligand (Fig 11b). Complete transfer of a proton from the acid to the basic binding site to form ion pairs in chloroform is unlikely, so the reversibility of the influence of the chiral acid ligand on the receptor carrying the basic binding site is likely to be due to competitive hydrogen-bonding interactions of different strengths between the various acidic and basic species in solution.



**Figure 11:** (a) A  $^{13}\text{C}$ -labelled dynamic foldamer bearing an N-terminal basic binding site, and a chiral phosphoric acid as a ligand. (b) The hydrogen-bonded complex between the phosphoric acid and the foldamer induces a preferred screw-sense preference that may be turned off and on by addition of ammonia and HCl.

By choosing acids with identifiably characteristic effects on the conformation of the receptor (strongly left-handed, strongly right-handed, or weakly left-handed) we found that we could force a receptor's conformation to cycle between these different preferences either by adding increasingly strong acids (a carboxylic acid, a phosphoric acid, or a triflamide) to the receptor (Fig 12).<sup>93</sup> Even more intriguingly, a mixture of acids of different strengths could also be selectively silenced or activated by addition of increasing amounts of base, or reactivated by acid. Fig 12 shows how a four-component mixture of receptor and three acids behaves as a chemical system capable of 'counting' the number of protons available to it and reporting an output (a left- or right-handed screw-sense preference) as a result.

The uncomplexed receptor (Fig 12a) responds to the addition of a chiral carboxylic acid (in blue) by adopting a left-handed screw-sense preference (Fig 12b). Adding a stronger acid, the phosphoric acid shown in orange, displaces the carboxylic acid from the binding site and induces a switch in screw-sense preference (Fig 12c). An even stronger acid still, the triflamide shown in red, further displaces the phosphoric acid and results in a second inversion to a weakly left-handed screw-sense (Fig 12d).



**Figure 12:** (a) A dynamic foldamer carrying a  $^{13}\text{C}$  NMR reporter of conformational preference and functionalised with a basic pyridine binding site. (b) The dynamic foldamer forms a hydrogen-bonded complex in which a chiral carboxylic acid (blue) induces a conformational preference for left-handed screw sense. (c) A chiral phosphoric acid (orange) induces a right-handed screw sense. (d) A chiral triflamide (red) restores a weak preference for left-handed screw sense. (e) Ammonia disrupts the hydrogen-bonded complex firstly of the most acidic triflamide ligand, then (f) of the phosphoric acid ligand and then finally (g) of the carboxylic acid ligand, returning the receptor to its native form.

Now, addition of ammonia provides an alternative partner for the most acidic proton available, and thus disrupts the hydrogen bonded complex between the basic foldamer and the most acidic (red) triflamide, leaving the phosphoric acid (orange) free to reassociate with the dynamic foldamer, introducing again a right-handed screw-sense preference (Fig 12e). A second equivalent of ammonia likewise removes the

proton from the phosphoric acid-foldamer complex, and allows the carboxylic acid to restore a preference for left-handed screw sense (Fig 12f). Finally a third equivalent of ammonia prevents association of any of the three acids with the dynamic foldamer, which thus reverts to its conformationally racemic native form (Fig 12g). The final three changes can each be reversed by readdition of HCl, which returns a proton to the system and neutralising the effect of the ammonia. The ability of the dynamic foldamer to report on the relative strengths of the different possible interactions between the components of complex mixture has analogies with the network of competitive interactions experienced by proteins in a cell. Future developments could lie in the direction of complex artificial control networks based on a type of molecular computation in which information is encoded in conformation.<sup>94</sup>

### Photoswitching foldamer conformation

Closely related in structure and function to G protein-coupled receptors are the opsins – the proteins of vision. Human vision is mediated by rhodopsin, a modified G protein-coupled receptor that carries a covalently attached ligand in the form of retinal. Photochemical switching of retinal between the *E* and *Z* configurations at one particular double bond leads to a conformational change that is communicated through the molecule and ultimately leads to the propagation of the nerve signal. Feringa has shown that a photochemical switch can lead to inversion of helical screw sense of polymeric polyisocyanate,<sup>52</sup> although because detection of screw-sense preference used circular dichroism, no data was obtained on how far the information encoded in the conformational switch is communicated. We showed that photochemical stimulation may lead to an inversion of screw sense that propagates through the helical oligomer and consequently inverts the sense of asymmetry induced by a catalytic thiourea unit<sup>95</sup> located at the remote terminus (Fig 13).<sup>89</sup> The photochemical inversion made use of the same concept that had been applied to the directed deprotonation of the oligomer shown in Fig 9. The starting tertiary amide contains no C terminal NH group, and as a result the C-terminal L-Ala residue (in blue) induces the oligomer to adopt a global left-handed helical structure, placing the locally achiral thiourea catalytic site in a chiral environment and allowing it to induce asymmetry in the catalytic addition of malonate to nitrostyrene to give an *S* product in 37:63 e.r. The transamidation of the (green) tertiary amide to an (orange) secondary amide by photochemical cleavage of the 5-bromo-8-nitroindolinamide<sup>96</sup> in the presence of isopropylamine allows a new hydrogen bond to form, reorganizing the terminal portion of the helix into a right-handed screw-sense preference. This resulting screw-sense switched foldamer also catalyses the Michael addition of malonate to nitrostyrene, with the sense of selectivity reversed, giving the *R* product in 77:23 e.r.




Fig 13. A photolabile tertiary amide function allows global screw-sense inversion of a dynamic foldamer to reverse the enantioselectivity of a remote catalytic site.

## Conclusions

Compounds that have a well defined but dynamic conformational structure offer opportunities to use synthetic scaffolds to mimic the features of natural molecules. The principal feature that divides biology from chemistry is information: no artificial chemical system has ever come even close to the informational complexity of even the simplest cell.<sup>97</sup> Nonetheless, dynamic foldamers allow spatial communication of information over distances that extend into the multi-nanometre range, in principle allowing communication across cell membranes (whose typical thickness is in the region of 4 nm). Future extensions of this work could allow artificial receptors to be built that function in a membrane environment. Such dynamic foldamers, potentially incorporated into cell membranes, would allow the construction of synthetic, controllable cellular compartments, providing a bridge between synthetic chemistry and synthetic biology. Dynamic foldamers also display the selective responses to competing interactions in a mixture of components, and further developments in this area could see information not only communicated by foldamers but processed by them as well.<sup>94</sup>

## Acknowledgements

Our work in this area has been supported by the EPSRC, the BBSRC and the ERC.

## Notes and references

1 A. Yonath, *Angew. Chem. Int. Ed.*, 2010, **49**, 4340–4354.

- 2 R. Dawkins, *The extended phenotype*, OUP (Oxford) 1982.
- 3 M. Berthelot, *La synthèse chimique*, Germer Baillière, Paris 1876.
- 4 K. C. Nicolaou and S. A. Snyder, *Proc. Natl. Acad. Sci.*, 2004, **101**, 11929–11936.
- 5 G. M. Whitesides, *Interface Focus*, 2015, **5**, 20150031–20150031.
- 6 S. H. Gellman, *Acc. Chem. Res.*, 1998, **31**, 173–180.
- 7 D. J. Hill, M. J. Mio, R. B. Prince, T. S. Hughes and J. S. Moore, *Chem. Rev.*, 2001, **101**, 3893–4012.
- 8 I. Huc, *Eur. J. Org. Chem.*, 2004, **2004**, 17–29.
- 9 S. Hecht and I. Huc, *Foldamers: Structure, Properties and Applications*, Wiley-VCH, Weinheim, 2007.
- 10 I. Saraogi and A. D. Hamilton, *Chem. Soc. Rev.*, 2009, **38**, 1726–1743.
- 11 H. Juwarker, J.-M. Suk and K.-S. Jeong, *Chem. Soc. Rev.*, 2009, **38**, 3316–3325.
- 12 R. B. Prince, S. A. Barnes and J. S. Moore, *J. Am. Chem. Soc.*, 2000, **122**, 2758–2762.
- 13 D.-W. Zhang, X. Zhao and Z.-T. Li, *Acc. Chem. Res.*, 2014, **47**, 1961–1970.
- 14 Q. Gan, Y. Ferrand, C. Bao, B. Kauffmann, A. Grélard, H. Jiang and I. Huc, *Science*, 2011, **331**, 1172–1175.
- 15 V. Azzarito, K. Long, N. S. Murphy and A. J. Wilson, *Nature Chem.*, 2013, **5**, 161–173.
- 16 M. Reggelin, S. Doerr, M. Klussmann, M. Schultz and M. Holbach, *Proc. Natl. Acad. Sci.*, 2004, **101**, 5461–5466.
- 17 R. A. Smaldone and J. S. Moore, *J. Am. Chem. Soc.*, 2007, **129**, 5444–5450.
- 18 M. M. Müller, M. A. Windsor, W. C. Pomerantz, S. H. Gellman and D. Hilvert, *Angew. Chem.*, 2009, **121**, 940–943.
- 19 C. A. Smith, D. Ban, S. Pratihari, K. Giller, C. Schwiegk, B. L. de Groot, S. Becker, C. Griesinger and D. Lee, *Angew. Chem. Int. Ed.*, 2015, **54**, 207–210.
- 20 J. Habchi, P. Tompa, S. Longhi and V. N. Uversky, *Chem. Rev.*, 2014, **114**, 6561–6588.
- 21 M. Fuxreiter, Á. Tóth-Petróczy, D. A. Kraut, A. T. Matouschek, R. Y. H. Lim, B. Xue, L. Kurgan and V. N. Uversky, *Chem. Rev.*, 2014, **114**, 6806–6843.
- 22 M. F. Perutz, *Q. Rev. Biophys.*, 1989, **22**, 139–236.
- 23 S. Granier and B. Kobilka, *Nature Chemical Biology*, 2012, **8**, 670–673.
- 24 A. Manglik and B. Kobilka, *Curr. Opin. Cell Biol.*, 2014, **27**, 136–143.
- 25 T. Nakano and Y. Okamoto, *Chem. Rev.*, 2001, **101**, 4013–4038.
- 26 J. J. L. M. Cornelissen, A. E. Rowan, R. J. M. Nolte and N. A. J. M. Sommerdijk, *Chem. Rev.*, 2001, **101**, 4039–4070.
- 27 D. Pijper and B. L. Feringa, *Soft Matter*, 2008, **4**, 1349–1372.
- 28 E. Yashima, K. Maeda, H. Iida, Y. Furusho and K. Nagai, *Chem. Rev.*, 2009, **109**, 6102–6211.
- 29 N. Chouaieb, A. Goriely and J. H. Maddocks, *Proc. Natl. Acad. Sci.*, 2006, **103**, 9398–9403.
- 30 R. A. Brown, T. Marcelli, M. De Poli, J. Solà and J. Clayden, *Angew. Chem. Int. Ed.*, 2012, **51**, 1395–1399.
- 31 J. D. Dunitz, *Angew. Chem. Int. Ed.*, 2001, **40**, 4167–4173.
- 32 B. Brodsky and A. V. Persikov, in *Fibrous Proteins: Coiled-Coils, Collagen and Elastomers*, Elsevier, Amsterdam, 2005, vol. 70, pp. 301–339.
- 33 M. Ishikawa, K. Maeda, Y. Mitsutsuji and E. Yashima, *J. Am.*

- Chem. Soc.*, 2004, **126**, 732–733.
- 34 T. Qi, V. Maurizot, H. Noguchi, T. Charoenraks, B. Kauffmann, M. Takafuji, H. Ihara and I. Huc, *Chem. Commun.*, 2012, **48**, 6337.
- 35 J. Clayden, L. Lemiègre, G. A. Morris, M. Pickworth, T. J. Snape and L. H. Jones, *J. Am. Chem. Soc.*, 2008, **130**, 15193–15202.
- 36 K. Maeda and E. Yashima, in *Supramolecular Chirality*, Springer-Verlag, Berlin/Heidelberg, 2006, vol. 265, pp. 47–88.
- 37 K. Ute, Y. Fukunishi, S. K. Jha, K. S. Cheon, B. Muñoz, K. Hatada and M. M. Green, *Macromolecules*, 1999, **32**, 1304–1307.
- 38 C. Dolain, H. Jiang, J.-M. Léger, P. Guionneau and I. Huc, *J. Am. Chem. Soc.*, 2005, **127**, 12943–12951.
- 39 J. Solà, G. A. Morris and J. Clayden, *J. Am. Chem. Soc.*, 2011, **133**, 3712–3715.
- 40 C. Toniolo, M. Crisma, F. Formaggio and C. Peggion, *Biopolymers*, 2001, **60**, 396–419.
- 41 Y. Inai, K. Tagawa, A. Takasu, T. Hirabayashi, T. Oshikawa and M. Yamashita, *J. Am. Chem. Soc.*, 2000, **122**, 11731–11732.
- 42 B. V. Prasad and P. Balaram, *CRC Crit. Rev. Biochem.*, 1984, **16**, 307–348.
- 43 C. Toniolo, G. M. Bonora, V. Barone, A. Bavoso, E. Benedetti, B. Di Blasio, P. Grimaldi, F. Lejl, V. Pavone and C. Pedone, *Macromolecules*, 1985, **18**, 895–902.
- 44 C. Toniolo and E. Benedetti, *Trends Biochem. Sci.*, 1991, **16**, 350–353.
- 45 J. E. Jones, V. Diemer, C. Adam, J. Raftery, R. E. Ruscoe, J. T. Sengel, M. I. Wallace, A. Bader, S. L. Cockroft, J. Clayden and S. J. Webb, *J. Am. Chem. Soc.*, 2016, jacs.5b12057.
- 46 W. Zhang, W. Jin, T. Fukushima, T. Mori and T. Aida, *J. Am. Chem. Soc.*, 2015, **137**, 13792–13795.
- 47 S. Azeroual, J. Surprenant, T. D. Lazzara, M. Kocun, Y. Tao, L. A. Cuccia and J.-M. Lehn, *Chemical Communications*, 2012, **48**, 2292–2294.
- 48 F. Dumitru, Y.-M. Legrand, A. Van der Lee and M. Barboiu, *Chemical Communications*, 2009, **0**, 2667–2669.
- 49 Circular Dichroism: Principles and Applications, ed. K. Nakanishi, N. Berova and R. W. Woody, Wiley-VCH, New York, 2nd ed., 2000; Circular Dichroism and the Conformational Analysis of Biomolecules, ed. G. D. Fasman, Plenum Press, New York, 1996.
- 50 R. P. Hummel, C. Toniolo and G. Jung, *Angew. Chem. Int. Ed.*, 1987, **26**, 1150–1152.
- 51 M. M. Green, N. C. Peterson, T. Sato, A. Teramoto, R. Cook and S. Lifson, *Science*, 1995, **268**, 1860–1866.
- 52 D. Pijper and B. L. Feringa, *Angew. Chem. Int. Ed. Engl.*, 2007, **46**, 3693–3696.
- 53 E. Yashima, K. Maeda and T. Nishimura, *Chem. Eur. J.*, 2004, **10**, 42–51.
- 54 E. Yashima and K. Maeda, *Macromolecules*, 2008, **41**, 3–12.
- 55 B. Pengo, F. Formaggio, M. Crisma, C. Toniolo, G. M. Bonora, Q. B. Broxterman, J. Kamphuis, M. Saviano, R. Iacovino, F. Rossi and E. Benedetti, *J. Chem. Soc., Perkin Trans. 2*, 1998, 1651–1657.
- 56 J. Clayden, A. Castellanos, J. Solà and G. A. Morris, *Angew. Chem. Int. Ed.*, 2009, **48**, 5962–5965.
- 57 M. Crisma, M. De Zotti, F. Formaggio, C. Peggion, A. Moretto and C. Toniolo, *J. Pept. Sci.*, 2015, **21**, 148–177.
- 58 N. Ousaka, Y. Takeyama and E. Yashima, *Chem. Eur. J.*, 2013, **19**, 4680–4685.
- 59 N. Ousaka, Y. Takeyama, H. Iida and E. Yashima, *Nature Chem.*, 2011, **3**, 856–861.
- 60 Y. Inai, Y. Kurokawa, A. Ida and T. Hirabayashi, *Bull. Chem. Soc. Jpn.*, 1999, **72**, 55–61.
- 61 Y. Inai, S. Ashitaka and T. Hirabayashi, *Polym J*, 1999, **31**, 246–253.
- 62 Y. Inai, Y. Kurokawa and T. Hirabayashi, *Biopolymers*, 1999, **49**, 551–564.
- 63 Y. Inai, Y. Ishida, K. Tagawa, A. Takasu and T. Hirabayashi, *J. Am. Chem. Soc.*, 2002, **124**, 2466–2473.
- 64 Y. Inai, N. Ousaka and T. Okabe, *J. Am. Chem. Soc.*, 2003, **125**, 8151–8162.
- 65 H. Komori and Y. Inai, *J. Org. Chem.*, 2007, **72**, 4012–4022.
- 66 N. Ousaka and Y. Inai, *J. Org. Chem.*, 2009, **74**, 1429–1439.
- 67 For other examples in which small changes of changes of foldamer structure or solvent lead to significant reorganisations of conformational preference, see: F. Mamiya, N. Ousaka and E. Yashima, *Angew. Chem.*, 2015, **54**, 14442–14446; F. Formaggio, G. Ballano, A. Moretto, C. Peggion, M. Crisma, R. Lettieri, E. Gatto, N. Venanzi and C. Toniolo, *Biopolymers*, 2011, **96**, 436–436; R. Lettieri, M. Bischetti, E. Gatto, A. Palleschi, E. Ricci, F. Formaggio, M. Crisma, C. Toniolo and M. Venanzi, *Biopolymers*, 2013, **100**, 51–63.
- 68 M. De Poli, L. Byrne, R. A. Brown, J. Solà, A. Castellanos, T. Boddaert, R. Wechsel, J. D. Beadle and J. Clayden, *J. Org. Chem.*, 2014, **79**, 4659–4675.
- 69 B. A. F. Le Bailly and J. Clayden, *Chem. Commun.*, 2014, **50**, 7949–7952.
- 70 Y. Ito, T. Miyake, S. Hatano, R. Shima, T. Ohara and M. Suginome, *J. Am. Chem. Soc.*, 1998, **120**, 11880–11893.
- 71 M. Suginome, S. Collet and Y. Ito, *Org. Lett.*, 2002, **4**, 351–354.
- 72 M. De Poli and J. Clayden, *Org. Biomol. Chem.*, 2014, **12**, 836–843.
- 73 L. Byrne, J. Solà, T. Boddaert, T. Marcelli, R. W. Adams, G. A. Morris and J. Clayden, *Angew. Chem. Int. Ed.*, 2014, **53**, 151–155.
- 74 S. J. Pike, M. De Poli, W. Zawodny, J. Raftery, S. J. Webb and J. Clayden, *Org. Biomol. Chem.*, 2013, **11**, 3168–3176.
- 75 B. A. F. Le Bailly, PhD thesis, University of Manchester 2015.
- 76 B. A. F. Le Bailly, L. Byrne, V. Diemer, M. Foroozandeh, G. A. Morris and J. Clayden, *Chem. Sci.*, 2015, **6**, 2313–2322.
- 77 Inai has nonetheless made use of chromophores with distinctive Cotton effects to establish the preferred screw sense of a localised part of an extended oligomer: see ref. 66.
- 78 In order to limit the number of compounds we needed to make, we incorporated three doubly <sup>13</sup>C-labelled Aib residues into each oligomer, and identified the individual labelled residues in each chain by enriching different positions of the chain with different % <sup>13</sup>C. The three labels in each oligomer were 'frame-shifted' by one residue relative to its neighbours in the oligomer series.
- 79 T. Boddaert, J. Solà, M. Helliwell and J. Clayden, *Chem. Commun.*, 2012, **48**, 3397–3399.
- 80 M. Tomsett, B. A. F. Le Bailly, S. Bijvoets, I. Maffucci, A. Contini, and J. Clayden, manuscript in preparation.
- 81 C. Toniolo and H. Brückner, *Peptaibiotics*, HCA / Wiley-VCH, Zürich, 2009.

## ARTICLE

Journal Name

- 82 G. B. Bunk czi, M. Schiell, L. S. V rtesy and G. M. Sheldrick, *J. Peptide Sci.*, 2003, **9**, 745–752.
- 83 M. De Poli, M. De Zotti, J. Raftery, J. A. Aguilar, G. A. Morris and J. Clayden, *J. Org. Chem.*, 2013, **78**, 2248–2255.
- 84 U. Orcel, M. De Poli, M. De Zotti and J. Clayden, *Chem. Eur. J.*, 2013, **19**, 16357–16365.
- 85 S. P. Fletcher, J. Solà, D. Holt, R. A. Brown and J. Clayden, *Beilstein. J. Org. Chem.*, 2011, **7**, 1304–1309.
- 86 M. De Zotti, B. Biondi, M. Crisma, C. U. Hjørringgaard, A. Berg, H. Brückner and C. Toniolo, *Biopolymers*, 2012, **98**, 36–49.
- 87 J. Solà, S. P. Fletcher, A. Castellanos and J. Clayden, *Angew. Chem. Int. Ed.*, 2010, **49**, 6836–6839.
- 88 E. Kolomiets, V. Berl, I. Odriozola, A.-M. Stadler, N. Kyritsakas and J.-M. Lehn, *Chem. Commun.*, 2003, 2868–2869.
- 89 B. A. F. Le Bailly, L. Byrne and J. Clayde, *Angew. Chem.* in press (doi 10.1002/ange.201510605)
- 90 J.-H. Ha and S. N. Loh, *Chem. Eur. J.*, 2012, **18**, 7984–7999.
- 91 T. D. James, M. D. Phillips and S. Shinkai, in *Boronic Acids in Saccharide Recognition*, Royal Society of Chemistry, 2006.
- 92 R. A. Brown, V. Diemer, S. J. Webb and J. Clayden, *Nature Chem.*, 2013, **5**, 853–860.
- 93 J. Brioché, S. J. Pike, S. Tshepelevitsh, I. Leito, G. A. Morris, S. J. Webb and J. Clayden, *J. Am. Chem. Soc.*, 2015, **137**, 6680–6691.
- 94 J. Lehn, *Chem. Eur. J.*, 2000, **6**, 2097–2102.
- 95 T. Okino, Y. Hoashi and Y. Takemoto, *J. Am. Chem. Soc.*, 2003, **125**, 12672–12673.
- 96 B. Amit, D. A. Ben-Efraim and A. Patchornik, *J. Am. Chem. Soc.*, 1976.
- 97 J.-M. Lehn, *Proc. Natl. Acad. Sci.*, 2002, **99**, 4763–4768.