ORGANIC CHEMISTRY

FRONTIERS

RESEARCH ARTICLE



View Article Online View Journal | View Issue

Cite this: Org. Chem. Front., 2014, 1, 661

Carbopalladation of bromoene-alkynylsilanes: mechanistic insights and synthesis of fused-ring bicyclic silanes and phenols[†]

Marie-Caroline A. Cordonnier,^a S. B. Jennifer Kan,^b Birgit Gockel,^a Shermin S. Goh^a and Edward A. Anderson*^a

The palladium-catalyzed cascade cyclization of silylated bromoenynes and alkenylstannanes provides a straightforward route to a range of bicyclic silylated cyclohexadienes. Mechanistic insights into aspects of carbopalladation and unusual palladium-mediated isomerizations have been obtained through the detection of reaction intermediates, the isolation of byproducts, and reaction monitoring by VT NMR spectroscopy. The utility of the bicyclic products is illustrated through oxidation to bicyclic enones and phenols.

Received 25th April 2014, Accepted 20th May 2014 DOI: 10.1039/c4go00123k

rsc.li/frontiers-organic

Introduction

Phenols represent one of the most important aromatic functional groups, featuring in numerous natural products and drugs.¹ Polycyclic phenols are of particular interest, as illustrated by highly bioactive compounds such as the anthracycline antibiotics, estrone, and morphine (Fig. 1).² As traditional methods for phenol preparation require harsh conditions that may not be compatible with sensitive functional groups,³ phenols are usually incorporated at an early stage of a synthesis, thus restricting synthetic planning.

Late-stage phenolation is therefore an attractive prospect that has driven the development of transition metal-catalyzed aryl halide hydroxylation and aryl C–H activation/oxidation methods.⁴ An alternative that avoids the prolonged heating often required by these processes is the use of a phenol surrogate, which could be revealed at a late stage of a synthetic route, but is stable to intermediate transformations. Here, arylboron derivatives have met with some success, particularly due to Molander's elegant work on the oxidation of aryl trifluoroborate salts,⁵ although the synthetic processability of these remains to be proven. In contrast, organosilanes – stalwarts of the protecting group field – display unrivalled tolerance towards multistep synthesis, and in recent work we have



Fig. 1 Fused-ring phenols in bioactive natural products.

reported the use of arylsilanes as a source of phenols through Tamao-type oxidation.^{6,7}

We have also disclosed a palladium-catalyzed cascade cyclization which prepares bi- or tricyclic products (1, Scheme 1) from the coupling of bromoenynes 2 with alkenyl- and dienylstannanes 3,⁸ a reaction pioneered and impressively explored by Suffert *et al.* for the synthesis of various polycycles.⁹ A feature of this work is the requirement for an internal alkyne (*i.e.* $\mathbb{R}^1 \neq \mathbb{H}$, Scheme 1) to avoid the formation of undesired triene isomers 4, which cannot undergo electrocyclization. One solution is to employ trimethylsilyl alkynes,¹⁰ and although this functionality indeed enables the desired cascade, it was clear that the resultant TMS-substituted cyclohexadienes would be of limited synthetic utility.

We realized that the use of a more functional silane would lead to a silicon-containing product capable of undergoing a range of further transformations, such as the aforementioned Tamao oxidation or Hiyama coupling.¹¹ Indeed, we have illus-

^aChemistry Research Laboratory, University of Oxford, 12 Mansfield Road, Oxford, OX1 3TA, UK. E-mail: edward.anderson@chem.ox.ac.uk; Fax: +44 (0)1865 285002; Tel: +44 (0)1865 285000

^bDepartment of Chemistry, University of Cambridge, Lensfield Road, Cambridge, CB2 1EW, UK

 $[\]dagger$ Electronic supplementary information (ESI) available: Copies of 1H and ^{13}C NMR spectra, and experimental details for NMR experiments. See DOI: 10.1039/c4q000123k



Scheme 1 Proposed synthesis of bicyclic phenols.

trated this principle in approaches to the CDE rings of rubriflordilactone A, where arylsilanes formed from the cyclizations of bromoenynes could be oxidized to give specific tricyclic phenols.^{8b,12} In this paper we wish to report methods for the synthesis of a range of silvlated bromoenynes, and their intermolecular coupling with alkenylstannanes to provide bicyclic silvlcyclohexadienes 5. The oxidation of these dienes and silane substituents leads to bicyclic arylsilanes (6) and phenols (7) that would be challenging to prepare by other routes, and offers a de novo approach to ring systems of this type.13 Alongside this synthetic work, we expand on the mechanistic pathways operative in the cascade cyclization, including two unusual palladium-mediated isomerization processes. The unexpected observation of an unprecedented formal 4-endotrig cyclization to afford fused ring cyclobutenes is also described.

Results and discussion

Synthesis of cascade cyclization substrates

We selected a range of linkers to connect the bromoalkene and silylalkyne, such that bicyclic products containing 5- to 7-membered rings, and heterocycles, could be obtained. Synthesis of carbon-tethered bromoenynes began with dimethylmalonate (Scheme 2), which following monoallylation with 2,3-dibromopropene (70%) was alkylated with alkynyl electrophiles 8a-c to give bromoenynes 9a-c (69-90%). The silylation of these alkynes was initially attempted using conditions employed in our arylsilane work.^{6a} Thus, deprotonation of 9a with *n*-butyllithium followed by trapping of the resultant alkynyllithium species with the moisture-sensitive (diethylamino)chlorodimethylsilane (Et2NMe2SiCl) gave an intermediate aminosilane which was not isolated, but immediately converted to the dimethylisopropoxysilane 10a (Route A, Scheme 2).¹⁴ Perhaps unsurprisingly, this silane proved to be rather unstable towards chromatography, however through rapid chromatographic purification using a solvent system buffered with triethylamine, 10a could be isolated in 75% yield. In order to address the chromatographic instability of



Scheme 2 Synthesis of malonate-derived silylbromoenynes.



Scheme 3 Synthesis of heteroatom-tethered silylbromoenynes.

this dimethylisopropoxysilane, we employed the analogous diethylsilane reagent $(Et_2N)SiEt_2SiCl$,¹⁵ which gave the alkynyl isopropoxydiethylsilanes **11a–c** in much improved yields following chromatographic purification (Route B). An alternative solution to the synthesis of **10a** proved to be the use of dimethylchlorosilane (HMe₂SiCl) as the acetylide trapping agent (Route C); the ensuing hydrolytically stable intermediate alkynylhydrosilane could be easily purified, and then smoothly oxidized to **10a** using Lee and Chang's elegant ruthenium-catalyzed dehydrogenative silylation methodology,¹⁶ which improved the yield of **10a** to 93%.

The syntheses of heteroatom-tethered silylbromoenynes could be achieved uneventfully using equivalent chemistry, with sulfonamide-tethered enyne **12** and ether-linked enyne **13** being prepared in reasonable yields over two steps from known sulfonamide **14** and propargyl alcohol, respectively (Scheme 3).

Cascade cyclizations of silylated bromoenynes

With a selection of silylated alkynes in hand, their palladiumcatalyzed cascade cyclizations were examined. These reactions



^{*a*} Isolated yield. ^{*b*} Isolated as an 80:13:7 ratio of **15f:16a:17a**. ^{*c*} Isolated as a 63:19:18 ratio of **15g:16b:17b**. See text for discussion.

were carried out according to our previously optimized conditions,^{8*a*} using $PdCl_2(PPh_3)_2$ as precatalyst in refluxing toluene (Table 1). For the purposes of this work, the propenyl and styrenyl stannanes **3a** and **3b** were selected as generic cross-coupling partners. The reaction of dimethylsilane **10a** was first investigated, which in spite of the relative instability of the alkynylsilane cyclized in excellent yields with both stannanes (92–97%, entries 1, 2). Notably, the hydrolytic stability of the silane increases markedly upon cyclization owing to the increased steric bulk of the cyclohexadienyl substituent in **15a** compared to the alkyne substituent of the starting material. The cyclizations of the (more robust) diethylsilane substrates proved more challenging (entries 3–5), with these reactions requiring longer reaction times, likely due to steric hindrance



Fig. 2 Cyclization byproducts **16a/b** and **17a/b**. Selected key ¹H NMR chemical shifts/multiplicities (in CDCl₃) for **16a** and **17a** are shown.

of the key transmetallation step following carbopalladation. This resulted in lower isolated yields, as exemplified by the malonate and sulfonamide tethered enynes **11a** and **12** which gave the corresponding bicycles **15c** and **15d** in 77 and 63% yields respectively. The oxygen-tethered substrate **13** proved particularly troublesome, delivering the dihydrofuran product **15e** in low yield, possibly due to instability of the allyl propargyl ether. Reaction of substrates featuring a longer tether also proved effective, giving the 6,6- and 7,6-bicyclic products **15f** and **15g** in good yields (entries 6, 7).

In the latter two cases, the desired bicyclic products were contaminated with significant quantities of partially separable isomeric byproducts 16a, b and 17a, b (Fig. 2), which were identified by careful analysis of ¹H-¹H COSY, and ¹H-¹³C HSQC and HMBC 2D NMR spectra;¹⁷ the assignment of one of these sets of byproducts (16a and 17a) is discussed here. Byproduct 16a contains an *exo*-methylene unit ($\delta_{\rm H}$ 5.01 and 4.85 ppm), and two vicinal alkene protons ($\delta_{\rm H}$ 6.00 ppm, dd, J = 15.5, 1.5 Hz; and 5.32 ppm, dq, J = 15.5 and 6.5 Hz), the latter being characteristic of the trans-alkene of the propenyl unit. These data, together with key HMBC correlations, are strongly suggestive of the formal anti-carbopalladation/ cross-coupling product 16a, the formation of which is entirely consistent with our earlier results.8a The identification of the second byproduct proved less straightforward. Key signals in the ¹H NMR spectrum at $\delta_{\rm H}$ 6.36 (d, J = 10.6 Hz), 6.31 (ddd, J = 16.5, 10.6 and 9.7 Hz), 5.26 (dd, J = 16.5 and 2.0 Hz) and 5.25 ppm (dd, I = 9.8 and 2.0 Hz) revealed a connectivity between four alkene protons and thus the surprising presence of a 1,3-butadienyl unit. A further significant piece of evidence was the detection of a methyl singlet at 1.50 ppm, to which the proton at 6.33 ppm showed an nOe enhancement. These combined observations, and ¹H-¹³C correlations, led us to propose structure 17a, which features an (E)-alkenylsilane as part of a conjugated triene.

A mechanistic hypothesis for the formation of these byproducts is depicted in Scheme 4. The formation of byproduct **16** is consistent with our earlier observation^{8a} that such undesired *anti*-trienes form in the course of the cyclization reaction, potentially *via* isomerization of the intermediate dienylpalladium complex **18** to its isomer **19**.¹⁸ The rate of transmetallation of complex **19** is likely to exceed that of **18**, so even small amounts of **19** may lead to significant quantities of *anti*-triene **16** (*i.e.*, a Curtin-Hammett situation). The formation of increased quantities of this triene for substrates **11b** and **11c**



Scheme 4 Mechanistic proposal for byproduct formation. E = CO_2Me , [Si] = SiEt₂Oi-Pr.

may reflect the increased flexibility of the larger tethering ring, which can distort to alleviate 1,3-allylic strain between the silvl substituent and exo-methylene in 19 - and therefore reduce the steric cost of placing a silane in this more hindered position. The formation of 17 may be explained by our second previous observation^{8a} that anti-trienes can isomerize to bicyclic products on prolonged exposure to the reaction conditions. We suggest that this process may proceed via oxidative addition of Pd(0) with diene 16 to give a palladacyclopentene 20.19 This could undergo a 1,3-allylic migration of the palladium atom to the 7-membered palladacycle 21, reductive elimination from which would afford the cyclohexadiene product 15.²⁰ The formation of byproduct 17 could be rationalised by β-hydride elimination from this common palladacycle intermediate 21, followed by reductive elimination of the resultant palladium(II) hydride species. We presume that the steric hindrance imposed by the methyl group in 17 prevents a 6π -electrocyclization of this compound.

As [4 + 1] oxidative additions of Pd(0) to dienes are rare,¹⁹ more concrete evidence to support these pathways was sought. Firstly, the formation of 16 via the intermediacy of dienylpalladium complex 19 was probed through the exposure of alkynylsilane 22 to one equivalent of $Pd(PPh_3)_4$ in d₈-toluene at 110 °C (Fig. 3). The characteristic methylene signals of 22 at $\delta_{\rm H}$ 5.41 and 5.71 ppm were rapidly converted (10 min) to two new sets of peaks: a prominent (apparent) singlet at 4.60 ppm, and two smaller singlets at 4.58 and 4.78 ppm. These were tentatively assigned as the exo-methylene peaks of the syn- and antidienylpalladium complexes 23 and 24 respectively, based on analysis by COSY and HSQC experiments. Support for this assignment was gained through the addition of 1.5 equivalents of tributylvinyltin to the NMR tube; further heating for 10 minutes led to exclusive formation of cyclohexadiene 25 and triene 26 in a ratio mirroring that of these intermediate species (84:16).

To explore the isomerization of **26** to **25**, a 1:0.32 mixture of these compounds, formed from reaction of silane **22** with



Fig. 3 Use of stoichiometric $Pd(PPh_3)_4/VT$ ¹H NMR spectroscopy to probe the putative dienylpalladium(II) intermediates.



Scheme 5 Exploration of conditions to effect the isomerization of *anti*-triene **26** to bicycle **25**.

vinyltributyltin for two hours, was purified by silica gel chromatography (Scheme 5); the triene **26** in this mixture would be expected to isomerize to give the **25** upon resubmission to the reaction conditions. Heating the mixture in toluene overnight at 110 $^{\circ}$ C in the absence of catalyst led to no conversion





Fig. 4 1 H NMR spectroscopic monitoring of the conversion of 26 to 25 catalyzed by Pd(PPh₃)₄. 1,4-Dimethoxybenzene was used as internal standard.

of 26 to the bicyclic product 25, with both compounds recovered unchanged after this period (thus ruling out a purely thermal process). Palladium(II) salts have previously been shown to promote alkene isomerisation,²¹ but subjection of the anti-triene to PdCl₂(PPh₃)₂ at 110 °C also led to no reaction. However, heating the mixture in the presence of Pd (PPh₃)₄ for 20 h led to complete consumption of *anti*-triene, and by performing this isomerization in d₈-toluene with monitoring by ¹H NMR spectroscopy in the presence of an internal standard (1,4-dimethoxybenzene), a smooth conversion of 26 to 25 was observed (Fig. 4). This clearly demonstrates that the isomerization of anti-triene to product is not a thermal process, and in fact requires a Pd(0) catalyst, thus offering some support to our proposed mechanism. At no point do we detect the formation of syn-triene, which lends some weight to our mechanistic hypothesis for the direct conversion of antitrienes to bicyclic products (Scheme 4, although we recognise that any syn-triene formed could electrocyclize rapidly).

Cyclization to 7,4-fused ring cyclobutenes

In the course of cyclization reactions to form seven-membered rings (including 11c), we had noticed the occasional formation of a different byproduct to those discussed thus far, the production of which seemed highly dependent on the reaction concentration, and quantity of stannane coupling partner. Although this byproduct was not observed under our optimized conditions, the use of <1.5 equivalents of stannane, or more dilute reaction conditions (i.e. such that transmetallation would be slowed) increased its formation. In fact, we had first observed such a species in the attempted 8π-electrocyclic coupling of bromoenyne 27 with stannane 28 (Fig. 5), which resulted in a surprising degree of apparent protodestannylation of 28 (leading to the known diene 29).²² We assigned the product formed in this reaction as the 7,4-fused cyclobutene 30 based on detailed analysis by 2D NMR experiments. Specifically, a complete set of HMBC correlations (Fig. 5) was observed between protons H1, H3 and H7, and carbons C2, C8



Fig. 5 Initial synthesis of 7,4-fused cyclobutene 30, and key HMBC spectra correlations. E = CO_2Me . • = strong correlation; • = weak correlation.



Scheme 6 Possible mechanisms for the formation of 7,4-fused cyclobutene 30. $E = CO_2 Me$.

and C9, together with long-range coupling between H1 and H7 (${}^{5}J$ = 2.7 Hz; this coupling was also observed in a ${}^{1}H{-}^{1}H$ COSY spectrum).

The formation of this product is not unreasonable if potential mechanisms for its formation are considered (Scheme 6). Following carbopalladation (**31**), one possibility would involve a 4-*endo-trig* carbopalladation (Path A), leading to cyclobutene



Scheme 7 Heck conditions for formation of 7,4-fused cyclobutenes.

32 – a pathway that might be favoured, in spite of ring strain, due to the formation of an allylpalladium intermediate. β -Hydride elimination would afford the observed product 30, and liberate a palladium(II) hydride species that could be reduced to palladium(0) by transmetallation with 28, thus leading to the 'protodestannylated' product 29. However, due to a lack of precedent for this mode of carbopalladation, we also consider an electrophilic palladation route feasible (Path B), in which attack by the exo-methylene on the proximal palladium(II) atom leads to palladacycle 33 - which may again be rendered possible by the formation of an allyl cation in this process. Now, loss of a proton generates palladacyclopentene 35, reductive elimination from which leads to 30.²³ The resemblance of intermediate 35 to those proposed in enyne cycloisomerization processes is clear;²⁴ the proton lost from this pathway could then effect protodestannylation of the coupling partner 28 to afford 29.

Whatever mechanism is operational, it was clear to us that this process overall corresponds to a Heck reaction in which regeneration of the palladium(0) catalyst is mediated by the stannane reagent. This suggested that an amine base might perform a similar role, and to our delight, the use of common Heck conditions (toluene, triethylamine) at just 5 mol% catalyst loading indeed led to a high-yielding cyclization of **27** to **30** (90%, Scheme 7). The more functional bromoenyne **11c** was also tested in this chemistry, which gave the corresponding 7,4-fused silylcyclobutene **36** in excellent yield (86%). This efficient process offers an alternative entry to this type of fused cyclobutene ring system.²⁵

Oxidations of silylcyclohexadienes

To illustrate the potential utility of this chemistry, we subjected the product dienylsilanes to a selection of oxidative transformations. Firstly, a direct Tamao oxidation of the dienylsilane was carried out,²⁶ which we hoped would deliver a bicyclic enone. In the case of the isopropoxydimethylsilane **15a**, this met with some success (Scheme 8), delivering **37** in moderate yield (43%). The successful formation of this product, which lacks any olefinic protons, could be confirmed by ¹³C NMR ($\delta_{\rm C}$ 194.9, 158.7, 134.8 ppm for the enone region) and IR spectroscopy ($\nu_{\rm max}$ 1667 cm⁻¹). Attempted oxidation of the equivalent diethylisopropoxysilanes proved unsuccessful, potentially due to competing nucleophilic epoxidation of the product enone, which highlights a reactivity benefit of the less-hindered dimethylalkoxysilane group.



Scheme 8 Bicyclic enone formation.



Scheme 9 Oxidation of the silylcyclohexadiene framework.

The silylcyclohexadiene frameworks could also be readily oxidized to the corresponding arylsilanes using manganese dioxide, conditions that we had successfully employed in other work²⁷ and which proved superior to the use of other oxidants such as DDQ. The resultant arylsilanes could generally be isolated in excellent yield; two examples are shown in Scheme 9 (**38**: 88%; **39**: 93%). These arylsilanes show potential for a range of transformations – but here, in keeping with our interests in the synthesis of phenols from arylsilanes,⁶ we chose to investigate Tamao oxidation. Under Tamao conditions (TBAF, KHCO₃, H₂O₂, 60 °C),²⁸ good yields of the corresponding phenols **40** and **41** were obtained, thus validating this approach to the synthesis of bi/polycyclic phenols.

Conclusions

In summary, we have developed efficient routes for the synthesis of functional alkynylsilanes and demonstrated their application in bicyclization cascade reactions. The resultant silvlated bicyclic cyclohexadienes are substrates for oxidation to bicyclic enones, arylsilanes, and phenols; the latter process thus affords bicyclic phenols from acyclic precursors in just three steps. Investigations into the mechanism of the cascade cyclization have revealed some unusual palladium-mediated isomerization pathways. Finally, 7,4-fused cyclobutene ring systems, arising from cyclization of a 7-membered exocyclic dienylpalladium complex, could be formed under standard 'Heck' type conditions. Together, these processes underline the rich reactivity – both expected and unexpected – that can be harvested from carbopalladation chemistry.

Experimental

General

Reagents were used as purchased, or purified by standard laboratory techniques. Reaction solvents were purified using an alumina column drying system, and reactions were performed under inert atmosphere unless otherwise stated.

Dimethyl 2-(2-bromoallyl)malonate. To a suspension of NaH (60% dispersion in mineral oil, 2.32 g, 58.0 mmol) in THF (50 mL) at 0 °C was added dropwise dimethyl malonate (5.52 mL, 48.4 mmol), and the mixture was stirred for 1 h. 2,3dibromopropene (80% tech. grade, 5.52 mL, 48.4 mmol) was added to the milky yellow mixture, and stirring was continued for 3 h until completion of the reaction as monitored by TLC. The reaction was quenched with NH₄Cl (sat., aq.). The aqueous phase was extracted with Et₂O and the combined organic extracts washed with brine and dried (MgSO₄). The solvent was evaporated in vacuo and the crude product purified by distillation at (101-106 °C, 1 mmHg) to afford the product as a colourless oil (8.50 g, 33.9 mmol, 70%). Rf 0.22 (5:1 pet. ether-ether); ¹H NMR (500 MHz, CDCl₃) $\delta_{\rm H}$ 5.69 (1H, dt, J = 2.0, 1.0 Hz), 5.48 (1H, d, J = 2.0 Hz), 3.89 (1H, t, J = 7.5 Hz), 3.76 (6H, s), 3.03 (2H, dd, J = 7.5, 1.0 Hz); ¹³C NMR (125 MHz, $CDCl_3$) δ_C 168.7, 129.2, 119.9, 52.7, 50.4, 40.5. Data in accordance with the literature.29

General procedure 1: alkylation of dimethyl 2-(2-bromoallyl)malonate

A solution of dimethyl 2-(2-bromoallyl)malonate (1 equiv.) in THF (2 mL mmol⁻¹) was added to a stirred suspension of NaH (60% dispersion in mineral oil, 1.2 equiv.) and the alkyne electrophile (1.2 equiv.) in THF (2 mL mmol⁻¹). The resulting mixture was stirred at room temperature under argon until completion of the reaction as monitored by TLC. The reaction was quenched by addition of NH₄Cl (sat., aq.). The aqueous layer was extracted with Et₂O and the combined organic extracts washed with brine, and dried (MgSO₄). The solvent was evaporated *in vacuo* and the crude product purified by flash chromatography to afford the bromoenyne product.

Dimethyl 2-(2-bromoallyl)-2-(prop-2-ynyl)malonate (9a). Synthesised from dimethyl 2-(2-bromoallyl)malonate (2.90 g, 11.7 mmol) and propargyl bromide (1.60 mL, 14.0 mmol) using general procedure 1. The crude product was purified by flash chromatography (5:1 pet. ether–ether) to afford **9a** as a colourless oil (2.91 g, 10.1 mmol, 86%). $R_{\rm f}$ 0.31 (5:1 pet. ether–ether); IR (thin film) $\nu_{\rm max}/{\rm cm}^{-1}$ 3289, 2954, 1737; ¹H NMR (500 MHz, CDCl₃) $\delta_{\rm H}$ 5.83 (1H, t, J = 0.6 Hz), 5.63 (1H, d, J = 1.6 Hz), 3.77 (6H, s), 3.31 (2H, br s), 2.93 (2H, d, J = 2.7 Hz), 2.05 (1H, t, J = 2.7 Hz); ¹³C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 169.5, 126.1, 122.9, 78.6, 72.1, 56.0, 53.0, 42.9, 22.2; HRMS (ES⁺) calcd for C₁₁H₁₃BrO₄ [M + H]⁺ 289.0074, found 289.0070. Data in accordance with the literature.^{8a}

Dimethyl 2-(2-bromoallyl)-2-(but-3-ynyl)malonate (9b). Synthesised from dimethyl 2-(2-bromoallyl)malonate (876 mg, 3.49 mmol) and but-3-ynyl trifluoromethanesulfonate (847 mg, 4.19 mmol) using general procedure 1. The crude product was

purified by flash chromatography (5 : 1 pet. ether–ether) to afford **9b** as a colourless oil (730 mg, 2.41 mmol, 69%). $R_{\rm f}$ 0.26 (5 : 1 pet. ether–ether); IR (thin film) $\nu_{\rm max}/{\rm cm}^{-1}$ 3285, 2954, 1730, 1624, 1433, 1201, 1150, 959, 649; ¹H NMR (500 MHz, CDCl₃) $\delta_{\rm H}$ 5.70–5.68 (1H, m), 5.61 (1H, d, J = 1.8 Hz), 3.75 (6H, s), 3.19 (2H, d, J = 0.5 Hz), 2.30–2.28 (2H, m), 2.21–2.19 (2H, m), 1.97 (1H, t, J = 2.8 Hz); ¹³C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 170.4, 126.7, 122.2, 82.8, 68.9, 56.4, 52.8, 43.3, 30.7, 14.0; HRMS (ES⁺) calcd for C₁₂H₁₅BrNaO₄ [M + Na]⁺ 325.0070, found 325.0062.

Dimethyl 2-(2-bromoallyl)-2-(pent-4-ynyl)malonate (9c). Synthesised from dimethyl 2-(2-bromoallyl)malonate (200 mg, 0.796 mmol) and pent-4-ynyl trifluoromethanesulfonate (207 mg, 0.957 mmol) using general procedure 1. The crude product was purified by flash chromatography (5 : 1 pet. ether-ether) to afford **9c** as a colourless oil (227 mg, 0.716 mmol, 90%). $R_{\rm f}$ 0.26 (5 : 1 pet. ether-ether); IR (thin film) $\nu_{\rm max}/{\rm cm^{-1}}$ 3291, 2952, 1731, 1625, 1434, 1278, 1150, 901, 640; ¹H NMR (500 MHz, CDCl₃) $\delta_{\rm H}$ 5.69–5.67 (1H, m), 5.59 (1H, d, J = 1.7 Hz), 3.75 (6H, s), 3.17 (2H, s), 2.21 (2H, dt, J = 7.0, 2.6 Hz), 2.12–2.10 (2H, m), 1.96 (1H, t, J = 2.6 Hz), 1.44–1.42 (2H, m); ¹³C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 170.8, 126.9, 121.9, 83.5, 68.8, 56.8, 52.7, 43.1, 30.8, 23.4, 18.5; HRMS (ES⁺) calcd for C₁₃H₁₇BrNaO₄ [M + Na]⁺ 339.0202, found 339.0200.

2-(2-bromoallyl)-2-(3-(isopropoxydimethylsilyl)-Dimethyl prop-2-ynyl)malonate (10a). Method A. To a solution of 9a (150 mg, 0.519 mmol) in dry THF (5 mL) at -78 °C was added *n*-BuLi (1.6 M in hexanes, 405 µL, 0.649 mmol, 1.25 equiv.), and the mixture was stirred for 20 min. Chloro(diethylamino)dimethylsilane (130 µL, 0.649 mmol, 1.25 equiv.) was added, and the reaction mixture was allowed to warm to rt. When complete as judged by TLC, the reaction was concentrated, diluted with dry pentane, and stirred for 15 min, then filtered through oven-dried Celite. The clear filtrate was evaporated to give the intermediate aminosilane. This was dissolved in CH₂Cl₂ (3 mL) and cooled to 0 °C, then imidazole (71 mg, 1.04 mmol, 2 equiv.), isopropanol (80 µL, 1.04 mmol, 2 equiv.) and DMAP (cat.) were added to the mixture, which was stirred at rt for 2 h. The reaction was quenched with NaHCO3 (aq., sat.), the aqueous layer was extracted with CH₂Cl₂, and the combined organic extracts were washed with brine, and dried (MgSO₄). The solvent was evaporated in vacuo and the product was purified by flash chromatography (10:1 pet. ether-ether with 1% Et₃N) to afford the 10a as a colourless oil (158 mg, 0.39 mmol, 75%). Rf 0.38 (5:1 pet. ether-ether); IR (thin film) $\nu_{\rm max}/{\rm cm}^{-1}$ 2360, 1658, 1568, 1382, 1369, 1290, 1030; ¹H NMR (400 MHz, CDCl₃) $\delta_{\rm H}$ 5.83–5.81 (1H, m), 5.63 (1H, d, J = 1.5 Hz), 4.09 (1H, septet, J = 6.1 Hz), 3.76 (6H, s), 3.30 (2H, s), 2.97 (2H, s), 1.18 (6H, d, J = 6.1 Hz), 0.22 (6H, s); ¹³C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 169.4, 126.2, 122.7, 100.8, 87.4, 65.9, 56.0, 53.0, 43.0, 25.4, 23.4, 0.6; HRMS (ES⁺) calcd for $C_{16}H_{25}BrNaO_5Si [M + Na]^+ 427.0533$, found 427.0547.

Dimethyl 2-(2-bromoallyl)-2-(3-(isopropoxydimethylsilyl)prop-2-ynyl)malonate (10a). Method B. To a solution of 9a (200 mg, 0.691 mmol) in THF (2 mL) at -78 °C was added LiHMDS (1 M in hexanes, 830 μ L, 0.83 mmol) and the mixture was stirred for 30 min. ClSiMe₂H (116 µL, 1.038 mmol) was then added, and the reaction was allowed to warm to rt, then stirred until completion as monitored by TLC. The reaction was quenched with H₂O, and the aqueous layer was extracted with ether. The combined organic extracts washed with brine and dried (MgSO₄). The solvent was evaporated in vacuo and the crude product was purified by flash chromatography (10:1 pet. ether-ether), to afford the intermediate alkynyl hydrosilane as a colourless oil (224 mg, 0.645 mmol, 93%). ¹H NMR (400 MHz, CDCl₃) $\delta_{\rm H}$ 5.65–5.58 (1H, m), 5.43 (1H, d, J = 1.5 Hz), 3.87 (1H, septet, J = 3.7 Hz), 3.56 (6H, s), 3.10 (2H, s), 2.76 (2H, s), 0.01 (6H, d, J = 3.7 Hz); ¹³C NMR (100 MHz, $CDCl_3$) δ_C 169.4, 126.1, 122.8, 102.5, 86.0, 56.0, 53.0, 42.9, 23.6, -3.0; HRMS (ES⁺) calcd for C₁₃H₁₉BrNaO₄Si [M + Na]⁺ 369.0128, found 369.0128. To a solution of this hydrosilane (30 mg, 0.086 mmol) in i-PrOH (100 µL) was added [RuCl₂-(p-cymene)]₂ (0.3 mg, 0.0004 mmol). The reaction was stirred at rt for 10 min, then concentrated. Pentane (1 mL) was added, and the crude product was filtered through Celite (ether eluent). Concentration of the filtrate gave 10a as a pale pink oil (35 mg, 0.086 mmol, 99%).

(Diethylamino)diethylchlorosilane. To a solution of dichlorodiethylsilane (20 mL, 134 mmol) and anhydrous triethylamine (20.6 mL, 147 mmol) in THF at 0 °C was added a solution of anhydrous diethylamine (13.8 mL, 137 mmol) in THF (6 mL) over 2 h. The resulting beige suspension was warmed to rt overnight with vigorous stirring maintained. The mixture was then filtered through Celite under N₂, the filter cake washed with anhydrous pentane, and the filtrate concentrated *in vacuo* to give an orange oil. This residue was purified by vacuum distillation (bp 92–94 °C, 36 mbar) to afford (diethylamino) diethylchlorosilane as a colourless oil (17.6 g, 90.8 mmol, 68%). ¹H NMR (250 MHz, CDCl₃) $\delta_{\rm H}$ 2.88 (4H, q, *J* = 7.0 Hz), 1.07–0.99 (12H, m), 0.90 (4H, q, *J* = 6.5 Hz). Data in accordance with literature.³⁰

General procedure 2: alkyne silylation with (diethylamino)diethylchlorosilane

To a solution of bromoenyne (1 equiv.) in THF (4.5 mL mmol⁻¹) at -78 °C under Ar was added LiHMDS (1 M in THF, 1.5 equiv.). The pale yellow solution was stirred at -78 °C for 20 min and (diethylamino)diethylchlorosilane (1.5 equiv.) was added dropwise to the reaction mixture. After stirring for 30 min, after which time the starting material had been consumed (as monitored by TLC), i-PrOH (4 equiv.), DMAP (cat.) and imidazole (4 equiv.) were added and the reaction was allowed to warm to rt. The reaction was quenched by addition of water, and the aqueous layer was extracted with ether. The combined organic extracts were washed with brine, dried (MgSO₄), and concentrated. The residue was purified by flash chromatography to afford the silylated bromoenyne.

Dimethyl 2-(2-bromoallyl)-2-(3-(diethyl(isopropoxy)silyl)prop-2-ynyl)malonate (11a). Synthesised from 9a (500 mg, 1.73 mmol), LiHMDS (2.60 mL, 2.60 mmol), $ClSiEt_2(NEt_2)$ (606 μ L, 2.60 mmol), i-PrOH (530 μ L, 6.90 mmol) and imidazole (471 mg, 6.92 mmol) using general procedure 2. The product was purified by flash chromatography (5 : 1 pet. etherether) to afford **11a** as a colourless oil (701 mg, 1.62 mmol, 94%). $R_{\rm f}$ 0.26 (5 : 1 pet. ether-ether); Anal. calcd for $C_{18}H_{29}BrO_5Si$: C, 49.88; H, 6.74, found: C, 49.96; H, 6.72; IR (thin film) $\nu_{\rm max}/{\rm cm}^{-1}$ 2956, 2178, 1742, 1626, 1025, 733; ¹H NMR (500 MHz, CDCl₃) $\delta_{\rm H}$ 5.84–5.82 (1H, m), 5.63 (1H, d, J = 1.7 Hz), 4.10 (1H, septet, J = 6.1 Hz), 3.76 (6H, s), 3.32 (2H, s), 3.00 (2H, s), 1.18 (6H, d, J = 6.1 Hz), 0.98 (6H, t, J = 7.9 Hz), 0.62 (4H, app. qd, J = 7.9 Hz, 2.2 Hz); ¹³C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 169.3, 126.2, 122.7, 101.6, 85.5, 66.0, 56.1, 53.0, 42.9, 25.4, 23.5, 6.8, 6.6; HRMS (ES⁺) calcd for $C_{18}H_{29}NaBrO_5Si$ [M + Na]⁺ 455.0860, found 455.0862.

Dimethyl 2-(2-bromoallyl)-2-(4-(diethyl(isopropoxy)silyl)but-3-ynyl)malonate (11b). Synthesised from 9b (545 mg, 1.80 mmol), LiHMDS (2.70 mL, 2.70 mmol), ClSiEt₂(NEt₂) (634 μL, 2.70 mmol), i-PrOH (550 μL, 7.20 mmol) and imidazole (490 mg, 7.20 mmol) using general procedure 2. The product was purified by flash chromatography (5:1 pet. ether-ether) to afford **11b** as a colourless oil (723 mg, 1.62 mmol, 90%); $R_{\rm f}$ 0.26 (5:1 Pet. ether-ether); Anal. calcd for C19H31BrO5Si: C, 51.00; H, 6.98, found: C, 50.92, H; 7.02; IR (thin film) $\nu_{\rm max}$ / cm⁻¹ 2956, 2173, 1735, 1625, 1435, 1088, 1027, 732; ¹H NMR (500 MHz, CDCl₃) $\delta_{\rm H}$ 5.84–5.82 (1H, m), 5.63 (1H, d, J = 1.7 Hz), 4.10 (1H, septet, J = 6.1 Hz), 3.76 (6H, s), 3.18 (2H, s), 2.31-2.29 (2H, m), 2.23-2.24 (2H, m), 1.18 (6H, d, J = 6.1 Hz), 0.98 (6H, t, J = 7.9 Hz), 0.62 (4H, app. qd, J = 7.9 Hz, 2.2 Hz); ¹³C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 170.4, 126.6, 122.2, 106.0, 81.9, 65.9, 56.5, 52.8, 43.4, 30.9, 25.5, 15.3, 6.8, 6.5; HRMS (ES⁺) calcd for $C_{19}H_{31}BrNaO_5Si [M + Na]^+$ 469.1016, found 469.1015.

Dimethyl 2-(2-bromoallyl)-2-(5-(diethyl(isopropoxy)silyl)pent-4-ynyl)malonate 11c). Synthesised from 9c (320 mg, 1.01 mmol), LiHMDS (1.50 mL, 1.50 mmol), ClSiEt₂(NEt₂) (355 µL, 1.50 mmol), i-PrOH (306 µL, 4.00 mmol) and imidazole (272 mg, 4.00 mmol) using general procedure 2. The product was purified by flash chromatography (5:1 pet. etherether) to afford 11c as a colourless oil (337 mg, 0.730 mmol, 72%); R_f 0.26 (5:1 pet. ether-ether); Anal. calcd for C₂₀H₃₃BrO₅Si: C, 52.05; H, 7.21, found: C, 51.94; H, 7.30; IR (thin film) $\nu_{\text{max}}/\text{cm}^{-1}$ 2955, 2172, 1735, 1625, 1434, 1278, 1026, 732; ¹H NMR (500 MHz, CDCl₃) $\delta_{\rm H}$ 5.68–5.66 (1H, m), 5.59 (1H, d, J = 1.5 Hz), 4.14 (1H, septet, J = 6.4 Hz), 3.75 (6H, s),3.18 (2H, s), 2.29 (2H, t, J = 7.3 Hz), 2.14-2.12 (2H, m), 1.49-1.47 (2H, m), 1.20 (6H, d, J = 3.4 Hz), 1.00 (6H, t, J = 6.4 Hz), 0.65–0.62 (4H, m); 13 C NMR (125 MHz, CDCl₃) δ_{C} 170.8, 127.0, 121.9, 106.8, 81.8, 65.9, 56.9, 52.9, 43.2, 30.9, 25.5, 23.4, 19.9, 6.9, 6.6; HRMS (ES⁺) calcd for C₂₀H₃₃BrNaO₅Si $[M + Na]^+$ 483.1173, found 483.1179.

N-(2-Bromoallyl)-4-methyl-*N*-(prop-2-ynyl)benzene sulfonamide. To a mixture of *N*-(2-bromoallyl)-4-methylbenzenesulfonamide (1.00 g, 3.45 mmol) and K_2CO_3 (953 mg, 6.89 mmol) in acetone (10 mL) was added propargyl bromide (80% in toluene, 460 µL, 4.14 mmol). The reaction mixture was stirred at 60 °C for 12 h, then it was cooled to rt and quenched with water. The aqueous layer was extracted with Et₂O, then the combined organic layers were dried (MgSO₄) and concentrated *in vacuo*. The crude product was purified by flash chromatography (2 : 1 pet. ether–ether) to afford the title compound as a colourless oil (903 mg, 2.75 mmol, 80%); $R_{\rm f}$ 0.35 (2 : 1 pet. ether–ether); IR (thin film) $\nu_{\rm max}/{\rm cm}^{-1}$ 3280, 1629, 1344, 1181, 1150, 892, 679; ¹H NMR (500 MHz, CDCl₃) $\delta_{\rm H}$ 7.75 (2H, d, J = 8.1 Hz), 7.31 (2H, d, J = 8.1 Hz), 5.97–5.95 (1H, m), 5.69–5.67 (1H, m), 4.13 (2H, d, J = 2.3 Hz), 4.08 (2H, s), 2.44 (3H, s), 2.05 (1H, t, J = 2.3 Hz); ¹³C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 143.9, 135.8, 129.6, 127.7, 126.9, 120.5, 76.0, 74.2, 53.8, 36.1, 21.6; HRMS (ES⁺) calcd for C₁₄H₁₈BrNNaO₃S [M + MeOH + Na]⁺ 382.0083, found 382.0077.

N-(2-Bromoallyl)-N-(3-(diethyl(isopropoxy)silyl)prop-2-ynyl)-4-methylbenzene sulfonamide (12). Synthesised from N-(2bromoallyl)-4-methyl-N-(prop-2-ynyl)benzene sulfonamide (150 mg, 0.457 mmol), LiHMDS (686 µL, 0.686 mmol), ClSiEt₂(NEt₂) (160 µL, 0.686 mmol), i-PrOH (140 µL, 1.80 mmol) and imidazole (125 mg, 1.84 mmol) using general procedure 2. The product was purified by flash chromatography (5:1 pet. ether-ether) to afford 12 as a colourless oil (107 mg, 0.226 mmol, 50%); R_f 0.26 (5:1 pet. ether-ether); Anal. calcd for C20H30BrNO3SSi: C, 50.84; H, 6.40; N, 2.96, found: C, 50.90; H, 6.32; N, 2.95. IR (thin film) $\nu_{\text{max}}/\text{cm}^{-1}$ 2960, 1353, 1163, 1119, 1092, 1002, 899, 734, 661; ¹H NMR (400 MHz, CDCl₃) $\delta_{\rm H}$ 7.77 (2H, d, J = 8.1 Hz), 7.33 (2H, d, J = 8.1 Hz), 5.98 (1H, d, J = 1.8 Hz), 5.72 (1H, d, J = 1.8 Hz), 4.23 (2H, s), 4.11 (2H, s), 3.95 (1H, sept, J = 6.0 Hz), 2.45 (3H, s),1.14 (6H, d, J = 6.0 Hz), 0.89 (6H, t, J = 7.8 Hz), 0.51 (4H, q, J = 7.8 Hz); ¹³C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 144.0, 136.3, 130.4, 130.1, 128.1, 119.7, 98.9, 66.5, 45.4, 41.0, 38.6, 26.2, 25.8, 7.8, 7.0; HRMS (ES⁺) calcd for $C_{20}H_{30}BrNNaO_3SSi [M + Na]^+$ 494.0791, found 494.0789.

2-Bromo-3-(prop-2-ynyloxy)prop-1-ene. To a solution of NaH (60% in mineral oil, 1.50 g, 35.8 mmol) in THF (10 mL) at 0 °C was added propargyl alcohol (1.00 mL, 17.0 mmol) and the reaction mixture was stirred for 30 min. 2,3-Dibromopropene (1.50 mL, 14.3 mmol) was slowly added, then the reaction mixture was allowed to warm to rt and stirred for 12 h. The reaction was quenched by addition of water and the aqueous layer was extracted with Et₂O; the combined organic extracts were washed with brine and dried (MgSO₄). The solvent was evaporated in vacuo and the crude product purified by flash chromatography (5:1 pet. ether-ether) to afford the 2-bromo-3-(prop-2-ynyloxy)prop-1-ene as a colourless oil (2.15 g, 12.3 mmol, 86%); Rf 0.26 (5:1 pet. ether-ether); IR (thin film) $\nu_{\rm max}/{\rm cm}^{-1}$ 3300, 2923, 2130, 1639, 1443, 1l62, 1086, 899, 668; ¹H NMR (400 MHz, CDCl₃) $\delta_{\rm H}$ 5.97–5.95 (1H, m), 5.67–5.65 (1H, m), 4.23–4.19 (4H, m), 2.47 (1H, t, J = 2.4 Hz); ¹³C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 128.4, 118.7, 78.8, 75.1, 73.3, 57.0; HRMS (ES^+) calcd for C₆H₇BrO $[M + H]^+$ 173.9681, found 173.9679. Data in accordance with the literature.³¹

(3-(2-Bromoallyloxy)prop-1-ynyl)diethyl(isopropoxy)silane (13). Synthesised from 2-bromo-3-(prop-2-ynyloxy) prop-1-ene (500 mg, 2.86 mmol), LiHMDS (4.30 mL, 4.30 mmol), ClSiEt₂-(NEt₂) (1.00 mL, 4.30 mmol), i-PrOH (875 μ L, 11.4 mmol) and imidazole (778 mg, 11.4 mmol) using general procedure 2. The product was purified by flash chromatography (5 : 1 pet. etherether) to afford **13** as a colourless oil (537 mg, 1.68 mmol, 59%); $R_{\rm f}$ 0.26 (5 : 1 pet. ether–ether); IR (thin film) $\nu_{\rm max}/{\rm cm}^{-1}$ 2969, 1461, 1380, 1240, 1122, 1088, 878, 734; ¹H NMR (400 MHz, CDCl₃) $\delta_{\rm H}$ 5.93 (1H, d, J = 1.7 Hz), 5.64–5.62 (1H, m), 4.23 (2H, s), 4.21 (2H, s), 4.12 (1H, septet, J = 6.1 Hz), 1.17 (6H, d, J = 6.1 Hz), 0.96 (6H, t, J = 7.8 Hz), 0.58 (4H, q, J =7.9 Hz); ¹³C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 128.7, 118.8, 101.3, 89.1, 73.3, 66.3, 57.9, 25.7, 6.8, 6.7; HRMS (ES⁺) calcd for C₁₃H₂₃BrNaO₂Si [M + Na]⁺ 341.0543, found 341.0531.

General procedure 3: cyclisation of bromoenynes to silylcyclohexyldienes

PdCl₂(PPh₃)₂ (0.1 equiv.) was added to a degassed solution of bromoenyne (1 equiv.) and vinyl stannane (1.5–1.7 equiv.) in toluene (20 mL mmol⁻¹) under argon. The reaction mixture was heated to reflux (110 °C) until completion as monitored by TLC and/or ¹H NMR spectroscopic analysis of an aliquot. After concentration, the crude mixture was purified by flash chromatography to obtain the bicyclic diene.

Dimethyl 7-(isopropoxydimethylsilyl)-5-methyl-4,5-dihydro-1H-indene-2,2(3H)-dicarboxylate (15a). Synthesised from 10a (23.9 mg, 0.0590 mmol) and (E)-tributyl(prop-1-enyl)stannane 3a (29.3 mg, 0.0885 mmol) using general procedure 3. The product was purified by flash chromatography (5:1 pet. etherether) to afford 15a as a colourless oil (20 mg, 0.0546 mmol, 92%); R_f 0.33 (5:1 pet. ether-ether); IR (thin film) $\nu_{\text{max}}/\text{cm}^{-1}$ 2358, 1737, 1252, 1024, 890, 822, 781; ¹H NMR (500 MHz, $CDCl_3$) δ_H 5.95 (1H, d, J = 3.3 Hz, H6), 3.95 (1H, septet, J =6.3 Hz, CHMe₂), 3.74 (3H, s, CO₂Me), 3.74 (3H, s, CO₂Me), 3.67-3.65 (1H, m, H5), 3.20 (2H, br s, H3), 3.03-3.01 (1H, m, H4), 2.96–2.94 (1H, m, H'4), 2.46–2.44 (1H, m, H3), 2.20–2.18 (1H, m, H'3), 1.13 (3H, d, J = 6.3 Hz, CHMe₂), 1.12 (3H, d, J = 6.3 Hz, $CHMe_2$, 1.04 (3H, d, J = 7.1 Hz, Me), 0.23 (6H, s, SiMe₂); ¹³C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 172.8, 172.8, 143.9, 132.1, 131.3, 131.1, 65.1, 58.5, 52.7, 52.7, 43.0, 41.7, 31.1, 30.0, 25.6, 20.1, -0.90; HRMS (ES⁺) calcd for C₁₉H₃₀NaO₅Si $[M + Na]^+$ 389.1743, found 389.1755.

Dimethyl 7-(isopropoxydimethylsilyl)-5-phenyl-4,5-dihydro-1H-indene-2,2(3H)-dicarboxylate (15b). Synthesised from 10a (26.6 mg, 0.0656 mmol) and (E)-tributyl(styryl)stannane 3b (38.8 mg, 0.0987 mmol) using general procedure 3. The product was purified by flash chromatography (5:1 pet. etherether) to afford 15b as a colourless oil (27.4 mg, 0.0639 mmol, 97%); $R_{\rm f}$ 0.30 (5:1 pet. ether-ether); IR (thin film) $\nu_{\rm max}/{\rm cm}^{-1}$ 2391, 1737, 1434, 1367, 1252, 1023, 782; ¹H NMR (500 MHz, $CDCl_3$) δ_H 7.30 (2H, m, ArH), 7.24 (3H, m, ArH), 6.10 (1H, d, J = 3.1 Hz, H6), 3.99 (1H, septet, J = 6.2 Hz, CHMe₂), 3.76 (3H, s, CO₂Me), 3.74 (3H, s, CO₂Me), 3.65-3.63 (1H, m, H5), 3.28-3.26 (2H, m, H1), 3.10 (1H, d, J = 17.4 Hz, H4), 2.94 (1H, d, J = 17.4 Hz, H'4), 2.45-2.43 (1H, m, H3), 2.29-2.27 (1H, m, H'3), 1.16 (3H, d, J = 6.2 Hz, CHMe₂), 1.14 (3H, d, J = 6.2 Hz, CHMe₂), 0.25 (6H, s, SiMe₂); ¹³C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 172.7, 145.4, 140.7, 133.4, 131.7, 131.1, 128.5, 127.6, 126.3, 65.2, 58.5, 52.7, 42.9, 41.2, 41.7, 32.0, 25.6, -0.9; HRMS (ES⁺) calcd for $C_{24}H_{32}NaO_5Si [M + Na]^+ 451.1892$, found 451.1911.

Dimethyl 7-(diethyl(isopropoxy)silyl)-5-phenyl-4,5-dihydro-1*H*-indene-2,2(3*H*)-dicarboxylate (15c). Synthesised from 11a (300 mg, 0.670 mmol) and (E)-tributyl(styryl)stannane 3b (435 mg, 1.11 mmol) using general procedure 3. The crude product was purified by flash chromatography (5:1 pet. etherether) to afford 15c as a colourless oil (237 mg, 0.519 mmol, 77%); $R_{\rm f}$ 0.42 (5:1 pet. ether-ether); IR (thin film) $\nu_{\rm max}/{\rm cm}^{-1}$ 2955, 1736, 1434, 1381, 1243, 1005, 874, 730, 699; ¹H NMR (400 MHz, CDCl3) δ_H 7.32-7.30 (2H, m, ArH), 7.27-7.23 (3H, m, ArH), 6.12-6.10 (1H, m, H6), 4.01 (1H, septet, J = 6.2 Hz, CHMe₂), 3.76 (3H, s, CO₂Me), 3.73 (3H, s, CO₂Me), 3.64-3.62 (1H, m, H5), 3.25 (2H, br s, H3), 3.10 (1H, d, J = 17.5 Hz, H1), 2.94 (1H, d, J = 17.5 Hz, H'1), 2.45-2.43 (1H, m, H4), 2.27-2.25 $(1H, m, H'4), 1.16 (3H, d, J = 6.2 Hz, CHMe_2), 1.15 (3H, d, J =$ 6.2 Hz, CHMe2), 1.00-0.95 (6H, m, SiCH2Me), 0.77-0.74 (4H, m, SiCH₂Me); ¹³C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 172.8, 172.7, 145.5, 141.3, 131.8, 131.5, 130.9, 128.5, 127.6, 126.3, 65.2, 58.5, 52.7, 52.7, 42.9, 42.2, 41.8, 32.1, 25.7, 25.6, 6.8, 5.0, 4.9; HRMS (ES^{+}) calcd for $C_{27}H_{40}NaO_6Si [M + MeOH + Na]^{+} 511.2486$, found 511.2432.

7-(Diethyl(isopropoxy)silyl)-5-methyl-2-tosyl-2,3,4,5-tetrahydro-1*H*-isoindole (15d). Synthesised from 12 (50.0 mg, 0.106 mmol) and (*E*)-tributyl(prop-1-enyl)stannane 3a (60.0 mg, 0.181 mmol) using general procedure 3. The product was purified by flash chromatography (5:1 pet. ether-ether) to afford 15d as a colourless oil (29.0 mg, 0.0670 mmol, 63%); $R_{\rm f}$ 0.29 (5:1 pet. ether-ether); IR (thin film) $\nu_{\rm max}/{\rm cm}^{-1}$ 2958, 1436, 1378, 1230, 1015, 874, 730, 699; ¹H NMR (500 MHz, $CDCl_3$) δ_H 7.72 (2H, d, J = 8.0 Hz, ArH), 7.31 (2H, d, J = 8.0 Hz, ArH), 5.93 (1H, d, J = 3.5 Hz, H6), 4.22–4.20 (2H, m, H1), 4.11-4.00 (2H, m, H3), 3.95 (1H, septet, J = 5.9 Hz, $CHMe_2$), 2.42 (4H, s, H5, ArMe), 2.15 (1H, dd, J = 17.0, 8.5 Hz, H5), 1.81-1.73 (1H, m, H'5), 1.13 (3H, d, J = 6.0 Hz, CHMe₂), 1.11 (3H, d, J = 6.0 Hz, CHMe₂), 1.02 (3H, d, J = 7.1 Hz, CHMe), 0.89 (6H, app. td, J = 8.0, 1.7 Hz, SiCH₂Me), 0.68-0.65 (4H, m, SiCH₂Me); ¹³C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 145.1, 143.2, 134.4, 129.9, 129.6, 128.1, 127.8, 127.5, 65.3, 56.3, 55.4, 29.8, 28.7, 25.7, 25.7, 21.5, 19.9, 6.8, 5.0, 5.0; HRMS (ES⁺) calcd for $C_{23}H_{35}NNaO_{3}SSi [M + Na]^{+} 456.1999$, found 456.2000.

Diethyl(isopropoxy)(6-methyl-1,3,6,7-tetrahydroisobenzofuran-4-yl)silane (15e). Synthesised from 13 (50.0 mg, 0.157 mmol) (E)-tributyl(prop-1-enyl)stannane and 3a (88.1)mg, 0.266 mmol) using general procedure 3. The crude product was purified by flash chromatography (5:1 pet. ether-ether) to afford 15e as a colourless oil (10.0 mg, 0.0356 mmol, 23%); $R_{\rm f}$ 0.29 (5:1 pet. ether-ether); IR (thin film) $\nu_{\text{max}}/\text{cm}^{-1}$ 2955, 1732, 1463, 1210, 866; ¹H NMR (500 MHz, $CDCl_3$) δ_H 5.99 (1H, d, J = 3.3 Hz, H5), 4.75-4.73 (2H, m, H1), 4.60-4.58 (2H, m, H3), 3.99 (1H, septet, J = 6.0 Hz, $CHMe_2$), 2.59–2.48 (1H, m, H8), 2.26 (1H, dd, J = 17.0, 9.0 Hz, H7), 1.92–1.84 (1H, m, H7), 1.16 (3H, d, J = 6.0 Hz, CHMe₂), 1.15 (3H, d, J = 6.0 Hz, CHMe₂), 1.09 (3H, d, J = 7.0 Hz, CHMe), 0.94 (6H, t, J = 7.9 Hz, SiCH₂Me), 0.75-0.65 (4H, m, SiCH₂Me); ¹³C NMR (125 MHz, CDCl₃) δ_C 144.6, 131.3, 129.7, 127.4, 76.3, 75.7, 65.2, 29.9, 27.7, 25.7, 20.1, 6.7, 4.9; HRMS (ES⁺) calcd for C₁₆H₂₈NaO₂Si $[M + Na]^+$ 303.1751, found 303.1749.

Dimethyl 5-(diethyl(isopropoxy)silyl)-7-methyl-3,4,7,8-tetrahydronaphthalene-2,2(1*H*)-dicarboxylate (15f). Synthesised from 11b (100 mg, 0.223 mmol) and (E)-tributyl(prop-1-enyl)stannane 3a (126 mg, 0.381 mmol) using general procedure 3. The product was purified by flash chromatography (5:1 pet. ether-ether) to afford 15f as a colourless oil (68.0 mg, 0.166 mmol, 75%); Rf 0.52 (5:1 pet. ether-ether); IR (thin film) $\nu_{\rm max}/{\rm cm}^{-1}$ 2956, 1735, 1435, 1250, 1170, 1121, 1083, 1018, 872, 731; ¹H NMR (500 MHz, CDCl_3) δ_{H} 5.99 (1H, d, J = 3.4 Hz, H6), 3.94 (1H, sept, J = 6.1 Hz, OCHMe₂), 3.72 (3H, s, CO₂Me), 3.71 (3H, s, CO₂Me), 2.60 (2H, m, H1), 2.31-2.29 (3H, m, H4 and H7), 2.24–2.20 (1H, m, H3), 2.12 (1H, dt, J = 13.4, 6.3 Hz, H3), 1.98 (1H, dd, J = 16.0, 7.5 Hz, H8), 1.80-1.69 (1H, m, H8), 1.12 (6H, t, J = 6.1 Hz, OCHMe₂), 1.01 (3H, d, J = 6.8Hz, C11), 0.95-0.90 (6H, m, SiCH₂Me), 0.75-0.72 (4H, m, SiCH₂Me); ¹³C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 172.4, 172.0, 144.9, 134.1, 127.8, 125.7, 65.2, 53.5, 52.7, 52.7, 36.4, 36.1, 29.5, 28.5, 25.8, 25.8, 25.2, 19.6, 7.0, 7.0, 5.7, 5.6; HRMS (ES⁺) calcd for $C_{22}H_{36}NaO_5Si [M + Na]^+$ 431.2224, found 431.2236. NMR data for 16a and 17a was obtained from a purified mixture of 16a and 17a (2.8:1 ratio).

Dimethyl (*Z*)-4-((*E*)-1-(diethyl(isopropoxy)silyl)but-2-en-1ylidene)-3-methylenecyclohexane-1,1-dicarboxylate (16a). ¹H NMR (500 MHz, CDCl₃) $\delta_{\rm H}$ 6.00 (1H, dd, *J* = 15.5, 1.5 Hz, H8), 5.32 (1H, dq, *J* = 15.5, 6.5 Hz, H9), 5.01 (1H, d, *J* = 2.3 Hz, H11), 4.87–4.83 (1H, m, H11), 4.00 (1H, sept, *J* = 6.0 Hz, CHMe₂), 3.72 (6H, s, CO₂*Me*), 2.81 (2H, s, H2), 2.58–2.52 (2H, m, H5), 2.11–2.06 (2H, m, H6), 1.74 (3H, dd, *J* = 6.5, 1.5 Hz, H10), 1.13 (6H, d, *J* = 6.0 Hz, CH*Me*₂), 0.91 (6H, t, *J* = 7.5 Hz, SiC*H*₂Me), 0.69 (4H, *J* = 7.5 Hz, SiCH₂*Me*); ¹³C NMR (126 MHz, CDCl₃) $\delta_{\rm C}$ 171.2, 151.58, 146.1, 132.1, 132.1, 127.6, 115.1, 65.2, 57.1, 52.8, 41.2, 32.3, 30.3, 25.9, 18.7, 7.3, 6.7.

Dimethyl (*E*)-3-(1-(diethyl(isopropoxy)silyl)buta-1,3-dien-1yl)-4-methylcyclopent-3-ene-1,1-dicarboxylate (17a). ¹H NMR (500 MHz, CDCl₃) $\delta_{\rm H}$ 6.36 (1H, d, *J* = 10.6 Hz, H8), 6.31 (1 H, ddd, *J* = 16.5, 10.6, 9.7 Hz, H9), 5.26 (1H, dd, *J* = 16.5, 2.0 Hz, H10), 5.13 (1H, dd, *J* = 9.7, 2.0, H10), 4.00 (1H, sept, *J* = 6.0 Hz, CHMe₂), 3.73 (6H, s, CO₂*Me*), 2.61 (1H, d, *J* = 17.2 Hz, H2), 2.45 (1H, d, *J* = 17.2 Hz, H2), 2.24–2.18 (2H, m, H6 and H5), 2.14–2.11 (1H, m, H6), 1.87–1.82 (1H, m, H5), 1.50 (3H, s, H11), 1.14 (6H, d, *J* = 6.0 Hz, CHMe₂), 0.91 (6H, t, *J* = 7.5 Hz, SiCH₂Me), 0.69 (4H, *J* = 7.5 Hz, SiCH₂Me); ¹³C NMR (126 MHz, CDCl₃) $\delta_{\rm C}$ 172.4, 172.2, 144.5, 140.2, 134.4, 130.5, 123.2, 118.9, 65.2, 53.9, 52.7, 52.7, 36.3, 28.3, 27.7, 26.0, 25.9, 20.3, 7.0, 6.8, 5.5, 5.3.

Dimethyl 4-(diethyl(isopropoxy)silyl)-2-methyl-5,6,7,9-tetrahydro-1*H*-benzo[7]annulene-8,8(2*H*)-dicarboxylate (15g). Synthesised from 11c (51.4 mg, 0.111 mmol) and (*E*)-tributyl-(prop-1-enyl)stannane 3a (55.0 mg, 0.166 mmol) using general procedure 3. The crude product was purified by flash chromatography (5:1 pet. ether–ether) to afford 15g as a colourless oil (27.5 mg, 0.0651 mmol, 58%, mixture with 16b and 17b); *R*_f 0.52 (5:1 pet. ether–ether); IR (thin film) ν_{max}/cm^{-1} 2954, 1736, 1455, 1228, 1172, 1007, 873, 729; ¹H NMR (500 MHz, CDCl₃) δ_H 5.98 (1H, d, *J* = 3.2 Hz, H3), 3.95 (1H, septet, *J* = 5.9 Hz, *CHM*e₂), 3.71 (3H, s, CO₂*Me*), 3.70 (3H, s, CO₂*Me*), 2.49–2.44 (2H, m, H9), 2.29–2.20 (2H, m, H5), 2.17–2.04 (3H, m, H5, H2, H1), 1.78–1.72 (1H, m, H1), 1.68–1.60 (2H, m, H7), 1.38–1.33 (2H, m, H6), 1.13 (3H, d, J = 6.0, CH Me_2), 1.13 (3H, d, J = 6.0, CH Me_2), 0.99 (3H, d, J = 7.0 Hz, CHMe), 0.96–0.87 (6H, m, SiCH₂Me), 0.76–0.60 (4H, m, SiCH₂Me); ¹³C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 172.3, 172.0, 145.3, 137.8, 135.6, 129.2, 65.1, 55.0, 52.3, 52.1, 39.7, 39.1, 37.2, 31.4, 29.8, 28.6, 26.7, 25.6, 25.4, 22.7, 19.4, 17.3, 13.6, 6.9, 6.8, 5.2, 5.1; HRMS (ES⁺) calcd for C₂₃H₃₈NaO₅Si [M + Na]⁺ 445.2381, found 445.2386.

Dimethyl 8-(trimethylsilyl)bicyclo[5.2.0]nona-1,7-diene-3,3dicarboxylate (30). To Pd(PPh₃)₄ (5.0 mg, 0.0045 mmol, 5 mol%) was added a degassed solution of bromoenyne 27^{8a} (35.0 mg, 0.0899 mmol) and Et₃N (25.0 µl, 0.180 mmol) in toluene (1.2 mL) under Ar, and the mixture was refluxed in a preheated oil bath (110 °C) until completion of the reaction (16 h). The mixture was concentrated, and the residue purified by flash chromatography (10:1 pet. ether-Et₂O) to obtain 30 as a colourless oil (25.0 mg, 0.0811 mmol, 90%); Rf 0.46 (2:1 pet. ether-Et₂O); IR (thin film) $\nu_{\text{max}}/\text{cm}^{-1}$ 2954, 1736, 1248, 840; ¹H NMR (500 MHz, MeOD) $\delta_{\rm H}$ 4.99 (1H, s, H2), 3.71 (6H, s, CO_2Me), 2.73 (2H, t, J = 2.9 Hz, H9), 2.38–2.32 (2H, m, H6), 2.29-2.25 (2H, m, H4), 1.67 (2H, app. qd, J = 6.0, 4.5 Hz, H5), 0.12 (9H, s, SiMe); ¹³C NMR (126 MHz, MeOD) $\delta_{\rm C}$ 173.1, 159.9, 150.9, 145.3, 110.9, 63.2, 53.3, 36.4, 35.7, 31.9, 23.6, -1.7; HRMS (ES+) calcd for $C_{16}H_{24}NaO_4Si [M + Na]^+ 331.1336$, found 331.1340.

8-(diethyl(isopropoxy)- λ^4 -sulfanyl)bicyclo[5.2.0]-Dimethyl nona-1,7-diene-3,3-dicarboxylate (36). To Pd(PPh₃)₄ (5.0 mg, 0.0045 mmol, 5 mol%) was added a degassed solution of bromoenyne 11c (32.0 mg, 0.0693 mmol) and Et₃N (20.0 µl, 0.144 mmol) in toluene (1.2 mL) under Ar, and the mixture was heated in a preheated oil bath (110 °C) until completion of the reaction (16 h). The mixture was concentrated, and the residue was purified by flash chromatography (10:1 pet. ether-Et2O) to obtain 36 as a colourless oil (23.0 mg, 0.0604 mmol, 87%); Rf 0.24 (9:1 pet. ether-Et₂O); IR (thin film) $\nu_{\text{max}}/\text{cm}^{-1}$ 2955, 1736, 1565, 1243, 1031; ¹H NMR (500 MHz, CDCl₃) $\delta_{\rm H}$ 5.09 (1H, s, H2), 3.99 (1H, sept, J = 6.0 Hz, SiOCHMe), 3.74 (6H, s, CO₂Me), 2.84 (2H, t, J = 2.9 Hz, H9), 2.39-2.35 (2H, m, H6), 2.33-2.30 (2H, m, H4), 1.71 (2H, dt, J = 10.8, 5.6 Hz, H5), 1.15 (6H, d, J = 6.0 Hz, SiOCHMe), 0.96 (6H, t, J = 7.9 Hz, SiCH₂Me), 0.68 (4H, q, J = 7.9 Hz, SiCH₂Me); ¹³C NMR (101 MHz, CDCl₃) $\delta_{\rm C}$ 171.8, 160.6, 146.9, 144.5, 110.5, 65.6, 62.0, 52.9, 36.9, 34.7, 31.5, 26.0, 22.6, 6.8, 5.7; HRMS (ESI⁺) calcd for $C_{20}H_{32}NaO_5Si [M + Na]^+$ 403.1911, found 403.1916.

Dimethyl 6-methyl-4-oxo-4,5,6,7-tetrahydro-1*H***-indene-2,2-**(*3H*)-**dicarboxylate (37). 15a** (16.8 mg, 0.0458 mmol) was dissolved in THF–MeOH (1 : 1, 400 µL). KF (5.3 mg, 0.090 mmol), KHCO₃ (9.2 mg, 0.090 mmol), H₂O₂ (30% aq., 5.2 µL, 0.050 mmol) were added and the reaction mixture was heated to 40 °C for 3 h. The reaction was allowed to cool to rt, then it was diluted with CH₂Cl₂ and water. The organic extract was washed with water, dried (MgSO₄), and concentrated. The product was purified by flash chromatography (2 : 1 pet. etherether) to afford 37 as a colourless oil (5.3 mg, 0.020 mmol, 43%); $R_{\rm f}$ 0.21 (2 : 1 pet. ether–ether); IR (thin film) $\nu_{\rm max}/{\rm cm}^{-1}$ 2349, 1734, 1667, 1434, 1262, 913, 745; ¹H NMR (400 MHz, CDCl₃) $\delta_{\rm H}$ 3.75 (6H, s, CO₂*Me*), 3.23–3.21 (4H, m, H1, H3), 2.42 (1H, dd, *J* = 17.0, 3.9 Hz, H5), 2.42–2.36 (1H, m, H7), 2.32–2.20 (1H, m, H6), 2.09 (1H, dd, *J* = 17.0, 12.0 Hz, H5), 2.07–2.00 (1H, m, H7), 1.07 (3H, d, *J* = 6.5 Hz, *Me*); ¹³C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 194.9, 172.0 (2C), 158.7, 134.8, 57.8, 52.5 (2C), 45.8, 45.2, 38.6, 33.8, 31.0, 21.0; HRMS (ES⁺) calcd for C₁₄H₁₉O₅ [M + H]⁺ 267.1227, found 267.1225.

Dimethyl 4-(diethyl(isopropoxy)silyl)-6-phenyl-1H-indene-2,2(3H)-dicarboxylate (38). To a solution of 15c (20.0 mg, 0.0438 mmol) in CH₂Cl₂ (1 mL) was added MnO₂ (19.0 mg, 0.220 mmol) and the reaction mixture was stirred at 30 °C for 12 h. The mixture was filtered through a pad of Celite and concentrated, and the residue was purified by flash chromatography (5:1 pet. ether-ether) to afford 38 as a colourless oil (17.0 mg, 0.0374 mmol, 85%); R_f 0.31 (5:1 pet. ether-ether); IR (thin film) $\nu_{\text{max}}/\text{cm}^{-1}$ 2923, 2853, 1740, 1460, 699; ¹H NMR $(500 \text{ MHz}, \text{CDCl}_3) \delta_H$ 7.60–7.40 (6H, m, H5, H7, ArH), 7.33 (1H, t, J = 7.3 Hz, ArH), 4.08 (1H, septet, J = 6.1 Hz, CHMe₂), 3.76 (6H, s, CO₂Me), 3.74 (2H, s, H1 or H3), 3.63 (2H, s, H1 or H3), 1.19 (6H, d, J = 6.1 Hz, CHMe₂), 1.03–0.93 (4H, m, SiCH₂Me); 13 C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 172.2, 144.6, 141.5, 139.9, 139.4, 132.7, 132.5, 128.7, 127.1, 127.0, 124.3, 65.5, 60.4, 52.9, 41.4, 40.1, 25.8, 6.8, 5.5; HRMS (ES⁺) calcd for C₂₆H₃₄NaO₅Si $[M + Na]^+$ 477.2068, found 477.2053.

4-(Diethyl(isopropoxy)silyl)-6-methyl-2-tosylisoindoline (39). To a solution of **15d** (29.0 mg, 0.0669 mmol) in CH₂Cl₂ (1 mL) was added MnO₂ (50.0 mg, 0.460 mmol) and the reaction mixture was stirred at 30 °C for 12 h. The mixture was filtered through a pad of Celite and concentrated, and the residue was purified by flash chromatography (5:1 pet. ether-ether) to afford **39** as a colourless oil (27.0 mg, 0.0625 mmol, 93%); $R_{\rm f}$ 0.30 (5:1 pet. ether-ether); IR (thin film) $\nu_{\text{max}}/\text{cm}^{-1}$ 2958, 1733, 1596, 1433, 1259, 699; ¹H NMR (500 MHz, CDCl₃) $\delta_{\rm H}$ 7.76 (2H, d, J = 8.3 Hz, ArH), 7.29 (2H, d, J = 8.3 Hz, ArH), 7.14 (1H, s, H5), 6.99 (1H, s, H7), 4.67 (2H, s, H3), 4.56 (2H, s, H1), 4.05 (1H, sept, J = 6.3 Hz, $CHMe_2$), 2.40 (3H, s, Ts-Me), 2.31 (3H, s, Me), 1.18 (6H, d, J = 6.3 Hz, CHMe₂), 0.92 (6H, t, J = 6.8 Hz, SiCH₂Me), 0.88–0.82 (4H, m, SiCH₂Me); ¹³C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 143.4, 138.8, 136.3, 135.6, 134.4, 133.7, 131.1, 129.7, 127.6, 124.2, 65.4, 54.6, 53.3, 25.7, 21.4, 21.2, 6.8, 5.5; HRMS (ES⁺) calcd for $C_{23}H_{33}NNaO_3SSi [M + Na]^+$ 454.1843, found 454.1845.

Dimethyl 4-hydroxy-6-phenyl-1*H*-indene-2,2(3*H*)-dicarboxylate (40). To a solution of 38 (17.0 mg, 0.0374 mmol) in THF-MeOH (1 : 1, 1 mL) was added and TBAF (1 M solution in THF, 13.0 μL, 0.013 mmol), KHCO₃ (7.5 mg, 0.080 mmol) and H₂O₂ (30% aq., 13 μL, 0.12 mmol). The reaction mixture was heated to 60 °C for 3 h, then it was cooled to rt, diluted with CH₂Cl₂ and water. The organic phase was washed with water, dried (MgSO₄), and concentrated. The residue was purified by flash chromatography (2 : 1 pet. ether–ether) to afford 40 (9.4 mg, 0.0288 mmol, 78%); $R_{\rm f}$ 0.15 (2 : 1 pet. ether–ether); IR (thin film) $\nu_{\rm max}/{\rm cm}^{-1}$ 2954, 1733, 1596, 1434, 1260, 1165, 864, 699; ¹H NMR (500 MHz, CDCl₃) $\delta_{\rm H}$ 7.52 (2H, d, *J* = 7.5 Hz, ArH), 7.41 (2H, t, *J* = 7.5 Hz, ArH), 7.32 (1H, t, *J* = 7.5 Hz, ArH), 7.01 (1H, s, H7), 6.85 (1H, s, H5), 3.78 (6H, s, CO₂*Me*), 3.66 (2H, s, H1), 3.60 (2H, s, H3); 13 C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 172.0, 151.8, 142.8, 142.5, 140.9, 128.7, 127.2, 127.1, 124.7, 115.6, 112.6, 60.3, 53.0, 40.9, 36.8; HRMS (ES⁻) calcd for C₁₉H₁₇O₅ [M]⁻ 325.1081, found 325.1081.

6-Methyl-2-tosylisoindolin-4-ol (41). To a solution of 39 (20.0 mg, 0.0463 mmol) in THF-MeOH (1:1, 1 mL) was added and TBAF (1 M solution in THF, 16.0 µL, 0.016 mmol), KHCO₃ (9.0 mg, 0.10 mmol) and H₂O₂ (30% aq., 17 µL, 0.15 mmol). The reaction mixture was heated to 60 °C for 3 h, then it was cooled to rt, diluted with CH₂Cl₂ and water. The organic phase was washed with water, dried (MgSO₄), and concentrated. The residue was purified by flash chromatography (2:1 pet. etherether) to afford 41 as a colourless oil (14.0 mg, 0.0461 mmol, 99%); $R_{\rm f}$ 0.26 (2:1 pet. ether-ether); IR (thin film) $\nu_{\rm max}/{\rm cm}^{-1}$ 2954, 1733, 1596, 1434, 1260, 1165, 864, 699; ¹H NMR $(500 \text{ MHz}, \text{CDCl}_3) \delta_H$ 7.76 (2H, d, J = 7.7 Hz, ArH), 7.31 (2H, d, J = 7.7 Hz, ArH), 6.54 (1H, s, H7), 6.47 (1H, s, H5), 4.57 (4H, s, H1, H3), 2.40 (3H, s, TsMe), 2.24 (3H, s, Me); ¹³C NMR (125 MHz, CDCl₃) $\delta_{\rm C}$ 150.6, 143.6, 139.8, 138.2, 133.6, 129.8, 127.6, 119.7, 115.2, 114.8, 54.0, 51.4, 21.5, 21.2; HRMS (ES⁺) calcd for $C_{16}H_{17}NNaO_3S [M + Na]^+$ 326.0821, found 326.0824.

Acknowledgements

We thank the EPSRC (Advanced Research Fellowship to E.A.A.; EP/E055273/1, studentship to M.-C.A.C.), the German Academic Exchange Service (DAAD) for a postdoctoral fellowship (B.G.), and A*STAR for a National Science Scholarship (S.S.G.).

Notes and references

- (a) J. H. P. Tyman, Synthetic and Natural Phenols, Elsevier, New York, 1996; (b) Z. Rappoport, The Chemistry of Phenols, Wiley-VCH, Weinheim, 2003; (c) I. P. Singh, S. B. Bharate and K. K. Bhutani, Curr. Sci., 2005, 89, 269–290.
- 2 (a) For examples of tetracycline and pyranonaphthoquinone natural products, see: A. G. Myers, D. A. Kummer, D. R. Li and A. Dion, *Chem. Sci.*, 2011, 2, 1710–1718;
 (b) A. G. Myers, C. X. Sun, Q. Wang, J. D. Brubaker, P. M. Wright, C. D. Lerner, K. Noson, M. Charest, D. R. Siegel and Y. M. Wang, *J. Am. Chem. Soc.*, 2008, 130, 17913–17927; (c) G. Minotti, P. Menna, E. Salvatorelli, G. Cairo and L. Gianni, *Pharmacol. Rev.*, 2004, 56, 185–229; (d) J. Sperry, P. Bachu and M. A. Brimble, *Nat. Prod. Rep.*, 2008, 25, 376–400.
- 3 For examples, see ref. 1*a* and: P. Hanson, J. R. Jones, A. B. Taylor, P. H. Walton and A. W. Timms, *J. Chem. Soc., Perkin Trans.* 2, 2002, 1135–1150.
- 4 (a) For reviews of C-X phenolation, see: M. C. Willis, Angew. Chem., Int. Ed., 2007, 46, 3402–3404; (b) S. Enthaler and A. Company, Chem. Soc. Rev., 2011, 40, 4912–4924; (c) For reviews of C-H phenolation, see: D. A. Alonso, C. Najera, I. M. Pastor and M. Yus, Chem. – Eur. J., 2010,

16, 5274–5284; (*d*) M. S. Sanford and T. W. Lyons, *Chem. Rev.*, 2010, **110**, 1147–1169.

- 5 (a) The equivalent oxidation of arylboranes can also provide phenols, but in most cases these derivative are less amenable to processing through multistep synthesis. See:
 R. E. Maleczka, F. Shi, D. Holmes and M. R. Smith, *J. Am. Chem. Soc.*, 2003, 125, 7792–7793; (b) G. A. Molander and L. N. Cavalcanti, *J. Org. Chem.*, 2011, 76, 623–630; (c) X. Y. Wang, J. M. Xu, C. W. Shao, D. Y. Su, G. L. Cheng and Y. F. Hu, *Org. Lett.*, 2010, 12, 1964–1967.
- 6 (*a*) S. Bracegirdle and E. A. Anderson, *Chem. Commun.*, 2010, **46**, 3454–3456; (*b*) E. J. Rayment, N. Summerhill and E. A. Anderson, *J. Org. Chem.*, 2012, 77, 7052–7060.
- 7 (a) For other examples of arylsilane oxidation, see:
 J. F. Hartwig and E. M. Simmons, J. Am. Chem. Soc., 2010,
 132, 17092–17095; (b) M. Suginome and H. Ihara, J. Am. Chem. Soc., 2009, 131, 7502–7503; (c) J. D. Sunderhaus,
 H. Lam and G. B. Dudley, Org. Lett., 2003, 5, 4571–4573; (d) K. Tamao, Proc. Jpn. Acad. B, Phys. Biol. Sci., 2008, 84, 123–133; (e) K. Tamao, T. Kakui, M. Akita, T. Iwahara,
 R. Kanatani, J. Yoshida and M. Kumada, Tetrahedron, 1983, 39, 983–990; (f) For a borodesilylation/oxidation sequence, see: V. Gevorgyan, C. H. Huang, N. Chernyak and
 A. S. Dudnik, Adv. Synth. Catal., 2011, 353, 1285–1305.
- 8 (a) S. B. J. Kan and E. A. Anderson, *Org. Lett.*, 2008, 10, 2323–2326; (b) M.-C. A. Cordonnier, S. B. J. Kan and E. A. Anderson, *Chem. Commun.*, 2008, 5818–5820.
- 9 (a) J. Suffert, B. Salem and P. Klotz, J. Am. Chem. Soc., 2001, 123, 12107–12108; (b) B. Salem, P. Klotz and J. Suffert, Org. Lett., 2003, 5, 845-848; (c) B. Salem, E. Delort, P. Klotz and J. Suffert, Org. Lett., 2003, 5, 2307-2310; (d) B. Salem and J. Suffert, Angew. Chem., Int. Ed., 2004, 43, 2826-2830; (e) C. Bour, G. Blond, B. Salem and J. Suffert, Tetrahedron, 2006, 62, 10567-10581; (f) For the most recent publication on this topic, and leading references, see: M. Charpenay, A. Boudhat, C. Hulot, G. Blond and J. Suffert, Tetrahedron, 2013, 69, 7568–7591; (g) For seminal work on alkyne carbopalladation followed by Stille cross-coupling, see: B. Burns, R. Grigg, P. Ratananukul, V. Sridharan, P. Stevenson, S. Sukirthalingam and T. Worakun, Tetrahedron Lett., 1988, 29, 5565-5568; (h) E. Negishi, Y. Noda, F. Lamaty and E. J. Vawter, Tetrahedron Lett., 1990, 31, 4393-4396; (i) J. M. Nuss, M. M. Murphy, R. A. Rennels, M. M. Heravi and B. J. Mohr, Tetrahedron Lett., 1993, 34, 3079-3082; (j) P. Fretwell, R. Grigg, J. M. Sansano, V. Sridharan, S. Sukirthalingam, D. Wilson and J. Redpath, Tetrahedron, 2000, 56, 7525-7539; (k) For a review of palladium-catalyzed cascade cyclizations, see: T. Vlaar, E. Ruijter and R. V. A. Orru, Adv. Synth. Catal., 2011, 353, 809-841; (l) For a general review of carbopalladation chemistry, see: E. Negishi, C. Coperet, S. Ma, S. Y. Liou and F. Liu, Chem. Rev., 1996, 96, 365-394.
- 10 Suffert has also noted the utility of silylalkynes in similar cascades. See: G. Blond, C. Bour, B. Salem and J. Suffert, *Org. Lett.*, 2008, **10**, 1075–1078.

- 11 (a) For recent reviews and leading references, see: Y. Nakao and T. Hiyama, *Chem. Soc. Rev.*, 2011, 40, 4893-4901;
 (b) S. E. Denmark and J. H. C. Liu, *Angew. Chem., Int. Ed.*, 2010, 49, 2978-2986.
- 12 S. S. Goh, H. Baars, B. Gockel and E. A. Anderson, *Org. Lett.*, 2012, **14**, 6278–6281.
- 13 (a) For other de novo routes to fused-ring phenols, see:
 A. S. K. Hashmi, T. Haffner, M. Rudolph and F. Rominger, *Chem. – Eur. J.*, 2011, 17, 8195–8201; (b) A. S. K. Hashmi,
 M. Rudolph, J. W. Bats, W. Frey, F. Rominger and T. Oeser, *Chem. – Eur. J.*, 2008, 14, 6672–6678; (c) A. S. K. Hashmi,
 R. Salathe and W. Frey, *Chem. – Eur. J.*, 2006, 12, 6991– 6996; (d) K. Hashmi, A. Stephen, T. M. Frost and J. W. Bats, *J. Am. Chem. Soc.*, 2000, 122, 11553–11554; (e) K. Tanaka,
 H. Hara and M. Hirano, *Org. Lett.*, 2009, 11, 1337–1340; (f) S. Akai, T. Ikawa, S. I. Takayanagi, Y. Morikawa,
 S. Mohri, M. Tsubakiyama, M. Egi, Y. Wada and Y. Kita, *Angew. Chem., Int. Ed.*, 2008, 47, 7673–7676; (g) K. Yoshida, R. Narui and T. Imamoto, *Chem. – Eur. J.*, 2008, 14, 9706– 9713.
- 14 L. H. Li, D. Wang and T. H. Chan, Organometallics, 1994, 13, 1757–1761.
- 15 (a) Diethylaminochlorodiethylsilane was prepared according to the procedure of Tamao: K. Tamao, A. Kawachi and Y. Ito, *Organometallics*, 1993, 12, 580–582; (b) For an alternative preparation, see: J. Beignet, P. J. Jervis and L. R. Cox, *J. Org. Chem.*, 2008, 73, 5462–5475.
- 16 (a) R. L. Miller, S. V. Maifeld and D. Lee, Org. Lett., 2004, 6, 2773–2776; (b) S. Chang, M. Lee and S. Ko, J. Am. Chem. Soc., 2000, 122, 12011–12012.
- 17 See the ESI† for NMR spectra of a purified mixture containing predominantly **16a** and **17a**.
- 18 (a) C. Amatore, S. Bensalem, S. Ghalem and A. Jutand, J. Organomet. Chem., 2004, 689, 4642–4646; (b) B. H. Lipshutz and A. Krasovskiy, Org. Lett., 2011, 13, 3818–3821.
- 19 M. Murakami, K. Itami and Y. Ito, J. Am. Chem. Soc., 1997, 119, 7163–7164.

- 20 A reductive elimination to the *syn*-triene followed by rapid electrocyclization cannot be ruled out; however, we have never observed such intermediates.
- 21 J. Q. Yu, M. J. Gaunt and J. B. Spencer, J. Org. Chem., 2002, 67, 4627–4629.
- 22 X. Wang, H. Chakrapani, J. W. Madine, M. A. Keyerleber and R. A. Widenhoefer, *J. Org. Chem.*, 2002, **67**, 2778– 2788.
- 23 An intermediate of this type has also been proposed in the mechanistically distinct palladium(II)-catalyzed cyclization of an allenynoate to give a fused cyclobutene. See: C. H. Oh, D. I. Park, S. H. Jung, V. R. Reddy, A. K. Gupta and Y. M. Kim, *Synlett*, 2005, 2092–2094.
- 24 (a) A. Furstner, A. Schlecker and C. W. Lehmann, *Chem. Commun.*, 2007, 4277–4279; (b) N. Saito, Y. Tanaka and Y. Sato, *Org. Lett.*, 2009, 11, 4124–4126.
- 25 (a) For a Mo-catalyzed route, see: Q. Shen and G. B. Hammond, J. Am. Chem. Soc., 2002, 124, 6534–6535;
 (b) For examples of syntheses via thermal [2 + 2] cycloaddition, see: K. M. Brummond and D. T. Chen, Org. Lett., 2005, 7, 3473–3475; (c) H. Ohno, T. Mizutani, Y. Kadoh, A. Aso, K. Miyamura, N. Fujii and T. Tanaka, J. Org. Chem., 2007, 72, 4378–4389; (d) C. Mukai, Y. Hara, Y. Miyashita and F. Inagaki, J. Org. Chem., 2007, 72, 4454–4461.
- 26 K. Tamao, M. Kumada and K. Maeda, *Tetrahedron Lett.*, 1984, 25, 321–324.
- 27 R. L. Greenaway, C. D. Campbell, O. T. Holton,
 C. A. Russell and E. A. Anderson, *Chem. Eur. J.*, 2011, 17, 14366–14370.
- 28 We found it possible to use a sub-stoichiometric quantity of TBAF to effect these oxidations. See ref. 6*b*.
- 29 M. G. Organ, J. T. Cooper, L. R. Rogers, F. Soleymanzadeh and T. Paul, *J. Org. Chem.*, 2000, **65**, 7959–7970.
- 30 F. Hoffmann, J. Wagler and G. Roewer, *Eur. J. Inorg. Chem.*, 2010, 1133–1142.
- 31 P. J. Parsons, P. Willis and S. C. Eyley, *Tetrahedron*, 1992, 48, 9461–9472.