



Cite this: *Catal. Sci. Technol.*, 2015, 5, 274

Solvent-free iridium-catalyzed CO₂ hydrosilylation: experiments and kinetic modeling†

E. A. Jaseer,^a Muhammad N. Akhtar,^a Mogahid Osman,^a A. Al-Shammari,^a Habeebullah B. Oladipo,^a Karin Garcés,^b Francisco J. Fernández-Alvarez,^{*b} Sulaiman Al-Khattaf^{*a} and Luis A. Oro^{*ab}

The iridium(III) complex [Ir(H)(CF₃SO₂)(NSiN)(coe)] (NSiN = bis(pyridine-2-yloxy)methylsilyl, coe = cyclooctene) has been demonstrated to be an active catalyst for the solvent-free hydrosilylation of CO₂ with 1,1,1,3,5,5,5-heptamethyltrisiloxane (HMTS) under mild reaction conditions (3 bar). The activity of this catalytic system depends on the reaction temperature. The best catalytic performance has been achieved at 75 °C. A kinetic study at variable temperature (from 25 °C to 75 °C) and constant pressure (3 bar) together with kinetic modeling has been carried out. The results from such a study show an activation energy of 73.8 kJ mol⁻¹ for the process.

Received 24th June 2014,
Accepted 30th July 2014

DOI: 10.1039/c4cy00815d

www.rsc.org/catalysis


1. Introduction

The development of sustainable processes for carbon dioxide (CO₂) reduction is a topic of general interest for the scientific community.¹ In particular, catalytic CO₂ hydrosilylation represents a thermodynamically favorable process for the transformation of CO₂ into value-added chemicals,² which also presents the advantages that some hydrosilanes are easy to handle, readily available and environmentally harmless.³ This CO₂ fixation method allows formation of silylformates which have been successfully used as C1 building blocks in the synthesis of formic acid,⁴ formamides,⁵ amines,⁶ and methane.⁷ Selectivity is a relevant parameter to take into account when new catalysts for CO₂ hydrosilylation to silylformates have to be designed. This is due to the reactivity of silylformates, which under the reaction conditions could be hydrosilylated to afford the corresponding bis(silyl)acetal, which could react further with excess hydrosilane present in the reaction mixture to give methoxysilane and (R₃Si)₂O (Scheme 1).⁸ Additionally, the reaction of the corresponding methoxysilane with one additional equivalent of hydrosilane could lead to the formation of methane and (R₃Si)₂O (Scheme 1).⁷ It is worth mentioning that some examples of transition metal catalysts

as well as organocatalysts which selectively catalyze the hydrosilylation of CO₂ to silylformates have been recently reported.^{4b,9-12} Among them, the phosphane-copper catalysts, using 1,4-dioxane as solvent, reported by Baba and co-workers are the most active known so far.⁴

On the other hand, the large-scale use of volatile organic solvents in industrial processes contributes to the impact of human activity on the environment.^{13,14} Therefore, the development of solvent-free synthetic and catalytic processes has become of wide interest for chemists.¹⁵ To the best of our knowledge only two solvent-free catalytic systems effective for the hydrosilylation of CO₂ to silylformate have been reported.^{12,16}

In this context, it has been previously communicated that the complex [Ir{SiMe(OSiMe₃)₂}(CF₃SO₃)(NSiN)(NCMe)] (**1**) (NSiN = bis(pyridine-2-yloxy)methylsilyl, coe = cyclooctene) catalyzes the selective formation of silylformate by solvent-free hydrosilylation of CO₂ with 1,1,1,3,5,5,5-heptamethyltrisiloxane (HMTS) under mild reaction conditions (r.t. and 3 bar). Using complex **1** as the catalyst precursor, an



Scheme 1 Possible transformations of *in situ* generated silylformate during catalytic CO₂ hydrosilylation processes.

^a Center of Research Excellence in Petroleum Refining & Petrochemicals, King Fahd University of Petroleum & Minerals, 31261 Dhahran, Saudi Arabia. E-mail: skhattaf@kfupm.edu.sa

^b Departamento de Química Inorgánica, Instituto de Síntesis Química y Catálisis Homogénea (ISQCH), Universidad de Zaragoza-CSIC, Facultad de Ciencias 50009, Zaragoza, Spain. E-mail: oro@unizar.es, paco@unizar.es

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c4cy00815d



activation step consisting of the replacement of the acetonitrile molecule by one molecule of HMTS to yield the corresponding active intermediate **2** containing a η^2 -(H-Si) coordinated HMTS molecule is required (Scheme 2).¹⁶ Theoretical studies at the DFT level showed that the Si-H bond activation is a triflate ligand-assisted process *via* TS-1 and that the activation of the CO₂ molecule proceeds *via* an outer-sphere mechanism through the transition state TS-2 (Scheme 3).¹⁶

It is worth noting that the yield of the reaction of the synthesis of **1** from **3** (Scheme 2) is relatively low due to purification difficulties. Subsequently, the use of an alternative catalyst precursor is of interest. In this regard, preliminary ¹H NMR studies of the reaction of **3** with HMTS in CD₂Cl₂ in the absence of acetonitrile showed the formation of cyclooctane together with a mixture of unidentified iridium-containing compounds. Interestingly, when the same reaction was carried out using a mixture of acetonitrile-d₃ and CD₂Cl₂ as solvents ¹H NMR studies evidenced the formation of cyclooctane and **1** (see the ESI[†]). These results are consistent with the formation of intermediate **4** which coordinates a molecule of acetonitrile to give **1** (Scheme 2). Therefore, it is reasonable to assume that under solvent-free conditions,

complex **3** could afford directly the active species **2** *via* η^2 -(H-Si) coordination of a molecule of silane to **4**.

The aforementioned background motivates us to explore the potential application of compound **3** as a catalyst precursor in solvent-free CO₂ hydrosilylation processes. Herein, we report a detailed kinetic study at variable temperature together with kinetic modeling of the catalytic hydrosilylation of CO₂ with 1,1,1,3,5,5,5-heptamethyltrisiloxane (HMTS) using complex **3** as a catalyst precursor.


2. Experimental

General information


All manipulations were performed with rigorous exclusion of air at an argon/vacuum manifold using standard Schlenk-tube techniques or in a dry box (MB-UNILAB). Catalytic experiments were carried out using a 25 mL Teflon-lined batch reactor. Gas chromatography-mass spectrometry (GC-MS) analyses were performed using an Agilent 7890A GC system, an Agilent 5975C inert MSD with a Triple-Axis Detector MS system, and an HP5-MS capillary column (30 m × 250 μm, film thickness 0.25 μm). Oven parameters: an initial temperature of 45 °C was held for 1.0 min, and then the temperature was increased up to 160 °C at a heating rate of 8.0 °C min⁻¹ and held for 1.0 min, then heated up again to 250 °C at a rate of 8.0 °C min⁻¹. Carrier gas flow was 1.2 mL min⁻¹ and the injection volume was 0.1 mL. Inlet temperature was 250 °C and the split ratio was 100 : 1. The iridium(III) complex **3** was prepared according to a method reported in the literature.¹⁶ NMR spectra were recorded on a Bruker ARX 300 or a Bruker Avance 300 MHz spectrometer. Chemical shifts (expressed in parts per million) were referenced to residual solvent peaks (¹H, ¹³C{¹H}). Coupling constants, *J*, were given in Hertz. 1,1,1,3,5,5,5-Heptamethyltrisiloxane (HMTS) and carbon dioxide (99.99 purity) were purchased from commercial sources. HMTS was dried over 4 Å molecular sieves.

Catalytic reactions at variable temperature and 3 bar

A 25 mL Teflon-lined batch reactor with Teflon lining inside was charged with dry HMTS (3 mL, 11.04 mmol). Then the reactor was closed, purged with CO₂ gas a few times and heated to the corresponding temperature. After stabilization to the corresponding temperature, the reactor was opened under argon and the iridium catalyst **3** (75 mg, 0.11 mmol), which was weighed in a glove box, was added. After closing the reactor, the system was purged a few times using vacuum and CO₂ gas. Then the pressure of CO₂ gas was adjusted to 3 bar. The liquid samples were taken periodically after releasing the CO₂ pressure, without opening the reactor, using a long needle through the sample withdrawal valve. The reactor was purged with vacuum and CO₂ several times after each sample withdrawal and pressure was returned to 3 bar using CO₂. For GC-MS sample preparation, the collected samples were diluted in dry tetrahydrofuran (0.5 mL) and analyzed by quantifiable GC-MS. The product yield was obtained by comparison of the area corresponding to the peaks assigned to



Scheme 2 Examples of the reaction of complex **3** with HSiMeR₂ (R = OSiMe₃).



Scheme 3 Energy profile for the outer-sphere mechanism of the hydrosilylation of CO₂ using complex **1** as the catalyst precursor ([Ir] = Ir(NSiN); R = Me; energy is in kJ mol⁻¹).



the reaction products with the area of the peak corresponding to HMTS.

Kinetic modeling of the catalytic reaction

A power law model was proposed as a descriptor of the kinetics of the first-order reaction¹⁷ of CO₂ with HMTS. The rate of the disappearance of HMTS (eqn (1)) and the rate of the formation of silylformate (eqn (2)) can be written as:

$$-\frac{V}{W_c} \frac{dC_{SX}}{dt} = (k_i C_{SX}) \quad (1)$$

$$\frac{V}{W_c} \frac{dC_F}{dt} = (k_i C_{SX}) \quad (2)$$

where C_{SX} and C_F are the molar concentrations of HMTS and silylformate in the system, respectively, V is the volume of the reactor, W_c is the weight of the catalyst, t is the time in hours and k_i is the rate constant. The reaction rate constants are related to the temperature through the Arrhenius equation.¹⁸ A nonlinear regression algorithm (MATLAB; ODE 45-4th order Runge–Kutta methods and LSQCURVEFIT) was used to evaluate the model equations and to obtain the kinetic parameters. The details of the regression analysis are similar to those previously described by Waziri *et al.*¹⁹

3. Results and discussion

Reaction of CO₂ with HMTS at 298 K using catalytic amounts of 3

We have found that the catalytic solvent-free reaction of CO₂ (298 K, 3.0 bar) with HMTS to produce the corresponding silylformate was faster when complex 3 (1.0 mol%) was used instead of 1 as the catalyst precursor. Indeed, using complex 3, the total consumption of HMTS to give a yellow oil identified by ¹H, ²⁹Si{¹H} NMR spectroscopy and GC-MS as a silylformate, HCOOSiMe(OSiMe₃)₂, (in approximately 89% yield) was observed after three and a half days of reaction, while 6 days were needed to achieve full conversion of HMTS using complex 1 (1.0 mol%) as the catalyst precursor under the same reaction conditions.

¹H NMR spectra of CD₂Cl₂ solutions of the reaction mixture confirm that at the end of the reaction HCOOSiMe(OSiMe₃)₂ is the major reaction product (see the ESI†). Interestingly, in these spectra we always observed the presence of a minor resonance at δ 1.54 ppm which shows a direct C–H bond correlation in the heteronuclear single quantum correlation (HSQC) with one signal that appears at δ 27.3 ppm in the ¹³C{¹H} NMR spectra of the reaction samples and subsequently has been assigned to the CH₂ moieties of cyclooctane. The formation of cyclooctane supports that the *in situ* generation of the active species 2 by reaction of 3 with HMTS could be proposed as the activation step of the catalytic process (Scheme 2). Furthermore, ²⁹Si{¹H} NMR spectra of CD₂Cl₂ solutions of the reaction product also confirm the

formation of HCOOSiMe(OSiMe₃)₂ as the major reaction product. These spectra show two single resonances at δ 11.0 ppm and at δ –58.4 ppm corresponding to the OSiMe₃ and HCOOSiMe silicon atoms, respectively.^{16,20} Additionally, traces of other unidentified silicon-containing minor species were observed (see Fig. 1 and S8 in the ESI†).

GC-MS spectra of the reaction samples taken after 3.5 days of reaction at 25 °C evidenced the full consumption of the starting HMTS and the presence of a peak with a retention time of 7.74 min corresponding to a major reaction product (close to 89% yield) (Fig. 2 and Table 1). The mass pattern found for that peak agrees with that expected for HCOOSiMe(OSiMe₃)₂ (see the ESI†). An additional peak due to {(Me₃SiO)₂MeSi}₂O appears at 14.02 min (9.4%). The presence of {(Me₃SiO)₂MeSi}₂O could be attributable to further reactions of silylformate and its derivatives with HMTS under the reaction conditions²¹ (Scheme 1) or to some iridium-catalyzed hydrolysis of the starting hydrosilane.²²

These results compare well with the work reported by Mizuno and co-workers using the catalytic system [Rh₂(OAc)₄] (0.25 mol%)/K₂CO₃.¹⁰




Fig. 1 ²⁹Si{¹H} NMR (79.5 MHz, CD₂Cl₂) spectrum of the product of the reaction of CO₂ (3 bar) with HMTS after 3.5 days at 25 °C using 3 (1.0 mol%) as the precatalyst.




Fig. 2 GC profile of the product of the reaction of CO₂ (3 bar) with HMTS after 3.5 days at 25 °C using 3 (1.0 mol%) as the precatalyst.



Table 1 Temperature dependence of the turnover frequency (TOF) value obtained for the catalytic hydrosilylation of CO₂ with HMTS using **3** (1.0 mol%) as the precatalyst

Temperature (°C)	Time (h)	Product (mmol) ^a	Yield (%) ^b	TOF (h ⁻¹)
25	80	9.8	88.8	1.1
35	41	10.0	90.6	2.2
45	23	10.0	90.6	3.9
55	9	9.6	87.0	9.7
65	5	9.5	86.1	17.3
75	3	9.6	87.0	29.1

^a Millimoles of silylformate obtained by quantitative GC-MS analysis.

^b [(mmol silylformate/initial mmol HMTS) × 100].

Kinetic study of the catalytic hydrosilylation of CO₂ with HMTS

Kinetic studies of the reaction of hydrosilylation of CO₂ with 1,1,1,3,5,5,5-heptamethyltrisiloxane (HMTS) in the presence of a catalytic amount of **3** (1.0 mol%) show that the rate of this process is temperature dependent. The results from the kinetic studies of the reaction are depicted in Fig. 3. The points in Fig. 3 represent the conversion of HMTS *versus* time. The dotted lines in Fig. 3 have no mathematical meaning and uniquely represent a continuous path between the experimental data points. These results evidenced that increasing the temperature of the reactor from 25 °C to 75 °C enhances the catalytic activity.

A higher catalytic activity was found at 75 °C. GC-MS studies of the reaction evidenced that after 3 hours at that temperature most of the starting HMTS has been consumed and HCOOSiMe(OSiMe₃)₂ (87.0%) is the major reaction product together with traces of {(Me₃SiO)₂MeSi}₂O (Fig. 4 and Table 1).

Table 1 shows that at temperatures higher than 45 °C there is a slight fall-off of the reaction yield. In addition, it has been observed that at these temperatures the silylformate is partially transformed into {(Me₃SiO)₂MeSi}₂O (see Fig. 4 and Table S1 in the ESI†). This observation agrees with the known reactivity of silylformates (see Scheme 1).²¹

The catalytic activity found for **3** in this work is higher than that previously reported for complex **1** and compares well with the catalytic activity of the zinc-based catalytic




Fig. 3 Representation of the experimental data obtained for HMTS conversion by GC-MS.





Fig. 4 Time dependence of the composition of the reaction samples taken from the solvent-free hydrosilylation of CO₂ (3.0 bar) with HMTS using **3** (1.0 mol%) as the catalyst precursor at 75 °C. Data obtained by quantitative GC-MS.

system reported by Sattler and Parkin for the solvent-free hydrosilylation of CO₂ with HSi(OEt)₃.¹² In addition, complex **3** shows better catalytic activity in the range of 55 °C and 75 °C (Table 1).

Kinetic modeling of the Ir-catalyzed CO₂ hydrosilylation

Kinetic modeling of the iridium-catalyzed hydrosilylation of CO₂ with HMTS has been developed based on the chemical equation presented in Scheme 4 and on the approximation that the rate of disappearance of HMTS (eqn (1)) is equal in absolute terms but opposite in sign to the rate of the formation of HCOOSiMe(OSiMe₃)₂ (eqn (2)).

A power law model¹⁷ was proposed as a descriptor of the kinetics of the first-order reaction shown in Scheme 4. The reaction rate constants are supposed to be related to temperature through the Arrhenius equation.¹⁸ The mole balance equations (eqn (1) and (2)) incorporating the Arrhenius relation between k_1 and E_a (see the ESI†) were evaluated by least-squares fitting of the kinetic parameters using the experimental data obtained from the GC-MS studies of the reaction of hydrosilylation of CO₂ with HMTS using **3** (1.0 mol%) as a precatalyst at different temperatures (see the ESI†). Experimental data were used to optimize the model parameters. In order to select the acceptable model the following criteria were considered. All of the kinetic parameters (specific reaction rates and activation energies) should be consistent with (i) physical principles, (ii) coefficient of determination (R^2), (iii) lower SSR (sum of the squares of the residuals), (iv) lower cross-correlation coefficient (γ), and (v) lower individual confidence intervals for the model parameters.



Scheme 4 Solvent-free catalytic CO₂ hydrosilylation with HMTS using **3** (1.0 mol%) as the catalyst precursor.






Fig. 5 Temperature dependence of the consumption of HMTS predicted by the kinetic model (continuous lines). HMTS conversion obtained from GC-MS experiments (points).

As a result of this, kinetic study values of $73.8 (\pm 2.9) \text{ kJ mol}^{-1}$ and of $3.1 \cdot 10^{-2} (\pm 0.03) \text{ s}^{-1}$ have been found for the activation energy of the process (E_a) and for the Arrhenius pre-exponential factor, respectively. This E_a value is consistent with the activation energy theoretically calculated at the DFT level for the outer-sphere mechanism proposed in Scheme 3.¹⁶

A comparison of the experimental data with the model predictions is plotted in Fig. 5. The temperature dependence of the consumption of HMTS predicted by our model (the continuous lines in Fig. 5) fits in an excellent manner with the experimental data obtained from the GC-MS studies of the reaction samples (data points in Fig. 5).

4. Conclusions

The iridium(III) complex $[\text{Ir}(\text{H})(\text{CF}_3\text{SO}_3)(\text{NSiN})(\text{coe})]$ (3) (NSiN = bis(pyridine-2-yloxy)methylsilyl, coe = cyclooctene) has been demonstrated to be an active catalyst precursor for the solvent-free hydrosilylation of CO_2 with 1,1,1,3,5,5,5-heptamethyltrisiloxane (HMTS) under mild reaction conditions (25 °C to 75 °C and 3 bar) allowing preparation of the corresponding silylformate on the gram scale and in high yields. The best catalytic performance has been achieved at 75 °C (TOF = 29.1 h^{-1}). A kinetic study at variable temperature together with kinetic modeling of the catalytic hydrosilylation of CO_2 with HMTS using complex 3 as a catalyst precursor has been carried out. The results from this study showed an E_a (activation energy) value of 73.8 kJ mol^{-1} for this process.

Acknowledgements

The authors express their appreciation to the Ministry of Higher Education, Saudi Arabia, for the establishment of the Center of Research Excellence in Petroleum Refining & Petrochemicals at King Fahd University of Petroleum &

Minerals (KFUPM) and the KFUPM-University of Zaragoza research agreement for their support. Financial support from MINECO/FEDER (projects CONSOLIDER INGENIO-2010 MULTICAT CSD2009-00050 and CTQ2011-27593) and DGA/FSE (group E07) is also acknowledged. Dr. F. J. Fernández-Alvarez thanks Dr. Ralte Lalrempuia for preliminary investigations.

Notes and references

- See for some recent examples; (a) P. G. Jessop, F. Joó and C.-C. Tai, *Coord. Chem. Rev.*, 2004, **248**, 2425–2442; (b) M. Aresta and A. Dibenedetto, *Dalton Trans.*, 2007, 2975–2992; (c) M. Peters, B. Köhler, W. Kuckshinrichs, W. Leitner, P. Markewitz and T. E. Müller, *ChemSusChem*, 2011, **4**, 1216–1240; (d) I. Omae, *Coord. Chem. Rev.*, 2012, **256**, 1384–1405; (e) F. J. Fernández-Alvarez, M. Iglesias, L. A. Oro and V. Polo, *ChemCatChem*, 2013, **5**, 3481–3494; (f) M. Aresta, A. Dibenedetto and A. Angelini, *Chem. Rev.*, 2014, **114**, 1709–1742; (g) T. Fujihara, K. Semba, J. Terao and Y. Tsuji, *Catal. Sci. Technol.*, 2014, **4**, 1699–1709; (h) C. Maeda, Y. Miyazaki and T. Ema, *Catal. Sci. Technol.*, 2014, **4**, 1482–1497; (i) Y.-N. Li, R. Ma, L.-N. He and Z.-F. Diao, *Catal. Sci. Technol.*, 2014, **4**, 1498–1512; (j) A. W. Kleij, *Catal. Sci. Technol.*, 2014, **4**, 1481–1481.
- (a) F. J. Fernández-Alvarez, A. M. Aitani and L. A. Oro, *Catal. Sci. Technol.*, 2014, **4**, 611–624; (b) W. Sattler and G. Parkin, *Catal. Sci. Technol.*, 2014, **4**, 1578–1584.
- (a) N. J. Lawrence, M. D. Drew and S. M. Bushell, *J. Chem. Soc., Perkin Trans. 1*, 1999, **1**, 3381–3391; (b) W.-S. Han, T.-J. Kim, S.-K. Kim, Y. Kim, Y. Kim, S.-W. Nam and S. O. Kang, *Int. J. Hydrogen Energy*, 2011, **36**, 12305–12312.
- (a) K. Motokura, D. Kashiwame, A. Miyaji and T. Baba, *Org. Lett.*, 2012, **14**, 2642–2645; (b) K. Motokura, D. Kashiwame, N. Takahashi, A. Miyaji and T. Baba, *Chem. – Eur. J.*, 2013, **19**, 10030–10037.
- (a) O. Jacquet, C. Das Neves Gomes, M. Ephritikhine and T. Cantat, *ChemCatChem*, 2013, **5**, 117–120; (b) C. Das Neves Gomes, O. Jacquet, C. Villiers, P. Thuéry, M. Ephritikhine and T. Cantat, *Angew. Chem., Int. Ed.*, 2012, **51**, 187–190; (c) O. Jacquet, C. Das Neves Gomes, M. Ephritikhine and T. Cantat, *J. Am. Chem. Soc.*, 2012, **134**, 2934–2937; (d) X. Frogneux, O. Jacquet and T. Cantat, *Catal. Sci. Technol.*, 2014, **4**, 1529–1533.
- (a) Y. Li, X. Fang, K. Junge and M. Beller, *Angew. Chem., Int. Ed.*, 2013, **52**, 9568–9571; (b) O. Jacquet, X. Frogneux, C. Das Neves Gomes and T. Cantat, *Chem. Sci.*, 2013, **4**, 2127–2131.
- (a) S. Park, D. Bézier and M. Brookhart, *J. Am. Chem. Soc.*, 2012, **134**, 11404–11407; (b) A. Berkefeld, W. E. Piers and M. Parvez, *J. Am. Chem. Soc.*, 2010, **132**, 10660–10661; (c) T. Matsuo and H. Kawaguchi, *J. Am. Chem. Soc.*, 2006, **128**, 12362–12363; (d) A. Berkefeld, W. E. Piers, M. Parvez, L. Castro, L. Maron and O. Eisenstein, *Chem. Sci.*, 2013, **4**, 2152–2162; (e) S. J. Mitton and L. Turculet, *Chem. – Eur. J.*, 2012, **18**, 15258–15262.
- T. C. Eiseenschmid and R. Eisenberg, *Organometallics*, 1989, **8**, 1822–1824.



- 9 (a) A. Jansen, H. Görls and S. Pitter, *Organometallics*, 2000, **19**, 135–138; (b) A. Jansen and S. Pitter, *J. Mol. Catal. A: Chem.*, 2004, **217**, 41–45; (c) P. Deglmann, E. Ember, P. Hofmann, S. Pitter and O. Walter, *Chem. – Eur. J.*, 2007, **13**, 2864–2879.
- 10 S. Itagaki, K. Yamaguchi and N. Mizuno, *J. Mol. Catal. A: Chem.*, 2013, **366**, 347–352.
- 11 L. Zhang, J. Cheng and Z. Hou, *Chem. Commun.*, 2013, **49**, 4782–4784.
- 12 W. Sattler and G. Parkin, *J. Am. Chem. Soc.*, 2012, **134**, 17462–17465.
- 13 M. Lancaster, *Green Chemistry: An introductory text*, RSC Paperbacks, Cambridge, 2002.
- 14 R. A. Sheldon, *Green Chem.*, 2005, **7**, 267–278.
- 15 (a) M. S. Singh and S. Chowdhury, *RSC Adv.*, 2012, **2**, 4547–4592; (b) M. A. P. Martins, C. P. Frizzo, D. N. Moreira, L. Buriol and P. Machado, *Chem. Rev.*, 2009, **109**, 4140–4182; (c) P. J. Walsh, H. Li and C. A. de Parrodi, *Chem. Rev.*, 2007, **107**, 2503–2545; (d) A. L. Garay, A. Pichon and S. L. James, *Chem. Soc. Rev.*, 2007, **36**, 846–855.
- 16 R. Lalrempuia, M. Iglesias, V. Polo, P. J. Sanz Miguel, F. J. Fernández-Alvarez, J. J. Pérez-Torrente and L. A. Oro, *Angew. Chem., Int. Ed.*, 2012, **51**, 12824–12827.
- 17 P. J. Crickmore, *Can. J. Chem. Eng.*, 1989, **67**, 392–396.
- 18 A. K. Agarwal and M. L. Brisk, *Ind. Eng. Chem. Process Des. Dev.*, 1985, **24**, 203–207.
- 19 Details about the regression analysis are described in S. Waziri, A. M. Aitani and S. Al-Khattaf, *Ind. Eng. Chem. Res.*, 2010, **49**, 6376–6387.
- 20 M. Chauhan, B. P. S. Chauhan and P. Boudjouk, *Org. Lett.*, 2000, **2**, 1027–1029.
- 21 Traces of CH₄ were detected in the GC/MS spectra.
- 22 K. Garcés, F. J. Fernández-Alvarez, V. Polo, R. Lalrempuia, J. J. Pérez-Torrente and L. A. Oro, *ChemCatChem*, 2014, **6**, 1691–1697.

