



CrossMark  
click for updates

Cite this: *Chem. Sci.*, 2015, 6, 254

Received 25th August 2014  
Accepted 22nd September 2014

DOI: 10.1039/c4sc02590c

www.rsc.org/chemicalscience

# Ligand design for Rh(III)-catalyzed C–H activation: an unsymmetrical cyclopentadienyl group enables a regioselective synthesis of dihydroisoquinolones†

Todd K. Hyster, Derek M. Dalton and Tomislav Rovis\*


We report the regioselective synthesis of dihydroisoquinolones from aliphatic alkenes and *O*-pivaloyl benzhydroxamic acids mediated by a Rh(III) precatalyst bearing sterically bulky substituents. While the prototypical Cp\* ligand provides product with low selectivity, sterically bulky Cp<sup>f</sup> affords product with excellent regioselectivity for a range of benzhydroxamic acids and alkenes. Crystallographic evidence offers insight as to the source of the increased regioselectivity.

C–H activation mediated processes have provided a unique retrosynthetic approach to access a variety of substituted heterocycles.<sup>1</sup> One tactic that has received increased attention is the coupling of  $\pi$ -components with heteroatom containing molecules.<sup>2</sup> A variety of transition metals are capable of catalyzing this type of transformation, providing access to dozens of heterocyclic motifs.<sup>1–3</sup> A challenge for these methods is controlling the regioselectivity of migratory insertion across alkenes and alkynes after the metallacycle forming C–H activation (eqn 1).


Steric and electronic effects are understood to control migratory insertion of unsymmetrical alkynes in Rh(III) catalyzed isoquinolone syntheses (eqn 1). When the substituents are electronically similar, the larger group resides  $\beta$ - to Rh in the metallacycle to avoid unfavorable steric interactions (selectivity is generally >10 : 1).<sup>4</sup> When the substituents are electronically different, the more electron-donating group prefers being  $\alpha$ - to rhodium in the metallacycle, presumably to stabilize the electron poor metal.<sup>5,6</sup> The type of C–H bond being activated also plays an important role in the regioselectivity of migratory insertion; aromatic substrates typically provide synthetically useful regioselectivities when electronically different alkynes are used (>10 : 1) but alkenyl C–H activation leads to products with lower regioselectivities, presumably due to minimal steric interactions during migratory insertion.<sup>7,8</sup> We found that sterically bulky di-*tert*-butylcyclopentadienyl ligand (Cp<sup>f</sup>) enhances the regioselectivity of the alkyne migratory insertion event in these cases, delivering regioselectivities (>10 : 1) modestly above those achievable by Cp\* ligated Rh complexes (<6 : 1). However, when the alkyne migratory

insertion was poorly selective with RhCp\* (<3 : 1), RhCp<sup>f</sup> complex was ineffective at providing synthetically useful levels of selectivity. Furthermore, the Cp<sup>f</sup> ligand was only effective with aryl substituted alkynes, presumably because of strong steric interactions between the ligand and alkyne in the insertion event.


## • Possible Migratory Insertion Products (1)




## • Previous Intermolecular Coupling with Alkenes (2) (ref 5h)



## • Previous Intramolecular Coupling with Alkenes (3) (ref 14a)



## • Proposed Ligand Controlled Alkene Migratory Insertion (4)



Migratory insertion of alkenes to access heterocycles using C–H activation chemistry is still relatively rare, with seminal studies by Glorius and Fagnou reporting the synthesis of dihydroisoquinolones.<sup>9–11</sup> Similar to alkynes, alkenyl electron-

Department of Chemistry, Colorado State University, Fort Collins, Colorado 80523, USA. E-mail: rovis@lamar.colostate.edu

† Electronic supplementary information (ESI) available: Experimental procedures and compound characterization. CCDC 1021286. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c4sc02590c



donating groups favor the position adjacent to the metal in the metallacycle delivering high regioselectivity. In contrast to alkynes, aliphatic alkenes afford product with poor regioselectivity (2 : 1) (eqn 2).<sup>5h,12</sup> We hypothesized competing steric and electronic effects cause the low regioselectivity, with steric effects favoring the formation of a 4-substituted product and electronics favoring the formation of a 3-substituted product.<sup>13</sup> As a temporary solution to this problem, our group and others have employed tethering strategies to increase the regioselectivity of the migratory insertion event (eqn 3).<sup>14,15</sup> Of course, regioselectivity controlled by the ligand on Rh would be the optimal solution to the selectivity problem (eqn 4).<sup>16</sup> Consequently, we focused our attention toward developing an intermolecular variant of this reaction that would provide product with improved regioselectivity.

As a model system, we explored the impact ligands have on the coupling of *O*-pivaloyl-benzhydroxamic acid **1a** with 1-decene **2a** to provide dihydroisoquinolones **3a** and **3a'**. When Cp\* is used as a ligand, the desired products are isolated in excellent yield but poor selectivity (2.4 : 1 **3a** : **3a'**) (Table 1, entry 1).

To determine the effect that ligand electronics have on product regioselectivity, we employed an electron deficient 1-trifluoromethyl-2,3,4,5-tetramethylcyclopentadienyl ligand originally developed by Gassman (Cp<sup>CF3</sup>)<sup>17</sup> and found that this catalyst provides **3a** and **3a'** products in good yield but without an increase in selectivity (2.4 : 1) (Table 1, entry 2).<sup>18,19</sup> Since ligand electronics did not appear to affect product regioselectivity, we tested an electron rich, sterically bulky di-phenyl-tri-methyl Cp ligand (Cp<sup>†</sup>) and were pleased to find a remarkable increase in selectivity from 2.4 : 1 to 12 : 1 (**3a** : **3a'**). Pleased by this improvement, we tested the sterically bulky di-*tert*-butyl Cp ligand Cp<sup>‡</sup> and were surprised to find that RhCp<sup>‡</sup> provides the desired product in 91% yield with exquisite regioselectivity (15 : 1) (Table 1, entry 4).

Having demonstrated that Cp<sup>‡</sup> is able to substantially increase the regioselectivity of alkene migratory insertion, we explored the scope of *O*-pivaloyl benzyhydroxamic acids

Table 1 Ligand optimization<sup>a</sup>

Entry	Catalyst	Yield (%)	Regioselectivity
1	[RhCp*Cl <sub>2</sub> ] <sub>2</sub>	90	2.4 : 1
2 <sup>b</sup>	[RhCp <sup>CF3</sup> Cl <sub>2</sub> ] <sub>2</sub>	85	2.4 : 1
3 <sup>c</sup>	[RhCp <sup>†</sup> Cl <sub>2</sub> ] <sub>2</sub>	82	12 : 1
4 <sup>d</sup>	[RhCp <sup>‡</sup> Cl <sub>2</sub> ] <sub>2</sub>	92	15 : 1

<sup>a</sup> Reaction conditions: **1a** (.2 mmol), 1-decene (.2 mmol), precatalyst (1 mol%), CsOAc (200 mol%), MeOH (0.1 M). <sup>b</sup> Cp<sup>CF3</sup> = 1-trifluoromethyl-2,3,4,5-tetramethylcyclopentadienyl. <sup>c</sup> Cp<sup>†</sup> = 1,2-di-phenyl-3,4,5-trimethylcyclopentadienyl. <sup>d</sup> Cp<sup>‡</sup> = 1,3-di-*t*-butylcyclopentadienyl.

amenable to this reaction. We were pleased to find *para*-substituted benzhydroxamic acids are well tolerated (50–76% yields) with both Cp\* and Cp<sup>†</sup> ligands. Notably, the low regioselectivity of <2.2 : 1 seen using [RhCp\*Cl<sub>2</sub>]<sub>2</sub> is dramatically improved with [RhCp<sup>†</sup>Cl<sub>2</sub>]<sub>2</sub> giving excellent regioselectivities of >14 : 1 (Table 2, entries 1–4). Electron rich amides derived from gallic acid provide product in excellent yield and regioselectivity (Table 2, entry 5). Finally, heterocyclic amides are both well tolerated under the reaction conditions and responsive to the sterically bulky ligand, providing product with excellent regioselectivity (Table 2, entries 6–7).

*meta*-Substituents also provide exquisite levels of regioselectivity for alkene migratory insertion when Cp<sup>†</sup> is used (>15 : 1) (Table 3). Interestingly, when Cp\* is used the C–H activation occurs exclusively at the 6-position; we suggest that this selectivity is the result of steric interactions between the 2-substituent and metal complex during concerted metallation deprotonation. Surprisingly, when Cp<sup>†</sup> is used, the regioselectivity of the C–H activation actually decreases to afford a mixture of 2- and 6-substituted products. While on the surface the decrease of selectivity appears counterintuitive, it can be explained by the uneven distribution of steric bulk in the Cp<sup>†</sup> ligand, which we will comment on subsequently. Not only is this decrease in selectivity mechanistically intriguing, it also offers









Table 2 Amide scope<sup>a</sup>

Entry	Starting material	Yield <sup>b</sup> (%)	Cp*	Cp <sup>†</sup>
1		50	1.5 : 1	19 : 1
2		76	2.2 : 1	19 : 1
3		70	1.9 : 1	16 : 1
4		75	1.7 : 1	14 : 1
5		95	1.9 : 1	15 : 1
6		84	2.5 : 1	19 : 1
7		88	1.8 : 1	19 : 1

<sup>a</sup> Reaction conditions: amide (.2 mmol), 1-decene (.2 mmol), precatalyst (1 mol%), CsOAc (200 mol%), MeOH (0.1 M). <sup>b</sup> Isolated yield of reaction using [RhCp<sup>†</sup>Cl<sub>2</sub>]<sub>2</sub> as a precatalyst.



Table 3 Influence of Cp<sup>t</sup> on C–H insertion selectivity<sup>a</sup>











			
<b>Catalyst</b>	<b>Regioselectivity</b>		
[RhCp*Cl <sub>2</sub> ] <sub>2</sub>	3:1	>19	: 1
[RhCp <sup>t</sup> Cl <sub>2</sub> ] <sub>2</sub> <sup>b</sup>	15:1	6	: 1
			
[RhCp*Cl <sub>2</sub> ] <sub>2</sub>	1.8:1	>19	: 1
[RhCp <sup>t</sup> Cl <sub>2</sub> ] <sub>2</sub> <sup>c</sup>	19:1	4	: 1
			
[RhCp*Cl <sub>2</sub> ] <sub>2</sub>	1.9:1	>19	: 1
[RhCp <sup>t</sup> Cl <sub>2</sub> ] <sub>2</sub> <sup>d</sup>	16:1	1	: 1
			
[RhCp*Cl <sub>2</sub> ] <sub>2</sub>	1.7:1	>19	: 1
[RhCp <sup>t</sup> Cl <sub>2</sub> ] <sub>2</sub> <sup>e</sup>	18:1	7.7	: 1

<sup>a</sup> Reaction conditions: amide (.2 mmol), 1-decene (.2 mmol), precatalyst (1 mol%), CsOAc (200 mol%), MeOH (0.1 M). isolated yield of reaction using [RhCp<sup>t</sup>Cl<sub>2</sub>]<sub>2</sub> as a precatalyst. <sup>b</sup> 67% yield. <sup>c</sup> 80% yield. <sup>d</sup> 85% yield. <sup>e</sup> 79% yield.

an exciting opportunity to potentially reverse the regioselectivity of C–H activation.

We next explored the alkene tolerance of the method. Allyl benzene **2b** furnishes a 1.6 : 1 ratio of dihydroisoquinolone with RhCp\* (Table 4, entry 1). When Cp<sup>t</sup> is employed, the regioselectivity increases to 4.5 : 1. Reducing the temperature to 0 °C further increases regioselectivity to 5.1 : 1. Placing an electron-donating group on the aromatic ring increases the selectivity of insertion (Table 4, entry 2). Aryl electron withdrawing substituents do not, however, provide a change in regioselectivity relative to phenyl (Table 4, entry 3). When the aromatic ring is moved further from the alkene the regioselectivity observed with Cp<sup>t</sup> increases and yields remain high (Table 4, entries 4 and 5). Ester and ketone functional groups are tolerated without a detrimental impact on regioselectivity (Table 4, entry 6, 9, and 10). Unfortunately, alkenes bearing stereocenters are ineffective at inducing diastereoselectivity with either ligand (Table 4, entries 6 and 8). Vinyl cyclopropanes participate with good levels of regioselectivity but poor diastereoselectivity (Table 4, entry 8). Unprotected alcohols provide product in excellent yield with no sign of oxidation (Table 4, entry 7).

Table 4 Impact of Cp<sup>t</sup> on alkene scope<sup>a</sup>

Entry	Alkene	Yield <sup>b</sup> (%)	Cp*	Cp <sup>t</sup>
1 <sup>c</sup>		85	1.6 : 1	5.1 : 1
2		68	1.6 : 1	9.4 : 1
3		70	1.3 : 1	5.5 : 1
4		95	2.3 : 1	14 : 1
5		85	1.6 : 1	8 : 1
6 <sup>d</sup>		92	1.2 : 1	7.2 : 1
7		80	1.4 : 1	12 : 1
8 <sup>e</sup>		93	1 : 1	11 : 1
9		89	2 : 1	14 : 1
10		94	3 : 1	14 : 1

<sup>a</sup> Reaction conditions: **1a** (.2 mmol), alkene (.2 mmol), precatalyst (1 mol%), CsOAc (200 mol%), MeOH (0.1 M). <sup>b</sup> Isolated yield of reaction using [RhCp<sup>t</sup>Cl<sub>2</sub>]<sub>2</sub> as a precatalyst. <sup>c</sup> Reaction conducted at 0 °C. <sup>d</sup> Products isolated as a 1 : 1 ratio of diastereomers. <sup>e</sup> Product isolated as a 2 : 1 ratio of diastereomers.

While it is desirable to achieve high regioselectivity for a single regioisomer, it is even more attractive to use a ligand to access alternate regioisomers. Currently, the only example of Rh(III)-catalyzed synthesis of 4-substituted dihydroisoquinolones is with potassium vinyltrifluoroborates where electronics are believed to control regioselectivity.<sup>20</sup> We found that when vinylcyclohexane was submitted to a reaction with [RhCp\*Cl<sub>2</sub>]<sub>2</sub> as the precatalyst,




Fig. 1 Impact of ligand on reaction of vinyl cyclohexane.



the 3-substituted dihydroisoquinolone **4a** was isolated in 90% yield with 11 : 1 regioselectivity (Fig. 1). However, when the same reaction was catalyzed by  $[\text{RhCp}^f\text{Cl}_2]_2$  the opposite isomer **4b** was isolated in 75% yield and 10 : 1 (**4b** : **4a**) regioselectivity. Given this unexpected discovery, we were interested in gaining insight into how  $\text{Cp}^f$  influences regioselectivity of alkene migratory insertion. A competition experiment between vinyl cyclohexane **2m** and 1-decene **2a** run to 10% conversion favored the formation of dihydroisoquinolone **3a** in >19 : 1 ratio as determined by  $^1\text{H}$  NMR. This experiment suggests that enhanced steric interactions between the substrate and ligand slow the rate of migratory insertion.


To investigate the steric differences between the  $\text{RhCp}^*$  and  $\text{RhCp}^f$  systems X-ray analysis was conducted on a 5-membered  $\text{RhCp}^f$  metallacycle. While we were unable to obtain a 5-membered rhodacycle from our system, Jones and coworkers previously characterized 5-membered rhodacycle **5a** from *N*-benzylidenemethanamine and  $[\text{RhCp}^*\text{Cl}_2]_2$ .<sup>21</sup> We found that a similar metallacycle **5b** derived from  $[\text{RhCp}^f\text{Cl}_2]_2$  could be obtained in crystalline form under identical conditions and was evaluated by single crystal X-ray diffraction.

A comparison of the bond lengths and angles reveals several notable differences between our  $\text{Cp}^f$  rhodacycle and the  $\text{Cp}^*$  rhodacycle reported by Jones (Fig. 2). The Rh–Cp centroid distance in **5b** is 0.011 Å longer than **5a** which is either the result of increased steric interactions, or an artifact of  $\text{Cp}^f$  being a less electron-donating ligand. While there are subtle differences in many bond lengths and angles, the most striking difference is the angle C3–Rh–Cl<sub>1</sub>, which is 98.03° in **5b** while only 90.09° in **5a**. The angle increase is likely the result of steric interactions caused by the *tert*-butyl moiety being situated directly over the Rh–Cl bond. As alkene exchange presumably occurs with Cl, we suggest that steric interactions between the *t*-butyl of the ligand

and the alkene substituent affect both the alkene coordination and 1,2-insertion events.

Based on the X-ray crystal structure and regioselectivity data, we propose the following model for regioselectivity of the 1,2-migratory insertion of alkenes, where steric contributions from the *t*-butyl groups influence both alkene coordination and insertion events to give high selectivity. With small alkyl alkenes, we propose that steric interactions from one *t*-butyl of  $\text{Cp}^f$  disfavor alkene coordination (**I**) and subsequent insertion to give the  $\beta$ -substituted product **3a'** (Fig. 3). Coordination of the alkene with the steric bulk oriented away from the *t*-butyl group finds minimized steric interactions during coordination (**II**). Subsequent migratory insertion from **II** places the alkyl substituent  $\alpha$  to Rh in the transition state, which we propose is able to stabilize a buildup of partial positive charge, making the  $\alpha$ -substituted product **3a** both sterically and electronically favored with  $\text{Cp}^f$ . In the case of the  $\text{Cp}^*$  ligand with small alkyl alkenes, neither steric nor electronic interactions dominate so low selectivity is observed.

However if the size of the alkene substituent is significantly increased, as in the case of vinyl cyclohexane, then  $\text{Cp}^f$  favors the opposite regioisomer. While certainly a puzzling result, we propose that the selectivity can be explained by  $\text{Cp}^f$  rotation such that the *t*-butyl groups both occupy the space above the metallacycle.  $\text{Cp}^f$  rotation gears the *O*-piv toward the alkene coordination site disfavoring alkene coordination to this side (**IV**) favoring the  $\alpha$ -substituted product **3a**. At the same time, alkene coordination (**III**) with the cyclohexyl opposite the *O*-piv minimizes steric interactions enabling insertion of the large alkene and preferential formation of  $\beta$ -substituted product **3a'**. While not conclusive, the observation that cyclohexyl alkene reacts significantly slower than *n*-octyl alkene suggests that migratory insertion of the cyclohexyl alkene proceeds through a higher energy and potentially highly ordered transition state, such as  $\text{Cp}^f$  rotation.



	Complex 5a	Complex 5b
Cp Centroid to Rh	1.834 Å	1.845 Å
Rh <sub>1</sub> –Cl <sub>1</sub>	2.384 Å	2.399 Å
Rh <sub>1</sub> –N <sub>1</sub>	2.095 Å	2.08 Å
Rh <sub>1</sub> –C <sub>14</sub>	2.029 Å	2.031 Å
C <sub>10</sub> –C <sub>3</sub> –Rh <sub>1</sub>	127.05°	130.09°
C <sub>3</sub> –Rh <sub>1</sub> –Cl <sub>1</sub>	90.09°	98.03°
C <sub>14</sub> –Rh <sub>1</sub> –Cl <sub>1</sub>	85.47°	85.69°
N <sub>1</sub> –Rh <sub>1</sub> –Cl <sub>1</sub>	86.69°	87.84°
N <sub>1</sub> –Rh <sub>1</sub> –C <sub>1</sub>	78.67°	78.76°

Fig. 2 X-Ray analysis.




Fig. 3 Rationale for selectivity.





## Conclusions

In conclusion, we found that sterically bulky di-*tert*-butyl cyclopentadienyl ligand (Cp<sup>t</sup>) is effective at increasing regioselectivity of alkene migratory insertion in the synthesis of dihydroisoquinolones by Rh(III) C–H activation catalysis. In contrast to previous cases where Cp\* delivers modest levels of regioselectivity of migratory insertion (with alkynes), Cp<sup>t</sup> renders previously non-selective reactions (with alkenes) highly regioselective. Furthermore, ligand control (Cp\* vs. Cp<sup>t</sup>) enables the highly selective synthesis of two regioisomeric products with vinylcyclohexanes. Finally, crystallographic evidence lends support for a possible explanation of the enhanced regioselectivity.

## Acknowledgements

We thank the NIGMS for generous support of this research (GM80442). T.R thanks Johnson Matthey for a generous loan of Rh salts. D.M.D. thanks the NIH Ruth Kirschstein predoctoral fellowship for funding.

## Notes and references

- (a) C. Zhu, R. Wang and J. R. Falck, *Chem.–Asian J.*, 2012, **7**, 1502; (b) F. W. Patureau, J. Wencel-Delord and F. Glorius, *Aldrichimica Acta*, 2012, **45**, 31; (c) D. A. Colby, R. G. Bergman and J. A. Ellman, *Chem. Rev.*, 2010, **110**, 624; (d) D. M. D'Souza and T. J. J. Müller, *Chem. Soc. Rev.*, 2007, **36**, 1095; (e) J. Wencel-Delord and F. Glorius, *Nat. Chem.*, 2013, **5**, 369.
- G. Song, F. Wang and X. Li, *Chem. Soc. Rev.*, 2012, **41**, 3651.
- (a) P. B. Arockiam, C. Bruneau and P. H. Dixneuf, *Chem. Rev.*, 2012, **112**, 5879; (b) L. Ackermann, *Acc. Chem. Res.*, 2014, **47**, 818.
- For early examples; see: (a) N. Guimond and K. Fagnou, *J. Am. Chem. Soc.*, 2009, **131**, 12050; (b) T. K. Hyster and T. Rovis, *J. Am. Chem. Soc.*, 2010, **132**, 10565; (c) D. R. Stuart, P. Alsabeh, M. Kuhn and K. Fagnou, *J. Am. Chem. Soc.*, 2010, **132**, 18326.
- For early examples; see: (a) K. Ueura, T. Satoh and M. Miura, *Org. Lett.*, 2007, **9**, 1407; (b) K. Ueura, T. Satoh and M. Miura, *J. Org. Chem.*, 2007, **72**, 5362; (c) D. R. Stuart, M. Bertrand-Laperle, K. M. N. Burgess and K. Fagnou, *J. Am. Chem. Soc.*, 2008, **130**, 16474; (d) S. Mochida, K. Hirano, T. Satoh and M. Miura, *J. Org. Chem.*, 2009, **74**, 6295; (e) N. Guimond, C. Gouliaras and K. Fagnou, *J. Am. Chem. Soc.*, 2010, **132**, 6908; (f) D. J. Schipper, M. Hutchinson and K. Fagnou, *J. Am. Chem. Soc.*, 2010, **132**, 6910; (g) S. Rakshit, F. W. Patureau and F. Glorius, *J. Am. Chem. Soc.*, 2010, **132**, 9585; (h) N. Guimond, S. I. Gorelsky and K. Fagnou, *J. Am. Chem. Soc.*, 2011, **133**, 6449.
- For rationale of alkyne electronics using Mayr's nucleophilicity scale; see: R. K. Friedman and T. Rovis, *J. Am. Chem. Soc.*, 2009, **131**, 10775.
- (a) T. K. Hyster and T. Rovis, *Chem. Sci.*, 2011, **2**, 1606; (b) T. K. Hyster and T. Rovis, *Chem. Commun.*, 2011, **47**, 11846.
- For additional examples of alkenyl Rh(III)-catalyzed C–H activation to synthesize heterocycles; see: (a) G. Song, D. Chen, C.-L. Pan, R. H. Crabtree and X. Li, *J. Org. Chem.*, 2010, **75**, 7487; (b) Y. Su, K. Han, G. Song and X. Li, *Org. Lett.*, 2010, **12**, 5462; (c) X. P. Zhang, D. Chen, M. A. Zhao, J. Zhao, A. Q. Jia and X. Li, *Adv. Synth. Catal.*, 2011, **353**, 719; (d) P. C. Too, T. Noji, Y.-J. Lim, X. Li and S. Chiba, *Synlett*, 2011, **22**, 2789.
- (a) S. Rakshit, C. Grohmann, T. Basset and F. Glorius, *J. Am. Chem. Soc.*, 2011, **133**, 2350; (b) Ref. 5h; (c) H. Wang and F. Glorius, *Angew. Chem., Int. Ed.*, 2012, **51**, 7318.
- For an enantioselective example using an artificial metalloenzyme; see: T. K. Hyster, L. Knörr, T. R. Ward and T. Rovis, *Science*, 2012, **338**, 500.
- For an enantioselective example using a chiral cyclopentadienyl ligand; see: (a) B. Ye and N. Cramer, *Science*, 2012, **338**, 504; (b) B. Ye and N. Cramer, *J. Am. Chem. Soc.*, 2013, **135**, 636; (c) B. Ye and N. Cramer, *Angew. Chem., Int. Ed.*, 2014, **53**, 7896.
- In contrast, 1-octyne provides a single regioisomer, see ref. 5h.
- (a) H. A. Malik, G. J. Sormunen and J. Montgomery, *J. Am. Chem. Soc.*, 2010, **132**, 6304; (b) T. Liu, J. Montgomery and K. N. Houk, *J. Am. Chem. Soc.*, 2011, **133**, 6959; (c) N. Wu, L. Deng, L. Liu, Q. Liu, C. Li and Z. Yang, *Chem.–Asian J.*, 2012, **8**, 65; (d) P. A. Wender and A. J. Dyckman, *Org. Lett.*, 1999, **1**, 2089.
- (a) T. A. Davis, T. K. Hyster and T. Rovis, *Angew. Chem., Int. Ed.*, 2013, **52**, 14181; (b) B. Ye, P. A. Donet and N. Cramer, *Angew. Chem., Int. Ed.*, 2014, **53**, 507; (c) Z. Shi, M. Boultradakis-Arapinis, D. C. Koester and F. Glorius, *Chem. Commun.*, 2014, **50**, 2650; (d) T. K. Hyster and T. Rovis, *Synlett*, 2013, **24**, 1842.
- For other tethering strategies; see: (a) X. Xu, Y. Liu and C.-M. Park, *Angew. Chem., Int. Ed.*, 2012, **51**, 9372; (b) N. Quiñones, A. Seoane, R. Garcia-Fandiño, J. L. Mascareñas and M. Gulias, *Chem. Sci.*, 2013, **4**, 2874.
- (a) M. Itoh, K. Hirano, T. Satoh, Y. Shibata, K. Tanaka and M. Miura, *J. Org. Chem.*, 2013, **78**, 1378; (b) K. Morimoto, M. Itoh, K. Hirano, T. Satoh, Y. Shibata, K. Tanaka and M. Miura, *Angew. Chem., Int. Ed.*, 2012, **51**, 5359; (c) Y. Shibata and K. Tanaka, *Angew. Chem., Int. Ed.*, 2011, **50**, 10922; (d) M. Shimizu, H. Tsurugi, T. Satoh and M. Miura, *Chem.–Asian J.*, 2008, **3**, 881.
- P. G. Gassman, J. W. Mickelson and J. R. Sowa, *J. Am. Chem. Soc.*, 1992, **114**, 6942.
- For catalytic uses of this ligand, see: (a) Ref. 7b; (b) J. M. Neely and T. Rovis, *J. Am. Chem. Soc.*, 2014, **136**, 2735.
- For Tanaka's electron deficient Cp ligand; see: (a) M. Fukui, Y. Hoshino, T. Satoh, M. Miura and K. Tanaka, *Adv. Synth. Catal.*, 2014, **356**, 1638; (b) Y. Hoshino, Y. Shibata and K. Tanaka, *Adv. Synth. Catal.*, 2014, **356**, 1577; (c) T. Piou and T. Rovis, *J. Am. Chem. Soc.*, 2014, **136**, 11292.
- M. Pisset, D. Oehlrich, F. Rombouts and G. A. Molander, *Org. Lett.*, 2013, **15**, 1528.
- L. Li, W. W. Brennessel and W. D. Jones, *J. Am. Chem. Soc.*, 2008, **130**, 12414.

