

CrossMark
click for updatesCite this: *Chem. Sci.*, 2015, 6, 1426

Bite-angle bending as a key for understanding group-10 metal reactivity of d^{10} -[M(NHC)₂] complexes with sterically modest NHC ligands†‡

Florian Hering,^a Jörn Nitsch,^a Ursula Paul,^a Andreas Steffen,^a
F. Matthias Bickelhaupt^{*bc} and Udo Radius^{*a}

Synthesis, characterization and investigations on the reactivity of the novel metal basic, yet isolable 14 VE NHC-complexes [M⁰(iPr₂Im)₂] (M = Pd **3**, Pt **4**; iPr₂Im = 1,3-di-isopropyl-imidazolin-2-ylidene; VE = valence electron; NHC = N-heterocyclic carbene) is reported and compared to the chemistry of the corresponding nickel complex. Quantum chemical analyses reveal that differences in the reactivity of group 10 NHC complexes are caused by differences in the rigidity and thus activation strain associated with bending the corresponding d^{10} -[M(NHC)₂] fragments during reaction. These results should have implications for the understanding of the fundamental steps in catalytic cycles, in which such complex fragments are employed.

Received 29th September 2014
Accepted 3rd December 2014

DOI: 10.1039/c4sc02998d

www.rsc.org/chemicalscience

Introduction

Transition metal-mediated activation and functionalization of element–element bonds is a vital area of homogeneous catalysis research. For low-valent, electron-rich transition metal complexes, bond activation is by and large achieved by either σ -bond metathesis or oxidative addition to a metal center, and especially complexes of d^8 transition metals play a crucial role in this chemistry aiming at catalysis.^{1,2} In the last two decades, N-heterocyclic carbenes (NHCs) have emerged as one of the most important classes of spectator ligands in organometallic chemistry.³ Due to their better σ -donor capabilities compared to tertiary phosphines, the employment of NHCs in transition metal complexes can create more electron-rich metal centers. Complexes stabilized with NHCs are expected to exceed the activity of related phosphine compounds for many applications, and NHC complexes of noble metals have already large impact in the field of catalysis.³ The isolation of unsaturated, metal basic complexes, such as 14 VE (valence electron) species d^{10} -[ML₂] or synthons thereof, is highly desirable for investigations

concerning the reactivity of NHC complexes and of elementary steps in catalytic cycles as well as for their use in catalysis.

Over the last years we have explored the manifold chemistry of the {Ni(iPr₂Im)₂} complex fragment (iPr₂Im = 1,3-di-isopropyl-imidazolin-2-ylidene), as provided by dinuclear [Ni₂(iPr₂Im)₄(COD)] (COD = 1,5-cyclooctadiene) or mononuclear [Ni(iPr₂Im)₂(η^2 -C₂H₄)] (see Scheme 1), in stoichiometric and catalytic reactions.⁴ In course of these studies we became more and more interested in synthesizing precursors for [M⁰(iPr₂Im)₂] (M = Pd, Pt; see Scheme 1), either complexes [M⁰(iPr₂Im)₂] themselves or suitable synthons of them. For M = Ni, a two coordinate complex [M(NHC)₂] of iPr₂Im is currently unknown.⁵ Investigations on isolated palladium(0) and especially on platinum(0) complexes bearing two NHC ligands are scarce and mainly involve NHCs with sterically demanding nitrogen substituents.^{6,7} For palladium, two major pathways for the synthesis of complexes [M(NHC)₂] have been established: (i) ligand substitution at [Pd(P(*o*Tol)₃)₂] (*o*Tol = 2-CH₃-C₆H₄) developed by Herrmann *et al.*^{6a} and (ii) ligand substitution at dimeric [(Pd(η^3 -C₄H₇)Cl)₂] in the presence of sodium dimethylmalonate, developed by Caddick and Cloke *et al.*^{6b} The latter


^aInstitut für Anorganische Chemie, Julius-Maximilians-Universität Würzburg, Am Hubland, D-97074 Würzburg, Germany. E-mail: u.radius@uni-wuerzburg.de

^bDepartment of Theoretical Chemistry and Amsterdam Center for Multiscale Modeling, VU University Amsterdam, De Boelelaan 1083, NL-1081 HV Amsterdam, The Netherlands

^cInstitute of Molecules and Materials, Radboud University Nijmegen, Heyendaalseweg 135, NL-6525 AJ Nijmegen, The Netherlands. E-mail: F.M.Bickelhaupt@vu.nl

† Electronic supplementary information (ESI) available: Additional figures and tables; experimental, crystallographic and computational details. CCDC 1025170 and 1025171. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c4sc02998d

‡ Dedicated to Prof. Dr. Reinhold Tacke on the occasion of his 65th birthday.



Scheme 1 {M(iPr₂Im)₂}-precursors: [Ni(iPr₂Im)₂(η^2 -C₂H₄)] **A**, [Pd(iPr₂Im)₂] and [Pt(iPr₂Im)₂].



procedure does not require the unappealing synthesis of the $[\text{Pd}(\text{P}\{\text{oTol}\}_3)_2]$ precursor,⁸ but the stoichiometry has to be controlled carefully due to equilibration in solution, since NHC dissociation from palladium seems to be more facile than generally envisioned.^{6c} However, using these methods, complexes such as $[\text{Pd}(\text{Dipp}_2\text{Im})_2]$ (Dipp = 2,6-diisopropylphenyl), $[\text{Pd}(\text{Mes}_2\text{Im})_2]$ (Mes = 2,4,6-trimethylphenyl), $[\text{Pd}(t\text{Bu}_2\text{Im})_2]$, $[\text{Pd}(\text{Ad}_2\text{Im})_2]$ (Ad = adamantyl) and of other NHCs with sterically demanding substituents at nitrogen have been prepared.⁶ Furthermore, synthesis of $[\text{Pd}(\text{Me}_2\text{Im})_2]$ *via* potassium reduction of $[\text{Pd}(\text{Me}_2\text{Im})_2\text{Cl}_2]$ and the characterization of this complex including X-ray analysis has been described lately.^{6f} In course of these studies, it turned out that complexes of the type $[\text{Pd}(\text{NHC})_2]$ are extremely air sensitive and form isolable peroxy or peroxocarbonate complexes. Isolation of $[\text{Pd}(\text{iPr}_2\text{Im})_2]$ has been claimed twice in the literature,^{6a,j} but the NMR data provided are in neither case in accordance with a D_{2d} or D_{2h} type structure of this complex.


The chemistry of isolated complexes $[\text{Pt}(\text{NHC})_2]$ is virtually unexplored.⁷ Arduengo *et al.*^{7a} and Nolan *et al.*^{7c} described the synthesis of $[\text{Pt}(\text{Mes}_2\text{Im})_2]$ and $[\text{Pt}(\text{Dipp}_2\text{Im})_2]$ starting from the NHC and $[\text{Pt}(\text{COD})_2]$ or $[\text{Pt}(\text{COD})(\text{Me})_2]$, respectively. Intra-molecular C–H activation of the ligand was observed for $[\text{Pt}(\text{Mes}_2\text{Im})_2]$ in the latter case.^{7c} This procedure seems to be further limited since the use of $t\text{Bu}_2\text{Im}$ leads to formation of $[\text{Pt}(t\text{Bu}_2\text{Im})(t\text{Bu}_2\text{Im})'(\text{H})(\text{Me})_2]$ (l' denotes coordination at the backbone as an "abnormal" NHC). Additionally, Braunschweig *et al.* reported the synthesis of $[\text{Pt}(\text{SIMes})_2]$ (SIMes = bis(mesityl)imidazolidin-2-ylidene) *via* ligand exchange at $[\text{Pt}(\text{PCy}_3)_2]$, whereas the same reaction with $t\text{Bu}_2\text{Im}$ led to mixed substituted $[\text{Pt}(t\text{Bu}_2\text{Im})(\text{PCy}_3)_2]$.^{7d}

Herein, we report the synthesis, spectroscopic and chemical characterization, and quantum chemical analysis of the complexes $[\text{Pd}(\text{iPr}_2\text{Im})_2]$ (3) and $[\text{Pt}(\text{iPr}_2\text{Im})_2]$ (4).

Results and discussion

Synthesis and characterization of $[\text{M}^0(\text{iPr}_2\text{Im})_2]$ (M = Pd, Pt)

All attempts to isolate the complexes $[\text{M}^0(\text{iPr}_2\text{Im})_2]$ (M = Pd, Pt) bearing NHC ligands with modest steric demand by the methods introduced above failed in our hands. However, we have found that complexes $[\text{M}(\text{iPr}_2\text{Im})_3\text{Cl}]^+\text{Cl}^-$ (M = Pd 1, Pt 2; see Scheme 2) are suitable precursors for the synthesis of 3 and 4. Complexes 1 and 2 have been prepared by the reaction of the metal halides MCl_2 and three equivalents (or an excess) of the NHC (see Scheme 2).⁹ The use of $\text{K}_2[\text{PtCl}_4]$ instead of PtCl_2 as platinum source also leads selectively to $[\text{Pt}(\text{iPr}_2\text{Im})_3\text{Cl}]^+\text{Cl}^-$ (2).



Scheme 2 Synthesis of $[\text{M}(\text{iPr}_2\text{Im})_3\text{Cl}]^+\text{Cl}^-$ (M = Pd 1, Pt 2) and $[\text{M}(\text{iPr}_2\text{Im})_2]$ (M = Pd 3, Pt 4).




Fig. 1 X-ray crystal structure of the cation of $[\text{Pd}(\text{iPr}_2\text{Im})_3\text{Cl}]^+\text{Cl}^-$ (1; left) and of $[\text{Pd}(\text{iPr}_2\text{Im})_2]$ (3; right) in the solid state (ellipsoids set at 50% probability level). Hydrogen atoms have been omitted for clarity. Selected bond lengths (Å) and angles (deg.) for $[\text{Pd}(\text{iPr}_2\text{Im})_3\text{Cl}]^+$: Pd–Cl1 2.3553(9), Pd–C1 2.0181(34), Pd–C2 2.0590(35), Pd–C3 2.0474(30); Cl1–Pd–C1 178.169(95), C2–Pd–C3 175.967(140). Selected bond lengths (Å) for 1: Pd–C1 2.0231 (18).

In some instances the formation of the imidazolium salt $\text{iPr}_2\text{Im} \cdot \text{HCl}$ as a side product was detected for this reaction. The imidazolium salt and complexes 1 and 2, respectively, are difficult to separate, since these compounds reveal similar solubility in common organic solvents. The NMR spectra of 1 and 2 are those typically observed for a T-shaped alignment of three iPr_2Im ligands and a X-ray analysis of 1 confirms the square planar structure with a T-shaped, propeller-like alignment of the three NHCs ligands of the 16 VE cation (see Fig. 1).

The reduction of 1 and 2 with potassium graphite leads in good yields to the two-coordinated complexes $[\text{Pd}(\text{iPr}_2\text{Im})_2]$ (3) and $[\text{Pt}(\text{iPr}_2\text{Im})_2]$ (4) (see Scheme 2), which were isolated as orange (3) or yellow (4) solids. The release of one equivalent iPr_2Im as well as the presence of minor amounts of $\text{iPr}_2\text{Im} \cdot \text{HCl}$ does not affect the reaction, since 3 and 4 do not react with additional iPr_2Im to afford $[\text{M}(\text{iPr}_2\text{Im})_3]$. For nickel, we have shown earlier that threefold coordination is feasible, as exemplified by the synthesis of $[\text{Ni}(\text{Me}_2\text{Im})_3]$ from the reaction of $[\text{Ni}(\text{COD})_2]$,^{4c} and we have also proof for the existence of $[\text{Ni}(\text{iPr}_2\text{Im})_3]$. For both, 3 and 4, the analytical data are in agreement with a coordination of the metal with two NHC ligands as proposed in Scheme 2. The ^1H NMR and $^{13}\text{C}\{^1\text{H}\}$ NMR spectra reveal one set of signals for the ligands of each complex comprising a doublet for the isopropyl methyl protons (3: 1.34 ppm, 4: 1.33 ppm), a septet for the isopropyl methine protons (3: 6.03 ppm, 4: 6.26 ppm), and a singlet for the olefinic protons of the backbone (3: 6.46 ppm, 4: 6.40 ppm) in the ^1H NMR spectrum. The ^{195}Pt NMR spectrum of 4 shows a resonance at $\delta = -5943$ ppm, which is close to ^{195}Pt NMR shifts reported for other $[\text{PtL}_2]$ complexes ($[\text{Pt}(\text{PCy}_3)_2]$: -6501 ppm, $[\text{Pt}(\text{SIMes})(\text{PCy}_3)]$: -6151 ppm, $[\text{Pt}(t\text{Bu}_2\text{Im})(\text{PCy}_3)]$: -6156 ppm, $[\text{Pt}(\text{SIMes})_2]$: -5462 ppm vs. $[\text{Pt}(\text{PPh}_3)_3]$: -4583 ppm).^{7a,d,10}

Single crystals of $[\text{Pd}(\text{iPr}_2\text{Im})_2]$ (3) suitable for X-ray diffraction have been obtained *via* slow diffusion of the solvent from a saturated solution of 3 in diethyl ether. The compound crystallizes in the space group $P2_1/c$ with the palladium atom located on a crystallographically imposed inversion center.

Both imidazole rings of 3 are coplanar arranged, which leads to a D_{2h} symmetric compound with Pd–C bond lengths of 2.0231(18) Å (Fig. 1). However, for steric reasons, a D_{2d}



symmetric structure with staggered NHC ligands should be expected. A search on all currently known X-ray crystal structures of d^{10} -[M(NHC)₂] complexes reveal that the dihedral angles between the NHC nitrogen and NHC carbene carbon atoms increases for bulky substituents.^{5a,6,7,11} Calculations on [M(H₂Im)₂] presented earlier¹² predict a D_{2d} type structure also for sterically less demanding NHCs. However, calculations on [M(iPr₂Im)₂] (M = Ni, Pd, Pt) performed at BP86/def2-TZVPP level (see ESI†) reveal a shallow potential for the rotation of the NHC ligands and differences in energy between a D_{2d} and D_{2h} type structure below 5 kJ mol⁻¹ in favor of a staggered alignment. Thus, we attribute the reason for experimentally observed D_{2h} type symmetry of **3** to packing forces.

Reactivity of [Pt(iPr₂Im)₂] (**4**)

We reported earlier stable NHC nickel η^2 -olefin, η^2 -alkyne, and η^2 -ketone complexes [Ni(iPr₂Im)₂(η^2 -C₂R₄)], [Ni(iPr₂Im)₂(η^2 -C₂R₂)], and [Ni(iPr₂Im)(η^2 -CO{-O=CR¹R²})].⁴ Due to the fact that complexes **3** and **4** did not react with additional NHC in solution, we became interested to investigate reactions of these two-coordinated complexes with π -acceptor ligands such as olefins, alkynes and ketones. In stark contrast to [Ni(iPr₂Im)₂], its Pd congener **3** merely shows weak interactions with ethylene giving [Pd(iPr₂Im)₂(η^2 -C₂H₄)] (**5**), which is very unstable even under an atmosphere of the olefin (see ESI†). However, if a solution of [Pt(iPr₂Im)₂] (**4**) is pressurized with 3 bars ethylene, [Pt(iPr₂Im)₂(η^2 -C₂H₄)] (**6**) is formed in quantitative yield and remains stable under an ethylene atmosphere (Scheme 3). This compound reveals one set of resonances for the NHC ligand (in C₆D₆ at 1.04, 5.48, 6.47 ppm), substantially shifted from the signals of **4** (1.33, 6.26, 6.40 ppm; see Fig. 2). More significantly, resonances of the η^2 -coordinated ethylene ligand were detected at 1.99 ppm with a ¹⁹⁵Pt-coupling constant ²J_{PtH} of 54.7 Hz in the ¹H NMR spectrum and at 13.3 ppm (¹J_{PtC} = 253.8 Hz) in the ¹³C NMR spectrum of **6**. If the ethylene pressure is reduced, complex **4** reforms in quantitative yield. This cycle can be





Fig. 2 ¹H NMR of [Pt(iPr₂Im)₂(η^2 -C₂H₄)] **6** in C₆D₆: (a) sample of [Pt(iPr₂Im)₂] **4** after pressurizing with ethylene to give **6**, (b) same sample after 14 d (revealing the stability of the complex in solution), (c) same sample after replacing the ethylene atmosphere with argon (revealing that **4** was formed from **6** after ethylene has been partially removed), (d) same sample after 5 h at room temperature; (e) sample after removal of all volatile material and addition of fresh C₆D₆ (= complex **4**).

repeated for several times without any loss of complex **4**. To our surprise, no reaction has been observed between **4** and more electron rich alkenes such as 1,5-cyclooctadiene or cyclooctene.


Accordingly, we were interested in the reactivity of complex **4** with respect to different alkynes such as phenyl and diphenyl acetylene, and 2-butyne (see Scheme 3). For nickel, we reported earlier the formation of stable nickel alkyne complexes [Ni(iPr₂Im)₂(η^2 -RC≡CR')].^{4a,c} The reaction of phenyl acetylene and diphenyl acetylene with **4** afforded the alkyne complexes [Pt(iPr₂Im)₂(η^2 -HC≡CPh)] (**7**) and [Pt(iPr₂Im)₂(η^2 -PhC≡CPh)] (**8**) as the main products.

If 2-butyne is used for the reaction with **4**, quantitative conversion cannot be observed (Scheme 3). For the reaction of **4** and 2-butyne in an equimolar stoichiometry an equilibrium was reached that contains approximately 33% of the reaction product [Pt(iPr₂Im)₂(η^2 -MeC≡CMe)] (**9**) and 66% of the starting material [Pt(iPr₂Im)₂] (**4**). After prolonged standing in solution the isomerization of the 2-butyne ligand to an 1,2-butadiene (allene) ligand in the coordination sphere of platinum can be observed. Complex [Pt(iPr₂Im)₂(η^2 -H₂C=C=C(H)Me)] (**10**) has been identified in solution in a mixture containing **4**, **9**, and **10** using ¹H and ¹³C NMR spectroscopy. The resonances of the allene ligand for complex **10** were detected at 2.06 ppm (CH₂), 6.02 ppm (CH) and 2.57 ppm (CH₃) in the proton NMR spectrum with ¹⁹⁵Pt coupling to the CH₂ (²J_{PtH} = 54.2 Hz) and CH (³J_{PtH} = 47.0 Hz) signals, which is in agreement with an 1,2-coordination of the allene ligand as proposed in Scheme 3. The assignment of the resonances was confirmed by a ¹H-¹H COSY experiment (see ESI†). The allene carbon atoms attached to platinum reveal resonances at -4.3 ppm and 158.4 ppm in the ¹³C{¹H} NMR spectrum. Complex **10** represents to our knowledge one of the few examples of an allene complex formed from the isomerization of an alkyne ligand in the coordination sphere of platinum.¹³

The reaction of **4** with acetophenone leads to an equilibrium between **4** and the α -C-H bond activation product *trans*-[Pt(iPr₂Im)₂(H)(-CH₂-C(O)Ph)] (**11**) (Scheme 3). Other likely reaction products such as a η^2 -ketone complex or a complex resulting from *ortho*-metallation of the phenyl ring of the



Scheme 3 Reactions of [Pt(iPr₂Im)₂] (**4**) with π -acceptor ligands.



ketone have not been observed. The formation of **11** is quantitative if an excess of acetophenone is used at elevated temperatures (80 °C). The hydride ligand of **11** gives rise to a resonance at -10.11 ppm ($^1J_{\text{PtH}} = 1133.3$ Hz) and the metal bound methylene unit to a resonance at 3.22 ppm ($^2J_{\text{PtH}} = 70.6$ Hz) in the proton NMR spectrum. Notably, benzene solutions of compound **11** are not stable and slowly decompose to starting material. We have demonstrated earlier that $[\text{Ni}_2(\text{iPr}_2\text{Im})_4(\text{COD})]$ reacts with ketones to give stable and thermally robust nickel ketone complexes $[\text{Ni}(\text{iPr}_2\text{Im})_2(\eta^2\text{-}\{\text{O}=\text{C}\text{R}^1\text{R}^2\})]$ (see Fig. 3). DFT calculations (BP86/def2-TZVPP) on the relative stabilities of complexes $[\text{M}(\text{iPr}_2\text{Im})_2(\eta^2\text{-}\{\text{O}=\text{C}\text{PhMe}\})]$ and *trans*- $[\text{M}(\text{iPr}_2\text{Im})_2(\text{H})(-\text{CH}_2-\text{C}(\text{O})\text{Ph})]$ for nickel and platinum (see Fig. 3 and ESI†) reveal a significant stabilization (-41.7 kJ mol $^{-1}$) of the platinum complex **11** with respect to an η^2 -ketone complex. A platinum η^2 -ketone complex would be unstable with respect to cleavage into **4** and acetophenone, according to these calculations. For nickel, the DFT calculations reveal the opposite trend, *i.e.* the η^2 -ketone complex is the global minimum, stabilized by -57.1 kJ mol $^{-1}$ with respect to $[\text{Ni}(\text{iPr}_2\text{Im})_2]$ and acetophenone and -60.9 kJ mol $^{-1}$ with respect to $[\text{Ni}(\text{iPr}_2\text{Im})_2(\text{H})(-\text{CH}_2-\text{C}(\text{O})\text{Ph})]$ (see Fig. 3).

Catalytic C–H bond activation reactions at positions α to the carbonyl group are currently of interest because they provide atom economic entries into the α -functionalization of organic carbonyl compounds.¹⁴ Compound **11** represents to our knowledge the first well characterized example of an α -CH bond activation product of a ketone at platinum (0) to give the corresponding hydrido α -keto-alkyl complex.^{15,16} Further investigations on the reactivity of **11** are underway.




Fig. 3 Top: contrary reactivity of $[\text{Ni}(\text{iPr}_2\text{Im})_2]$ (as provided by $[\text{Ni}_2(\text{iPr}_2\text{Im})_4(\text{COD})]$ and $[\text{Pt}(\text{iPr}_2\text{Im})_2]$ towards acetophenone: Formation of the η^2 -ketone complexes $[\text{Ni}(\text{iPr}_2\text{Im})_2(\eta^2\text{-}\{\text{O}=\text{C}\text{PhMe}\})]$ for nickel and the α -C–H bond activation product *trans*- $[\text{Pt}(\text{iPr}_2\text{Im})_2(\text{H})(-\text{CH}_2-\text{C}(\text{O})\text{Ph})]$ (**11**) for platinum. Bottom: results of DFT calculations (BP86/def2-TZVPP; see ESI†) on the relative stabilities of the α -C–H bond activation products *trans*- $[\text{M}(\text{iPr}_2\text{Im})_2(\text{H})(-\text{CH}_2-\text{C}(\text{O})\text{Ph})]$ and η^2 -ketone complexes $[\text{M}(\text{iPr}_2\text{Im})_2(\eta^2\text{-}\{\text{O}=\text{C}\text{PhR}^2\})]$ for $\text{M} = \text{Pt}$ (left side) and $\text{M} = \text{Ni}$ (right side).

Calculations on model systems

In order to understand the reasons for these differences in nickel, palladium and platinum chemistry, we have performed calculations on model systems $[\text{M}(\text{H}_2\text{Im})_2(\eta^2\text{-C}_2\text{H}_4)]$ using the ADF program at the ZORA-BLYP/TZ2P level.¹⁷ The calculated bond energies for the ethylene ligand dissociation in complexes $[\text{M}(\text{H}_2\text{Im})_2(\eta^2\text{-C}_2\text{H}_4)]$ into $[\text{M}(\text{H}_2\text{Im})_2]$ and ethylene clearly reveal the trend observed experimentally, *i.e.* that ethylene binds significantly more strongly to the nickel than to the palladium and platinum complexes: -104.2 kJ mol $^{-1}$ (Ni), -25.0 kJ mol $^{-1}$ (Pd) and -20.5 kJ mol $^{-1}$ (Pt; see Table 1, Fig. 4 and ESI†).

We have analyzed the metal–ethylene bonding in $[\text{M}(\text{H}_2\text{Im})_2(\eta^2\text{-C}_2\text{H}_4)]$ complexes using the activation strain model and ADF's energy decomposition analysis (EDA).^{17b,c} Thus, the bond energy ΔE is decomposed into the strain energy ΔE_{strain} associated with the deformation of the fragments $[\text{M}(\text{H}_2\text{Im})_2]$ and C_2H_4 during the bond formation reaction plus the mutual interaction ΔE_{int} between the deformed fragments. The latter is further decomposed into classical electrostatic attraction ΔV_{elstat} , steric (Pauli) repulsion ΔE_{Pauli} , and bonding orbital interactions ΔE_{oi} .^{17b} The results are given in Table 1. Note that the orbital interaction ΔE_{oi} is least stabilizing for the 4d metal palladium: -415.1 kJ mol $^{-1}$ (Ni); -334.5 kJ mol $^{-1}$ (Pd); -523.4 kJ mol $^{-1}$ (Pt).¹⁸ Our analyses show that, in line with the work of Dewar, Chatt and Duncanson,¹⁹ the bonding can be described as ligand to metal σ -donation and a π -back donation from the metal to the unoccupied π^* -orbital of ethylene. In agreement with studies presented earlier,^{20,21} the orbital interaction of π -symmetry has for each metal a larger impact compared to the orbital interaction of σ -symmetry and increases for $[\text{M}(\text{H}_2\text{Im})_2(\eta^2\text{-C}_2\text{H}_4)]$ from Pd (-249.6 kJ mol $^{-1}$,

Table 1 Energy decomposition analysis [kJ mol $^{-1}$] of the metal–ethylene bond in $[\text{M}(\text{H}_2\text{Im})_2(\eta^2\text{-C}_2\text{H}_4)]$ complexes^a

M	ΔV_{elstat}	ΔE_{Pauli}	ΔE_{steric}	ΔE_{oi}	ΔE_{int}	ΔE_{strain}	ΔE
Ni	-543.5	+741.8	+198.4	-417.1	-245.0	+140.8	-104.2
Pd	-521.3	+695.7	+174.4	-334.5	-184.2	+159.2	-25.0
Pt	-773.2	+1038.2	+265.0	-523.4	-284.1	+263.6	-20.5

^a Computed at ZORA-BLYP/TZ2P; for full data, see Table S1; $\Delta E = \Delta E_{\text{strain}} + \Delta E_{\text{int}}$; $\Delta E_{\text{int}} = \Delta V_{\text{elstat}} + \Delta E_{\text{Pauli}} + \Delta E_{\text{oi}} + \Delta E_{\text{disp}}$ (see text and ref. 17).




Fig. 4 Analysis of metal–ethylene bonding in $[\text{M}(\text{H}_2\text{Im})_2(\eta^2\text{-C}_2\text{H}_4)]$ complexes using the activation strain model and ADF's energy decomposition analysis (EDA) (see text).



75% of ΔE_{oi}) to Ni ($-317.4 \text{ kJ mol}^{-1}$, 76%) to Pt ($-369.0 \text{ kJ mol}^{-1}$, 71%). The interaction energy ΔE_{int} follows this trend in ΔE_{oi} : $-245.0 \text{ kJ mol}^{-1}$ (Ni); $-184.2 \text{ kJ mol}^{-1}$ (Pd); $-284.1 \text{ kJ mol}^{-1}$ (Pt).

We find that the overall bond energies ΔE is significantly weakened for palladium ($-25.0 \text{ kJ mol}^{-1}$) and platinum ($-20.5 \text{ kJ mol}^{-1}$) compared to nickel ($-104.2 \text{ kJ mol}^{-1}$). The main reason lies in an increase in strain energy ΔE_{strain} going from Ni to Pd and especially Pt. This strain ΔE_{strain} appears to stem mainly from the bending of the $[\text{M}(\text{H}_2\text{Im})_2]$ fragment from its preferred linear geometry towards smaller L–M–L bite angles. While the deformation energies of the ethylene ligand (for Pd: $+42.2$; Ni: $+48.6$ and Pt: $+74.3 \text{ kJ mol}^{-1}$) depend on the strength of π -back donation, the origin of the deformation energies for the $[\text{M}(\text{H}_2\text{Im})_2]$ fragment are more subtle and increase from Ni ($+92.2 \text{ kJ mol}^{-1}$) to Pd ($+117.0 \text{ kJ mol}^{-1}$) to Pt ($+189.3 \text{ kJ mol}^{-1}$). Bending of linear $[\text{M}(\text{H}_2\text{Im})_2]$ is costly in energy, and it has to be emphasized that this energy loss upon bending is by and large an electronic effect. In addition to steric factors, it has been shown earlier that fourteen-electron dicoordinate group-10 metal complexes $\text{d}^{10}\text{[ML}_2\text{]}$ adopt linear geometries mainly due to a significant destabilization of one of the occupied, in the linear case nonbonding d orbitals (d_{xz} ; for a schematic Walsh diagram see ESI†).²² An alternative but, ultimately equivalent way of understanding the resistance of $\text{d}^{10}\text{[ML}_2\text{]}$ complexes to L–M–L bending is that such bending destabilizes the interaction ΔE_{int} between ML and L through increasing Pauli repulsion between the ligands' lone-pair orbital lobes and a destabilization of the ML d_{z^2} hybrid orbital.^{22e} For the good σ -donating NHC ligands, the $\text{d}^{10}\text{[ML}_2\text{]}$ bending strain is relatively high in the case of the noble metals Pd and Pt and – in combination with the weak orbital interaction ΔE_{oi} term for Pd – is mainly responsible for the weak metal olefin bond. Note that NHC-coordinated palladium and platinum olefin complexes are known to be stable if one utilizes chelating bidentate NHC ligands, for which the linearization of the $[\text{M}(\text{NHC})_2]$ moiety is not possible.²³ The importance of $[\text{M}(\text{NHC})_2]$ -bending to this chemistry can also be illustrated with the aid of an analysis on the stability of the model complexes $[\text{M}(\text{H}_2\text{Im})_3]$ with respect to a cleavage into $[\text{M}(\text{H}_2\text{Im})_2]$ and H_2Im (Table 2). The deformation energies of the H_2Im ligand are small and lie between 1 and 3 kJ mol^{-1} (for Pd: $+1.0$; Pt: $+1.9$ and Ni: $+2.8 \text{ kJ mol}^{-1}$), while the strain energies ΔE_{strain} for the $[\text{M}(\text{H}_2\text{Im})_2]$ fragments are much higher and increase substantially going from Ni ($+70.7 \text{ kJ mol}^{-1}$) to Pd ($+90.5 \text{ kJ mol}^{-1}$) to Pt ($+130.4 \text{ kJ mol}^{-1}$). Thus, the energy gain ΔE_{int} obtained from the interaction of $[\text{M}(\text{H}_2\text{Im})_2]$

with H_2Im is almost used up for palladium and platinum due to the L–M–L bending strain of $[\text{M}(\text{H}_2\text{Im})_2]$. As a result, we calculate a significant overall binding energy for Ni ($-107.9 \text{ kJ mol}^{-1}$) and a much more reduced binding energy for the metals with larger radii Pd (-25.6) and Pt ($-26.5 \text{ kJ mol}^{-1}$).

Our DFT calculations show the same trend in bond energy ΔE for the full complexes $[\text{M}(\text{iPr}_2\text{Im})_2(\text{L})]$ as discussed above for the truncated models $[\text{M}(\text{H}_2\text{Im})_2(\text{L})]$ (see ESI†). In agreement with the experiment, these calculations reveal that addition of another iPr_2Im ligand to $[\text{M}(\text{iPr}_2\text{Im})_2]$ should be repulsive for palladium ($+4.4 \text{ kJ mol}^{-1}$) and platinum ($+13.1 \text{ kJ mol}^{-1}$). In addition, we calculated even smaller BDEs for 2-butyne complexes $[\text{M}(\text{iPr}_2\text{Im})_2(\eta^2\text{-C}_2\text{Me}_2)]$ than for the ethylene complexes $[\text{M}(\text{iPr}_2\text{Im})_2(\eta^2\text{-C}_2\text{H}_4)]$ of palladium and platinum.

Conclusions

We have reported the synthesis, characterization and first investigations concerning the reactivity of the novel complexes $[\text{M}^0(\text{iPr}_2\text{Im})_2]$ ($\text{M} = \text{Pd}$ **3**, Pt **4**), stabilized with an NHC ligand of low steric demand. These complexes represent rare examples of metal basic, yet isolable 14 VE complexes of the type $\text{d}^{10}\text{[ML}_2\text{]}$. Complexes **3** and **4** do not react with additional NHC and the reaction with selected π -acidic ligands leads in some cases, if at all, to π -complexes, which are much less stable compared to the corresponding nickel complexes. The reactions presented here are illustrative examples of the differences one can observe by changing the central atom in a complex going from a 3d metal to a 4d or 5d metal atom within the same ligand framework and clearly show that L–M–L bending strain of $\text{d}^{10}\text{[ML}_2\text{]}$ is a key to understand group 10 metal reactivity. This is a new appearance of the concept of bite-angle flexibility that was recently introduced in the context of metal-mediated C–H bond activation.^{22f}

Acknowledgements

This work was supported by the Julius-Maximilians-University Würzburg and the Deutsche Forschungsgemeinschaft (DFG). F.M.B. also thanks the Netherlands Organization for Scientific research (NWO-CW and NWO-EW) and the National Research School Combination-Catalysis (NRSC-C) for financial support.

Notes and references

- (a) J. Hartwig, *Organotransition Metal Chemistry – from Bonding to Catalysis*, University Science Books, Sausalito, 2010; (b) C. Elschenbroich, *Organometallics*, Weinheim, 3rd edn, 2006; (c) R. H. Crabtree, *The Organometallic Chemistry of the Transition Metals*, Wiley, Hoboken, 4th edn, 2005.
- (a) *Applied Homogeneous Catalysis with Organometallic Compounds*, ed. B. Cornils and W. A. Herrmann, Weinheim, 2nd edn, 2002; (b) D. Steinborn, *Grundlagen der metallorganischen Komplexkatalyse*, 2, Auflage, Vieweg + Teubner, Wiesbaden, 2010.
- (a) W. A. Herrmann, *Angew. Chem., Int. Ed.*, 2002, **41**, 1290; (b) *N-Heterocyclic Carbenes in Synthesis*, ed. S. P. Nolan, Wiley-VCH, Weinheim, 2006; (c) *N-Heterocyclic Carbenes*

Table 2 Energy decomposition analysis [kJ mol^{-1}] of the metal–carbene bond in $[\text{M}(\text{H}_2\text{Im})_3]$ complexes between $[\text{M}(\text{H}_2\text{Im})_2]$ and H_2Im^a

M	ΔE_{steric}	ΔE_{oi}	ΔE_{int}	ΔE_{strain}	ΔE
Ni	+123.4	–278.5	–181.3	+73.4	–107.9
Pd	+132.4	–224.7	–117.1	+91.5	–25.6
Pt	+225.8	–357.3	–158.8	+132.3	–26.5

^a Computed at ZORA-BLYP/TZ2P; for full data, see Table S2; $\Delta E = \Delta E_{\text{strain}} + \Delta E_{\text{int}}$; $\Delta E_{\text{int}} = \Delta V_{\text{elstat}} + \Delta E_{\text{Pauli}} + \Delta E_{\text{oi}} + \Delta E_{\text{disp}}$ (see text and ref. 17).



- in Transition Metal Catalysis, *Topics in Organometallic Chemistry*, ed. F. Glorius, Springer, Berlin, 2007, vol. 21; (d) M. C. Jahnke and F. E. Hahn, *Angew. Chem., Int. Ed.*, 2008, **47**, 3122; (e) S. Díez-Gonzalez, N. Marion and S. P. Nolan, *Chem. Rev.*, 2009, **109**, 3612; (f) T. Dröge and F. Glorius, *Angew. Chem., Int. Ed.*, 2010, **49**, 6940; (g) N-Heterocyclic Carbenes – From Laboratory Curiosities to Efficient Synthetic Tools, *RSC Catalysis Series*, ed. S. Díez-González, RSC, Cambridge, 2011; (h) H. D. Velazquez and F. Verpoort, *Chem. Soc. Rev.*, 2012, **41**, 7032; (i) R. H. Crabtree, *Coord. Chem. Rev.*, 2013, **257**, 755; (j) L.-A. Schaper, S. J. Hock, W. A. Herrmann and F. E. Kühn, *Angew. Chem., Int. Ed.*, 2013, **52**, 270; (k) K. Riener, S. Haslinger, A. Raba, M. P. Högerl, M. Cokoja, W. A. Herrmann and F. E. Kühn, *Chem. Rev.*, 2014, **114**, 5215; (l) J. A. Mata, F. E. Hahn and E. Peris, *Chem. Sci.*, 2014, **5**, 1723.
- 4 (a) T. Schaub and U. Radius, *Chem.–Eur. J.*, 2005, **11**, 5024; (b) T. Schaub, M. Backes and U. Radius, *J. Am. Chem. Soc.*, 2006, **128**, 15964; (c) T. Schaub, M. Backes and U. Radius, *Organometallics*, 2006, **25**, 4196; (d) T. Schaub, M. Backes and U. Radius, *Chem. Commun.*, 2007, 2037; (e) T. Schaub, C. Döring and U. Radius, *Dalton Trans.*, 2007, 1993; (f) T. Schaub, P. Fischer, A. Steffen, T. Braun, U. Radius and A. Mix, *J. Am. Chem. Soc.*, 2008, **130**, 9304; (g) T. Schaub, M. Backes and U. Radius, *Eur. J. Inorg. Chem.*, 2008, 2680; (h) T. Schaub, M. Backes, O. Plietzsch and U. Radius, *Dalton Trans.*, 2009, 7071; (i) T. Zell, M. Feierabend, B. Halfter and U. Radius, *J. Organomet. Chem.*, 2011, **696**, 1380; (j) T. Zell and U. Radius, *Z. Anorg. Allg. Chem.*, 2011, **637**, 1858; (k) T. Zell, T. Schaub, K. Radacki and U. Radius, *Dalton Trans.*, 2011, 1852; (l) T. Schaub, P. Fischer, T. Meins and U. Radius, *Eur. J. Inorg. Chem.*, 2011, 3122; (m) P. Fischer, K. Götz, A. Eichhorn and U. Radius, *Organometallics*, 2012, **31**, 1374; (n) T. Zell, P. Fischer, D. Schmidt and U. Radius, *Organometallics*, 2012, **31**, 5065; (o) B. Zarzycki, T. Zell, D. Schmidt and U. Radius, *Eur. J. Inorg. Chem.*, 2013, 2051; (p) T. Zell and U. Radius, *Z. Anorg. Allg. Chem.*, 2013, **639**, 334; (q) D. Schmidt, T. Zell, T. Schaub and U. Radius, *Dalton Trans.*, 2014, 10816.
- 5 Danopoulos *et al.* reported the formation of a new signal set in the proton NMR for the thermal decomposition reaction of *cis*-[Ni(iPr₂ImMe₂)₂Me₂] at 50 °C, which was attributed to the formation of the related complex [Ni(iPr₂ImMe₂)₂] (iPr₂ImMe₂ = 1,3-diisopropyl-4,5-dimethylimidazolin-2-ylidene). This complex was never characterized or mentioned thereafter: (a) A. A. Danopoulos and D. Pugh, *Dalton Trans.*, 2008, 30. Own investigations on the related complex *trans*-[Ni(iPr₂Im)₂Me₂] revealed stability with respect to a thermal decomposition to [Ni(iPr₂Im)₂]; (b) P. Fischer, T. Linder and U. Radius, *Z. Anorg. Allg. Chem.*, 2012, **638**, 1491.
- 6 (a) V. P. W. Böhm, C. W. K. Gstöttmayr, T. Weskamp and W. A. Herrmann, *J. Organomet. Chem.*, 2000, **595**, 186; (b) S. Caddick, F. G. N. Cloke, G. K. B. Clentsmith, P. B. Hitchcock, D. McKercher, L. R. Titcomb and M. R. V. Williams, *J. Organomet. Chem.*, 2001, **617–618**, 635; (c) L. R. Titcomb, S. Caddick, F. G. N. Cloke, D. J. Wilson and D. McKercher, *Chem. Commun.*, 2001, 1388; (d) K. Arentsen, S. Caddick, F. G. N. Cloke, A. P. Herring and P. B. Hitchcock, *Tetrahedron Lett.*, 2004, **45**, 3511; (e) C. W. K. Gstöttmayr, V. P. W. Böhm, E. Herdtweck, M. Grosche and W. A. Herrmann, *Angew. Chem., Int. Ed.*, 2002, **41**, 1363; (f) G. Altenhoff, R. Goddard, C. W. Lehmann and F. Glorius, *Angew. Chem., Int. Ed.*, 2003, **42**, 3690; (g) M. M. Konnick, I. A. Guzei and S. S. Stahl, *J. Am. Chem. Soc.*, 2004, **126**, 10212; (h) M. Yamashita, K. Goto and T. Kawashima, *J. Am. Chem. Soc.*, 2005, **127**, 7294; (i) S. Fantasia and S. P. Nolan, *Chem.–Eur. J.*, 2008, **14**, 6987; (j) E. Lee and D. V. Yandulov, *J. Organomet. Chem.*, 2011, **696**, 4095.
- 7 (a) A. J. Arduengo III, S. F. Gamper, J. C. Calabrese and F. Davidson, *J. Am. Chem. Soc.*, 1994, **116**, 4391; (b) P. L. Arnold, F. G. N. Cloke, T. Geldbach and P. B. Hitchcock, *Organometallics*, 1999, **18**, 3228; (c) G. C. Fortman, N. M. Scott, A. Linden, E. D. Stevens, R. Dorta and S. P. Nolan, *Chem. Commun.*, 2010, **46**, 1050; (d) J. Bauer, H. Braunschweig, P. Brenner, K. Kraft, K. Radacki and K. Schwab, *Chem.–Eur. J.*, 2010, **16**, 11985.
- 8 M. Matsumoto, H. Yoshioko, K. Nakatsu, T. Yoshida and S. Otsuka, *J. Am. Chem. Soc.*, 1974, **96**, 3322.
- 9 Details on the reaction of iPr₂Im with the metal halides MCl₂ (M = Pd, Pt) will be given elsewhere.
- 10 (a) P. S. Pregosin, Group 10: Nickel to Platinum, in *Transition Metal Nuclear Magnetic Resonance*, ed. P. S. Pregosin, Elsevier, Amsterdam, 1991, pp. 216–263; (b) R. Benn, H. M. Büch and R.-D. Reinhardt, *Magn. Reson. Chem.*, 1985, **23**, 559.
- 11 (a) S. Caddick, F. G. N. Cloke, P. B. Hitchcock and A. K. d. K. Lewis, *Angew. Chem., Int. Ed.*, 2004, **43**, 5824.
- 12 (a) U. Radius and F. M. Bickelhaupt, *Organometallics*, 2008, **27**, 3410; (b) U. Radius and F. M. Bickelhaupt, *Coord. Chem. Rev.*, 2009, **253**, 678.
- 13 The other example we are aware of involves methylation of an alkynylphosphine ligand: M. A. Bennett, L. Kwan, A. D. Rae, E. Wenger and A. C. Willis, *Dalton Trans.*, 2002, 226.
- 14 (a) F. Bellina and R. Rossi, *Chem. Rev.*, 2010, **110**, 1082; (b) C. C. C. Johansson and T. J. Colacot, *Angew. Chem., Int. Ed.*, 2010, **49**, 676.
- 15 In many cases the formation of stable η²-ketone complexes or orthometallation of the aryl ring is the dominant path for the reaction of benzophenone with a suitable transition metal precursor; for the latter see for example: (a) F. Kakiuchi and S. Murai, *Acc. Chem. Res.*, 2002, **35**, 826.
- 16 Some platinum hydrido α-keto-alkyl complexes are known but have been prepared using other routes: (a) A. Albinati, G. Bracher, D. Carmona, J. H. P. Jans, W. T. Klooster, T. F. Koetzle, A. Macchioni, J. S. Ricci, R. Thouvenot and L. M. Venanzi, *Inorg. Chim. Acta*, 1997, **265**, 255; (b) C. E. Shuchart, R. R. Willis, A. Wojcicki, A. L. Rheingold and B. S. Haggerty, *Inorg. Chim. Acta*, 2000, **307**, 1; (c) T. F. Vaughan, D. J. Koedyk and J. L. Spencer, *Organometallics*, 2011, **30**, 5170.



- 17 (a) G. te Velde, F. M. Bickelhaupt, E. J. Baerends, C. Fonseca Guerra, S. J. A. van Gisbergen, J. G. Snijders and T. Ziegler, *J. Comput. Chem.*, 2001, **22**, 931; (b) F. M. Bickelhaupt and E. J. Baerends, in *Reviews in Computational Chemistry*, ed. K. B. Lipkowitz and D. B. Boyd, Wiley-VCH, New York, 2000, vol. 15, pp. 1–86; (c) I. Fernandez and F. M. Bickelhaupt, *Chem. Soc. Rev.*, 2014, **43**, 4953.
- 18 See, for example: (a) T. Ziegler and J. Autschbach, *Chem. Rev.*, 2005, **105**, 2695; (b) J. Li and T. Ziegler, *Organometallics*, 1996, **15**, 3844; (c) S. Dapprich and G. Frenking, *Angew. Chem., Int. Ed.*, 1995, **34**, 354.
- 19 (a) M. J. S. Dewar, *Bull. Soc. Chim. Fr.*, 1951, 71; (b) J. Chatt and L. A. Duncanson, *J. Chem. Soc.*, 1953, 2939.
- 20 G. Frenking and N. Fröhlich, *Chem. Rev.*, 2000, **100**, 717.
- 21 (a) K. Kitaura, S. Sakaki and K. Morokuma, *Inorg. Chem.*, 1981, **20**, 2292; (b) T. Ziegler, *Inorg. Chem.*, 1985, **24**, 1547; (c) J. Li, G. Schreckenbach and T. Ziegler, *Inorg. Chem.*, 1995, **34**, 3245.
- 22 (a) T. A. Albright, J. K. Burdett and M. H. Whangbo, *Orbital Interactions in Chemistry*, Wiley & Sons, New York, 1985; (b) K. Tatsumi, R. Hoffmann, A. Yamamoto and J. K. Stille, *Bull. Chem. Soc. Jpn.*, 1981, **54**, 1857; (c) M.-D. Su and S.-Y. Chu, *Inorg. Chem.*, 1998, **37**, 3400; (d) M. A. Carvajal, J. J. Novoa and S. Alvarez, *J. Am. Chem. Soc.*, 2004, **126**, 1465; For consequences for the reactivity, see: (e) L. P. Wolters and F. M. Bickelhaupt, *ChemistryOpen*, 2013, **2**, 106; (f) L. P. Wolters, W. J. van Zeist and F. M. Bickelhaupt, *Chem.–Eur. J.*, 2014, **20**, 11370.
- 23 See for example: (a) M. Brendel, C. Braun, F. Rominger and P. Hofmann, *Angew. Chem., Int. Ed.*, 2014, **53**, 8741; (b) S. N. Sluijter, S. Warsink, M. Lutz and C. J. Elsevier, *Dalton Trans.*, 2013, 7365.

