

CrossMark  
click for updatesCite this: *Chem. Sci.*, 2015, 6, 3354

# Accurate molecular weight determination of small molecules *via* DOSY-NMR by using external calibration curves with normalized diffusion coefficients†‡

Roman Neufeld and Dietmar Stalke\*

Determination of the aggregation and solvation numbers of organometallic complexes in solution is an important task to increase insight in reaction mechanisms. Thus knowing which aggregates are formed during a reaction is of high interest to develop better selectivity and higher yields. Diffusion-ordered spectroscopy (DOSY), which separates NMR signals according to the diffusion coefficient, finds increasing use to identify species in solution. However, there still is no simple relationship between diffusion coefficient and molecular weight (MW). Some methods have been developed to estimate the MW but still with a significant error of  $\pm 30\%$ . Here we describe a novel development of MW-determination by using an external calibration curve (ECC) approach with normalized diffusion coefficients. Taking the shape of the molecules into account enables accurate MW-predictions with a maximum error of smaller than  $\pm 9\%$ . Moreover we show that the addition of multiple internal references is dispensable. One internal reference (that also can be the solvent) is sufficient. If the solvent signal is not accessible, 16 other internal standards (aliphatics and aromatics) are available to avoid signal overlapping problems and provide flexible choice of analytes. This method is independent of NMR-device properties and diversities in temperature or viscosity and offers an easy and robust method to determine accurate MWs in solution.

Received 23rd February 2015  
Accepted 18th March 2015

DOI: 10.1039/c5sc00670h

www.rsc.org/chemicalscience

## Introduction

Chemists have always had a vital interest in the size of molecules. Especially the aggregation and solvation numbers of organometallic intermediates play a dominant role in reaction mechanisms and product forming.<sup>1</sup> Therefore the knowledge of reactive aggregates is a prerequisite to understand how molecules react and why they form which products.<sup>2</sup> Apart from colligative property measurements, such as cryoscopy,<sup>3</sup> diffusion-ordered spectroscopy (DOSY) has gained increasing importance in this area.<sup>4</sup> The physical observable that can be derived from the diffusion NMR experiment is the diffusion coefficient  $D$  that is sensitive to size and shape of the molecular species.<sup>5</sup> A number of empirical methods for relating diffusion coefficients to the molecular weight (MW) have been proposed.<sup>6</sup> Often the Stokes–Einstein equation<sup>7</sup> and its modifications<sup>8</sup> are useful and enable molecular size estimation of large particles

that are much larger than the solvent.<sup>9</sup> Besides that, especially the empirically derived power law (eqn (1)) is probably the most powerful class of relations which correlates the MW and the diffusion coefficient according to

$$D = KMW^\alpha \quad (1)$$

This power law<sup>10</sup> gives good results but is restricted to a specific class of compounds.<sup>11</sup> Especially the polymer community applied it to estimate the MW distribution of polymer solutions like *e.g.* globular proteins,<sup>11</sup> oligosaccharides,<sup>12</sup> polyethylene-oxides<sup>13</sup> and denatured peptides<sup>14</sup> in variable solvents. Even small molecules correlate to the power law, as demonstrated in the work of Crutchfield and Harris.<sup>15</sup> Unfortunately this work only allows a MW estimation with a relatively high error of  $\pm 30\%$ . For small organometallic molecules Li and Williard *et al.* have used an analogue approach of the power law by introducing at least three internal references to one NMR sample in order to get an internal calibration curve (ICC).<sup>16</sup> For small molecules this method gives much better results. They were able to characterize for example THF-solvated LDA to be dimeric<sup>17</sup> and a 1 : 1 mixture of LiHMDS with HMPA (hexamethylphosphoramide) to adopt a disolvated dimeric structure in TOL- $d_8$  solution.<sup>18</sup> The same ICC-method was employed by

Institut für Anorganische Chemie, Georg-August-Universität, Tammannstrasse 4, Göttingen, Germany. E-mail: dstalke@chemie.uni-goettingen.de

† In memory to Professor Paul von Ragué Schleyer, the protagonist of organolithium chemistry.

‡ Electronic supplementary information (ESI) available: Includes detailed information about the calibration curves, the model compounds and the calculation of the molar van der Waals-density. See DOI: 10.1039/c5sc00670h



Armstrong and Mulvey *et al.* to characterize diisopropylamide and 2,2,6,6-tetramethylpiperidine (TMP) turbo-Grignard reagents in THF solution.<sup>19</sup> Unfortunately the ICC-method has some important disadvantages: on the one hand the ICC employs just a few references (mostly 3) and is often based on a small molecular weight distribution. On the other hand each ICC is only useful for one NMR sample. Additionally all of the internal references are limited to a lot of prerequisites: (a) they should be inert to the analyte in solution; (b) the NMR signals should not overlap with other components; (c) the internal references should have no coordinating ability to the analyte; (d) they should be well soluble in the solvent and finally (e) the internal references should have a well set molecular weight distribution.<sup>16</sup> Therefore it is a big challenge to choose the appropriate internal references, because often at least one of the above-mentioned exigencies would not be met.

In this article we describe the use of an external calibration curve (ECC) approach with fixed diffusion coefficients. This method allows highly accurate MW-determination without the necessity of multiple references that disturb the measurements. Furthermore we show for the first time how solvent temperature, concentration, the shape and the mass density of analytes influence the power law derived MW-determination. And finally we use ECCs to determine the aggregation of Na-indenide in THF-*d*<sub>8</sub> solution.

## Results and discussions

### Application of a normalized reference system with fixed diffusion coefficients

DOSY spectra are frequently affected by various sources of errors like *e.g.* diversities in temperature, fluctuation, convection, viscosity and concentration effects. Additionally the NMR-device constants like *e.g.* gradient strength and pulse duration influence the absolute diffusion coefficient. So it becomes clear that DOSYs only can be compared under *ceteris paribus* conditions. To overcome the complications of these effects and to enable tabulated diffusion coefficients, the use of an internal standard is necessary. Those standards provide more resilient diffusion coefficients for any changes in the NMR sample.<sup>20</sup> This ratio, often termed as relative diffusivity  $D_{\text{rel}}$  is defined as:

$$D_{\text{rel}} = D_{\text{x}}/D_{\text{ref}} \quad (2)$$

where ref and x correspond to the reference and analyte, respectively. This approach reduces the impact of viscosity and temperature and provides more robust data.<sup>21</sup> Besides the above-mentioned advantages this method has a disadvantage. Eqn (2) produces relative diffusion values that always depend on the one reference molecule it has been referenced for. This reference has no strict or defined diffusion value. We found that it is advantageous to employ relative diffusion coefficients with fixed diffusion values. Subsequently we show that this approach allows not only one molecule to act as a reference rather than every compound that is part of the calibration curve to act as an internal reference. We realized that the logarithmic diffusion values are connected approximately linearly, independent of

gradient and magnetic field strength, gyromagnetic ratio, gradient pulse duration, and changes in temperature or viscosity. That is why the linear eqn (3) was empirically derived:

$$\log D_{\text{x,norm}} = \log D_{\text{ref,fix}} - \log D_{\text{ref}} + \log D_{\text{x}} \quad (3)$$

where  $\log D_{\text{ref,fix}}$  is the fixed value of the reference,  $\log D_{\text{ref}}$  the measured diffusion coefficient of the reference,  $\log D_{\text{x}}$  the diffusion coefficient of analyte x and  $\log D_{\text{x,norm}}$  the relative diffusion value of the analyte x normalized to the reference. Eqn (3) ensures that all molecules get a normalized diffusion coefficient relative to the internal reference. For our measurements we picked for all TOL-*d*<sub>8</sub> solvates adamantane (ADAM) and for all THF-*d*<sub>8</sub> solvates 2,2,3,3-tetramethylbutane (TMB) as internal references (see Table 1). The validity of eqn (3) was proved by measuring DOSY spectra of various model compounds on two different NMR spectrometer devices, where one spectrometer had calibrated and the other uncalibrated gradients. Although differences in gradient calibration, temperature and concentration automatically give deviations in absolute diffusion (see ESI, S-Fig. 1†), eqn (3) provides excellent results for the normalized diffusion coefficient. It is independent of the NMR spectrometer or sample diversity with an average standard deviation of only  $\sigma(\log D_{\text{x,norm}}) = 0.003$  in TOL-*d*<sub>8</sub> and 0.002 in THF-*d*<sub>8</sub> (ESI, S-Table 1 and S-Table 2†). This experiment demonstrates the robustness of eqn (3) and the normalized diffusion coefficients.

### External calibration curves and internal references

The MW-determination developed by Li and Williard *et al.* relies on an internal calibration curve (ICC) that has been generated by a single DOSY measurement where all references are present in the same NMR sample. The calibration curves which are presented in this article were generated by measuring 28 model compounds in independent NMR samples. That is the reason why we name these calibration curves “external”. The calibration curves have been plotted the common way by linearizing<sup>22</sup> the power law (2) with taking the logarithm of both sides (see ESI,† chapter IV).<sup>23</sup> Plotting  $\log D$  versus  $\log \text{MW}$  of all model molecules in one diagram gives a linear correlation but with a significant deviation especially for the very low and higher weight molecules that prevents accurate MW-determination (Fig. 1). To find a better correlation between diffusion coefficient and the MW we generated three dimensional, shape

Table 1  $\log D_{\text{ref,fix}}$  values of the used internal references

Internal reference	$\log D_{\text{ref,fix}}^a$
ADAM in TOL- <i>d</i> <sub>8</sub> <sup>b</sup>	−8.8454
TMB in THF- <i>d</i> <sub>8</sub> <sup>c</sup>	−8.7749

<sup>a</sup> Each diffusion coefficient was estimated by using the middle  $\log D$  value of 15 DOSY measurements of 15 mM solutions at 25 °C.

<sup>b</sup> ADAM has two signals in the <sup>1</sup>H-NMR-spectrum. For determining the diffusion coefficient, we always used the signal of the −CH<sub>2</sub> groups with the highest intensity. <sup>c</sup> The chemical shift of one ADAM signal is very close to the THF-*d*<sub>7</sub> signal at 1.73 ppm. Therefore we used TMB as internal reference for all THF-*d*<sub>8</sub> solvates.






Fig. 1  $\log D$  versus  $\log MW$  in TOL- $d_8$ . All compounds were normalized to  $\log D_{\text{ref,fix}}(\text{ADAM}) = -8.8454$ , see Table 4.

optimized models of all model compounds. By comparing the diffusion coefficient of each molecule with its shape and compactness features, one can see a significant disparity that prompted us to separate the molecules into three different types: (1) compact spheres (CS), (2) dissipated spheres and ellipsoids (DSE) and (3) expanded discs (ED) (Fig. 1).<sup>24</sup> Of course the transitions between those types occur across a foggy line but there are clear systematic trends that can be rationalized. From Fig. 2 it is obvious that CS have nearly the same radius in all dimensions with a high-density space like for example the compact molecules ADAM or  $\text{Si}(\text{SiMe}_3)_4$ .

However, the majority of small molecules diffuse like DSE. These are either spherical-like molecules that are less compact than CS (e.g. compounds with dative bonds) and/or ellipsoidal molecules like e.g. tetramethoxypropane or 2,2-bis(diphenylphosphino)-1,1-binaphthyl (BINAP). Small aromatic compounds like toluene, indene or naphthalene with  $MW < 160 \text{ g mol}^{-1}$  diffuse also like DSE molecules. The significance of two dimensional geometries begins approximately at  $MW > 178 \text{ g mol}^{-1}$ . This is the region where the type of the ED begins, including molecules like anthracene ( $178 \text{ g mol}^{-1}$ ) or tetraphenyl-naphthalene (TPhN,  $433 \text{ g mol}^{-1}$ ). Depending on these different molecular types<sup>25</sup> we plotted their  $\log D$  versus  $\log MW$  in one separate plot. These give excellent linear fits with a small error and very high  $R^2$  values of  $R^2 \approx 1$  (see ESI, S-Fig. 2 and S-Fig. 3†). Those fits result in six different ECCs:  $\text{ECC}_{\text{CS}}^S$ ,  $\text{ECC}_{\text{DSE}}^S$

and  $\text{ECC}_{\text{ED}}^S$ , each for every solvent ( $S = \text{TOL-}d_8$  or  $\text{THF-}d_8$ ). The linear fit parameters are summarized in Table 2. We also merged all calibration curves to generate a merged  $\text{ECC}_{\text{merge}}^S$ .

$\alpha$  is related to the Flory exponent that comes from the fractal theory and can be described as a measure of compactness of a molecular shape. A Flory exponent of  $-\alpha = 0.33$  notes that the space is totally filled and no holes are left. On the other hand a Flory exponent of  $-\alpha = 1$  means that the molecule is completely one-dimensional and extends linearly like a rigid rod.<sup>11</sup> The estimated  $\alpha$ -values from the ECCs stay in very good agreement with the Flory exponent. The ECCs for compact spheres have a low  $\alpha$ -value ( $-\alpha \approx 0.5$ ) and the ECCs for the less compact spheres and ellipsoids have a higher  $\alpha$ -value ( $-\alpha \approx 0.6$ ). The two dimensional huge discs have the highest Flory exponent ( $-\alpha \approx 0.8$ ) like expected.

When internal calibration curves (ICC) are used, then all references have to be in the same sample. The diffusion coefficients of those internal references show a linear dependency. In our external calibration curves (ECC), (where each model compound has been measured with ADAM or TMB as internal reference), we also see a linear behavior. These compounds behave as they were all measured in the same NMR sample. Therefore the idea occurred that beside ADAM and TMB, basically all model compounds could act as internal references for the ECCs, according to:

$$\log D_{x,\text{norm}} = \log D_{\text{ref,fix}}^* \quad (4)$$

Consequently we measured DOSYs with some model compounds (e.g. ADAM +  $\text{Si}(\text{SiMe}_3)_4$  + naphthalene in TOL- $d_8$ ) in the same NMR sample and used every compound as an internal reference by applying eqn (4). In fact it is possible to determine accurate MW of all compounds by using the normalized  $\log D_{x,\text{norm}}$  value as fixed reference (Table 3). Utilizing e.g. TOL- $d_7$  as internal reference, ADAM,  $\text{Si}(\text{SiMe}_3)_4$  and naphthalene can be determined with an average deviation of  $\pm 5\%$ . That means that the “real” molecular weight was missed by only 5%, although we used a calibration curve on the basis of many references that were not present in this NMR sample. With the ECC it is possible to simulate a bench of internal references by adding just one of them to the NMR sample. All 28 compounds behave like they were all measured in the same NMR sample. This interrelation has the colossal advantage that it is unnecessary to introduce all references into the same NMR sample. Signal overlapping, analyte-reference interaction problems,




Fig. 2 Example molecules that were classified in our calibration curves as CS, DSE and ED like molecules.

Table 2 Linear fit parameter for the four ECCs each for TOL- $d_8$  and THF- $d_8$  solutions

S	TOL- $d_8$		THF- $d_8$	
	$-\log K$	$-\alpha$	$-\log K$	$-\alpha$
$\text{ECC}_{\text{CS}}^S$	7.7581	0.5018	7.7427	0.4943
$\text{ECC}_{\text{DSE}}^S$	7.5197	0.6098	7.5360	0.5824
$\text{ECC}_{\text{ED}}^S$	7.1008	0.7836	7.1205	0.7519
$\text{ECC}_{\text{merge}}^S$	7.4595	0.6318	7.4664	0.6095



**Table 3** Mixed composition of compounds (each 15 mM) in TOL- $d_8$  acting themselves as internal reference for the ECC<sup>TOL</sup>-MW-determination

Analyte	MW [g mol <sup>-1</sup> ]	Ref 1 TOL- $d_7$		Ref 2 ADAM		Ref 3 Si(SiMe <sub>3</sub> ) <sub>4</sub>		Ref 4 naphthalene	
		MW <sub>det</sub> [g mol <sup>-1</sup> ]	$\Delta$ MW [%]	MW <sub>det</sub> [g mol <sup>-1</sup> ]	$\Delta$ MW [%]	MW <sub>det</sub> [g mol <sup>-1</sup> ]	$\Delta$ MW [%]	MW <sub>det</sub> [g mol <sup>-1</sup> ]	$\Delta$ MW [%]
TOL- $d_7^b$	99	96	3	97	2	96	3	97	2
ADAM <sup>a</sup>	136	144	-6	147	-8	144	-6	145	-7
Si(SiMe <sub>3</sub> ) <sub>4</sub> <sup>a</sup>	321	304	5	309	4	303	5	305	5
Naphthalene <sup>b</sup>	128	122	5	124	3	122	5	122	5

<sup>a</sup> ECC<sup>TOL</sup><sub>CS</sub> was used to calculate the MW. <sup>b</sup> ECC<sup>TOL</sup><sub>DSE</sub> was used to calculate the MW.

wasting chemicals and deuterated solvents can be avoided, *i.e.* saved.

Due to the normalized diffusion coefficients everyone can use the ECCs, independent of the NMR device, without always recording new calibration curves. Inert compounds that are suitable to act as internal reference and their log  $D_{\text{ref,fix}}^*$  values are summarized in Table 4. In the next section the use of the residual solvent signal of TOL- $d_8$  and THF- $d_8$  (that arises from the proton of isotopomers containing one less deuterium atom than the perdeuterated solvent)<sup>26</sup> as internal reference is examined in detail. In the following those isotopomers are referred to as TOL- $d_7$  and THF- $d_7$ , respectively.

#### Quality of TOL- $d_7$ and THF- $d_7$ as internal reference

To compare the quality of the ECCs we calculated the MWs of all model compounds by using the log  $D_{\text{ref,fix}}^*$  values of TOL- $d_7$  and THF- $d_7$  as internal references. The MWs of the model compounds were determined by using the appropriate ECCs. The MW of cyclopentane (Fig. 3, very left point,  $\Delta$ MW = 7%) was

*e.g.* determined by using ECC<sub>CS</sub> and TPhN (Fig. 3, very right point,  $\Delta$ MW = 0%) by using ECC<sub>ED</sub>. Fig. 3 shows that the quality of MW-determination is reference-independent. It does not matter if ADAM/TMB or TOL- $d_7$ /THF- $d_7$  were used as internal reference. Both give excellent MW-predictions with a standard deviation of 4%. The maximum error in both solvents is  $\pm 9\%$  so we postulate that this is the maximum resolution of this DOSY method.

#### Influence of the shape

There are several modifications of the Stokes–Einstein equation which take the molecule's shape into account (by adding *e.g.* correction factors).<sup>6,8</sup> The power law derived MW-determinations distinguish mostly between compound classes,<sup>11</sup> large<sup>9</sup> and small molecules,<sup>15</sup> but not directly between different shapes within those molecular classes. In this section we will demonstrate that the accuracy of the power law derived MW-prediction is highly affected, also for small molecules, by the analyte's shape. To validate this issue we determined the MWs of all

**Table 4** Overview of all ECC-adapted references that fulfill the requirement of internal references for 15 mM solutions. All TOL- $d_8$  solvates were normalized to ADAM and all THF- $d_8$  solvates were normalized to TMB

MW [g mol <sup>-1</sup> ]	Compound <sup>a</sup>	TOL- $d_8$ log $D_{\text{ref,fix}}^*$	THF- $d_8$ log $D_{\text{ref,fix}}^*$
70	Cyclopentane	-8.6694	-8.6437
79	THF- $d_7^b$	—	-8.6335
88	TMS	-8.7445	-8.7018
92	TOL	—	-8.6742
99	TOL- $d_7^d$	-8.7289	—
114	TMB	-8.7963	-8.7749 <sup>c</sup>
116	Indene	-8.7698	-8.7325
128	Naphthalene	-8.7932	-8.7461
136	ADAM <sup>e</sup>	-8.8454 <sup>c</sup>	—
178	Diphenylacetylene	-8.9095	-8.8535
178	Anthracene	-8.8574	-8.8129
192	9-Methylanthracene	-8.8824	-8.8440
202	Pyrene	-8.8960	-8.8457
204	1-Phenylnaphthalene	-8.9184	-8.8812
228	Triphenylene	-8.9552	-8.8869
321	Si(SiMe <sub>3</sub> ) <sub>4</sub>	-9.0038	-8.9773
433	Tetraphenylnaphthalene	-9.1660	-9.1054

<sup>a</sup> When a compound had more than one <sup>1</sup>H signal, the average diffusion coefficient was used. <sup>b</sup> Due to the very high access of the solvent, the signal of THF- $d_7$  can be used as internal reference, but a higher MW<sub>det</sub> error can occur, when the solvent is coordinating to *e.g.* a metal. <sup>c</sup> The “original” log  $D_{\text{ref,fix}}^*$  values that were used for all ECCs. <sup>d</sup> We calculated the middle diffusion coefficients of the three aromatic protons. The final diffusion coefficient was calculated by middling this value with the diffusion coefficient of the methyl group at 1.73 ppm. <sup>e</sup> ADAM has two signals in the <sup>1</sup>H-spectrum. For determining the diffusion coefficient, we always used the signal of the -CH<sub>2</sub> groups with the highest intensity.








Fig. 3 Using (A) ADAM/TOL- $d_7$  or (B) TMB/THF- $d_7$  as internal reference (15 mM) gives a good MW-determination with a standard deviation of  $\sigma = 4\%$ . All of these model compounds were used to derive the ECCs.

compounds using *e.g.* the  $ECC_{ED}^S$  for expanded discs or  $ECC_{DSE}^S$  for ellipsoidal model molecules, *etc.* (Fig. 4). When for example the  $ECC_{CS}^S$  (that is for compact spherical molecules) is used on expanded disc like molecules, the determined MW will have a massive error especially for big molecules (Fig. 4A). Anticipating for example TPhN (an expanded flat disc,  $433 \text{ g mol}^{-1}$ ) in TOL- $d_8$  to be spherical would produce a  $MW_{det}$  of  $639 \text{ g mol}^{-1}$  that is a 48% overestimated mass (Fig. 4A). Using *e.g.* the  $ECC_{ED}^S$  for non-oblate molecules would produce especially for small molecules  $<170 \text{ g mol}^{-1}$  a large error (Fig. 4C). The merged calibration curve  $ECC_{merge}^S$  (Fig. 4D) determines MWs in a range of  $\Delta MW = \pm 23\%$ . But the deviation is much smaller in the region of approximately  $120\text{--}200 \text{ g mol}^{-1}$ . On one hand that means that in this MW-region all molecules diffuse more or less independently from their shape. On the other hand the MW-determination (and the self-diffusion) of molecules that are outside that region, is increasingly influenced by the analyte's shape. Using the wrong calibration curve (or for example wrong molecules for an ICC) can produce highly erroneous MW-values. This is the reason why other calibration curves<sup>8a,15,27</sup> that correlate a bundle of different molecules without considering the right shape, lead to bigger deviations from the correct MWs in the range of  $\Delta MW = \pm 30\%$ . By taking the correct shape into account it is possible to determine more accurate MWs with a deviation of  $\Delta MW < \pm 9\%$  (see Fig. 3).




Fig. 4 Using exclusively (A)  $ECC_{CS}^S$ , (B)  $ECC_{DSE}^S$ , (C)  $ECC_{ED}^S$  and (D)  $ECC_{merge}^S$  on all model compounds (15 mM).



### Influence of the concentration on TOL- $d_8$ solvates

All above-mentioned ECCs were derived from 15 mM solutions. We wanted to test how good the ECC-MW-determination works when the concentration is much higher than 15 mM. Therefore DOSYs of some model compounds have been measured at concentrations of 120 mM. The results are shown in Fig. 5. The Stokes–Einstein equation is only valid for infinite diluted solutions. Therefore the error should be much bigger with high concentration solutions. However, the average deviation of the estimated MW is only a little higher and most of the compounds are still in the  $\pm 9\%$  range. The biggest error arises probably due to intermolecular interactions that result in higher estimated MWs. Especially  $\pi$ -interactions of big aromatic systems, at high concentrations could significantly increase the error of estimated MWs. Anyway, all compounds without aromatic rings have been determined within  $\pm 9\%$  deviation. The power law seems to be valid even for higher concentrations, if inter- or intramolecular interactions can be excluded.<sup>28</sup>

### Influence of the temperature

According to the Stokes–Einstein equation the self-diffusion should be inversely proportional to the viscosity. Indeed, increasing the solvent viscosity by cooling the NMR sample from room temperature to  $-75^\circ\text{C}$  leads to an increase of the diffusion coefficient by almost two magnitudes! Thanks to the internal reference, the ECC-MW-determination of  $\text{Si}(\text{SiMe}_3)_4$  (TTS,  $321\text{ g mol}^{-1}$ ) in the full range of  $-75^\circ\text{C}$  to  $+100^\circ\text{C}$  still gives good results (Fig. 6). The internal reference is able to compensate for viscosity changes in the solvent.<sup>23</sup> Notably the MW-determination is still possible at temperatures close to the boiling point of the solvent. This would give the opportunity to observe species that are forming during reactions at elevated temperatures. Furthermore, it is obvious that the signal of TOL- $d_7$  is a useful internal reference for both high and low temperatures. But using more polar THF- $d_7$  below  $-50^\circ\text{C}$  can get problematic. This probably results from solvent–solvent interactions, *i.e.* hydrogen bonding. Anyhow, it is advisable to use non-polar references and low concentrations for low temperature measurements.<sup>28</sup>




Fig. 5 ECC-MW-determination of a few model compounds at a concentration of 120 mM at  $25^\circ\text{C}$ . ADAM was used in equimolar concentration as internal reference.




Fig. 6 ECC-MW-determination of TTS (15 mM) in (A) TOL- $d_8$  and in (B) THF- $d_8$  at different temperatures.

### Influence of halides and molecular density

In the Stokes–Einstein equation the diffusion coefficient relies on the shape and on the hydrodynamic radius of the particle. The latter can be described by the volume of the surrounding solvent molecules, the electron cloud and also the volume of the atoms. One has to keep in mind that the volume of an atom is not proportional to its atom weigh. Especially halides, compared to their atomic radii, have very high atomic masses and therefore a high mass density. For instance a potassium cation has an ion radius of 138 pm and an atom weight of  $39\text{ g mol}^{-1}$ . An iodine atom has almost the same radius of 133 pm but an atom weight of  $127\text{ g mol}^{-1}$  that is 320% bigger than that of the  $\text{K}^+$  cation.<sup>29</sup> Our ECCs were elaborated with references that consist mostly of hydrocarbons with some heteroatoms of the third period like silicon, phosphorus and sulfur. Therefore especially compounds containing heavy halides will be underestimated in MW. While chlorine-containing compounds are estimated with good accuracies the much denser bromides show bigger deviations from the correct MWs (see Table 5). Especially an increasing halide/carbon ratio leads to bigger errors. For example triphenylmethylbromide with one bromine atom is underestimated by 12%. But 9,10-dibromoanthracene with two bromides, a small carbon backbone and therefore a very high molar density is underestimated in MW by 42%. That means that the power law depends heavily on the molecular density.



Table 5 ECC-MW-determination of molecules with halides

Compound	MW [g mol <sup>-1</sup> ]	MW <sub>det</sub> <sup>a</sup> [g mol <sup>-1</sup> ]	ΔMW [%]	MD <sub>w</sub> /10 <sup>29</sup> [g mol <sup>-1</sup> m <sup>-3</sup> ]
1-Hexylchloride	120	117	2	5.49
1-Octylchloride	149	143	4	5.29
1-Decylchloride	177	176	1	5.13
1-Propylbromide	123	82	34	9.66
Triphenylmethylbromide	323	283	12	6.45
9,10-Dibromoanthracene	336	194 <sup>b</sup>	42	8.71
1-Butyliodide	184	102	45	11.15

<sup>a</sup> ECC<sub>DSE</sub> was used to determine the MW. <sup>b</sup> ECC<sub>ED</sub> was used to determine the MW.

There are more or less extensive ways to calculate the density of a molecule. We decided to derive a simple but robust equation that correlates the MW to the approximated volume of a compound. Therefore eqn (5) was derived, where MW is the molecular weight and  $V_w$  is the van der Waals-volume of an atom. In respect to this equation we calculated the van der Waals-volumes of all atoms of a compound and summed them up to one single van der Waals-sphere ( $V_{sph}$ ) (see ESI, S-Table 11†).<sup>30</sup> Of course this method is just an approximation without considering the real covalent bond-bond distances and the shape of the compounds. But the ratio between the MW and the sum of all van der Waals-volumes ( $V_w$ ) leads to a value that represents approximately a molar van der Waals-density (MD<sub>w</sub>) in a unit of g mol<sup>-1</sup> m<sup>-3</sup>:

$$MD_w = MW / \sum V_w = MW / V_{sph} \quad (5)$$

Plotting MD<sub>w</sub> against MW give for our model compounds an average density distribution of around  $5.2 \times 10^{29}$  g mol<sup>-1</sup> m<sup>-3</sup> (Fig. 7).

Obviously the ECCs presented in this article work well with molecules with a molar density between  $4.3 \times 10^{29}$  g mol<sup>-1</sup> m<sup>-3</sup> and  $5.9 \times 10^{29}$  g mol<sup>-1</sup> m<sup>-3</sup>. Higher molar van der Waals-densities like for example 9,10-dibromoanthracene [MD<sub>w</sub> =  $8.7 \times 10^{29}$  g mol<sup>-1</sup> m<sup>-3</sup>] will be underestimated and lower MD<sub>w</sub>

values will be overestimated in MW. To obtain accurate MWs for molecules with high densities it is necessary to measure a new calibration curve with references of comparable molar densities and geometries. Additional studies are in progress.

### Deuterated compounds

The MW-estimation of the residual solvent peak (THF-*d*<sub>7</sub>, 79 g mol<sup>-1</sup>) gives a MW of 63 g mol<sup>-1</sup> that would be underestimated in MW (ΔMW = 20%). The determined MW fits much better to the protonated THF-*h*<sub>8</sub> species (72 g mol<sup>-1</sup>, ΔMW = 8%). This is congruent to the nearly similar atomic radius of D compared to H. According to the above-mentioned correlation of the atomic volume and the corresponding diffusion coefficient, it is clear that deuterated molecules diffuse approximately like their protonated counterparts, although they have slightly bigger MWs. In the case of TOL-*d*<sub>7</sub> (99 g mol<sup>-1</sup>) this effect is less pronounced (MW<sub>det</sub> = 96 g mol<sup>-1</sup>, ΔMW = 3% rel. to TOL-*d*<sub>7</sub>, ΔMW = -5% rel. to TOL-*h*<sub>8</sub>) due to the relative higher MW of toluene. Moreover, especially in the case of multiple THF-*d*<sub>8</sub>-coordination it is advisable to use the molecular weight of THF-*h*<sub>8</sub> to have an accurate MW-determination.

### Determining the MW of alkali organometallics

On the one hand organometallic compounds tend to aggregate *via* coordinative bonds that are significantly longer than covalent bonds. Additionally solvent molecules can associate and dissociate in solution.<sup>31</sup> Therefore we anticipate that the space between all atoms is less packed than in the “sigma bonded-compact spheres” model. On the other hand alkaline organometallics frequently adopt spherical and ellipsoidal shapes, according to the ring-stacking principle.<sup>32</sup> This is why we think that the ECC<sub>DSE</sub><sup>S</sup> for “dissipated spheres and ellipsoids” is the best calibration curve for s-block organometallic compounds. As a proof of principle we made an ECC<sub>DSE</sub><sup>THF</sup>-MW-determination of LDA in THF-*d*<sub>8</sub> solution<sup>33</sup> (15 mM). THF solvated LDA is known to exist exclusively as a disolvated dimer (MW = 358 g mol<sup>-1</sup>).<sup>34</sup> In fact, using ECC<sub>DSE</sub><sup>THF</sup> estimates a MW of MW<sub>det</sub> = 347 g mol<sup>-1</sup> with a deviation of only 3% (see Table 6).

Recently we showed that *ortho* lithium dimethylaniline (Me<sub>2</sub>NC<sub>6</sub>H<sub>4</sub>Li) crystallizes in the presence of *tert*-butyl lithium (*t*-BuLi) as a separated lithium organic aggregate [*t*-BuLi]<sub>4</sub>·4 [Me<sub>2</sub>NC<sub>6</sub>H<sub>4</sub>Li]<sub>4</sub> in the same crystal.<sup>41</sup> Dissolving those crystals in




Fig. 7 MD<sub>w</sub> distribution in the model compounds and molecules with heavy atoms.



**Table 6** ECC-MW-determination of various lithium organics in solution. ECC<sub>DSE</sub> was used to determine the MWs

Species	MW	MW <sub>det</sub>	ΔMW
[(Me <sub>2</sub> CH) <sub>2</sub> NLi(THF)] <sub>2</sub>	358	347 <sup>a</sup>	3
[Me <sub>2</sub> NC <sub>6</sub> H <sub>4</sub> Li] <sub>4</sub>	508	527 <sup>b</sup>	−4
[(Me <sub>2</sub> NC <sub>6</sub> H <sub>4</sub> Li) <sub>3</sub> ( <sup>t</sup> BuLi)]	445	435 <sup>b</sup>	2
[(Me <sub>2</sub> NC <sub>6</sub> H <sub>4</sub> Li) <sub>2</sub> ( <sup>t</sup> BuLi) <sub>2</sub> ]	382	367 <sup>b</sup>	4
[(Me <sub>2</sub> NC <sub>6</sub> H <sub>4</sub> Li)( <sup>t</sup> BuLi) <sub>3</sub> ]	319	316 <sup>b</sup>	1
[ <sup>t</sup> BuLi] <sub>4</sub>	256	244 <sup>b</sup>	5

<sup>a</sup> 1-Phenylnaphthalene was used as internal reference. <sup>b</sup> TOL-*d*<sub>7</sub> was used as internal reference.

TOL-*d*<sub>8</sub> resulted in an unexpectedly complicated <sup>7</sup>Li NMR spectrum that shows five relatively sharp distinguishable signals over a range of nearly 2.5 ppm. These compounds were analysed by <sup>7</sup>Li-DOSY experiments and were anticipated to belong to homo and heteroleptic tetramers of *o*-lithium anilide and <sup>t</sup>BuLi. The ECC<sub>DSE</sub><sup>THF</sup>-MW-determination confirms those results (see Table 6).

### Determining the MW of Na-indenide in THF

Alkali metal indenides are important precursors for the synthesis of metallocenes of the main group and transition metals. Without donating ligands like THF, dimethoxyethane (DME) or chelating crown ethers *etc.* they build up polymeric stack structures.<sup>35</sup> The solid-state structure of base-free Na-indenide is unknown. With donating ligands Li- and Na-indenide with PMDETA or crown ethers form contact ion pairs (CIP)<sup>36</sup> and with ammonia solvent separated ion pairs (SSIP).<sup>37</sup> There are no investigations about the aggregation of Na-indenide in solution. One reason for that may be the relatively bad NMR properties of the sodium nucleus that has a spin of 3/2.<sup>38</sup> This quadrupole results in broad lines that get even broader with asymmetry of the environment. The <sup>1</sup>H-DOSY experiment is independent of that nucleus. Therefore Na-indenide is an interesting candidate for discovering its aggregation in solution. The most feasible species are the THF solvated monomers (**M1–M4**) and the dimers **D1–D2** (Fig. 8). The molar density for all species is between MD = 5.07 and 5.43 × 10<sup>29</sup> g mol<sup>−1</sup> m<sup>−3</sup> which ensures that those aggregates are suitable for our calibration curves (see Fig. 7).

At room temperature (RT) the ECC<sub>DSE</sub><sup>THF</sup>-MW-determination estimates a MW of MW<sub>det</sub> = 331 g mol<sup>−1</sup>. The comparison of that MW with the most likely Na-indenide species (1–7) is shown in Table 7. Both dimers with two- (**D1**: ΔMW = 21%) and

**Table 7** ECC-MW-determination of Na-indenide (15 mM) in THF-*d*<sub>8</sub> at various temperatures. TMB (15 mM) was used as internal reference and ECC<sub>DSE</sub><sup>THF</sup> to determine the MWs

Species	<i>n</i>	MW [g mol <sup>−1</sup> ]	ΔMW [%]		
			−50 °C	+25 °C	+60 °C
<b>M1</b>	1	210	−84	−57	−36
<b>M2</b>	2	282	−37	−17	−1
<b>M3</b>	3	354	−9	7	19
<b>M4</b>	4	426	10	22	33
<b>D1</b>	1	420	8 <sup>a</sup>	21	32
<b>D2</b>	2	564	32	41	49
Indene	111	111	4	2	−30
(H)HMDS	161	161	−5	−5	−1


<sup>a</sup> The disolvated dimer **D1** (ΔMW = 8%) would also fit to the estimated MW, but this aggregation makes in this context chemically not much sense.

four THF molecules (**D2**: ΔMW = 41%) can be excluded. The same is true for the mono- (**M1**: ΔMW = −57%) and disolvated monomers (**M2**: ΔMW = −17%). The tri-solvated monomer (**M3**: ΔMW = 7%) gives the best match. Such a three-fold THF coordination fits perfectly many crystal structures of THF solvated sodium cyclopentadienide derivatives.<sup>39</sup> We can also identify without difficulty the signals of remaining indene (ΔMW = 2%) and hexamethyldisilazane ((H)HMDS, ΔMW = −5%). Those very accurate MWs indicate that the exchange of the latter with Na-indenide **M3** is very slow or not present. Otherwise the estimated MWs of indene or (H)HMDS should be much higher.

At −50 °C it is obvious that the equilibrium of Na-indenide changes to a higher MW of MW<sub>det</sub> = 386 g mol<sup>−1</sup>. That MW is right in between three- (**M3**: ΔMW = −9%) and four-fold (**M4**: ΔMW = 10%) THF-coordinated Na-indenide, indicating that a fourth THF coordination is attractive at low temperatures. Again, indene (ΔMW = 4%) and (H)HMDS (ΔMW = −5%) are not involved in that Na-indenide–THF equilibrium. By warming up the THF solution to +60 °C the opposite trend is evident. The ECC-MW-determination estimates for Na-indenide a much lower MW of MW<sub>det</sub> = 286 g mol<sup>−1</sup> that would fit to a disolvated Na-indenide monomer (**M2**: ΔMW = −1%) but additionally the MW of indene rises significantly to (MW<sub>det</sub> = 158 g mol<sup>−1</sup>, ΔMW = −37%). That indicates that there is a rapid exchange of Na-indenide and indene at high temperatures producing a merged MW for both. Anyway, HMDS is still not involved in that equilibrium (ΔMW = −1%) perhaps due to its higher basicity and steric demand, compared with indene (p*K*<sub>s</sub> = 26 vs. 20).<sup>40</sup>

## Conclusions

In this article we described how to determine very accurately and reliably molecular weights from DOSY measurements. We derived the equation (log *D*<sub>x,norm</sub> = log *D*<sub>ref,fix</sub> − log *D*<sub>ref</sub> + log *D*<sub>x</sub>) that facilitates defining fixed and normalized diffusion values. This approach enables the use of only one internal reference to determine accurate molecular weights within a deviation of only

**Fig. 8** Most plausible Na-indenide species.



$\Delta MW < \pm 9\%$ . We showed that the ECC-MW-determination is valid at a wide temperature range, enabling to monitor reactions and intermediates at low and high temperatures. Further we showed how the shape and the molar density of compounds influence the power law derived MW-determination. We derived an equation that calculates a term of a molar van der Waals-density that helps to approximate, whether molecules are valid for specified calibration curves or not. In the end we used the ECC-approach to characterize the aggregations of Na-indenide at various temperatures. The ECC-method could easily be extended to other compounds, complexes or solvents giving the opportunity to develop a database of external calibration curves and internal references that would enable access to accurate MW-determination for everyone. An Excel spreadsheet that allows calculating MWs from diffusion coefficients is available for download from the authors' web site (see ESI†).

## Experimental

TOL- $d_8$  and THF- $d_8$  (99.8%, Aldrich, with little amounts of water) were used for the calibration curves. In the case of the Na-indenide, dry THF- $d_8$  was used that was kept with 4 Å molecular sieves under argon. All samples were prepared by adding adamantane (ADAM, in TOL- $d_8$ ) respectively 2,2,3,3-tetramethylbutane (TMB, in THF- $d_8$ ) and analyte (each 15 mM, when not mentioned otherwise) in an equimolar ratio. The diffusion coefficients of the model compounds in TOL- $d_8$  were normalized to the ADAM signal with a fixed value of  $\log D_{\text{ref,fix}}(\text{ADAM}) = -8.8454$ . THF solvates were normalized to the TMB signal to the fixed value of  $\log D_{\text{ref,fix}}(\text{TMB}) = -8.7749$ . NMR experiments were recorded on two devices: (1) Bruker Avance 400 spectrometer equipped with an observe broadband probe with z-axis gradient coil with maximum gradient strength of 57 G cm<sup>-1</sup> and (2) Bruker Ascend 400 spectrometer equipped with an inverse broadband probe with z-axis gradient coil with maximum gradient strength of 51 G cm<sup>-1</sup>. All spectra were acquired in 5 mm NMR tubes. Sample spinning was deactivated during the measurements. All DOSY experiments were performed using a double stimulated echo sequence with bipolar gradient pulses and three spoil gradients with convection compensation (dstebpgp3s).<sup>41</sup> The diffusion time was  $\Delta = 0.1$  s. The duration of the magnetic field pulse gradients  $\delta/2$  was adjusted for each temperature in a range of 400–3500  $\mu\text{s}$ . The delay for gradient recovery was 0.2 ms and the eddy current delay 5 ms. For each DOSY-NMR experiment, a series of 16 spectra on 32 K data points were collected. The pulse gradients ( $g$ ) were incremented from 2 to 98% of the maximum gradient strength in a linear ramp with a total experiment time of 45 min. The temperature was set and controlled at 298 K with an air flow of 400 l h<sup>-1</sup> in order to avoid any temperature fluctuations due to sample heating during the magnetic field pulse gradients. After Fourier transformation and baseline correction, the diffusion dimension was processed with the Topspin 3.1 software. Diffusion coefficients, processed with a line broadening of 2 Hz, were calculated by Gaussian fits with the T1/T2 software of Topspin.

## Acknowledgements

We thank the Danish National Research Foundation (DNRF93) funded Center for Materials Crystallography (CMC) for support.

## Notes and references

- 1 P. F. Godenschwager and D. B. Collum, *J. Am. Chem. Soc.*, 2008, **130**, 8726–8732.
- 2 (a) E. Carl and D. Stalke, in *Lithium Compounds in Organic Synthesis – From Fundamentals to Applications*, ed. R. Luisi and V. Capriati, Wiley-VCH, Weinheim, 2014, pp. 1–31; (b) H. J. Reich, *Chem. Rev.*, 2013, **113**, 7130–7178; (c) T. Stey and D. Stalke, in *The Chemistry of Organolithium Compounds*, ed. Z. Rappoport and I. Marek, John Wiley & Sons, New York, 2004, pp. 47–120; (d) I. Pavel, W. Kiefer and D. Stalke, in *The Chemistry of Organolithium Compounds*, ed. Z. Rappoport and I. Marek, John Wiley & Sons, New York, 2004, pp. 227–265; (e) K. Sorger, P. v. R. Schleyer, R. Fleischer and D. Stalke, *J. Am. Chem. Soc.*, 1996, **118**, 6924–6933; (f) C. Lambert and P. v. R. Schleyer, *Angew. Chem.*, 1994, **106**, 1187–1199; *Angew. Chem., Int. Ed. Engl.*, 1994, **33**, 1129.
- 3 (a) H. P. S. Chauhan, J. Carpenter and S. Joshi, *Appl. Organomet. Chem.*, 2014, **28**, 605–613; (b) L. E. N. Allan, G. G. Briand, A. Decken, J. D. Marks, M. P. Shaver and R. G. Wareham, *J. Organomet. Chem.*, 2013, **736**, 55–62; (c) M. G. Davidson, R. Snaith, D. Stalke and D. S. Wright, *J. Org. Chem.*, 1993, **58**, 2810–2816; (d) D. R. Armstrong, D. Barr, W. Clegg, S. M. Hodgson, R. E. Mulvey, D. Reed, R. Snaith and D. S. Wright, *J. Am. Chem. Soc.*, 1989, **111**, 4719–4727; (e) D. Reed, D. Barr, R. E. Mulvey and R. Snaith, *J. Chem. Soc., Dalton Trans.*, 1986, 557–564; (f) W. Bauer and D. Seebach, *Helv. Chim. Acta*, 1984, **67**, 1972–1988; (g) H. L. Lewis and T. L. Brown, *J. Am. Chem. Soc.*, 1970, **92**, 4664–4670.
- 4 (a) M. Sebban, L. Guilhaudis and H. Oulyadi, in *Lithium Compounds in Organic Synthesis*, ed. R. Luisi and V. Capriati, Wiley-VCH, Weinheim, 2014; (b) A. R. Kennedy, S. M. Leenhouts, J. J. Liggat, A. J. Martinez-Martinez, K. Miller, R. E. Mulvey, C. T. O'Hara, P. O'Keefe and A. Steven, *Chem. Commun.*, 2014, **50**, 10588–10591; (c) A. R. Kennedy, R. E. Mulvey, R. I. Urquhart and S. D. Robertson, *Dalton Trans.*, 2014, **43**, 14265–14274; (d) C. Su, J. Guang, W. Li, K. Wu, R. Hopson and P. G. Williard, *J. Am. Chem. Soc.*, 2014, **136**, 11735–11747; (e) C. Su, R. Hopson and P. G. Williard, *J. Am. Chem. Soc.*, 2013, **135**, 12400–12406; (f) C. Su, R. Hopson and P. G. Williard, *Eur. J. Inorg. Chem.*, 2013, **2013**, 4136–4141; (g) C. Su, R. Hopson and P. G. Williard, *J. Org. Chem.*, 2013, **78**, 7288–7292; (h) C. Su, R. Hopson and P. G. Williard, *J. Am. Chem. Soc.*, 2013, **135**, 14367–14379; (i) D. R. Armstrong, A. R. Kennedy, R. E. Mulvey and S. D. Robertson, *Dalton Trans.*, 2013, **42**, 3704–3711; (j) P. Pregosin, *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.*, 2013, **69**, 1433–1436; (k) S. E. Baillie, W. Clegg, P. García-Álvarez, E. Hevia, A. R. Kennedy, J. Klett and



- L. Russo, *Organometallics*, 2012, **31**, 5131–5142; (l) A.-C. Pöppler, M. M. Meinholz, H. Faßhuber, A. Lange, M. John and D. Stalke, *Organometallics*, 2011, **31**, 42–45; (m) M. Granitzka, A.-C. Pöppler, E. K. Schwarze, D. Stern, T. Schulz, M. John, R. Herbst-Irmer, S. K. Pandey and D. Stalke, *J. Am. Chem. Soc.*, 2011, **134**, 1344–1351; (n) W. Li, G. Kagan, R. Hopson and P. G. Williard, *ARKIVOC*, 2011, 5, 180–187; (o) D. R. Armstrong, W. Clegg, P. García-Alvarez, M. D. McCall, L. Nuttall, A. R. Kennedy, L. Russo and E. Hevia, *Chem.-Eur. J.*, 2011, **17**, 4470–4479; (p) P. García-Alvarez, R. E. Mulvey and J. A. Parkinson, *Angew. Chem.*, 2011, **123**, 9842–9845; *Angew. Chem., Int. Ed.*, 2011, **50**, 9668–9671; (q) I. Fernández, E. Martínez-Viviente, F. Breher and P. S. Pregosin, *Chem.-Eur. J.*, 2005, **11**, 1495–1506; (r) I. Keresztes and P. G. Williard, *J. Am. Chem. Soc.*, 2000, **122**, 10228–10229; (s) K. W. Henderson, D. S. Walther and P. G. Williard, *J. Am. Chem. Soc.*, 1995, **117**, 8680–8681; (t) K. F. Morris and C. S. Johnson, *J. Am. Chem. Soc.*, 1992, **114**, 3139–3141.
- 5 P. S. Pregosin, P. G. A. Kumar and I. Fernández, *Chem. Rev.*, 2005, **105**, 2977–2998.
- 6 (a) H. C. Chen and S. H. Chen, *J. Phys. Chem.*, 1984, **88**, 5118–5121; (b) A. Polson, *J. Phys. Colloid Chem.*, 1950, **54**, 649–652.
- 7  $D = kT/(\delta\pi\eta R_h)$ , where  $R_h$  is the hydrodynamic radius,  $D$  the diffusion coefficient,  $k$  the Boltzmann constant,  $T$  the absolute temperature and  $\eta$  the solvent viscosity.
- 8 (a) R. Evans, Z. Deng, A. K. Rogerson, A. S. McLachlan, J. J. Richards, M. Nilsson and G. A. Morris, *Angew. Chem.*, 2013, **125**, 3281–3284; *Angew. Chem., Int. Ed.*, 2013, **52**, 3199–3202; (b) A. Macchioni, G. Ciancaleoni, C. Zuccaccia and D. Zuccaccia, *Chem. Soc. Rev.*, 2008, **37**, 479–489.
- 9 S. Floquet, S. Brun, J.-F. Lemonnier, M. Henry, M.-A. Delsuc, Y. Prigent, E. Cadot and F. Taulelle, *J. Am. Chem. Soc.*, 2009, **131**, 17254–17259.
- 10 Where  $K$  is a molecule dependent constant and  $\alpha$  is a coefficient that depends highly on the shape of the particle.
- 11 S. Augé, P.-O. Schmit, C. A. Crutchfield, M. T. Islam, D. J. Harris, E. Durand, M. Clemancey, A.-A. Quoineaud, J.-M. Lancelin, Y. Prigent, F. Taulelle and M.-A. Delsuc, *J. Phys. Chem. B*, 2009, **113**, 1914–1918.
- 12 S. Viel, D. Capitani, L. Mannina and A. Segre, *Biomacromolecules*, 2003, **4**, 1843–1847.
- 13 A. Chen, D. Wu and C. S. Johnson, *J. Am. Chem. Soc.*, 1995, **117**, 7965–7970.
- 14 D. K. Wilkins, S. B. Grimshaw, V. Receveur, C. M. Dobson, J. A. Jones and L. J. Smith, *Biochemistry*, 1999, **38**, 16424–16431.
- 15 C. A. Crutchfield and D. J. Harris, *J. Magn. Reson.*, 2007, **185**, 179–182.
- 16 D. Li, I. Keresztes, R. Hopson and P. G. Williard, *Acc. Chem. Res.*, 2008, **42**, 270–280.
- 17 D. Li, R. Hopson, W. Li, J. Liu and P. G. Williard, *Org. Lett.*, 2008, **10**, 909–911.
- 18 G. Kagan, W. Li, R. Hopson and P. G. Williard, *Org. Lett.*, 2009, **11**, 4818–4821.
- 19 (a) R. E. Mulvey and S. D. Robertson, *Angew. Chem.*, 2013, **125**, 11682–11700; *Angew. Chem., Int. Ed.*, 2013, **52**, 11470–11487; (b) D. R. Armstrong, P. García-Alvarez, A. R. Kennedy, R. E. Mulvey and J. A. Parkinson, *Angew. Chem.*, 2010, **122**, 3253–3256; *Angew. Chem., Int. Ed.*, 2010, **49**, 3185–3188.
- 20 (a) E. J. Cabrita and S. Berger, *Magn. Reson. Chem.*, 2001, **39**, 142–148; (b) S. Yao, G. Howlett and R. Norton, *J. Biomol. NMR*, 2000, **16**, 109–119.
- 21 (a) E. Durand, M. Clemancey, J.-M. Lancelin, J. Verstraete, D. Espinat and A.-A. Quoineaud, *J. Phys. Chem. C*, 2009, **113**, 16266–16276; (b) J. Jones, D. Wilkins, L. Smith and C. Dobson, *J. Biomol. NMR*, 1997, **10**, 199–203.
- 22  $\log D_{x,\text{norm}} = \log K + \alpha \log \text{MW}$ .
- 23 D. Li, G. Kagan, R. Hopson and P. G. Williard, *J. Am. Chem. Soc.*, 2009, **131**, 5627–5634.
- 24 The full classification of all 28 compounds is available in the ESI†.
- 25 A fourth type of molecule shape is also thinkable that should distinguish from the already mentioned ones: rod shaped molecules. But in this article we focus on CS, DSE and ED molecules.
- 26 G. R. Fulmer, A. J. M. Miller, N. H. Sherden, H. E. Gottlieb, A. Nudelman, B. M. Stoltz, J. E. Bercaw and K. I. Goldberg, *Organometallics*, 2010, **29**, 2176–2179.
- 27 M. A. Jacobson, I. Keresztes and P. G. Williard, *J. Am. Chem. Soc.*, 2005, **127**, 4965–4975.
- 28 At lower temperatures inter- or intramolecular interactions increase. Therefore the error of MW-determination will also increase for high concentrated samples. We recommend to use low concentrated samples (<20 mM) for low temperature measurements.
- 29 P. Paetzold, *Chemie: Eine Einführung*, Walter de Gruyter, Berlin, 2009.
- 30 A. Bondi, *J. Phys. Chem.*, 1964, **68**, 441–451.
- 31 D. Stern, N. Finkelmeier, K. Meindl, J. Henn and D. Stalke, *Angew. Chem.*, 2010, **122**, 7021–7024; *Angew. Chem., Int. Ed.*, 2010, **49**, 6869–6872.
- 32 (a) R. E. Mulvey, *Chem. Soc. Rev.*, 1998, **27**, 339–346; (b) K. Gregory, P. v. R. Schleyer and R. Snaith, *Adv. Inorg. Chem.*, 1991, **37**, 47–142; (c) R. E. Mulvey, *Chem. Soc. Rev.*, 1991, **20**, 167–209; (d) D. R. Armstrong, D. Barr, R. Snaith, W. Clegg, R. E. Mulvey, K. Wade and D. Reed, *J. Chem. Soc., Dalton Trans.*, 1987, 1071–1081.
- 33 R. Neufeld and D. Stalke, *Angew. Chem., Int. Ed. Engl.*, 2015, submitted.
- 34 A. S. Galiano-Roth and D. B. Collum, *J. Am. Chem. Soc.*, 1989, **111**, 6772–6778.
- 35 R. E. Dinnebier, S. Neander, U. Behrens and F. Olbrich, *Organometallics*, 1999, **18**, 2915–2918.
- 36 (a) R. Michel, R. Herbst-Irmer and D. Stalke, *Organometallics*, 2011, **30**, 4379–4386; (b) T. Kähler, U. Behrens, S. Neander and F. Olbrich, *J. Organomet. Chem.*, 2002, **649**, 50–54; (c) R. E. Dinnebier, U. Behrens and F. Olbrich, *Organometallics*, 1997, **16**, 3855–3858.
- 37 R. Michel, T. Nack, R. Neufeld, J. M. Dieterich, R. A. Mata and D. Stalke, *Angew. Chem.*, 2013, **125**, 762–766; *Angew. Chem., Int. Ed.*, 2013, **52**, 734–738.



- 38 L. L. Tomasevich and D. B. Collum, *J. Am. Chem. Soc.*, 2014, **136**, 9710–9718.
- 39 (a) T. Nishinaga, D. Yamazaki, H. Stahr, A. Wakamiya and K. Komatsu, *J. Am. Chem. Soc.*, 2003, **125**, 7324–7335; (b) Z. Hou, Y. Zhang, H. Tezuka, P. Xie, O. Tardif, T.-A. Koizumi, H. Yamazaki and Y. Wakatsuki, *J. Am. Chem. Soc.*, 2000, **122**, 10533–10543.
- 40 R. R. Fraser, T. S. Mansour and S. Savard, *J. Org. Chem.*, 1985, **50**, 3232–3234.
- 41 (a) A. Jerschow and N. Müller, *J. Magn. Reson.*, 1997, **125**, 372–375; (b) A. Jerschow and N. Müller, *J. Magn. Reson.*, 1996, **123**, 222–225.

