



CrossMark
click for updates

Cite this: *Chem. Sci.*, 2016, 7, 85

Received 14th August 2015
Accepted 26th October 2015

DOI: 10.1039/c5sc03025k

www.rsc.org/chemicalscience

Visible light-mediated gold-catalysed carbon(sp²)-carbon(sp) cross-coupling†

Suhong Kim,^a Jaime Rojas-Martin^{ab} and F. Dean Toste^{*a}

A dual photoredox and gold-catalysed cross-coupling reaction of alkynyltrimethylsilanes and aryldiazonium tetrafluoroborates is described. The reaction proceeds through visible-light mediated oxidative addition of aryldiazoniums, transmetalation of alkynyltrimethylsilanes and aryl-alkynyl reductive elimination. Exclusive selectivity for silyl-substituted alkynes is observed, with no reactivity observed for terminal alkynes.

Introduction


For over a decade, homogeneous gold catalysis has been investigated because it provides access to novel modes of reactivity and enables rapid generation of complex molecular architectures.¹ However, in contrast to other low-valent late transition metal catalysts, the majority of gold(I) complexes are unreactive towards the oxidative addition of aryl and vinyl halides and pseudohalides.² Consequently, most gold-catalysed cross-coupling reactions have required sacrificial oxidants to access the +3 oxidation state.³ For example, this reactivity platform has been exploited in a gold-catalysed oxidative coupling of organosilanes and 1,2-alkene functionalizations (Scheme 1A). As an alternative, visible light-mediated oxidative addition of

aryldiazoniums has emerged as a method for the generation of the requisite gold(III) intermediates *via* photoredox catalysis, thereby obviating the need for sacrificial oxidants.⁴ These gold(III) intermediates have been intercepted by a variety of nucleophiles in dual catalytic photoredox-gold reactions (Scheme 1B).

Although aryldiazoniums and photoredox catalysts provide an efficient method to generate gold(III) intermediates, several challenges must be addressed in order to facilitate carbon-carbon cross-coupling *via* these species. First, aryldiazonium salts are highly electrophilic and reactive towards many classes of nucleophilic coupling partners. Aryldiazoniums are generally decomposed under Kumada and Negishi coupling conditions and do not tolerate ligand additives and stoichiometric bases employed in the majority of Suzuki, Sonogashira and Hiyama coupling reactions.⁵ Second, as visible light-mediated oxidative addition of aryldiazoniums is proposed to proceed through radical intermediates, nucleophilic coupling partners that can participate in single electron transfer, such as organotin and organotrifluoroborates compounds, are problematic.⁶ Inspired by the oxidative coupling of organosilanes, we envisioned that the gold(III) complexes generated *via* photoredox catalysis might undergo transmetalation with organosilanes, generating an intermediate poised for carbon-carbon bond formation through reductive elimination.^{7,8}

Results and discussions

While aryldiazonium tetrafluoroborates have been shown to react with halo-, azido-, allyl- and thiotrimethylsilanes, their reactions with alkynyltrimethylsilanes has not been previously reported.⁹ Additionally, previous reports of organosilane transmetalation with gold lead us to hypothesize that the transmetalation of alkynyltrimethylsilanes to gold(III) intermediates might result in productive cross-coupling. On the basis of this hypothesis, we examined the combined gold/photoredox coupling of 1 equivalent of aryldiazonium salt **1a** and 1 equivalent of alkynyltrimethylsilane **2**.



Scheme 1 Photoredox catalyst and visible light-mediated Au^I/Au^{III} catalysis.

^aDepartment of Chemistry, University of California, Berkeley, CA 94720, USA. E-mail: fdtoste@berkeley.edu

^bDepartamento de Química Orgánica (Módulo 01), Universidad Autónoma de Madrid, C/Francisco Tomás y Valiente 7, Cantoblanco, 28049 Madrid, Spain


† Electronic supplementary information (ESI) available. See DOI: 10.1039/c5sc03025k



Evaluation of reaction conditions showed that the combination of Ar_3PAuCl and $\text{Ru}(\text{bpy})_3(\text{PF}_6)_2$, in acetonitrile gave the highest yield of aryl alkyne **3a** (Table 1). Changing either the solvent or the photoredox catalyst proved detrimental to the yield of the desired product. With respect to the ligands on the gold catalyst, both electronic and steric factors impacted the efficiency of the cross-coupling. For example, electron-deficient triarylphosphine ligands resulted in significantly lower yield of **3a** (Table 1, entry 10). The sterics of the ligand showed an even more dramatic effect on reaction yield. The reaction conducted with (*p*-tol) $_3\text{PAuCl}$ provided 72% yield of the desired product, while the much more hindered (*o*-tol) $_3\text{PAuCl}$ -catalysed reaction only provided 17% yield of **3a** (Table 1, entry 11 and 12). The yield of product was lower when tricyclohexylphosphinegold(i) chloride was used as catalyst instead of triphenylphosphinegold(i) chloride (Table 1, entry 13). Gold(i) complexes of dialkylbiarylphosphine were also examined, but all showed less than 5% yield due to the combined effect of dialkyl groups and the steric effect of biaryl groups. Finally, no coupling product was observed in the absence of the ruthenium catalyst (entry 14) or when the reaction was conducted in the dark (entry 15).

With an optimized catalyst system in hand, the scope with respect to the aryldiazonium coupling partner was examined (Table 2). The yields of alkynylation reactions of electron-poor aryldiazonium salts were generally higher than those of reactions with electron-rich aryldiazonium coupling partners. It should be noted that the intrinsic instability of *ortho*-substituted aryldiazonium salts limited their use in this transformation (Table 2, entry **3k** and **3l**). It is also noteworthy that all halogen substitutions on the aryldiazonium coupling partner

Table 2 Scope of aryldiazonium tetrafluoroborates



Reaction conditions: **1a-1** (0.24 mmol, 1.2 equiv.), **2** (0.2 mmol, 1.0 equiv.), Ph_3PAuCl (10 mol%), $\text{Ru}(\text{bpy})_3(\text{PF}_6)_2$ (2 mol%), acetonitrile (2 mL), N_2 atmosphere, visible light, rt for 3 h. Isolated yields.

Table 1 Optimization of reaction conditions

Reaction scheme showing the synthesis of **3a** from **1a** and **2** using a gold catalyst and photoredox catalyst in a solvent under visible light at room temperature.

Entry	Gold cat.	Photoredox cat.	Solvent	Yield ^a
1	Ph_3PAuCl	$\text{Ru}(\text{bpy})_3(\text{PF}_6)_2$	CH_3CN	70
2	Ph_3PAuCl	$\text{Ru}(\text{bpy})_3(\text{PF}_6)_2$	Acetone	39
3	Ph_3PAuCl	$\text{Ru}(\text{bpy})_3(\text{PF}_6)_2$	CH_3NO_2	15
4	Ph_3PAuCl	$\text{Ru}(\text{bpy})_3(\text{PF}_6)_2$	DMF	10
5	Ph_3PAuCl	$\text{Ru}(\text{bpy})_3(\text{PF}_6)_2$	MeOH	18
6	Ph_3PAuCl	$\text{Ru}(\text{bpy})_3(\text{PF}_6)_2$	EtOH	4
7	Ph_3PAuCl	$\text{Ir}(\text{ppy})_3$	CH_3CN	21
8	Ph_3PAuCl	$\text{Ir}(\text{ppy})_2(\text{dtbbpy})\text{PF}_6$	CH_3CN	45
9	(<i>p</i> -MeOPh) $_3\text{PAuCl}$	$\text{Ru}(\text{bpy})_3(\text{PF}_6)_2$	CH_3CN	73
10	(<i>p</i> -CF $_3$ Ph) $_3\text{PAuCl}$	$\text{Ru}(\text{bpy})_3(\text{PF}_6)_2$	CH_3CN	43
11	(<i>p</i> -tol) $_3\text{PAuCl}$	$\text{Ru}(\text{bpy})_3(\text{PF}_6)_2$	CH_3CN	72
12	(<i>o</i> -tol) $_3\text{PAuCl}$	$\text{Ru}(\text{bpy})_3(\text{PF}_6)_2$	CH_3CN	17
13	Cy_3PAuCl	$\text{Ru}(\text{bpy})_3(\text{PF}_6)_2$	CH_3CN	48
14	Ph_3PAuCl	—	CH_3CN	0
15 ^b	Ph_3PAuCl	$\text{Ru}(\text{bpy})_3(\text{PF}_6)_2$	CH_3CN	0

Reaction conditions: 4-fluorobenzenediazonium tetrafluoroborate **1a** (0.2 mmol, 1 equiv.), 1-trimethylsilyl-2-phenylacetylene **2** (0.2 mmol, 1 equiv.), gold catalyst (5 mol%), photoredox catalyst (1 mol%), solvent (2 mL), N_2 atmosphere, visible light, rt for 6 h.^a GC yields. ^b The reaction was run in the dark. bpy = 2,2'-bipyridine, ppy = 2-phenylpyridine, dtbbpy = 4,4'-di-*tert*-butyl-2,2'-bipyridine.

were preserved during the coupling reaction. Bromo- and iodoaryldiazonium salts were readily coupled and leaving halides intact for use in further reactions (Table 2, **3b** and **3j**); this chemoselectivity is challenging under typical palladium-catalysed Sonogashira coupling reactions.¹⁰


The difference in reactivity of electron-rich and -poor aryldiazoniums may be the result of competing reaction pathways. Oxidative addition is believed to proceed through single electron reduction of aryldiazoniums and this process is preferred with electron-poor aryldiazonium salts.⁴ On the other hand, the generation of aryl cations, through loss of dinitrogen, is facile with electron-rich aryldiazoniums. Correspondingly, we observed trace amount of desired product when benzenediazonium tetrafluoroborate was heated at 60 °C for 9 hours in the presence of 1-trimethylsilyl-2-(4-methoxy)phenylacetylene; however the same reactivity was not observed with 4-fluorophenyldiazonium salts on the same conditions.

The scope of alkynyltrimethylsilanes was also investigated (Table 3). Both aryl- and alkylethynyltrimethylsilanes were coupled with modest to high product yields. *o*-Isopropylphenyl, biphenyl and naphthylalkynyltrimethylsilanes participated in the gold-catalysed coupling (Table 3, **3m**, **3n** and **3o**). Potentially sensitive benzylic and propargylic C-H and C-X bonds were well tolerated under the reaction conditions (Table 3, **3q**, **3aa** and **3s-v**). Additionally, alkynyltrimethylsilane **3w** was prepared in 78% yield from the coupling of aryldiazonium salt **1d** and bis(trimethylsilyl) acetylene, providing a compliment to using trimethylsilylacetylene in a traditional Sonogashira coupling (Table 3, **3w**).

Another interesting feature of this reaction is the effect of the silyl group on yield (Table 4). When trimethylsilylacetylene was



Table 3 Scope of alkynyltrimethylsilanes



Reaction conditions: **1d** (0.24 mmol, 1.2 equiv.), **2m-aa** (0.2 mmol, 1.0 equiv.), Ph₃PAuCl (10 mol%), Ru(bpy)₃(PF₆)₂ (2 mol%), acetonitrile (2 mL), N₂ atmosphere, visible light, rt for 3 h. Isolated yields. Ar = *p*-nitrophenyl. ^a 5 equiv. of bis(trimethylsilyl)acetylene was used with 1 equiv. of **1d**. ^b GC yields.


coupled with **1d** to generate terminal alkyne **3x**, the yield was 15% with none of the corresponding silylacetylene **3w** (Table 3). This observation suggests that, under the current base free

Table 4 Effect of counteranion and silyl group

Reaction conditions: aryl diazonium salt (0.2 mmol, 1 equiv.), phenylethynylsilane (0.2 mmol, 1 equiv.), Ph₃PAuCl (5 mol%), Ru(bpy)₃(PF₆)₂ (1 mol%), acetonitrile (2 mL), N₂ atmosphere, visible light, rt for 6 h. ^a GC yields.

Entry	Counteranion (X)	Silyl group (Y)	Yield ^a
1	BF ₄ ⁻	TMS	70
2	TsO ⁻	TMS	68
3	PF ₆ ⁻	TMS	24
4	BF ₄ ⁻	TES	61
5	BF ₄ ⁻	TBDMS	10
6	BF ₄ ⁻	TIPS	10
7	BF ₄ ⁻	TBDPS	1
8	BF ₄ ⁻	H	3

Reaction conditions: aryl diazonium salt (0.2 mmol, 1 equiv.), phenylethynylsilane (0.2 mmol, 1 equiv.), Ph₃PAuCl (5 mol%), Ru(bpy)₃(PF₆)₂ (1 mol%), acetonitrile (2 mL), N₂ atmosphere, visible light, rt for 6 h. ^a GC yields.



Scheme 2 Reaction with allenyltrimethylsilanes.

reaction conditions, the silyl group plays a critical role in the coupling. To investigate this role, a set of reactions was performed by varying counteranions of aryl diazonium salts and silyl group identities of alkynylsilanes. The yield of coupling product was diminished as the steric hindrance of silyl group was increased (Table 4, entry 1 and entries 4–7). While both tetrafluoroborate and tosylate diazonium salts proved efficient coupling partners, the yield was diminished when hexafluorophosphate salts were used.¹¹ Moreover, when the corresponding terminal alkyne was used instead of alkynyltrimethylsilanes, only 3% of the coupling product was observed (Table 4, entry 8).¹² Taken together, these results suggest that transmetalation from the organosilane is critical for high efficiency of the current coupling reaction.

Other organotrimethylsilanes were tested under the optimized reaction conditions. While the reactions of aryl and vinylsilanes were complicated by competing reactions,^{13,14} the gold/photoredox-catalysed coupling of allenyltrimethylsilane **4** provided propargylic compound **5** as the sole coupling product (Scheme 2). The product from aryl–allenyl reductive elimination and the other aryl–propargylic coupling isomer were prepared independently (Scheme 2, **6** and **7**). We examined whether **6** was an intermediate that underwent isomerization to **5** facilitated by visible light and the photoredox catalyst. However, no isomerization was observed of **6** to either **5** or **7** under the reaction conditions. Therefore, the formation of **5** is best rationalized by transmetalation of **4** to the gold(III) intermediate, followed by 1,3-migration and aryl–propargyl reductive elimination.

Conclusions

In summary, we have demonstrated that trimethylsilylalkynes and aryl diazonium tetrafluoroborates can be coupled *via* dual photoredox and gold catalysis strategy. The reaction proceeds under mild conditions and shows excellent functional group tolerance, including aryl halides that may be reactive under traditional coupling conditions. This process complements the Sonogashira coupling, especially when aryl–alkynyl coupling is required under non-basic conditions and may present an advantage when TMS-alkynes are available rather than corresponding terminal alkynes.

Acknowledgements

We are grateful for financial support from the NIHGMS (RO1 GM073932). Jaime Rojas-Martin thanks MINECO for the EEBB-



FPI fellowship. We thank Mark D. Levin, William J. Wolf and Matthew S. Winston for helpful discussions.

Notes and references

- For recent reviews on gold catalysis: (a) A. S. K. Hashimi, *Acc. Chem. Res.*, 2014, **47**, 864; (b) C. Obradors and A. M. Echavarren, *Chem. Commun.*, 2014, **50**, 16; (c) D. Garayalde and C. Nevado, *ACS Catal.*, 2012, **2**, 1462; (d) M. Rudolph and A. S. K. Hashmi, *Chem. Soc. Rev.*, 2012, **41**, 2448; (e) L. P. Liu and G. B. Hammond, *Chem. Soc. Rev.*, 2012, **41**, 3129; (f) N. D. Shapiro and F. D. Toste, *Synlett*, 2010, 675.
- (a) T. Lauterbach, M. Livendahl, A. Rosellón, P. Espinet and A. M. Echavarren, *Org. Lett.*, 2010, **12**, 3006. For recent breakthroughs, (b) M. D. Levin and F. D. Toste, *Angew. Chem., Int. Ed.*, 2014, **53**, 6211; (c) M. S. Winston, W. J. Wolf and F. D. Toste, *J. Am. Chem. Soc.*, 2014, **136**, 7777; (d) J. Guenther, S. Mallet-Ladeira, L. Estevez, K. Miqueu, A. Amgoune and D. Bourissou, *J. Am. Chem. Soc.*, 2014, **136**, 1778; (e) M. Joost, A. Zeineddine, L. Estevez, S. Mallet-Ladeira, K. Miqueu, A. Amgoune and D. Bourissou, *J. Am. Chem. Soc.*, 2014, **136**, 14654.
- For early publications on gold-catalysed oxidative coupling, (a) A. Kar, N. Mangu, H. M. Kaiser, M. Beller and M. K. Tse, *Chem. Commun.*, 2008, 386; (b) G. Zhang, Y. Peng, L. Cui and L. Zhang, *Angew. Chem., Int. Ed.*, 2009, **48**, 3112; (c) A. D. Melhado, W. E. Brenzovich, A. D. Lackner and F. D. Toste, *J. Am. Chem. Soc.*, 2010, **132**, 8885; (d) W. E. Brenzovich, D. Benitez, A. D. Lackner, H. P. Shunatona, E. Tkatchouk, W. A. Goddard and F. D. Toste, *Angew. Chem., Int. Ed.*, 2010, **49**, 5519; (e) W. E. Brenzovich, J. F. Brazeau and F. D. Toste, *Org. Lett.*, 2010, **12**, 4728; (f) G. Zhang, L. Cui, Y. Wang and L. Zhang, *J. Am. Chem. Soc.*, 2010, **132**, 1474; (g) T. de Haro and C. Nevado, *J. Am. Chem. Soc.*, 2010, **132**, 1512; (h) J. P. Brand, J. Charpentier and J. Waser, *Angew. Chem., Int. Ed.*, 2009, **48**, 9346; (i) J. P. Brand and J. Waser, *Angew. Chem., Int. Ed.*, 2010, **49**, 7304; (j) M. N. Hopkinson, A. Tessier, A. Salisbury, G. T. Giuffredi, L. E. Combettes, A. D. Gee and V. Gouverneur, *Chem.–Eur. J.*, 2010, **16**, 4739; (k) L. T. Ball, G. C. Lloyd-Jones and C. A. Russell, *Org. Lett.*, 2010, **12**, 4724; (l) L. T. Ball, G. C. Lloyd-Jones and C. A. Russell, *Science*, 2012, **337**, 1644.
- For visible-light mediated oxidative addition of aryldiazoniums on gold catalysis, (a) B. Sahoo, M. N. Hopkinson and F. Glorius, *J. Am. Chem. Soc.*, 2013, **135**, 5505; (b) M. N. Hopkinson, B. Sahoo and F. Glorius, *Adv. Synth. Catal.*, 2014, **356**, 2794; (c) X. Z. Shu, M. Zhang, Y. He, H. Frei and F. D. Toste, *J. Am. Chem. Soc.*, 2014, **136**, 5844; (d) Y. He, H. Wu and F. D. Toste, *Chem. Sci.*, 2015, **6**, 1194.
- A. Roglans, A. Pla-Quintana and M. Moreno-Manas, *Chem. Rev.*, 2006, **106**, 4622.
- For single-electron transmetalation of organotrifluoroborates (a) J. C. Tellis, D. N. Primer and G. A. Molander, *Science*, 2014, **345**, 433; (b) D. N. Primer, I. Karakaya, J. C. Tellis and G. A. Molander, *J. Am. Chem. Soc.*, 2015, **137**, 2195.
- W. J. Wolf, M. S. Winston and F. D. Toste, *Nat. Chem.*, 2013, **6**, 159.
- For cross-coupling reactions of alkynyltrimethylsilanes, (a) S. A. Lermontov, I. M. Rakov, N. S. Zefirov and P. J. Stang, *Tetrahedron Lett.*, 1996, **37**, 4051; (b) G. W. Kabalka, L. Wang and R. M. Pagni, *Tetrahedron*, 2001, **57**, 8017; (c) S. Kang, H. Ryu and Y. Hong, *J. Chem. Soc., Perkin Trans. 1*, 2007, 736; (d) C. Yang and S. P. Nolan, *Organometallics*, 2002, **21**, 1020; (e) L. K. Hwang, Y. Na, J. Lee, Y. Do and S. Chang, *Angew. Chem., Int. Ed.*, 2005, **44**, 6166; (f) W. J. Sommer and M. Weck, *Adv. Synth. Catal.*, 2006, **348**, 2101; (g) N. Sakai, R. Komatsu, N. Uchida, R. Ikeda and T. Konakahara, *Org. Lett.*, 2010, **12**, 1300; (h) Y. Nishihara, S. Noyori, T. Okamoto, M. Suetsugu and M. Iwasaki, *Chem. Lett.*, 2011, **40**, 972; (i) Y. Nishihara, D. Ogawa, S. Noyori and M. Iwasaki, *Chem. Lett.*, 2012, **41**, 1503; (j) L. Peng, F. Xu, Y. Suzuma, A. Orita and J. Otera, *J. Org. Chem.*, 2013, **78**, 12802.
- For reactions of aryldiazonium tetrafluoroborate with halo-, azido-, allyl- and thiotrimethylsilanes, (a) T. Keumi, T. Umeda, Y. Inoue and H. Kitajima, *Bull. Chem. Soc. Jpn.*, 1989, **62**, 89; (b) H. Mayr and K. Grimm, *J. Org. Chem.*, 1992, **57**, 1057; (c) G. K. S. Prakash, D. Hoole, D. S. Ha, J. Wilkinson and G. A. Olah, *ARKIVOC*, 2002, **13**, 50.
- For Sonogashira cross-coupling of aryldiazonium salts, (a) S. Darses, G. Michaud and J. Genet, *Eur. J. Org. Chem.*, 1999, 1875; (b) G. Fabrizi, A. Goggiamani, A. Sferrazza and S. Cacchi, *Angew. Chem., Int. Ed.*, 2010, **49**, 4067; (c) B. Panda and T. K. Sarkar, *Chem. Commun.*, 2010, **46**, 3131; (d) X. Wu, H. Neumann and M. Beller, *Chem. Commun.*, 2011, **47**, 7959; (e) This challenge was overcome by employing *o*-benzenedisulfonimide as a counteranion.; M. Barbero, S. Cadamuro and S. Dughera, *Eur. J. Org. Chem.*, 2014, 598. (f) While preparing a manuscript, gold-catalysed cross-coupling of aryldiazonium salts with terminal alkynes and boronic acids was published.; R. Cai, M. Lu, E. Y. Aguilera, Y. Xi, N. G. Akhmedov, J. L. Petersen, H. Chen and X. Shi, *Angew. Chem., Int. Ed.*, 2015, **54**, 8772.
- After the completion of coupling reaction with tetrafluoroborate salts, stoichiometric amounts of trimethylsilyl fluoride were observed by ¹H NMR.
- To examine the robustness of this reaction towards terminal alkynes, 1 equiv. of phenylacetylene and 1 equiv. of phenylethynyltrimethylsilane were reacted with 1 equiv. **1a** surprisingly, less than 5% of **3a** was detected.
- When (*Z*)-styryltrimethylsilane was used the corresponding *Z*-product was obtained from gold-catalysed cross coupling and *E*-product was obtained from uncatalysed addition of the aryldiazonium to the vinylsilane (see: A. Benkeser, E. W. Bennet and R. A. Hickner, *J. Am. Chem. Soc.*, 1957, **79**, 6253.) After consumption of the starting material, photoredox promoted isomerization of *E*-stilbene to *Z*-stilbene occurred.
- In the case aryltrimethylsilanes, further optimization is required due to the formation of significant amounts of competing homo-coupling reactions.

