



CrossMark
click for updates

Cite this: *Chem. Sci.*, 2016, 7, 3689

Supramolecular reactions of metallo-architectures: Ag_2 -double-helicate/ Zn_4 -grid, Pb_4 -grid/ Zn_4 -grid interconversions, and Ag_2 -double-helicate fusion†

Adrian-Mihail Stadler,^{*ab} Juan Ramírez,^c Jean-Marie Lehn^a and Bruno Vincent^d

Supramolecular reactions are of importance in many fields. We report herein three examples where complexes of hydrazone-based ligands are involved. A Ag_2 -double-helicate was converted, by treatment with $\text{Zn}(\text{OTf})_2$, into a Zn_4 -grid (exchange of metal ions and change of the nature of the initial complex). A Pb_4 -grid was converted, upon reaction with ZnCl_2 or ZnBr_2 , into a Zn_4 -grid (exchange of metal ions, but conservation of the nature of the initial complex). The reverse conversions were also achieved. The fusion of a Ag_2 -double-helicate with another Ag_2 -double-helicate was performed (exchange of ligands, but conservation of the nature of the complexes) and resulted in a mixture of three helicates (two homostranded ones and one heterostranded one).

Received 17th November 2015
Accepted 3rd February 2016

DOI: 10.1039/c5sc04403k

www.rsc.org/chemicalscience

Introduction

Like covalent molecules, supramolecular¹ assemblies may participate in various reactions. The understanding of supramolecular reactions is of much interest because they are involved in many areas such as complex chemical systems and networks,² adaptive³ and stimuli-responsive⁴ chemical systems, fabrication of nanodevices and materials,⁴ biomolecular processes. Thus, in the complexity and diversity of supramolecular chemistry, the reactivity of supramolecules plays a crucial role. It includes the processes:

(a) of (self)assembly (*i.e.* formation of supramolecular architectures through assembly, but also their participation, as subunits, in more complex assemblies), and correlatively, partial or total disassembly;

(b) of partial or total reorganization or exchange (at the supramolecular and, additionally and possibly, at the covalent level), that involves the breaking of several or all of the initial supramolecular connections and formation of new ones;

(c) without breaking or formation of new supramolecular connections (*e.g.* covalent modifications after self-assembly⁵).

Amongst supramolecular architectures, double helices and helicates,⁶ as well as grids⁷ arouse much interest and work. For example, DNA⁸ and the ion channel generated by gramicidine⁹

have a double helical structure, and there are double helical complexes that act as molecular machines¹⁰ or catalysts.¹¹ Grid-like complexes have been studied for their electrochemical and magnetic properties,⁷ for their capacity to encapsulate ions¹² or as starting materials for building more complex architectures (*e.g.* a Solomon link¹³), amongst other things. However, supramolecular interconversions of grids and helicates have not, except several examples,^{14,15} been much explored.

With these ideas in mind – and using principles such as the displacement of an equilibrium through precipitation, and the preference of Ag^+ for tetrahedral and of Zn^{2+} for octahedral coordination – we designed, as reported herein, three supramolecular reactions¹⁶ of reorganization and exchange (Fig. 1) involving grids and double helicates. They are related through the ligands¹⁷ (which are pyrimidine-bis-hydrazones;¹⁸ Fig. 2)




Fig. 1 Stylized representation of the three types of supramolecular reactions reported herein.

^aUniversité de Strasbourg, CNRS, UMR 7006, ISIS, 8 Allée G. Monge, Strasbourg, France. E-mail: mstadler@unistra.fr

^bInstitute of Nanotechnology (INT), Karlsruhe Institute of Technology (KIT), 76344, Eggenstein-Leopoldshafen, Germany

^cInstitut Pasteur Paris, 28 Rue du Docteur Roux, 75015 Paris, France

^dService de RMN, Faculté de Chimie, 1 Rue B. Pascal, Strasbourg, France

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c5sc04403k






Fig. 2 Structural formulae and stylized representations of ligands 1 and 2, and of grids and double helicites.

that produce the supramolecular complexes, as well as through the nature of complexes, and occur due to the dynamic character of the present metal–ligand connections. These reactions (Fig. 1) can be seen as:




Fig. 3 Stylized representation of: (a) the conversion of Ag_21_2-DH double helicite into the Zn_41_4-G grid; (b) the interconversion Ag_21_2-DH/Zn_41_4-G ; (c) the conversion of Ag_22_2-DH double helicite into the Zn_42_4-G grid; (d) the conversion of Zn_42_4-G grid into Ag_22_2-DH double helicite. Charges and stoichiometric coefficients are omitted for simplicity.




Fig. 4 Stylized representation of: (a) reaction of Pb_41_4-G grid with $ZnCl_2$, $ZnBr_2$, and $Zn(OTf)_2$ and $Bu_4P^+Br^-$ (solvent CD_3CN); (b) Zn_41_4-G/Pb_41_4-G and Pb_41_4-G/Zn_41_4-G conversions. Charges and stoichiometric coefficients are omitted for simplicity.




Fig. 5 Stylized representation of the fusion (conproportionation) reaction of double helicites Ag_21_2-DH and Ag_22_2-DH with formation of heterostrand species $Ag_2(1)(2)-DH$. Charges are omitted for simplicity.

(i) a change of the nature of the supramolecular architecture, from a Ag^+ dinuclear double helicite (DH) into a Zn^{2+} tetranuclear grid (G), induced by replacement of Ag^+ by Zn^{2+} (Fig. 3). In this reaction, not only the nature of the complex and that of the metal ion change, but also the conformation of the ligand (helical → unfolded), the charge ($2^+ \rightarrow 8^+$) and the nuclearity of the complex ($2 \rightarrow 4$) and the number of ligands per complex ($2 \rightarrow 4$). In regard to this last change, this process can be compared with the conversion or the equilibrium between supramolecular dimer and tetramer of bioactive proteins,¹⁹ or between other homo-oligomers²⁰ with influence on the protein functions.

(ii) a substitution²¹ (metal ion exchange or transmetallation), in a sole operation, of the four Pb^{2+} ions of a grid-like²² complex by Zn^{2+} ions (Fig. 4);

(iii) a fusion (conproportionation)²³ between two Ag^+ double helicites²⁴ (Fig. 5).

While in case (ii) the equilibrium is shifted towards the Zn^{2+} grid through the precipitation of Pb^{2+} as its halides (chloride and bromide), in cases (i) and (iii), the conversions can be done without precipitation.

Results and discussion

(i) The conversion $Ag_2L_2-DH \rightarrow Zn_4L_4-G$ ($L = 1, 2$) through transmetallation is a dramatic reorganization of the nature of the metallo-supramolecular architecture induced by the replacement of Ag^+ by Zn^{2+} (Fig. 3a and c): $2 Ag_2L_2-DH + 4 Zn^{2+} \rightarrow Zn_4L_4-G + 4 Ag^+$. Ag^+ prefers a tetrahedral coordination geometry which is, in the case of ligands 1 and 2, achieved from 2 two- Nsp^2 -atom bidentate pyridine-hydrazone sites. In this

way, Ag^+ induces the formation of double helicates with ligands **1** and **2**. Zn^{2+} prefers an octahedral coordination environment that results from **2** three- Nsp^2 -atom tridentate sites of type pyridine-hydrazone-pyrimidine, thus generating a grid.

Reaction of 1 equiv. of $\text{Ag}_2\text{L}_2\text{-DH}^{15c}$ with 2 equiv. of $\text{Zn}(\text{OTf})_2$ ($\text{OTf}^- = \text{CF}_3\text{SO}_3^-$) produces – without the need to precipitate Ag^+ as a halide – the corresponding grid $\text{Zn}_4\text{L}_4\text{-G}^{15b,22a}$ (solvent: CD_3NO_2 with 6–14% CD_3CN ; ESI, pp. S9–S11†). Where ZnCl_2 is used in the reaction with $\text{Ag}_2\text{L}_2\text{-DH}$, two equivalents of AgOTf per equiv. of DH are required according to the equation (ESI p. S8†):



On treatment of the double helicate $\text{Ag}_2\text{L}_2\text{-DH}$ in CD_3NO_2 with 2 equiv. of $\text{Zn}(\text{OTf})_2$ – added as a solution in a small volume of CD_3CN , or as a solid – the grid $\text{Zn}_4\text{L}_4\text{-G}$ was obtained. When the double helicate $\text{Ag}_2\text{L}_2\text{-DH}$ in CD_3NO_2 was treated with 2 equiv. of $\text{Zn}(\text{OTf})_2$, added as a solution in a small volume of CD_3CN (about 6–14% of the CD_3NO_2 volume), the grid $\text{Zn}_4\text{L}_4\text{-G}$ was obtained. When $\text{Zn}(\text{OTf})_2$ was added as a solid, without CD_3CN , was obtained a mixture without the $\text{Zn}_4\text{L}_4\text{-G}$ grid; addition of a small volume of CH_3CN (about 6–14% of the CD_3NO_2 volume) to this mixture produced the expected grid $\text{Zn}_4\text{L}_4\text{-G}$. A possible explanation could be that, in the case of the reaction $\text{Ag}_2\text{L}_2\text{-DH} \rightarrow \text{Zn}_4\text{L}_4\text{-G}$, the CH_3CN acts as a coordinating species for the Ag^+ ions and so contributes to the displacement of the equilibrium from the double helix towards the grid. The grid $\text{Zn}_4\text{L}_4\text{-G}$ should be – due to the π -stacking aromatic interaction between a phenyl ring and the two ligands between which that phenyl is located within the grid – more stable than the grid $\text{Zn}_4\text{L}_4\text{-G}$. This stability may be sufficient to make possible the formation of the grid $\text{Zn}_4\text{L}_4\text{-G}$ from the corresponding double helicate without, unlike in the case of the grid $\text{Zn}_4\text{L}_4\text{-G}$, the assistance of CH_3CN .

DOSY NMR was also used to study the conversion $\text{Ag}_2\text{L}_2\text{-DH} \rightarrow \text{Zn}_4\text{L}_4\text{-G}$ ($\text{L} = \text{1, 2}$). As expected, the volume of the grid species obtained from double helicates on treatment with $\text{Zn}(\text{OTf})_2$ was found in agreement with that of the grid prepared from the free ligands **L** and $\text{Zn}(\text{OTf})_2$.

The reverse conversion $\text{Zn}_4\text{L}_4\text{-G} \rightarrow \text{Ag}_2\text{L}_2\text{-DH}$ can be done as follows: after treatment of the grid with KOH , the solvent (CD_3CN or CD_3NO_2) is removed, and the ligand is extracted with CDCl_3 and separated from the solid residue (by centrifugation or filtration); after removal of CDCl_3 , CD_3NO_2 is added, then AgOTf is added to form the helicate. In order to simplify the procedure, we used ligand **2** and a mixture of CDCl_3 and CD_3NO_2 where ligand **2**, as well as the corresponding grid and double helicate were soluble. After precipitation of Zn^{2+} with KOH , the mixture was centrifuged (the ligand **2** being soluble in the mixture of solvents), and to the recovered liquid phase AgOTf was added to produce the $\text{Ag}_2\text{L}_2\text{-DH}$ (ESI, p. S13†).

In a pH-dependent system (Fig. 3b), the interconversion between $\text{Ag}_2\text{L}_2\text{-DH}$ and $\text{Zn}_4\text{L}_4\text{-G}$ was achieved as follows (ESI, p. S10†): the grid was generated from the double helicate by reaction with Zn^{2+} ; then, Zn^{2+} was complexed with hexacyclen, and the double helicate was regenerated; partial protonation of

hexacyclen with TfOH caused release of Zn^{2+} and formation of the grid (incomplete yield); finally, addition of triethylamine reactivated the hexacyclen that again encapsulated Zn^{2+} and resulted in the reformation of the double helicate.

(ii) The $\text{Pb}_4\text{L}_4\text{-G} \rightarrow \text{Zn}_4\text{L}_4\text{-G}$ conversion (Fig. 4a) can formally be seen as a substitution of Pb^{2+} by Zn^{2+} ions, although the real mechanism, involving breaking and formation of supramolecular bonds, must be more complex. Reaction of $\text{Pb}_4\text{L}_4\text{-G}^{15b}$ with 4 equiv. of $\text{Zn}(\text{OTf})_2$ produces a mixture which no longer contains the grid-like species $\text{Pb}_4\text{L}_4\text{-G}$ or $\text{Zn}_4\text{L}_4\text{-G}$ (ESI p. S2†). This suggests that the affinity of Zn^{2+} for the ligand, as well as its preference for octahedral coordination are not sufficient to displace the equilibrium towards $\text{Zn}_4\text{L}_4\text{-G}$. We considered that the involvement of Pb^{2+} ions in a weakly dissociating or sparingly soluble compound should displace the equilibrium. Indeed, addition of Br^- (as $\text{Bu}_4\text{P}^+\text{Br}^-$) to the above mixture, or treatment of $\text{Pb}_4\text{L}_4\text{-G}$ with four equivalents of ZnBr_2 or ZnCl_2 produced – along with the formation of PbX_2 ($\text{X} = \text{Br, Cl}$) which precipitates and, doing so, shifts the equilibrium – the expected $\text{Zn}_4\text{L}_4\text{-G}$ grid (solvent: CD_3CN ; ESI pp. S3–S4†): $\text{Pb}_4\text{L}_4\text{-G} + 4 \text{ZnX}_2 \rightarrow \text{Zn}_4\text{L}_4\text{-G} + 4 \text{PbX}_2$.

The reverse conversion $\text{Zn}_4\text{L}_4\text{-G} \rightarrow \text{Pb}_4\text{L}_4\text{-G}$ grid was achieved in several steps (Fig. 4b). Treatment of $\text{Zn}_4\text{L}_4\text{-G}$ (in CD_3CN) with KOH led to the precipitation of Zn^{2+} (as $\text{Zn}(\text{OH})_2$ or $\text{K}_2[\text{Zn}(\text{OH})_4]$), as well as of the free ligand **1**. After removal of CD_3CN , the free ligand **1** was extracted with CDCl_3 and used further for the preparation of $\text{Pb}_4\text{L}_4\text{-G}$ (see ESI, p. S5†).

Thus, in addition to its self-assembly from Zn^{2+} and a ligand, the same Zn^{2+} grid, $\text{Zn}_4\text{L}_4\text{-G}$, can be obtained, in reactions (i) and (ii), from a Ag^+ dinuclear double helicate or from a Pb^{2+} tetranuclear grid (exchange of metal ions and reorganization of the architectures).

(iii) The fusion (conproportionation) reaction of double helicate $\text{Ag}_2\text{L}_2\text{-DH}^{15c}$ with 1 equiv. of $\text{Ag}_2\text{L}_2\text{-DH}$ (Fig. 5) according to the equation



produces a mixture that contains each of the three helicates, namely two homoleptic (homost stranded) ones and one heteroleptic (heterost stranded) one. Ligands **1** and **2** equally participate to homo- and heteroleptic helicates, and so the observed molar percentages are of approximately 25% for $\text{Ag}_2\text{L}_2\text{-DH}$, 25% for $\text{Ag}_2\text{L}_2\text{-DH}$, and 50% for $\text{Ag}_2(\text{1})(\text{2})\text{-DH}$. For characterization of the new compound $\text{Ag}_2\text{L}_2\text{-DH}$, see ESI pp. S14–S20;† for ^1H , ^{13}C and DOSY of the mixture of three helicates, see ESI pp. S25–S31.†

For the reactions described above it might appear necessary, in practice, to slightly (2–10%) increase the amounts of reagents with respect to those theoretically calculated.

Experimental

For experimental details, see the ESI.†

Conclusions

To summarize, three supramolecular reactions were investigated: (i) a $\text{Ag}_2\text{L}_2\text{-DH}$ double-helicate into $\text{Zn}_4\text{L}_4\text{-G}$ grid



conversion, where the exchange of metal ions changes the nature of the metallo-supramolecular architecture, (ii) a $\text{Zn}_4\text{1}_4\text{-G}$ grid into $\text{Pb}_4\text{1}_4\text{-G}$ grid conversion driven by a halide-induced precipitation and where the nature of the metallo-supramolecular architecture is conserved, and (iii) a double exchange of ligands during the fusion of two double helicates.

The grid/grid and double-helicate/grid conversions were made reversible by precipitation of Zn^{2+} with KOH and subsequent reaction of the free ligand with Ag^+ or Pb^{2+} , or, for one DH/G interconversion, in a pH-dependent way.

In perspective, such ligands could be introduced in larger and more complex, suitably decorated, architectures where such supramolecular reactions can act as actuators of various properties (charge, volume, multivalency).

Acknowledgements

We thank Prof. Jack Harrowfield for helpful discussions, and Stéphanie Kouaho for mass spectrometry analyses.

Notes and references

- 1 J.-M. Lehn, *Angew. Chem., Int. Ed. Engl.*, 1988, **27**, 89–121; J.-M. Lehn, *Angew. Chem., Int. Ed. Engl.*, 1990, **29**, 1304–1319.
- 2 R. F. Ludlow and S. Otto, *Chem. Soc. Rev.*, 2008, **37**, 101–108.
- 3 D. G. Kurth, *Sci. Technol. Adv. Mater.*, 2008, **9**, 014103.
- 4 J.-M. Lehn, *Angew. Chem., Int. Ed.*, 2013, **52**, 2836–2850.
- 5 K. C.-F. Leung, C.-P. Chak, C.-M. Lo, W.-Y. Wong, S. Xuan and C. H. K. Cheng, *Chem.-Asian J.*, 2009, **4**, 364–381; X. Yan, F. Wang, B. Zheng and F. Huang, *Chem. Soc. Rev.*, 2012, **41**, 6042–6065; A. J. McConnell, C. S. Wood, P. P. Neelakandan and J. R. Nitschke, *Chem. Rev.*, 2015, **115**, 7729–7793.
- 6 See, for example: M. Wang, W.-J. Lan, Y.-R. Zheng, T. R. Cook, H. S. White and P. J. Stang, *J. Am. Chem. Soc.*, 2011, **133**, 10752–10755.
- 7 For a selection of reviews, see: C. Piguet, G. Bernardinelli and G. Hopfgartner, *Chem. Rev.*, 1997, **97**, 2005–2062; C. Piguet, *J. Inclusion Phenom. Macrocyclic Chem.*, 1999, **34**, 361–391; M. Albrecht, *Chem. Rev.*, 2001, **101**, 3457–3497; M. J. Hannon and L. J. Childs, *Supramol. Chem.*, 2004, **16**, 7–22 For reviews, see: M. Ruben, J. Rojo, F. J. Romero-Salguero, L. H. Uppadine and J.-M. Lehn, *Angew. Chem., Int. Ed.*, 2004, **43**, 3644–3662; L. N. Dawe, T. S. M. Abedin and L. K. Thompson, *Dalton Trans.*, 2008, 1661–1675; L. N. Dawe, K. V. Shuvaev and L. K. Thompson, *Chem. Soc. Rev.*, 2009, **38**, 2334–2359; A.-M. Stadler, *Eur. J. Inorg. Chem.*, 2009, 4751–4770; J. Hardy, *Chem. Soc. Rev.*, 2013, **42**, 7881–7899.
- 8 J. D. Watson and F. H. Crick, *Nature*, 1953, **171**, 737–738.
- 9 B. M. Burkhardt, N. Li, D. A. Langs, W. A. Pangborn and W. L. Duax, *Proc. Natl. Acad. Sci. U. S. A.*, 1998, **95**, 12950–12955.
- 10 K. Miwa, Y. Furusho and E. Yashima, *Nat. Chem.*, 2010, **2**, 444–449.
- 11 C.-T. Yeung, L.-H. Yeung, C.-S. Tsang, W.-Y. Wong and H.-L. Kwong, *Chem. Commun.*, 2007, 5203–5205.
- 12 B. R. Manzano, F. A. Jalón, I. M. Ortiz, M. L. Soriano, F. Gómez de la Torre, J. Elguero, M. A. Maestro, K. Mereiter and T. D. W. Claridge, *Inorg. Chem.*, 2008, **47**, 413–428.
- 13 J. E. Beves, J. J. Danon, D. A. Leigh, J.-F. Lemonnier and I. J. Vitorica-Yrezabal, *Angew. Chem., Int. Ed.*, 2015, **54**, 7555–7559.
- 14 For assembly/disassembly of double helicates, see: J.-P. Gisselbrecht, M. Gross, J.-M. Lehn, J.-P. Sauvage, R. Ziessel, C. Piccinni-Leopardi, J. M. Arrieta, G. Germain and M. V. Meersche, *Nouv. J. Chim.*, 1984, **8**, 661–667; Y. Yao, M. W. Perkovic, D. P. Rillema and C. Woods, *Inorg. Chem.*, 1992, **31**, 3956–3962; K. T. Potts, M. Keshavarz-K, F. S. Tham, H. D. Abruña and C. R. Arana, *Inorg. Chem.*, 1993, **32**, 4422–4435; V. Amendola, L. Fabbrizzi, L. Linati, C. Mangano, P. Pallavicini, V. Pedrazzini and M. Zema, *Chem.-Eur. J.*, 1999, **5**, 3679–3688; V. Amendola, L. Fabbrizzi, L. Gianelli, C. Maggi, C. Mangano, P. Pallavicini and M. Zema, *Inorg. Chem.*, 2001, **40**, 3579–3587; V. Amendola, L. Fabbrizzi, P. Pallavicini, E. Sartirana and A. Taglietti, *Inorg. Chem.*, 2003, **42**, 1632–1636; M. Boiocchi and L. Fabbrizzi, *Chem. Soc. Rev.*, 2014, **43**, 1835–1847. See also: M. Barboiu, G. Vaughan, N. Kyritsakas and J.-M. Lehn, *Chem.-Eur. J.*, 2003, **9**, 763–769.
- 15 (a) Grid/double-cross/stick: M. Barboiu, G. Vaughan, R. Graff and J.-M. Lehn, *J. Am. Chem. Soc.*, 2003, **125**, 10257–10265; (b) Grid/stick: A.-M. Stadler, N. Kyritsakas, R. Graff and J.-M. Lehn, *Chem.-Eur. J.*, 2006, **12**, 4503–4522; (c) Double helicate/stick: A.-M. Stadler, N. Kyritsakas, G. Vaughan and J.-M. Lehn, *Chem.-Eur. J.*, 2007, **13**, 59–68; (d) Grid/pincer: J. Ramirez, A. M. Stadler, N. Kyritsakas and J.-M. Lehn, *Chem. Commun.*, 2007, 237–239; (e) Pincer/stick: J. Ramirez, A.-M. Stadler, L. Brelot and J.-M. Lehn, *Tetrahedron*, 2008, **64**, 8402–8410; (f) Double helicate/stick: D. J. Hutchinson, S. A. Cameron, L. R. Hanton and S. C. Moratti, *Inorg. Chem.*, 2012, **51**, 5070–5081; (g) Double-helicate/grid: A.-M. Stadler, C. Burg, J. Ramirez and J.-M. Lehn, *Chem. Commun.*, 2013, **49**, 5733–5735; (h) see also: P. N. W. Baxter, R. G. Khoury, J.-M. Lehn, G. Baum and D. Fenske, *Chem.-Eur. J.*, 2000, **6**, 4140–4148; (i) C. S. Campos-Fernandez, B. L. Schottel, H. T. Chifotides, J. K. Bera, J. Bacsá, J. M. Koomen, D. H. Russell and K. R. Dunbar, *J. Am. Chem. Soc.*, 2005, **127**, 12909–12923.
- 16 For examples of supramolecular interconversions, see: D. P. Funeriu, J.-M. Lehn, K. M. Fromm and D. Fenske, *Chem.-Eur. J.*, 2000, **6**, 2103–2111; K. Kumazawa, Y. Yamanoi, M. Yoshizawa, T. Kusukawa and M. Fujita, *Angew. Chem., Int. Ed.*, 2004, **43**, 5936–5940; A. M. Brown, M. V. Ovchinnikov, C. L. Stern and C. A. Mirkin, *J. Am. Chem. Soc.*, 2004, **126**, 14316–14317; O. Mamula, M. Lama, H. Stoeckli-Evans and S. Shova, *Angew. Chem., Int. Ed.*, 2006, **45**, 4940–4944; K. Harano, S. Hiraoka and M. Shionoya, *J. Am. Chem. Soc.*, 2007, **129**, 5300–5301; J. Heo, Y.-M. Jeon and C. A. Mirkin, *J. Am. Chem. Soc.*,



- 2007, **129**, 7712–7713; L. Zhao, B. H. Northrop and P. J. Stang, *J. Am. Chem. Soc.*, 2008, **130**, 11886–11888; S. Hiraoka, Y. Sakata and M. Shionoya, *J. Am. Chem. Soc.*, 2008, **130**, 10058–10059; H. Dong, J. Yang, X. Liu and S. Gou, *Inorg. Chem.*, 2008, **47**, 2913–2915; P. J. Lusby, P. Müller, S. J. Pike and A. M. Z. Slawin, *J. Am. Chem. Soc.*, 2009, **131**, 16398–16400; K. Parimal, E. H. Witlicki and A. H. Flood, *Angew. Chem., Int. Ed.*, 2010, **49**, 4628–4632; S. Chen, L.-J. Chen, H.-B. Yang, H. Tian and W. Zhu, *J. Am. Chem. Soc.*, 2012, **134**, 13596–13599; X. Yan, J.-F. Xu, T. R. Cook, F. Huang, Q.-Z. Yang, C.-H. Tung and P. J. Stang, *Proc. Natl. Acad. Sci. U. S. A.*, 2014, **111**, 8717–8722; X. Lu, X. Li, K. Guo, T.-Z. Xie, C. N. Moorefield, C. Wesdemiotis and G. R. Newkome, *J. Am. Chem. Soc.*, 2014, **136**, 18149–18155; Y.-F. Han, L. Zhang, L.-H. Weng and G.-X. Jin, *J. Am. Chem. Soc.*, 2014, **136**, 14608–14615; Y. Li, Z. Jiang, J. Yuan, D. Liu, T. Wu, C. N. Moorefield, G. R. Newkome and P. Wang, *Chem. Commun.*, 2015, **51**, 5766–5769; D. Preston, A. Fox-Charles, W. K. C. Lo and J. D. Crowley, *Chem. Commun.*, 2015, **51**, 9042–9045; M. Han, Y. Luo, B. Damaschke, L. Gýmez, X. Ribas, A. Jose, P. Peretzki, M. Seibt and G. H. Clever, *Angew. Chem., Int. Ed.*, 2016, **55**, 445–449.
- 17 J.-L. Schmitt, A.-M. Stadler, N. Kyritsakas and J.-M. Lehn, *Helv. Chim. Acta*, 2003, **86**, 1598–1624.
- 18 X. Su and I. Arahamian, *Chem. Soc. Rev.*, 2014, **43**, 1963–1981.
- 19 See for example the cases of (a) spectrin: S.-C. Liu and J. Palek, *Nature*, 1980, **285**, 586–588; E. Ungewickell and W. Gratzner, *Eur. J. Biochem.*, 1978, **88**, 379–385. (b) the clock protein KaiB: T. Iida, R. Mutoh, K. Onai, M. Morishita, Y. Furukawa, K. Namba and M. Ishiura, *Genes Cells*, 2015, **20**, 173–190; (c) 6-phosphogluconate dehydrogenase, see: S. Hanau, L. Proietti d'Empaire, I. Capone, S. Alberighi, R. Montioli and F. Dallochio, *Biochim. Biophys. Acta*, 2013, **1834**, 2647–2652; (d) Bovine IF1, see: E. Cabezon, P. J. G. Butler, M. J. Runswick and J. E. Walker, *J. Biol. Chem.*, 2000, **275**, 25460–25464; (e) phosphofructokinase, see: J. Xu, T. Oshima and M. Yoshida, *J. Mol. Biol.*, 1990, **215**, 597–606.
- 20 For the case of (a) uracil phosphoribosyltransferase from *Escherichia coli*, see: K. F. Jensen and B. Mygind, *Eur. J. Biochem.*, 1996, **240**, 637–645; (b) mushroom tyrosinase, see: R. L. Jolley Jr, D. A. Robb and H. S. Mason, *J. Biol. Chem.*, 1969, **244**, 1593–1599; (c) D-amino acid oxidase, see: E. Antonini, M. Brunori, R. Bruzzesi, E. Chiancone and V. Massey, *J. Biol. Chem.*, 1966, **241**, 2358–2366 See also: (d) E. K. Jaffe, *Open Conf. Proc. J.*, 2010, **1**, 1–6; T. Selwood and E. K. Jaffe, *Arch. Biochem. Biophys.*, 2012, **519**, 131–143.
- 21 For an example of supramolecular substitution reaction, see: J. Santamaria, T. Martín, G. Hilmersson, S. L. Craig and J. Rebek Jr, *Proc. Natl. Acad. Sci. U. S. A.*, 1999, **96**, 8344–8347.
- 22 Pb²⁺ or Zn²⁺ hydrazone-based grids: (a) M. Barboiu, M. Ruben, G. Blasen, N. Kyritsakas, E. Chacko, M. Dutta, O. Radekovich, K. Lenton, D. J. R. Brook and J.-M. Lehn, *Eur. J. Inorg. Chem.*, 2006, 784–789; (b) A. R. Stefankiewicz, G. Rogez, J. Harrowfield, M. Drillon and J.-M. Lehn, *Dalton Trans.*, 2009, 5787–5802; (c) M. Dutta, M. Movassat, D. J. R. Brook, A. Oliver and D. Ward, *Supramol. Chem.*, 2011, **23**, 630–640; (d) D. J. Hutchinson, L. R. Hanton and S. C. Moratti, *Inorg. Chem.*, 2011, **50**, 7637–7649.
- 23 M. L. Saha, S. Pramanik and M. Schmittel, *Chem. Commun.*, 2012, **48**, 9459–9461.
- 24 For other examples of Ag⁺ hydrazone-based double helicates, see: D. J. Hutchinson, P. M. James, L. R. Hanton and S. C. Moratti, *Inorg. Chem.*, 2014, **53**, 2122–2132; D. J. Hutchinson, L. R. Hanton and S. C. Moratti, *Dalton Trans.*, 2014, **43**, 8205–8218.

