

## CORRECTION

[View Article Online](#)  
[View Journal](#) | [View Issue](#)Cite this: *Chem. Sci.*, 2016, 7, 5596**Correction: Essential role of hydride ion in ruthenium-based ammonia synthesis catalysts**Masaaki Kitano,<sup>a</sup> Yasunori Inoue,<sup>b</sup> Hiroki Ishikawa,<sup>b</sup> Kyosuke Yamagata,<sup>b</sup>  
Takuya Nakao,<sup>b</sup> Tomofumi Tada,<sup>a</sup> Satoru Matsuishi,<sup>a</sup> Toshiharu Yokoyama,<sup>ac</sup>  
Michikazu Hara<sup>\*bcd</sup> and Hideo Hosono<sup>\*abcd</sup>

DOI: 10.1039/c6sc90039a

[www.rsc.org/chemicalscience](http://www.rsc.org/chemicalscience)Correction for 'Essential role of hydride ion in ruthenium-based ammonia synthesis catalysts' by Masaaki Kitano *et al.*, *Chem. Sci.*, 2016, 7, 4036–4043.

The authors regret that in the original article incorrect units were used to define both the number of surface Ru atoms ( $N_s$ ) and the ammonia synthesis rate ( $r_{\text{NH}_3}$ ) in columns 6 and 7 of Table 1. In both cases, ' $\mu\text{mol}$ ' should have been used instead of ' $\text{mmol}$ '. Therefore, the correct units for  $N_s$  and  $r_{\text{NH}_3}$  are ' $\mu\text{mol g}^{-1}$ ' and ' $\mu\text{mol g}^{-1} \text{h}^{-1}$ ', respectively.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

<sup>a</sup>Materials Research Center for Element Strategy, Tokyo Institute of Technology, 4259 Nagatsuta, Midori-ku, Yokohama 226-8503, Japan. E-mail: [hosono@msl.titech.ac.jp](mailto:hosono@msl.titech.ac.jp)<sup>b</sup>Laboratory for Materials and Structures, Tokyo Institute of Technology, 4259 Nagatsuta, Midori-ku, Yokohama 226-8503, Japan. E-mail: [mhara@msl.titech.ac.jp](mailto:mhara@msl.titech.ac.jp)<sup>c</sup>ACCEL, Japan Science and Technology Agency, 4-1-8 Honcho, Kawaguchi, Saitama 332-0012, Japan<sup>d</sup>Frontier Research Center, Tokyo Institute of Technology, 4259 Nagatsuta, Midori-ku, Yokohama 226-8503, Japan