



Cite this: *RSC Adv.*, 2017, 7, 1646

# Carbon metabolism and transcriptional variation in response to salt stress in the genome shuffled *Candida versatilis* and a wild-type salt tolerant yeast strain†

Wei Qi,<sup>abc</sup> Wen-Tao Zhang<sup>d</sup> and Fu-Ping Lu<sup>\*abc</sup>

The carbon metabolism and molecular mechanisms of adaptation response when exposed to conditions causing osmotic stress in strains of a wild-type of *Candida versatilis* (WT) and S3–5 were investigated. The levels of glucose, glycerol and ethanol, as well as transcriptional variation of eight genes that encoded hexokinase (*HXK1*), mitogen-activated protein kinase (*HOG1*), glycerol-3-phosphate dehydrogenase (*GPD1*), aquaglyceroporin (*FPS1*), pyruvate kinase (*PYK1*), pyruvate decarboxylase (*PDC1*), aldehyde dehydrogenase (*ALD6*), alcohol dehydrogenase (*ADH1*) in short-term response to salt stress were determined. The results showed that an increase in salt concentration negatively affected glucose consumption and ethanol production in *C. versatilis*. In contrast, glycerol produced by *C. versatilis* was positively affected by salt addition. The results showed that *HXK1* contributes to rerouting glycolytic flux towards higher glycolysis. The glycerol production generated by S3–5 was higher as compared to WT, which is ascribed to the up-regulated expression of *GPD1* and *HOG1* genes and down-regulated expression of the *FPS1* gene after the initiation of stress treatment.

Received 13th October 2016  
Accepted 10th November 2016

DOI: 10.1039/c6ra25188a

www.rsc.org/advances

## Introduction

*Candida versatilis*, a salt-tolerant yeast, is widely used in the fermentation of soy sauce, soybean paste and miso.<sup>1</sup> It could contribute in a positive manner to the flavour development though a series of metabolic activities, such as formation of 4-hydroxy-2(or 5)-ethyl-5(or 2)-methyl-3(2*H*)-furanone (HEMF) and 4-ethyl guaiacol (4-EG).<sup>2</sup> During the fermentation process, yeast is subjected to a high salt osmotic stress which would affect its viability and fermentation efficiency. Dynamic changes occur in the complex biological networks which would provide insight into the link between the genome, gene-expression, proteins and metabolites, etc.

Various studies have been performed to investigate salt-tolerant yeast responses and adaptation stress mechanisms by which cells adapt to 'adverse' conditions in the fermentation,<sup>3–6</sup> including (1) its transport systems for adjusting ion homeostasis and counteracting the toxic effects caused by sodium ions,<sup>7</sup> such as plasma membrane H<sup>+</sup>-ATPase<sup>8,9</sup> and Na<sup>+</sup>/H<sup>+</sup> antiporter<sup>9</sup>; (2) synthesis and accumulation compatible solutes, such as glycerol,<sup>10,11</sup> trehalose,<sup>12</sup> mannitol,<sup>3,4</sup> which stabilize the activity and structure of diverse proteins and membranes<sup>13</sup>; (3) regulation membrane fluidity and permeability by changing lipid components<sup>14</sup>; (4) multiple signalling pathways to perceive and to quickly respond to altered osmolarity, such as high osmolarity glycerol (HOG) response pathway.<sup>15,16</sup>

Cells adapt to rapid and drastic environmental changes by turning 'on' and 'off' specific subsets of genes and proteins. From a molecular point of view, it is important to obtain information of signal transduction and the transcriptional response on the yeast adaptation during stress-responsive phase. The cellular mechanisms of adaptation to salt stress in strains of a wild type of *C. versatilis* (WT) and S3–5, genome shuffling strains of *C. versatilis* with improving tolerance to salt, has been well conducted in the previous research.<sup>3,4,6</sup> However, there have been very few studies on this topic by rapidly shifting *Candida versatilis* cells from a medium without salt to stressful conditions.<sup>17</sup> *C. versatilis* growing exponentially on glucose produces glycerol and ethanol, the both compounds can be used as single carbon and energy sources.<sup>4</sup> Moreover, glycerol

<sup>a</sup>Key Laboratory of Industrial Fermentation Microbiology, Tianjin University of Science & Technology, Ministry of Education, No. 29, 13th Road, Tianjin Economy-and-Technology Development Area, Tianjin 300457, P. R. China. E-mail: qfweisiming@126.com; Fax: +86-22-60601332; Tel: +86-22-60601428

<sup>b</sup>National Engineering Laboratory for Industrial Enzymes, Tianjin University of Science & Technology, Tianjin Economy-and-Technology Development Area, Tianjin 300457, P. R. China

<sup>c</sup>Tianjin Key Laboratory of Industrial Microbiology, Tianjin University of Science & Technology, Tianjin Economy-and-Technology Development Area, Tianjin 300457, P. R. China

<sup>d</sup>Key Laboratory of Food Nutrition and Safety, Tianjin University of Science & Technology, Ministry of Education, Tianjin 300457, P. R. China

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c6ra25188a






Fig. 1 Carbon metabolism of yeast.

have been implicated as potential stress protectants that accumulate in yeasts during various stress conditions.<sup>17</sup> In the present study, glucose, glycerol and ethanol metabolism as well as expression of related genes (underlined in Fig. 1) was analyzed by the treatment of NaCl. The investigation of such molecular responses will help us to understand the molecular mechanisms by which cells adapt to fermentation conditions.

## Materials and methods

### Chemicals

Analytical grade chemicals and solvents were purchased from Tianjin Chemical Reagent Research Institute (Tianjin, China). Chromatographic grade (HPLC) acetonitrile were obtained from Merck (Darmstadt, Germany). Double-distilled water for the dilution of samples was purified using a Milli-Q system (Millipore, Bedford, MA, USA).

### Yeast strains

Yeast strains *C. versatilis* wild-type (WT) and S3-5 used in this work were kept in our laboratory. The genome shuffled strain S3-5 was derived from the WT yeast strain *C. versatilis*, initially isolated for improved tolerance to salt.<sup>18</sup>

### Media and growth conditions

*C. versatilis* cells were grown in 10 ml YPD at 30 °C with rotary shaking (at 180 rpm). After 24 h incubation, the culture was transferred to 250 ml flask containing 100 ml of YPD (1% yeast extract, 2% peptone, 2% glucose). The yeast was incubated to its early stationary phase (OD 1.0 (600 nm)). Cells in early stationary phase were inoculated at 10% (v/v) in YPD containing 0, 2 M and 4 M NaCl and incubated at 30 °C.

### Analytical methods and data analysis

5 ml culture was undertaken at 0, 0.5, 1, 2, 4, 6 h and filtered using a 0.45 µm membrane filter (Millipore, France) under a pump-applied vacuum. The clarified supernatant was used for determination of extra-cellular glucose, glycerol and ethanol, and the filtrate was used for dry biomass assay.

### Dry biomass

The filtrate was oven-dried at 105 °C for 4 h to constant mass.<sup>19</sup>

### Intra- and extra-cellular concentrations of glucose and glycerol

Cells (5 ml culture) were collected by centrifugation (5000 rpm, 4 °C, 5 min) and washed twice with double-distilled water and heated in a constant temperature water bath at 95 °C for 15 min. After centrifugation as mentioned previously, the supernatant was used to measure the internal glucose and glycerol.

The glucose and glycerol was performed using a Shimadzu 20AB (Shimadzu, Japan), with a refractive index detector after separation in an Aminex HPX-87H column. The column oven and RID were set to 30 °C and 35 °C respectively. The acetonitrile–water (70%, v/v) mix was used as the eluent at a flow rate of 1 ml min<sup>-1</sup>. The intra-cellular glucose and glycerol content was expressed as a percentage of dry biomass.

### The content of external ethanol

The ethanol was analyzed by headspace solid phase micro-extraction (HS-SPME) gas chromatography spectrometry as De Martinis described.<sup>20</sup> The GC analysis was performed using a Shimadzu GC with flame ionization detector (GC-FID) and equipped with a DB-5MS fused silica column (30 m × 0.25 mm × 0.25 µm, J&W Scientific, Folsom, CA). The oven temperature was maintained at 60 °C for 10 min and increased to 240 °C at a rate of 5 °C min<sup>-1</sup>. The injector and transfer-line temperatures were 250 °C and 280 °C, respectively. Helium was used as carrier gas (about 47 ml min<sup>-1</sup>, 30 kPa) and the compounds were detected *via* a flame-ionization detector.

### RNA isolation and real-time PCR analysis

Total RNA prepared using the MiniBEST Universal RNA Extraction Kit (Takara, Dalian, China). Three biological replicates were prepared for each accession and salt combination. RNA was quantified by Nanodrop (ND-2000, NanoDrop Technologies, USA); quality was assessed by agarose gel analyses and 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA). Ten micrograms of total RNA from each sample was converted to double-stranded cDNA using the PrimeScript™ RT reagent Kit (Perfect Real Time) (TaKaRa, Dalian, China).

Primers (Table S1†) were designed using primer premier 5.0 (Premier, Canada) and provided by Sangon Biotech Co., Ltd. (Shanghai, China).

Real-time PCR analysis was performed in Eppendorf mastercycler ep realplex<sup>4</sup> (Eppendorf, Germany) using SYBR Green. Reactions were performed in a volume of 20 µl containing 2 µl cDNA, 0.8 µl forward and reverse primers, and 10 µl SYBR®



Premix Ex Taq™ II (Takara, Dalian, China). Duplicates of each sample and negative controls were included in the analysis. The amplification conditions were as follow: 95 °C for 30 s; 40 cycles of 95 °C for 5 s, 55 °C for 30 s. Each sample was repeated three times. Data were analyzed by the  $2^{-\Delta\Delta C_T}$  method.

### Statistical analysis

Statistical significance was determined using the SAS statistical analysis program, Version 8.01 (North Carolina State University, USA).

## Results and discussion

### The level of glucose of the salt-tolerant yeast *Candida versatilis* under stress conditions

Glucose was transported by facilitated diffusion *via* carrier protein transporter in *Candida versatilis*.<sup>3</sup> Extra-cellular glucose was absorbed and consumed by *C. versatilis* to maintain energy metabolism and central carbon metabolism (Fig. 2B and D). The level of intra-cellular glucose of *C. versatilis* after NaCl treatment was higher than that in cells treated without stressing agent (Fig. 2A and C). Our result also indicated that glucose was uptaken and utilized by S3-5 was higher as compare to WT (Table 1). The transient growth arrest following hyperosmotic

shock further adds to redirecting almost all glycolytic flux from biomass towards glycerol production.<sup>21</sup>

Prior research pointed out that three genes, *SNF3*, *HXT1* and *HXT2*, encode three different glucose transporters, and the rate of sugar uptake in yeast cells is controlled by changes in affinity of the corresponding transporters as well as by an irreversible inactivation that affects their  $V_{max}$ .<sup>22</sup> But the mechanisms involved in glucose transport processes of *Candida versatilis* are unknown at present.

### Glycerol accumulation of the salt-tolerant yeast *Candida versatilis* under stress conditions

As shown in Fig. 3A and C, intra-cellular glycerol after treatment of NaCl was obviously more than control group, moreover, the intra-cellular glycerol by the treatment of 2 M NaCl was higher than that by 4 M NaCl. The similar phenomenon has also been observed.<sup>3</sup> Salt stress was accompanied by an increase in the intracellular level of glycerol,<sup>23</sup> which could decrease the intracellular water potential, restore water influx and consequently restore cell volume and turgor pressure.<sup>24</sup> Osmoadaptation is robust to loss of individual adaptation pathways because of the existence and upregulation of alternative routes of glycerol accumulation.<sup>21</sup> In yeasts, salt-tolerant yeast *Debaryomyces hansenii* and *Saccharomyces cerevisiae*, exposed to strong osmotic stress the intracellular glycerol concentration can reach molar levels.<sup>25</sup>




Fig. 2 The content of glucose of the salt-tolerant yeast *Candida versatilis*: (A) intracellular glucose of WT; (B) extracellular glucose of WT; (C) intracellular glucose of S3-5; (D) extracellular glucose of S3-5.

Table 1 Summary of salt-tolerant yeast *Candida versatilis*: expressing the carbon metabolism enzymes in response to salt stress

Strains	NaCl (g L <sup>-1</sup> )	Dry cell mass (g L <sup>-1</sup> )	Glucose consumption rate (mg L <sup>-1</sup> h <sup>-1</sup> )	Glycerol productivity (g L <sup>-1</sup> h <sup>-1</sup> )	Ethanol productivity (g L <sup>-1</sup> h <sup>-1</sup> )
WT	0	3.84	3.49	0.28	11.40
	90	3.02	2.05	1.42	9.08
	180	1.49	2.81	2.04	1.65
S3-5	0	4.46	4.63	0.59	11.75
	90	3.20	3.77	2.26	9.21
	180	1.73	3.47	3.81	1.88






Fig. 3 The content of glycerol of the salt-tolerant yeast *Candida versatilis*: (A) intracellular glycerol of WT; (B) extracellular glycerol of WT; (C) intracellular glycerol of S3-5; (D) extracellular glycerol of S3-5.




Fig. 4 The content of ethanol of the salt-tolerant yeast *Candida versatilis*: (A) WT; (B) S3-5.

Extra-cellular glycerol level of *C. versatilis* after treatment of NaCl was significantly higher than salt-free group, and extra-cellular glycerol increased with increasing salt concentration

in the medium (Fig. 3B and D). The intra-, extra-cellular glycerol of S3-5 was higher than that of WT. The level of intra-cellular glycerol produced by S3-5 increased 17.2% than WT at 6 h,




Fig. 5 Expression level of genes related to glucose metabolism of the salt-tolerant yeast *Candida versatilis* by quantitative real-time PCR analysis: (A) expression level of *HXK1* gene after 0 M NaCl treatment; (B) expression level of *HXK1* gene after 2 M NaCl treatment; (C) expression level of *HXK1* gene after 4 M treatment; (D) relative expression level of *HXK1* (S3-5/WT).






Fig. 6 Expression level of genes related to glycerol metabolism of the salt-tolerant yeast *Candida versatilis* by quantitative real-time PCR analysis: (A) expression level of *HOG1* after 0 M NaCl treatment; (B) expression level of *HOG1* after 2 M NaCl treatment; (C) relative expression level of *HOG1* after 4 M NaCl treatment; (D) relative expression level of *HOG1* (S3-5/WT); (E) relative expression level of *GPD1* after 0 M NaCl treatment; (F) relative expression level of *GPD1* after 2 M NaCl treatment; (G) relative expression level of *GPD1* after 4 M NaCl treatment; (H) relative expression level of *GPD1* (S3-5/WT); (I) relative expression level of *FPS1* after 0 M NaCl treatment; (J) relative expression level of *FPS1* after 2 M NaCl treatment; (K) relative expression level of *FPS1* after 4 M NaCl treatment; (L) relative expression *FPS1* (S3-5/WT).

and extra-cellular glycerol level of S3-5 increased by 57.6% than WT at the same time.

There have two different types of active-transport systems for glycerol transport,  $\text{Na}^+$ /glycerol and  $\text{H}^+$ /glycerol symports,<sup>26</sup> and glycerol of *Candida versatilis* transported by proton symport.<sup>3</sup> The previous conclusion that the salt tolerance of S3-5 was higher than WT ascribed to the higher glycerol production, have verified by the experimental results.<sup>6</sup>

#### Ethanol accumulation of the salt-tolerant yeast *Candida versatilis* under stress conditions

The extra-cellular ethanol was affected by the level of salt (Fig. 4). At the beginning process, the level of ethanol was relatively lower under stress cultivated, which because intermediate of glucose metabolism was consumed to yield glycerol. The ethanol became increased until the yeast adapt to the salt stress. The level of ethanol produced by S3-5 increased at 2 h, while in that of WT elevated until 4 h.






Fig. 7 Expression level of genes related to ethanol metabolism of the salt-tolerant yeast *Candida versatilis* by quantitative real-time PCR analysis: (A) relative expression level of *PDC1* after 0 M NaCl treatment; (B) relative expression level of *PDC1* after 2 M NaCl treatment; (C) relative expression level of *PDC1* after 4 M NaCl treatment; (D) relative expression level of *PDC1* (S3–5/WT); (E) relative expression level of *PYK1* after 0 M NaCl treatment; (F) relative expression level of *PYK1* after 2 M NaCl treatment; (G) relative expression level of *PYK1* after 4 M NaCl treatment; (H) relative expression level of *PYK1* (S3–5/WT); (I) relative expression level of *ADH1* after 0 M NaCl treatment; (J) relative expression level of *ADH1* after 2 M NaCl treatment; (K) relative expression level of *ADH1* after 4 M NaCl treatment; (L) relative expression level of *ADH1* (S3–5/WT); (M) relative expression level of *ALD6* after 0 M NaCl treatment; (N) relative expression level of *ALD6* after 2 M NaCl treatment; (O) relative expression level of *ALD6* after 4 M NaCl treatment; (P) relative expression level of *ALD6* (S3–5/WT).

The results obtained by analysis of glucose, glycerol and ethanol showed that salt pressure contributes to rerouting glycolytic flux towards higher glycolysis and osmoadaptation prioritizes the redox and energy balance in glycolysis while rerouting flux from biomass to glycerol production.<sup>21</sup>

### The glucose metabolic pathway and the stress response

Glucose uptake occurs which is mediated by hexose transporters.<sup>27</sup> The main pathway of yeast carbon metabolism is Embden–Meyerhof–Parnas pathway (EMP), in which glucose was catalyzed by hexokinase encoded by *HXK1* gene (hexose transporter gene) to glucose-6-phosphate. The expression level of *HXK1* gene increased with increasing salt concentration in the medium (Fig. 5A–C), the expression level of the *HXK1* gene of S3–5 was higher than that of WT (Fig. 5D).

### The glycerol metabolic pathway and the stress response

Glycerol, as a primary osmoregulatory solute, is formed from the glycolytic intermediate dihydroxyacetone phosphate in two steps: firstly, dihydroxyacetone phosphate is reduced to glycerol-3-phosphate as catalysed by glycerol-3-phosphate dehydrogenase (GPD) encoded by the *GPD1* gene (there is only one *C. versatilis* *GPDH* gene<sup>5</sup>), and glycerol-3-phosphate then yields glycerol. After a hyperosmotic challenge, intracellular glycerol levels are increased, as a combined result of the diminished efflux and the enhanced synthesis,<sup>17</sup> an additional Hog1-independent mechanism retains intracellular glycerol for adaptation.

The expression of the *GPD1* gene (Fig. 6A–D) and *HOG1* gene (Fig. 6E–H) were induced by NaCl. The expression levels of *GPD1* gene increased rapidly after the initiation of stress treatment, similar to the expression pattern of *HOG1*. The previous research suggested that the response of yeast to osmotic stress



is controlled by HOG pathway, whose important target gene is *GPD1*,<sup>5</sup> and the transcription of *GPD1* in *C. versatilis* cells was stimulated by high concentrations of NaCl.<sup>5</sup> Moreover, the result was also observed by Silva-Graca that glycerol-3-phosphate dehydrogenase of *Candida versatilis* exhibited a very significant increase with increasing salt concentrations.<sup>4</sup> Because rapid transcriptional response under osmotic stress is accompanied by a delay in phosphorylation and nuclear translocation of Hog1p,<sup>4,28</sup> therefore, extra-glycerol begin to increase at 1 h. The transcription levels of the *GPD1* and *HOG1* gene of S3-5 were higher than that of WT which is the key reason that S3-5 has higher glycerol production and stronger tolerance to salt stress compare to WT.

Intra-cellular glycerol level are not only affected by yield but also regulated by transmembrane transport through the glycerol facilitator, Fps1p.<sup>29</sup> Under hyperosmolarity condition, together with increased glycerol synthesis, yeast cells accumulate intracellular glycerol content *via* closing Fps1p. Its functions as a turgor valve during cell fusion.<sup>28,30</sup> Hyperosmotic stress also leads to rapid closure of the glycerol facilitator Fps1 preventing glycerol outflow.<sup>21</sup> Down-regulated expression of *FPS1* gene was observed (Fig. 6I–L). The results showed that the expression of *FPS1* gene of S3-5 was lower than WT, which might be leading to S3-5 has higher level of glycerol production.

### The ethanol metabolic pathway and the stress response

Ethanol is formed from the glycolytic intermediate glyceraldehyde-3-phosphate in three steps: firstly, glyceraldehyde-3-phosphate metabolize to pyruvic acid catalysed by pyruvate kinase (PYK) encoded by the *PYK1* gene; then pyruvic acid reduced to aldehyde catalysed by pyruvate dehydrogenase complex (PDC) encoded by the *PDC1* gene; thirdly, aldehyde reduced to ethanol catalysed by alcohol dehydrogenase (ADHs) encoded by the *ADH* gene, moreover, acetic acid formed from aldehyde catalysed by acetaldehyde dehydrogenase (ALD) encoded by the *ALD6* gene. As for the ethanol production, several ADHs, including but not limited to *ADH1*, *ADH2*, *ADH3*, *ADH4*, *ADH5*, *ADH6* and *ADH7*, catalyze the NAD(P)H dependent reduction of acetaldehyde to ethanol.<sup>31</sup> Here, *ADH1* was chosen the target gene to reveal the flux to ethanol when cultured on glucose.<sup>32</sup>

The expression of *PDC1* gene was down-regulated by the treatment of salt stress (Fig. 7A–D).<sup>25</sup> On the other hand, the expression levels of *PYK1* gene increased with increasing salt concentration (Fig. 7E–H). The expression levels of *ADH1* gene decreased after NaCl treatment (Fig. 7I–L), the decreased ADH activity led to enhanced production of glycerol was observed in *C. versatilis*.<sup>31</sup> While expression levels of *ALD6* gene was up-regulated by the treatment of salt stress (Fig. 7M–P), which was consistent with the prior work that Ald6p, a protein related to acetate biosynthesis, is induced under conditions of high osmotic pressure.<sup>33</sup> Moreover, *ALD6* overexpression increases cell growth and ethanol production.<sup>34</sup> Several studies pointed out that the activity of alcohol dehydrogenase decreases during high glycerol production under high osmotic pressure conditions, whereas the activity of aldehyde dehydrogenase (and

probably acetate formation) increases, taking into account the demand of NADH for glycerol formation.<sup>33</sup>

Analysis of the data from *PYK1*, *PDC1*, *ADH1* and *ALD6* genes involved in ethanol metabolism of *C. versatilis* indicated that *PDC1* and *ADH1* genes were transcribed at low levels after salt stress, and *PYK1* and *ALD6* genes at high levels by the treatment of NaCl, which is in accordance with the extra-cellular ethanol detected previously.

Kinetic parameters of salt-tolerant yeast *Candida versatilis* expressing the carbon metabolism enzymes in response to salt stress were summarized in Table 1. For cell growth, final dry cell mass was decreased by 20% and 60% after 2 M and 4 M NaCl treatment as compare to control, respectively.

Glycerol productivity of S3-5 by the treatment of 2 M and 4 M NaCl were 59% and 86% higher than WT. During glucose fermentation, most of the metabolic flux goes to ethanol known as the Crabtree effect, our data showed that improved the concentration of stressing agent could reroute glycolytic flux towards glycerol metabolism.

## Conclusions

Analysis of carbon metabolism and transcriptional variation in response to salt stress in the genome shuffled *C. versatilis* 3-5 and a wild-type salt tolerant yeast strain. The level of glucose consumption, glycerol and ethanol production was affected by salt. The expression level of *HXX1* gene of S3-5 was higher than WT, which contributes to rerouting glycolytic flux towards higher glycolysis. Genome shuffling strains *C. versatilis* 3-5 with improving tolerance to salt might be caused by glycerol accumulation, as the genes involved in glycerol synthesis were found to be highly expressed under these stress conditions, such as up-regulated expression of *GPD1* gene and *HOG1* gene, and down-regulated expression of *FPS1* gene.

## Acknowledgements

This work was supported by grants from National Natural Science Foundation of China (31501449, 2015IM102).

## References

- 1 Y. Zhu and J. Tramper, *Biotechnol. Adv.*, 2013, **31**, 1448–1457.
- 2 C. van der Sluis, J. Tramper and R. H. Wijffels, *Trends Food Sci. Technol.*, 2001, **12**, 322–327.
- 3 M. Silva-Garca and C. Lucas, *FEMS Yeast Res.*, 2003, **3**, 247–260.
- 4 M. Silva-Graca, L. Neves and U. Lucas, *FEMS Yeast Res.*, 2003, **3**, 347–362.
- 5 Y. Watanabe, K. Nagayama and Y. Tama, *Yeast*, 2008, **25**, 107–116.
- 6 W. Qi, Z. C. Fan, C. L. Wang, L. H. Hou, X. H. Wang, J. F. Liu and X. H. Cao, *Eur. Food Res. Technol.*, 2014, **238**, 675–682.
- 7 A. Rodríguez-Navarro and B. Benito, *Biochim. Biophys. Acta, Biomembr.*, 2010, **1798**, 1841–1853.
- 8 Y. Watanabe, M. Yamaguchi, J. Sakamoto and Y. Tamai, *Yeast*, 1993, **9**, 213–220.



- 9 Y. Watanabe, H. Akita, Y. Higuchi, R. Tsujimatsu, T. Kaneta and Y. Tamai, *Biosci., Biotechnol., Biochem.*, 2008, **72**, 1005–1014.
- 10 L. Pribylova, J. de Montigny and H. Sychrova, *Yeast*, 2007, **24**, 171–180.
- 11 M. Lenassi, J. Zajc, C. Gostincar, A. Gorjan, N. Gunde-Cimerman and A. Plemenitas, *Fungal Biol.*, 2011, **115**, 959–970.
- 12 H. B. Kwon, E. T. Yeo, S. E. Hahn, S. C. Bae, D. Y. Kim and M. O. Byun, *FEMS Yeast Res.*, 2003, **3**, 433–440.
- 13 D. Araiza-Olivera, J. G. Sampedro, A. Mujica, A. Pena and S. Uribe-Carvajal, *FEMS Yeast Res.*, 2010, **10**, 282–289.
- 14 T. C. Dakal, L. Solieri and P. Giudici, *Int. J. Food Microbiol.*, 2014, **185**, 140–157.
- 15 J. H. Parmar, S. Bhartiya and K. V. Venkatesh, *Mol. BioSyst.*, 2011, **7**, 1138–1148.
- 16 S. Galafassi, M. Toscano, I. Vigentini, J. Piskur and C. Compagno, *Food Microbiol.*, 2013, **36**, 316–319.
- 17 L. Li, Y. Ye, L. Pan, Y. Zhu, S. Zheng and Y. Lin, *Biochem. Biophys. Res. Commun.*, 2009, **387**, 778–783.
- 18 X. Cao, L. Hou, M. Lu and C. Wang, *Int. J. Food Sci. Technol.*, 2010, **45**, 17–22.
- 19 F. Mo, H. Zhao, H. Lei and M. Zhao, *Appl. Biochem. Biotechnol.*, 2013, **171**, 1339–1350.
- 20 B. S. De Martinis and C. C. S. Martin, *Forensic Sci. Int.*, 2002, **128**, 115–119.
- 21 E. Petelenz-Kurdziel, C. Kuehn, B. Nordlander, D. Klein, K. K. Hong, T. Jacobson, P. Dahl, J. Schaber, J. Nielsen, S. Hohmann and E. Klipp, *PLoS Comput. Biol.*, 2013, **9**, e1003084.
- 22 R. Lagunas, *FEMS Microbiol. Rev.*, 1993, **10**, 229–242.
- 23 N. A. Abdel and E. I. Moghaz, *Adv. Biomed. Res.*, 2010, **1**, 169–176.
- 24 T. You, P. Ingram, M. D. Jacobsen, E. Cook, A. McDonagh, T. Thorne, M. D. Lenardon, A. P. de Moura, M. C. Romano, M. Thiel, M. Stumpf, N. A. Gow, K. Haynes, C. Grebogi, J. Stark and A. J. Brown, *BMC Res. Notes*, 2012, **5**, 258.
- 25 M. J. Taherzadeh, L. Adler and G. Lidén, *Enzyme Microb. Technol.*, 2002, **31**, 53–66.
- 26 F. Lages, M. Silva-Graca and C. Lucas, *Microbiology*, 1999, **145**, 2577–2585.
- 27 E. Vaudano, A. Costantini, O. Noti and E. Garcia-Moruno, *Food Microbiol.*, 2010, **27**, 802–808.
- 28 S. Hohmann, *Microbiol. Mol. Biol. Rev.*, 2002, **66**, 300–372.
- 29 X. M. Tang, G. Kayingo and B. A. Pior, *Yeast*, 2005, **22**, 571–581.
- 30 J. Philips and I. Herskowitz, *J. Cell Biol.*, 1997, **138**, 961–974.
- 31 O. de Smidt, J. C. du Preez and J. Albertyn, *FEMS Yeast Res.*, 2012, **12**, 33–47.
- 32 J. Lian, T. Si, N. U. Nair and H. Zhao, *Metab. Eng.*, 2014, **24**, 139–149.
- 33 T. Hirasawa, K. Yamada, K. Nagahisa, T. N. Dinh, C. Furusawa, Y. Katakura, S. Shioya and H. Shimizu, *Process Biochem.*, 2009, **44**, 647–653.
- 34 S. E. Park, H. M. Koo, Y. K. Park, S. M. Park, J. C. Park, O. K. Lee, Y. C. Park and J. H. Seo, *Bioresour. Technol.*, 2011, **102**, 6033–6038.

