# **RSC Advances**



View Article Online

View Journal | View Issue

# PAPER



Cite this: RSC Adv., 2017, 7, 11233

Received 27th December 2016 Accepted 7th February 2017

DOI: 10.1039/c6ra28744a

rsc.li/rsc-advances

## Introduction

#### The direct carboxylation of alkenes<sup>1,2</sup> is a formidable challenge despite its high potential as a practical method for the preparation of unsaturated carboxylic acid derivatives. A well-known process for carboxylation of alkenes is the transition-metalcatalyzed C-C bond formation with CO<sub>2</sub>, which has become a fundamental synthetic transformation in recent years.<sup>2</sup> Most of those methodologies require the use of rare and expensive metals.3 More recently, Tanaka's group reported that alkenes underwent carboxylation with CO2 in the presence of EtAlCl2 and 2,6-dibromopyridine to afford the corresponding $\alpha,\beta$ - or $\beta,\gamma$ unsaturated carboxylic acids, but the substrate scope was limited to α-arylalkenes and trialkyl-substituted alkenes (Scheme 1).<sup>4</sup> The radical-mediated approach can be a complementary route to the carboxylation of alkenes because radicals have high activity and efficiency.<sup>5</sup> By virtue of this radical carboxylation strategy, several terminal alkenes, which are difficult to carboxylation with CO<sub>2</sub>, have been successfully transformed into the corresponding $\alpha_{\beta}$ unsaturated carboxylic acids.

In the last decade, visible-light driven photoredox catalysis has developed into a versatile tool for organic synthesis that is especially attractive as a sustainable chemical process.<sup>6</sup> This is because solar energy (visible light) is clean and renewable. However, most of the reported such reactions often employ ruthenium or iridium polypyridyl complexes which are expensive and potentially toxic. Therefore, the organic dyes, eosin Y,<sup>7</sup> would be an appropriate alternative to transition metal photocatalysts.<sup>8</sup>

# Carboxylation of styrenes with CBr<sub>4</sub> and DMSO *via* cooperative photoredox and cobalt catalysis<sup>†</sup>

Cai-xia Song,<sup>a</sup> Ping Chen<sup>\*bc</sup> and Yu Tang<sup>\*abc</sup>

Cooperative photoredox and cobalt catalyzed carboxylation of styrenes with  $CBr_4$  to afford the corresponding  $\alpha,\beta$ -unsaturated carboxylic acids has been realized through radical addition and Kornblum (DMSO) oxidation. DMSO serves as the oxidant, oxygen source and solvent under these photocatalytic conditions.



R<sup>1</sup> = aryl R<sup>2</sup> = H, alkyl, aryl

Scheme 1 Carboxylation of styrenes

Traditionally, carboxylation of styrenes<sup>1*a*-*c*</sup> was though addition of CX<sub>4</sub> and acid-catalyzedhydrolysis reaction for two steps. In contrast to traditional methods we design a new method for the synthesis of  $\alpha$ , $\beta$ -unsaturated carboxylic acids directly. We herein report the successful development of a carboxylation of styrenes employing CBr<sub>4</sub> and DMSO *via* cooperative photoredox and cobalt catalysis (CoI<sub>2</sub>)<sup>9</sup> (Scheme 1).

# **Results and discussion**

At first, the carboxylation of alkene was performed using styrene **1a** and  $CBr_4$  with eosin Y disodium (10 mol%) as a photocatalyst under irradiation from a fluorescent bulb (18 W) and under a nitrogen atmosphere for 15 h in DMSO at 50 °C as the model reaction (Table 1).<sup>10</sup> In contrast, Xia group reported the bromination of styrene in the similar photoredox conditions but for 120 hours.<sup>11</sup> Subsequently, several catalysts were evaluated to give rise to target compound **2a** (entries 1–4), of which eosin Y disodium provided the highest yield of the target product **2a** and the intermediate **3** (entry 1). Atom-transfer radical additions of organic halides to alkenes giving the intermediate **3** 

<sup>&</sup>quot;School of Pharmaceutical Science and Technology, Key Laboratory for Modern Drug Delivery & High-Efficiency, Tianjin University, Tianjin, 300072 P. R. China. E-mail: tangyu@ouc.edu.cn

<sup>&</sup>lt;sup>b</sup>Key Laboratory of Marine Drugs, Chinese Ministry of Education, School of Medicine and Pharmacy, Ocean University of China, 5 Yushan Road, Qingdao, P. R. China. E-mail: chenping8315@126.com

<sup>&</sup>lt;sup>c</sup>Laboratory for Marine Drugs and Bioproducts, Qingdao National Laboratory for Marine Science and Technology, Qingdao 266237, P. R. China

<sup>†</sup> Electronic supplementary information (ESI) available. See DOI: 10.1039/c6ra28744a

Table 1 Optimization of visible light driven carboxylation of styrene

+ CX <sub>4</sub> catalyst, additive 1a halide visible light 2a 3								
Entry <sup>a</sup>	Halide	Catalyst	Additive	$2\mathbf{a}^{b}$ (%)	$3^{b}$ (%)			
1	$CBr_4$	C1	None	15	28			
2	$CBr_4$	C3	None	7	19			
3	$CBr_4$	C4	None	11	23			
4	$CBr_4$	C2	None	10	26			
5 <sup>c</sup>	$CBr_4$	C1	None	13	17			
$6^d$	$CBr_4$	C1	None	8	27			
7	BrCCl <sub>3</sub>	C1	None	13	25			
8	$CCl_4$	C1	None	n.r.	n.r.			
9	$CBr_4$	C1	$Co(acac)_2$	n.r.	n.r.			
10	$CBr_4$	C1	$Co(OAc)_2 \cdot 4H_2O$	Trace	n.d.			
11	$CBr_4$	C1	CoCO <sub>3</sub>	n.r.	n.r.			
12	$CBr_4$	C1	$CoF_2$	28	12			
13	$CBr_4$	C1	$CoCI_2$	19	8			
14	$CBr_4$	C1	CoBr <sub>2</sub>	Trace	n.d.			
15	$CBr_4$	C1	CoI <sub>2</sub>	70	19			
16	$CBr_4$	C1	$I_2$	Trace	34			
$17^e$	$CBr_4$	C1	CoI <sub>2</sub>	42	19			
18 <sup>f</sup>	$CBr_4$	C1	CoI <sub>2</sub>	65	15			
$19^g$	$CBr_4$	C1	CoI <sub>2</sub>	42	24			
$20^{h}$	$CBr_4$	C1	CoI <sub>2</sub>	51	39			
21 <sup>i</sup>	CBr <sub>4</sub>	C1	CoI <sub>2</sub>	72	9			
$22^j$	$CBr_4$	C1	CoI <sub>2</sub>	70	12			

<sup>*a*</sup> Reaction conditions: all reactions were carried out with **1a** (0.50 mmol), halide (1.00 mmol), 10 mol% catalyst and additive (0.50 mmol) in solvent (2.00 mL) at 50 °C for 15 h irradiation by 18 W fluorescent lamp under nitrogen atmosphere unless otherwise stated. <sup>*b*</sup> Isolated yield; n.r. = no reaction; n.d. = no detected. <sup>*c*</sup> 5 mol% eosin Y disodium was used. <sup>*d*</sup> 20 mol% eosin Y disodium was used. <sup>*e*</sup> The reaction was conducted at rt. <sup>*f*</sup> The reaction was conducted at 80 °C. <sup>*g*</sup> CoI<sub>2</sub> (0.50 equiv.) was used. <sup>*h*</sup> CoI<sub>2</sub> (2.00 equiv.) was used. <sup>*i*</sup> The reaction was conducted using 1.00 mL DMSO. <sup>*f*</sup> Green LEDs was used for irradiation.



were previously reported by groups of Stephenson,<sup>11*a*-*c*</sup> Reiser<sup>11*d*</sup> and Melchiorre<sup>11*e*</sup>. Different concentrations of eosin Y disodium were tested. Unfortunately, the yield of **2a** could not be improved (entries 5 and 6). We also investigated BrCCl<sub>3</sub> (ref. 11*a* and *b*) but the yield did not increase (entry 7). No conversion could be observed when CCl<sub>4</sub> was used (entry 8).<sup>12</sup> This phenomenon may be attributed to the carbon–chlorine bond more stable (relative to a carbon–bromine bond). Furthermore, we examined different solvent types, such as the protic solvent MeOH; non-polar solvent THF; dipolar solvents CH<sub>3</sub>CN, CH<sub>3</sub>NO<sub>3</sub> and DMF. However, **2a** was not observed in any of the cases (not shown). Cobalt catalysts in various oxidation states have recently attracted particular attention in the development

of applicable transformations due to its economy and low toxicity.<sup>9</sup> Subsequently, different cobalt catalysts were tested, including Co(acac)<sub>2</sub>, Co(OAc)<sub>2</sub>·4H<sub>2</sub>O, CoCO<sub>3</sub>, CoF<sub>2</sub>, CoCI<sub>2</sub>, CoBr<sub>2</sub> and CoI<sub>2</sub> (entries 9–15). These investigations identified CoI<sub>2</sub> as the most efficient catalyst, which provided the desired product **2a** in 70% isolated yield. We also explored the effect of iodide and observed that the addition of I<sub>2</sub> resulted in a low yield (entry 16). However, lower or higher reaction temperature led to lower yields (entries 17–18). Next, we varied the amount of the CoI<sub>2</sub> from 0.50 equiv. to 2.00 equiv. (entries 19–20). These experiments revealed that 1.0 equiv. of CoI<sub>2</sub> gave the best yield (entry 15). In addition, a study of the reaction concentration indicated that increased concentration (*i.e.*, 0.25 mol L<sup>-1</sup> *vs.* 0.50 mol L<sup>-1</sup>) slightly improved the yield (entries 15 *vs.* 21). The use of green LEDs (entry 22) was almost effective as white LEDs.

After determining the optimized conditions for the synthesis of 2a, we examined a diverse range of styrenes (Table 2). Initially, we evaluated the electronic effect of substrates with substituents on the aryl ring. Arylalkenes with methyl substituents at the *m* and *p* positions on the aryl ring furnished the target products in good yields (Table 2, 2c and 2d), while the omethyl substituted arylalkene had a low yield of 45% due to the steric effect (Table 2, 2b). The styrenes bearing the electronwithdrawing groups (F, Cl, and Br) were tolerated and provided moderate yields (Table 2, 2e-2g). For comparison, the yields were well tolerated when the styrenes contained electrondonating substituents such as methoxy and tert-butyl (Table 2, 2h, 2i and 2k). However, when dimethoxy and phenyl groups were introduced to the phenyl ring, the corresponding  $\alpha,\beta$ unsaturated acid products (Table 2, 2j and 2l) were isolated in 53% and 40% yields, respectively. It is noteworthy that 4-(trifluoromethyl)styrene gave the target product in good yield (Table 2, 2m). Nevertheless, the styrenes bearing the nitro and nitrile groups failed to provide the target products (Table 2, 2n and 20). These result indicated strong electron withdrawing groups impaired the reactivity to give the target products. Moreover, we were delighted to find that 1-naphthylalkene could also afford the corresponding  $\alpha$ ,  $\beta$ -unsaturated acid **2p** in vield of 62%.

Next, we turned our attention to exam the scope of arylalkenes with various substituents on the  $\alpha$ - or  $\beta$ -positions under the optimized conditions (Table 3). Unfortunately, the desired product was not obtained when (E)-1,2-diphenylethene was conducted under the optimized reaction conditions. Then we increased the reaction temperature to 100 °C and obtained the target product. In addition, both (Z)-and (E)-1,2-diphenylethene produced product 5a in 15% and 36% yields. The reaction of stilbenes 4b and 4c bearing substituents such as trifluoromethyl and nitro groups furnished the regioselective products 5b and 5c, respectively, in 22% and 32% yields. Moreover, any any lakene with the methyl group on the  $\beta$ -position (4d) furnished the corresponding  $\alpha,\beta$ -unsaturated acid product in a low yield (Table 3, 5d). These results indicated that arylalkenes with various substituents on the  $\beta$ -positions gave low yields which may be because of the steric effect. It is worth mentioning that 1,1-diphenylethylene and indene also proceeded efficiently with yields of 68% and 63% (Table 3, 5e and





<sup>*a*</sup> Alkene (**1a-1p** 0.50 mmol), 10 mol% eosin Y disodium, CBr<sub>4</sub> (1.00 mmol), CoI<sub>2</sub> (0.50 mmol), DMSO (1.00 mL), 50 °C, N<sub>2</sub>. <sup>*b*</sup> Isolated yield. <sup>*c*</sup> Afforded the intermediate 1-nitro-4-(1,3,3,3-tetrabromopropyl) benzene (**3n**) and 4-(1,3,3,3-tetrabromopropyl)benzonitrile (**3o**) in 25% and 35% yields, respectively.

 Table 3
 Carboxylation of arylvinyls<sup>a,b</sup>



<sup>*a*</sup> Alkene (**4a–4f** 0.50 mmol), 10 mol% eosin Y disodium, CBr<sub>4</sub> (1.00 mmol), CoI<sub>2</sub> (0.50 mmol), DMSO (1.00 mL), 100 °C, N<sub>2</sub>, 18 W white LED. <sup>*b*</sup> Isolated yield. <sup>*c*</sup> Using (*Z*)-1,2-diphenylethene. <sup>*d*</sup> Using (*E*)-1,2-diphenylethene. <sup>*e*</sup> Reaction run at 50 °C.

**5f**). Notably, 2-vinylthiophene was also a suitable substrate for the reaction, affording the product 5 g in 58% yield. However, no desired products were observed when other vinyl-substituted heterocycles such as 2-vinylpyrrole, 2-vinylpyridine and 3-vinylindole were used.

Several control experiments were designed to elucidate the plausible reaction mechanism for this carboxylation of styrenes (Scheme 2a). Firstly, the reaction of **1a** and  $CBr_4$  in the absence of eosin Y disodium, visible light or  $CoI_2$  under the standard reaction conditions was investigated. **3a** and **2a** were obtained in moderate yields respectively (Scheme 2a, entries 1 and 2). These results indicated that eosin Y disodium, visible light and  $CoI_2$  were all very important to achieve this reaction. In comparison, 30% yield of **3a** with trace amount of **2a** were detected without eosin Y disodium, which was essential factor in the second step reaction (Scheme 2a, entries 2 *vs.* 3). Moreover, TEMPO (2.00 equiv.), a well-known radical scavenger, was found to inhibit this reaction process (Scheme 2a, entry 4), which suggested that the first step of the reaction may occur through a radical mechanism.

To gain insight into mechanism of the second step reaction, a proposed intermediate **3a** was tested under the standard conditions resulting in target product **2a** in a yield of 76% (Scheme 2b, entry 1), which indicated that **3a** was a possible intermediate for this reaction. Besides, in the absence of eosin Y disodium and light or CoI<sub>2</sub>, the reaction generated **2a** in the yields of 53% and 60% respectively (Scheme 2b, entries 2 and 3), showing that eosin Y disodium and CoI<sub>2</sub> all play important roles in this step of the reaction mechanism. **3a** failed to be transformed into **2a** in the presence of eosin Y disodium, visible light and TEMPO. These results suggested that the mechanism may contain a radical process (Scheme 2b, entries 3 *vs.* 4). In comparison, almost the similar yield of **2a** was generated (Scheme 2a, entries 2, 5 and 6). Addition of TEMPO had no obvious effect on the reaction which suggested that this

a) + $CBr_4 = \frac{eosin Y disodium}{Col_2, DMSO, 50 °C, 15h}$ + $COOH$								
1a		visible light		2a		3a		
ontru <sup>a</sup>	CBr	eosin Y	visible	Col	TEMPO	<b>2</b> a <sup>b</sup>	3a <sup>b</sup>	
entry	CDI4	disodium	light		LIVIFO	(%)	(%)	
1	+	-	-	+	-	47	19	
2	+	+	+	-	-	15	28	
3	+	-	+	-	-	trace	30	
4	+	+	+	+	+	n.r.	n.r.	

<sup>*o*</sup> Reaction conditions: all reactions were carried out with **1a** (0.50 mmol), CBr<sub>4</sub> (1.00 mmol), 10 mol % eosin Y disodium, Col<sub>2</sub> (0.50 mmol) and TEMPO (1.00 mmol) in DMSO (1.00 mL) at 50 °C for 15 h irradiation by 18 W fluorescent lamp under nitrogen atmosphere unless otherwise stated; <sup>*b*</sup> Isolated yield; n.r. = no reaction.

	Br Cl 3a	Br <sub>3</sub> eosin N Col <sub>2</sub> , DMS visib	∕ disodium <mark>O</mark> , 50 ºC, 15h le light	CO 2a	ОН
entry <sup>a</sup>	eosin Y disudium	visible light	$Col_2$	TEMPO	2a (%) <sup>b</sup>
1	+	+	+	-	76
2	-	-	+	-	53
3	+	+	-	-	60
4	+	+	-	+	trace
5	-	-	+	+	51
6	+	+	+	+	47
a n	· · · · · · · · · · · · · · · · · · ·		and a second	1 1 2 (0	20

<sup>*a*</sup> Reaction conditions: all reactions were carried out with **3a** (0.20 mmol), 10 mol % eosin Y disodium, Col<sub>2</sub> (0.20 mmol) and TEMPO (0.80 mmol) in DMSO (1.00 mL) at 50 °C for 15 h irradiation by 18 W fluorescent lamp under nitrogen atmosphere unless otherwise stated; <sup>*b*</sup> Isolated yield.

Scheme 2 Mechanistic studies of cascade reactions.

b)



mechanism may contain a CoI<sub>2</sub>-catalyzed hydrolysis process (Scheme 2b, entries 5 and 6). The above results strongly suggested that this transformation may be achieved *via* cooperative photoredox and CoI<sub>2</sub>.

To further understand the mechanism, <sup>18</sup>O-labeled  $H_2O$  and DMSO experiments were performed between styrene **1a** and CBr<sub>4</sub> under optimized conditions (Scheme 3). When the reaction of **1a** was conducted in the presence of 10.00 equiv. of  $H_2^{18}O$  under the standard conditions, only **2a** and <sup>18</sup>O-**2a** were obtained in 51% yield with the ratio of 26.67 : 1.00 based on MS analysis. And no <sup>18</sup>O<sub>2</sub>-**2a** was observed (Scheme 3a). In contrast, when the reaction was performed with <sup>18</sup>O-labeled DMSO, the three products could also be obtained with the ratio 1.00 : 10.19 : 31.54 (Scheme 3b). These experimental results indicated that DMSO served as the source of oxygen in this transformation.

Based on the above results and previous reports, a plausible mechanism for the carboxylation is shown (Scheme 4). The



Scheme 4 Proposed reaction mechanism.

This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence. Dpen Access Article. Published on 13 February 2017. Downloaded on 8/9/2025 4:30:16 AM.

excited [eosin Y disodium]\* or Co(II) transfers an electron to the halide, generating a radical<sup>13,14</sup> that adds to the alkene under electronic and steric control. The resulting radical combines with the halide with concurrent electron transfer back to [eosin Y disodium]<sup>+</sup> or Co(III), thus, regenerating the catalyst or Co(II).<sup>15</sup> Then, the intermediate **3a** is eliminated, giving compound **C**, which undergoes SET (single electron transfer) reduction to generate the alkyl radical.<sup>16</sup> Oxidation of the alkyl radical by [eosin Y disodium]<sup>+</sup> generates carbocation **E**, accompanied by formation of the catalyst. Nucleophilic attack of DMSO to carbocation **E** affords alkoxysulfonium **F**,<sup>17</sup> which also may be obtained from **3a** by  $CoI_2$ -catalyzed hydrolysis reaction. Then, a reaction similar to Kornblum (DMSO) oxidation proceeds to give compound **G**. Finally, nucleophilic attack by DMSO leads to the desired product **2a**.

## Conclusions

In summary, we have developed a methodology that combines the photoredox/cabolt-catalyzed radical addition of halides onto alkenes and Kornblum (DMSO) oxidation of the dibromosubstituted carbocation, which enabled the transformation of alkenes with a variety of functional groups into the corresponding  $\alpha$ , $\beta$ -unsaturated carboxylic acids. The preliminary mechanistic studies suggested that DMSO as the oxidant, oxygen source and solvent plays a key roles in this reaction.

# Experimental

#### General information

All reactions were carried out under nitrogen and stirred magnetically. DMSO was distilled prior to use. <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on 500 MHz or 600 MHz spectrometers using CDCl<sub>3</sub> with TMS or residual solvent as standard unless otherwise noted. Melting points were determined using a Laboratory Device WRS-3 and are uncorrected/calibrated. TLC analysis was performed using Kangbino glass-backed plates (60 Å, 250  $\mu$ m) and visualized using UV and phosphomolybdic acid stains. High-resolution mass spectra were obtained using an Agilent 1290 Infinity II-6530B in the ESI mode. Low-resolution mass spectra were obtained using an Agilent 1290 Infinity II-6460 in the ESI mode.

#### Preparation of substrates (derivatives of styrene)

2-Methylstyrene (1b).<sup>18</sup> 2-Methylbenzaldehyde (1.00 g, 8.30 mmol, 1.00 equiv.) was added to a solution of methyltriphenylphosphonium bromide (3.5 g, 10.00 mmol, 1.20 equiv.) and K<sub>2</sub>CO<sub>3</sub> (1.80 g, 13.00 mmol, 1.5 equiv.) in dioxane (7.00 mL). The mixture was refluxed at 110 °C over 10 h. After being cooled to room temperature, excess solvent was removed by rotary evaporation. The residue was diluted with petroleum ether and filtered through celite and concentrated under vacuum. The crude product was purified by flash column chromatography (petroleum ether) to obtain the corresponding product as a colorless oil; 0.50 g, 4.23 mmol; 51% yield. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>),  $\delta$  7.51–7.47 (m, 1H), 7.21–7.14 (m, 3H), 6.96 (dd, J = 17.4, 11.0 Hz, 1H), 5.65 (dd, J = 17.4, 1.4 Hz, 1H), 5.30 (dd, J = 11.0, 1.4 Hz, 1H), 2.37 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>),  $\delta$  136.8, 135.4, 134.8, 130.2, 127.6, 126.1, 125.3, 115.1, 19.7.

3-Methylstyrene (1c).<sup>18</sup> *m*-Tolualdehyde (1.00 g, 8.30 mmol, 1.00 equiv.) was added to a solution of methyltriphenylphosphonium bromide (3.57 g, 10.00 mmol, 1.20 equiv.) and K<sub>2</sub>CO<sub>3</sub> (1.70 g, 13.00 mmol, 1.5 equiv.) in dioxane (7.00 mL). The mixture was refluxed at 110 °C over 10 h. After being cooled to room temperature, excess solvent was removed by rotary evaporation. The residue was diluted with petroleum ether and filtered through celite and concentrated under vacuum. The crude product was purified by flash column chromatography (petroleum ether) to obtain the corresponding product as a colorless oil; 0.80 g, 6.80 mmol; 82% yield. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.27–7.23 (m, 3H), 7.12–7.10 (m, 1H), 6.73 (dd, *J* = 17.6, 10.9 Hz, 1H), 5.78 (d, *J* = 17.6 Hz, 1H), 5.25 (d, *J* = 10.9 Hz, 1H), 2.38 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  138.1, 137.5, 137.0, 128.6, 128.4, 126.9, 123.4, 113.6, 21.4.

3-Methoxystyrene (1i).<sup>19</sup> 3-Methoxybenzaldehyde (1.00 g, 7.40 mmol, 1.00 equiv.) was added to a solution of methyltriphenylphosphonium bromide (3.20 g, 8.90 mmol, 1.20 equiv.) and K<sub>2</sub>CO<sub>3</sub> (1.60 g, 11.80 mmol, 1.60 equiv.) in THF (10.00 mL). The mixture was refluxed at 75 °C over 10 h. After being cooled to room temperature, excess solvent was removed by rotary evaporation. The residue was diluted with petroleum ether and filtered through celite and concentrated under vacuum. The crude product was purified by flash column chromatography (petroleum ether) to obtain the corresponding product as a colorless oil; 0.79 g, 6.00 mmol; 81% yield. <sup>1</sup>H NMR  $(500 \text{ MHz}, \text{CDCl}_3), \delta$  7.27 (t, J = 7.9 Hz, 1H), 7.03 (m, 1H), 6.98 (s, 100 Hz)1H), 6.86–6.83 (m, 1H), 6.72 (dd, *J* = 17.6, 10.9 Hz, 1H), 5.77 (d, J = 17.6 Hz, 1H), 5.28 (d, J = 10.9 Hz, 1H), 3.84 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>), δ 159.8, 139.0, 136.8, 129.5, 118.9, 114.1, 113.4, 111.5, 55.2.

3,4-(Dimethoxy)styrene (1j).<sup>19</sup> 3,4-Dimethoxybenzaldehyde (1.00 g, 6.00 mmol, 1.00 equiv.) was added to a solution of methyltriphenylphosphonium bromide (2.60 g, 7.20 mmol, 1.20 equiv.) and K<sub>2</sub>CO<sub>3</sub> (1.20 g, 9.00 mmol, 1.50 equiv.) in THF (10.00 mL). The mixture was refluxed at 75 °C over 10 h. After being cooled to room temperature, excess solvent was removed by rotary evaporation. The residue was diluted with petroleum ether and filtered through celite and concentrated under vacuum. The crude product was purified by flash column chromatography (petroleum ether) to obtain the corresponding product as a colorless oil; 0.83 g, 4.98 mmol; 83% yield. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>),  $\delta$  6.99–6.94 (m, 2H), 6.83 (d, J = 8.2 Hz, 1H), 6.66 (dd, J = 17.5, 10.9 Hz, 1H), 5.62 (d, J = 17.5 Hz, 1H), 5.16 (d, J = 10.9 Hz, 1H), 3.91 (s, 3H), 3.89 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>), *δ* 149.0, 149.0, 136.5, 130.7, 119.4, 1118, 111.0, 108.5, 55.9, 55.8.

**4-Phenylstyrene (11)**.<sup>18</sup> 4-Biphenylcarboxaldehyde (1.00 g, 5.49 mmol, 1.00 equiv.) was added to a solution of methyltriphenylphosphonium bromide (2.40 g, 6.70 mmol, 1.20 equiv.) and  $K_2CO_3$  (1.20 g, 8.77 mmol, 1.50 equiv.) in THF (7.00 mL). The mixture was refluxed at 75 °C over 10 h. After being cooled to room temperature, excess solvent was removed by rotary evaporation. The residue was diluted with petroleum ether and filtered through celite and concentrated under vacuum. The crude product was purified by flash column chromatography (petroleum ether) to obtain the corresponding product as a white solid; 0.60 g, 3.34 mmol; 61% yield; mp 119–120 °C (Lit. 120–122 °C), <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>),  $\delta$  7.63–7.57 (m, 4H), 7.50 (d, *J* = 8.2 Hz, 2H), 7.45 (t, *J* = 7.5 Hz, 2H), 7.36 (t, *J* = 7.5 Hz, 1H), 6.78 (dd, *J* = 17.6, 10.9 Hz, 1H), 5.81 (d, *J* = 17.6 Hz, 1H), 5.29 (d, *J* = 11.0 Hz, 1H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>),  $\delta$  140.7, 140.6, 136.6, 136.4, 128.8, 127.3, 127.2, 127.0, 126.6, 113.9.

4-(Trifluoromethyl)styrene (1m).18 4-(Trifluoromethyl)benzaldehyde (1.00 g, 5.70 mmol, 1.00 equiv.) was added to a solution of methyltriphenylphosphonium bromide (2.40 g, 6.84 mmol, 1.20 equiv.) and K<sub>2</sub>CO<sub>3</sub> (1.18 g, 8.55 mmol, 1.50 equiv.) in THF (8.00 mL). The mixture was refluxed at 75 °C over 10 h. After being cooled to room temperature, excess solvent was removed by rotary evaporation. The residue was diluted with petroleum ether and filtered through celite and concentrated under vacuum. The crude product was purified by flash column chromatography (petroleum ether) to obtain the corresponding product as a colorless oil; 0.66 g, 3.88 mmol; 67% yield. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>),  $\delta$  7.59 (d, J = 8.2 Hz, 2H), 7.51 (d, J = 8.1 Hz, 2H), 6.76 (dd, J = 17.6, 10.9 Hz, 1H), 5.86 (d, J = 17.6 Hz, 1H), 5.40 (d, J = 10.9 Hz, 1H); <sup>13</sup>C NMR (125 MHz,  $CDCl_3$ ),  $\delta$  140.9, 135.6, 129.8 (q, J = 32.3 Hz), 126.4, 125.5 (q, J =3.8 Hz), 123.1, 116.4; <sup>19</sup>F NMR (470 MHz,  $CDCl_3$ )  $\delta - 62.5$  (s, 3F)

**4-Nitrostyrene** (10).<sup>18</sup> 4-Nitrobenzaldehyde (1.00 g, 6.60 mmol, 1.00 equiv.) was added to a solution of methyltriphenylphosphonium bromide (2.80 g, 7.90 mmol, 1.20 equiv.) and K<sub>2</sub>CO<sub>3</sub> (1.40 g, 9.90 mmol, 1.50 equiv.) in THF (10.00 mL). The mixture was refluxed at 75 °C over 10 h. After being cooled to room temperature, excess solvent was removed by rotary evaporation. The residue was diluted with petroleum ether and filtered through celite and concentrated under vacuum. The crude product was purified by flash column chromatography (petroleum ether) to obtain the corresponding product as a yellow oil; 0.65 g, 4.36 mmol; 66% yield. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>),  $\delta$  8.16 (d, *J* = 8.5 Hz, 2H), 7.52 (d, *J* = 8.7 Hz, 2H), 6.77 (dd, *J* = 17.6, 10.9 Hz, 1H), 5.92 (d, *J* = 17.6 Hz, 1H), 5.49 (d, *J* = 10.9 Hz, 1H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>),  $\delta$  147.1, 143.8, 134.9, 126.8, 123.9, 118.6.

**4-Cyanostyrene** (1p).<sup>18</sup> 4-Cyanobenzaldehyde (0.50 g, 3.82 mmol, 1.00 equiv.) was added to a solution of methyltriphenylphosphonium bromide (1.60 g, 4.58 mmol, 1.20 equiv.) and K<sub>2</sub>CO<sub>3</sub> (0.83 g, 6.04 mmol, 1.50 equiv.) in dioxane (4.00 mL). The mixture was refluxed at 110 °C over 10 h. After being cooled to room temperature, excess solvent was removed by rotary evaporation. The residue was diluted with petroleum ether and filtered through celite and concentrated under vacuum. The crude product was purified by flash column chromatography (petroleum ether) to obtain the corresponding product as a colorless oil; 0.48 g, 3.70 mmol; 98% yield. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>),  $\delta$  7.62 (d, J = 8.2 Hz, 2H), 7.49 (d, J = 8.2 Hz, 2H), 6.73 (dd, J = 17.6, 10.9 Hz, 1H), 5.88 (d, J = 17.3 Hz, 1H), 5.45 (d, J = 10.9 Hz, 1H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>),  $\delta$  141.9, 135.4, 132.4, 126.7, 118.9, 117.7, 111.1.

2-Vinylnaphthalene (1p).<sup>20</sup> 2-Naphthaldehyde (1.00 g, 6.40 mmol, 1.00 equiv.) was added to a solution of methyltriphenylphosphonium bromide (2.70 g, 7.69 mmol, 1.20 equiv.) and K<sub>2</sub>CO<sub>3</sub> (1.30 g, 9.60 mmol, 1.50 equiv.) in THF (12.00 mL). The mixture was refluxed at 75 °C over 10 h. After being cooled to room temperature, excess solvent was removed by rotary evaporation. The residue was diluted with petroleum ether and filtered through celite and concentrated under vacuum. The crude product was purified by flash column chromatography (petroleum ether) to obtain the corresponding product as a white solid; 0.96 g, 6.20 mmol; 97% yield; mp 63-64 °C (Lit. 62–66 °C), <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>), δ 7.85–7.80 (m, 3H), 7.77 (s, 1H), 7.69-7.64 (m, 1H), 7.51-7.44 (m, 2H), 6.91 (ddd, J = 17.6, 10.9, 2.0 Hz, 1H), 5.90 (dd, J = 17.6, 2.0 Hz, 1H), 5.36 (dd, J = 10.8, 2.0 Hz, 1H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>), δ 136.9, 135.0, 133.6, 133.2, 128.2, 128.0, 127.7, 126.4, 126.2, 125.2, 123.2, 114.2.

(E/Z)-1-Styryl-4-(trifluoromethyl)benzene (4b/4b').<sup>18</sup> A solution of benzyltriphenylphosphonium bromide (1.52 g, 3.50 mmol, 1.20 equiv.) and LiOH · H<sub>2</sub>O (0.36 g, 8.60 mmol, 3.00 equiv.) in isopropanol (10.00 mL) was stirred at room temperature for 30 min, and then 4-(trifluoromethyl)benzaldehyde (0.50 g, 2.87 mmol, 1.00 equiv.) was added. The reaction mixture was stirred at 75 °C for 6 h. After being cooled to room temperature, excess solvent was removed by rotary evaporation. The residue was quenched with water and extracted with ethyl acetate and the combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under reduced pressure. The crude product was purified by flash column chromatography (petroleum ether) to obtain the corresponding product (E/Z)38/62) as a white solid; 0.64 g, 2.58 mmol; 90% yield. <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>, signals corresponding to (*E*)-isomer)  $\delta$  7.60 (s, 4H), 7.46 (d, J = 8.1 Hz, 1H), 7.38 (t, J = 7.6 Hz, 2H), 7.30 (t, J = 7.4 Hz, 1H), 7.19 (d, J = 16.2 Hz, 1H), 7.12 (d, J = 16.3 Hz, 1H); <sup>13</sup>C NMR (150 MHz, CDCl<sub>3</sub>, signals corresponding to (E)isomer), δ 140.8, 136.6, 131.2, 129.4, 128.3, 128.8, 127.1, 126.8, 126.6, 125.7 (q, J = 3.8 Hz), 123.3; <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>, representative signals corresponding to (Z)-isomer),  $\delta$  7.46 (d, J = 8.1 Hz, 2H), 7.33 (d, J = 8.1 Hz, 2H), 7.25–7.21 (m, 5H), 6.72 (d, J = 12.2 Hz, 1H), 6.59 (d, J = 12.2 Hz, 1H); <sup>13</sup>C NMR (150 MHz,  $CDCl_3$ , representative signals corresponding to (Z)isomer) & 140.9, 136.7, 132.3, 129.2, 128.8, 128.8, 128.4, 127.6, 125.2 (q, J = 3.7 Hz), 123.3, 121.5.

(*E*)-1-Nitro-4-styrylbenzene (4c).<sup>18</sup> A solution of benzyltriphenylphosphonium bromide (5.20 g, 12.01 mmol, 1.20 equiv.) and LiOH·H<sub>2</sub>O (1.24 g, 29.76 mmol, 3.00 equiv.) in 2-propanol (40.00 mL) was stirred at room temperature for 30 min, and then 4-nitrobenzaldehyde (1.50 g, 9.93 mmol, 1.00 equiv.) was added. The reaction mixture was stirred at 75 °C for 6 h. After being cooled to room temperature, excess solvent was removed by rotary evaporation. The residue was quenched with water and extracted with ethyl acetate and the combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under reduced pressure. The crude product was purified by flash column chromatography (petroleum ether) to obtain the corresponding product: (*E*)-isomer; yellow solid; 1.16 g, 5.10 mmol; 52% yield; mp 154–155 °C (Lit. 155–158 °C), <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>),  $\delta$  8.22 (d, J = 8.6 Hz, 2H), 7.64 (d, J = 8.6 Hz, 2H), 7.55 (d, J = 7.6 Hz, 2H), 7.40 (t, J = 7.6 Hz, 2H), 7.34 (t, J = 7.3 Hz, 1H), 7.27 (d, J = 16.5 Hz, 1H), 7.15 (d, J = 16.3 Hz, 1H), <sup>13</sup>C NMR (150 MHz, CDCl<sub>3</sub>), 146.8, 143.9, 136.2, 133.3, 128.9, 128.9, 127.0, 126.9, 126.3, 124.2.

(Z/E)- $\beta$ -Methylstyrene (4d/4d').<sup>18</sup> Benzaldehyde (0.50 g, 4.72 mmol, 1.00 equiv.) was added to a solution of ethyltriphenylphosphonium bromide (2.00 g, 5.39 mmol, 1.14 equiv.) and K<sub>2</sub>CO<sub>3</sub> (1.00 g, 7.25 mmol, 1.50 equiv.) in dioxane (4.00 mL). The mixture was refluxed at 110 °C over 10 h. After being cooled to room temperature, excess solvent was removed by rotary evaporation. The residue was diluted with petroleum ether and filtered through celite and concentrated under vacuum. The crude product was purified by flash column chromatography (petroleum ether) to obtain the corresponding product (E/Z = 44/56) as a colorless oil; 0.33 g, 3.00 mmol; 65% yield. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>, signals corresponding to (Z)isomer) & 7.41-7.27 (m, 4H), 7.27-7.20 (m, 1H), 6.51-6.43 (m, 1H), 5.88–5.81 (m, 1H), 1.95 (dd, J = 7.2, 1.8 Hz, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>, signals corresponding to (Z)-isomer)  $\delta$  137.7, 129.9, 128.9, 128.1, 126.8, 126.4, 14.6; <sup>1</sup>H NMR (500 MHz,  $CDCl_3$ , representative signals corresponding to (E)-isomer) δ 7.41-7.27 (m, 4H), 7.27-7.20 (m, 1H), 6.51-6.43 (m, 1H), 6.32-6.25 (m, 1H), 1.93 (dd, J = 6.6, 1.6 Hz, 3H); <sup>13</sup>C NMR (125 MHz,  $CDCl_3$ , representative signals corresponding to (*E*)-isomer) δ 138.0, 131.1, 128.5, 123.8, 125.9, 125.7, 18.5.

**2-Vinylthiophene (4g).**<sup>21</sup> Thiophene-2-carbaldehyde (1.00 g, 8.92 mmol, 1.00 equiv.) was added to a solution of ethyltriphenylphosphonium bromide (3.80 g, 10.70 mmol, 1.20 equiv.) and K<sub>2</sub>CO<sub>3</sub> (1.85 g, 13.38 mmol, 1.50 equiv.) in dioxane (10.00 mL). The mixture was refluxed at 110 °C over 10 h. After being cooled to room temperature, excess solvent was removed by rotary evaporation. The residue was diluted with petroleum ether and filtered through celite and concentrated under vacuum. The crude product was purified by flash column chromatography (petroleum ether) to obtain the corresponding product as a colorless oil; 0.59 g, 5.35 mmol; 60% yield. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>),  $\delta$  7.19 (d, *J* = 5.0 Hz, 1H), 7.00 (m, 2H), 6.85 (dd, *J* = 17.5, 11.0 Hz, 1H), 5.60 (d, *J* = 12.5 Hz, 1H), 5.17 (d, *J* = 10.5 Hz, 1H), <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>),  $\delta$  143.1, 129.9, 127.3, 125.8, 124.3, 113.2.

**2-Vinylpyrrole** (**4h**).<sup>21</sup> Pyrrole-2-carbaldehyde (1.00 g, 10.52 mmol, 1.00 equiv.) was added to a solution of ethyltriphenylphosphonium bromide (4.51 g, 12.63 mmol, 1.20 equiv.) and K<sub>2</sub>CO<sub>3</sub> (2.18 g, 15.78 mmol, 1.50 equiv.) in dioxane (10.00 mL). The mixture was refluxed at 110 °C over 10 h. After being cooled to room temperature, excess solvent was removed by rotary evaporation. The residue was diluted with 9 : 1 petroleum ether/ethyl acetate and filtered through celite and concentrated under vacuum. The crude product was purified by flash column chromatography (19 : 1 petroleum ether/EtOAc) to obtain the corresponding product as a colorless oil; 0.55 g, 5.79 mmol; 55% yield. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>),  $\delta$  8.24 (s, 1H), 6.78 (s, 1H), 6.61 (dd, *J* = 18.0, 11.0 Hz, 1H), 6.32–6.23 (m, 2H), 5.31 (d, *J* = 17.5 Hz, 1H), 5.05 (d, *J* = 11.5 Hz, 1H), <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>),  $\delta$  130.9, 127.2, 118.8, 109.5, 108.0.

This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.

#### Paper

3-Vinylindole (4j).<sup>22</sup> To a stirred suspension of methyltriphenylphosphonium bromide (2.96 g, 8.30 mmol, 1.2 equiv.) in THF (20 mL), cooled to -50 °C, *n*-BuLi (2.90 mL, 2.4 M in hexanes, 6.90 mmol, 1.00 equiv.) was slowly added. The resulting yellow suspension was stirred and allowed to warm to 0 °C in approximately 45 minutes. After cooling to -30 °C, a premixed solution of an indole 3-carboxaldehyde (6.90 mmol, 1 equiv.) and LiHMDS (6.90 mL, 1.0 M in THF, 6.90 mmol, 1.00 equiv.), in THF (8.00 mL) was added. The resulting suspension was then stirred at room temperature for 1 h, then poured onto  $H_2O$  and extracted with EtOAc (2×). The combined organic phases were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated. The crude residue was then purified by a short chromatography on silica gel (petroleum ether/ethyl acetate 9:1) to obtain the corresponding product as a white solid; 0.90 g, 6.28 mmol; 91% yield; mp 84-86 °C (Lit. 80-81 °C), <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>),  $\delta$  11.19 (s, 1H), 7.80 (d, J = 8.5 Hz, 1H), 7.48 (s, 1H), 7.38 (d, J =8.1 Hz, 1H), 7.12 (t, J = 7.0 Hz, 1H), 7.05 (t, J = 8.0 Hz, 1H), 6.86 (dd, J = 18.0, 11.5 Hz, 1H), 5.61 (d, J = 17.5 Hz, 1H), 5.04 (d, J = 11.5 Hz, 1H). <sup>13</sup>C NMR (126 MHz, DMSO- $d_6$ ),  $\delta$  137.3, 130.7, 125.9, 125.6, 120.0, 119.97, 119.95, 114.3, 112.2, 109.4.

#### Carboxylation of alkenes

Procedures for the carboxylation of arylvinyls. Styrene (52 mg, 0.50 mmol, 1.00 equiv.), CBr<sub>4</sub> (331 mg, 1.00 mmol, 2.00 equiv.), 10 mol% eosin Y disodium (31 mg, 0.05 mmol) and CoI<sub>2</sub> (156 mg, 0.50 mmol, 1.00 equiv.) and 1 mL DMSO were combined in dried flask. The solution was stirred at 50 °C for 15 h on the condition of visible light (18 W fluorescent lamp) under nitrogen. After cooling to room temperature, the reaction mixture was extracted with ethylacetate (20.00 mL × 3) and then washed with brine (20.00 mL). The organic layer was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> followed by filtration and then condensation. The residue was purified by silica gel column chromatography to afford corresponding product **2a** and intermediate **3a**.

**Cinnamic acid (2a).**<sup>23</sup> Following the general procedure: white solid; 53 mg, 0.36 mmol; 72% yield; mp 134–136 °C (Lit. 133–135 °C);  $R_{\rm f} = 0.51$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01); <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ ),  $\delta$  12.39 (brs, 1H), 7.67–7.64 (m, 2H), 7.59 (d, J = 16.0 Hz, 1H), 7.42–7.37 (m, 3H), 6.52 (d, J = 16.0 Hz, 1H); <sup>13</sup>C NMR (125 MHz, DMSO- $d_6$ ),  $\delta$  168.1, 144.4, 134.7, 130.7, 129.4, 128.7, 119.7; MS (ESI): m/z (% relative intensity) = 147.0 (100) (M – H)<sup>-</sup>.

(*E*)-3-(*o*-Tolyl)acrylic acid (2b).<sup>24</sup> Following the general procedure using 2-methylstyrene in place of styrene: white solid; 62 mg, 0.23 mmol; 45% yield; mp 184–185 °C (Lit. 175–176 °C);  $R_{\rm f} = 0.46$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  8.10 (d, J = 15.6 Hz, 1H), 7.59 (d, J = 7.8 Hz, 1H), 7.31 (t, J = 7.8 Hz, 1H), 7.26–7.22 (m, 2H), 6.39 (d, J = 16.2 Hz, 1H), 2.46 (s, 3H); <sup>13</sup>C NMR (150 MHz, CDCl<sub>3</sub>)  $\delta$  172.3, 144.7, 138.0, 133.0, 130.9, 130.5, 126.65, 126.5, 118.2, 19.8; MS (ESI): m/z (% relative intensity) = 161.1 (100) (M – H)<sup>-</sup>.

(*E*)-3-(*m*-Tolyl)acrylic acid (2c).<sup>25</sup> Following the general procedure using 3-methylstyrene in place of styrene: white solid; 102 mg, 0.37 mmol; 74% yield; mp 115–117  $^{\circ}$ C (Lit.

116–118 °C);  $R_{\rm f} = 0.42$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>),  $\delta$  7.77 (d, J = 15.6 Hz, 1H), 7.36 (d, J = 7.2 Hz, 2H), 7.30 (t, J = 7.8 Hz, 1H), 7.23 (d, J = 7.2 Hz, 1H), 6.45 (d, J = 16.2 Hz, 1H), 2.38 (s, 3H); <sup>13</sup>C NMR (150 MHz, CDCl<sub>3</sub>),  $\delta$  172.6, 147.3, 138.7, 134.0, 131.6, 129.0, 128.9, 125.6, 117.1, 21.3; MS (ESI): m/z (% relative intensity) = 161.0 (100) (M – H)<sup>-</sup>.

(*E*)-3-(*p*-Tolyl)acrylic acid (2d).<sup>24</sup> Following the general procedure using 4-methylstyrene in place of styrene: white solid; 92 mg, 0.34 mmol; 67% yield; mp 190–192 °C (Lit. 199–200 °C);  $R_{\rm f} = 0.45$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01); <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ ),  $\delta$  7.56–7.52 (m, 3H), 7.20 (d, J = 8.0 Hz, 2H), 6.44 (d, J = 16.0 Hz, 1H), 2.30 (s, 3H); <sup>13</sup>C NMR (125 MHz, DMSO- $d_6$ ),  $\delta$  168.1, 144.4, 140.6, 131.9, 129.9, 128.6, 118.5, 21.4; MS (ESI): m/z (% relative intensity) = 161.1 (100) (M – H)<sup>-</sup>.

(*E*)-3-(4-Fluorophenyl)acrylic acid (2e).<sup>24</sup> Following the general procedure using 4-fluorostyrene in place of styrene: white solid; 38 mg, 0.23 mmol; 46% yield; mp 198–200 °C (Lit. 205–207 °C);  $R_{\rm f} = 0.41$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01); <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.38 (brs, 1H), 7.78–7.73 (m, 2H), 7.58 (d, J = 16.0 Hz, 1H), 7.24 (t, J = 9.0 Hz, 2H), 6.49 (d, J = 16.0 Hz, 1H); <sup>13</sup>C NMR (125 MHz, DMSO- $d_6$ ),  $\delta$  168.0, 163.6 (d,  ${}^2J_{\rm CF} = 246.3$  Hz), 143.1, 131.3 (d, J = 3.1 Hz), 131.0 (d, J = 8.5 Hz), 119.6 (d, J = 2.4 Hz), 116.3 (d, J = 21.2 Hz); <sup>19</sup>F NMR (470 MHz, DMSO- $d_6$ ),  $\delta - 110.44$  (m, 1F); MS (ESI): m/z (% relative intensity) = 165.0 (100) (M – H)<sup>-</sup>.

(*E*)-3-(4-Chlorophenyl)acrylic acid (2f).<sup>25</sup> Following the general procedure using 1-chlorostyrene in place of styrene: white solid; 56 mg, 0.30 mmol; 60% yield; mp 241–243 °C (Lit. 247–249 °C);  $R_{\rm f} = 0.38$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01); <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>),  $\delta$  12.50 (brs, 1H), 7.76 (d, J = 8.5 Hz, 2H), 7.62 (d, J = 16.0 Hz, 1H), 7.51 (d, J = 8.5 Hz, 2H), 6.59 (d, J = 16.0 Hz, 1H); <sup>13</sup>C NMR (125 MHz, DMSO-*d*<sub>6</sub>),  $\delta$  167.8, 143.0, 135.1, 133.7, 130.4, 129.4, 120.5; MS (ESI): *m/z* (% relative intensity) = 181.0 (100) (M – H)<sup>-</sup>.

(*E*)-3-(4-Bromophenyl)acrylic acid (2g).<sup>26</sup> Following the general procedure using 4-bromostyrene in place of styrene: white solid; 56 mg, 0.30 mmol; 60% yield; mp 255–256 °C (Lit. 264–265 °C);  $R_{\rm f} = 0.43$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01); <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ ),  $\delta$  7.65 (d, J = 8.5 Hz, 2H), 7.60 (d, J = 8.5 Hz, 2H), 7.55 (d, J = 16.0 Hz, 1H), 6.55 (d, J = 16.0 Hz, 1H); <sup>13</sup>C NMR (125 MHz, DMSO- $d_6$ ),  $\delta$  167.8, 143.1, 134.0, 132.3, 130.6, 124.0, 120.6; MS (ESI): m/z (% relative intensity) = 226.9 (100) (M – H)<sup>-</sup>

(*E*)-3-(4-Methoxyphenyl)acrylic acid (2h).<sup>24</sup> Following the general procedure using 4-methoxy styrene in place of styrene: white solid; 68 mg, 0.38 mmol; 76% yield; mp 170–171 °C (Lit. 175–176 °C);  $R_{\rm f} = 0.40$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01), <sup>1</sup>H NMR (600 MHz, DMSO- $d_6$ )  $\delta$  12.23 (brs, 1H), 7.64 (d, J = 9.0 Hz, 2H), 7.55 (d, J = 16.2 Hz, 1H), 6.97 (d, J = 8.4 Hz, 2H), 6.38 (d, J = 16.2 Hz, 1H), 3.80 (s, 3H); <sup>13</sup>C NMR (150 MHz, DMSO- $d_6$ ),  $\delta$  167.8, 160.9, 143.7, 129.9, 126.8, 116.5, 114.3, 55.3; MS (ESI): m/z (% relative intensity) = 177.0 (100) (M – H)<sup>-</sup>.

(*E*)-3-(3-Methoxyphenyl)acrylic acid (2i).<sup>25</sup> Following the general procedure using 3-methoxystyrene in place of styrene: white solid; 60 mg, 0.34 mmol; 67% yield; mp 110–112  $^{\circ}$ C (Lit.

117–119 °C);  $R_{\rm f} = 0.43$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>),  $\delta$  7.77 (d, J = 16.2 Hz, 1H), 7.32 (t, J = 7.8 Hz, 1H), 7.15 (d, J = 7.2 Hz, 1H), 7.07 (s, 1H), 6.97 (dd, J = 7.8, 1.8 Hz, 1H), 6.45 (d, J = 15.6 Hz, 1H), 3.84 (s, 3H); <sup>13</sup>C NMR (150 MHz, CDCl<sub>3</sub>),  $\delta$  172.3, 159.9, 147.1, 135.4, 130.0, 121.1, 117.6, 116.7, 113.2, 55.4; MS (ESI): m/z (% relative intensity) = 177.0 (100) (M – H)<sup>-</sup>.

(*E*)-3-(3,4-Dimethoxyphenyl)acrylic acid (2j).<sup>27</sup> Following the general procedure using 3,4-dimethoxytyrene in place of styrene: white solid; 55 mg, 0.27 mmol; 53% yield; mp 189–191 °C (Lit. 191 °C);  $R_{\rm f} = 0.34$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>),  $\delta$  7.73 (d, J = 16.0 Hz, 1H), 7.15 (dd, J = 8.0, 1.5 Hz, 1H), 7.08 (d, J = 1.5 Hz, 1H), 6.89 (d, J = 8.5 Hz, 1H), 6.33 (d, J = 16.0 Hz, 1H), 3.93 (s, 6H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>),  $\delta$  172.2, 151.5, 149.3, 147.0, 127.0, 123.1, 114.8, 111.0, 109.8, 56.0, 55.9; MS (ESI): m/z (% relative intensity) = 206.9 (100) (M – H)<sup>-</sup>.

(*E*)-3-(4-(*tert*-Butyl)phenyl)acrylic acid (2k).<sup>26</sup> Following the general procedure using 4-(*tert*-butyl)styrene in place of styrene: white solid; 83 mg, 0.34 mmol; 68% yield; mp 202–203 °C (Lit. 202–204 °C);  $R_{\rm f} = 0.46$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>),  $\delta$  7.79 (d, J = 15.9 Hz, 1H), 7.50 (d, J = 8.2 Hz, 2H), 7.43 (d, J = 8.2 Hz, 2H), 6.43 (d, J = 15.9 Hz, 1H), 1.33 (s, 9H); <sup>13</sup>C NMR (150 MHz, CDCl<sub>3</sub>),  $\delta$  172.7, 154.4, 147.0, 131.3, 128.3, 126.0, 116.4, 35.0, 31.2; MS (ESI): m/z (% relative intensity) = 203.0 (100) (M – H)<sup>-</sup>.

(*E*)-3-([1,1'-Biphenyl]-4-yl)acrylic acid (2l). Following the general procedure using 4-phenylstyrene in place of styrene: white solid; 50 mg, 0.20 mmol; 40% yield; mp 215–217 °C;  $R_f = 0.39$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01); <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ ),  $\delta$  7.77 (d, J = 8.0 Hz, 2H), 7.73–7.69 (m, 4H), 7.63 (d, J = 16.0 Hz, 1H), 7.47 (t, J = 7.5 Hz, 2H), 7.38 (t, J = 8.0 Hz, 1H), 6.57 (d, J = 16.0 Hz, 1H); <sup>13</sup>C NMR (125 MHz, DMSO- $d_6$ ),  $\delta$  168.0, 143.9, 142.2, 139.7, 133.8, 129.5, 129.3, 128.4, 127.5, 127.1, 119.6; MS (ESI): m/z (% relative intensity) = 223.0 (100) (M – H)<sup>-</sup>, HRMS (ESI): m/z calcd for C<sub>15</sub>H<sub>12</sub>O<sub>2</sub> (M – H)<sup>-</sup> 223.0765, found 223.0764.

(*E*)-3-(4-(Trifluoromethyl)phenyl)acrylic acid (2m).<sup>28</sup> Following the general procedure using 4-trifluoromethylstyrene in place of styrene: white solid; 66 mg, 0.31 mmol; 61% yield; mp 206–208 °C;  $R_{\rm f} = 0.43$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01); <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ ),  $\delta$  12.59 (brs, 1H), 7.91 (d, J = 8.0 Hz, 2H), 7.75 (d, J = 8.5 Hz, 2H), 7.65 (d, J = 16.0 Hz, 1H); 6.67 (d, J = 16.0 Hz, 1H); <sup>13</sup>C NMR (125 MHz, DMSO- $d_6$ ),  $\delta$  167.6, 142.5, 138.7, 130.3 (q, J = 31.7), 129.3, 126.1 (q, J = 3.7 Hz), 124.5 (d, J = 271.3 Hz), 122.6; <sup>19</sup>F NMR (470 MHz, DMSO- $d_6$ ),  $\delta - 61.28$  (s, 3F); MS (ESI): m/z (% relative intensity) = 215.0 (100) (M - H)<sup>-</sup>.

(*E*)-3-(Naphthalen-2-yl)acrylic acid (2p). Following the general procedure using 2-vinylnaphthalene in place of styrene: white solid; 61 mg, 0.31 mmol; 62% yield; mp 203–204 °C;  $R_f$  = 0.41 (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01); <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ ),  $\delta$  12.44 (brs, 1H), 8.17 (s, 1H), 7.96–7.91 (m, 3H), 7.88–7.84 (m, 1H), 7.74 (d, J = 16.0 Hz, 1H), 7.58–7.53 (m, 2H), 6.65 (d, J = 15.5 Hz, 1H); <sup>13</sup>C NMR (125 MHz, DMSO- $d_6$ ),  $\delta$  168.1, 144.4, 134.1, 133.3, 132.3, 130.1, 129.0, 128.9, 128.1, 127.7, 127.2, 124.4, 120.0; MS (ESI): m/z (% relative intensity) =

197.0 (100) (M – H)<sup>–</sup>; HRMS (ESI): m/z calcd for  $C_{13}H_{10}O_2$  (M – H)<sup>–</sup> 197.0608, found 197.0605.

(1,3,3,3-Tetrabromopropyl)benzene (3a).<sup>29</sup> Following the general procedure: white solid; 19 mg, 0.05 mmol; 9% yield; mp 55–57 °C;  $R_{\rm f} = 0.63$  (hexane); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>),  $\delta$  7.53 (d, J = 7.3 Hz, 2H), 7.40 (t, J = 7.4 Hz, 2H), 7.34 (t, J = 7.3 Hz, 1H), 5.38 (dd, J = 7.8, 4.0 Hz, 1H), 4.19–4.07 (m, 2H), <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>),  $\delta$  140.8, 129.0, 128.9, 128.2, 66.5, 50.2, 35.3. MS (ESI): m/z (% relative intensity) = 437.0 (100) (M + H)<sup>+</sup>.

**1-Nitro-4-(1,3,3,3-tetrabromopropyl)benzene (3n).** Following the general procedure: yellow oil; 60 mg, 0.13 mmol; 25% yield;  $R_{\rm f} = 0.59$  (petroleum ether : ethyl acetate = 9 : 1); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>),  $\delta$  8.24 (d, J = 8.5 Hz, 2H), 7.69 (d, J = 8.5 Hz, 2H), 5.38 (dd, J = 8.5, 3.5 Hz, 1H), 4.18–4.02 (m, 2H), <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  147.9, 147.5, 129.3, 124.1, 66.0, 47.4, 34.0. MS (ESI): m/z (% relative intensity) = 477.7 (100) (M + H)<sup>+</sup>; HRMS (ESI): m/z calcd for C<sub>13</sub>H<sub>10</sub>O<sub>2</sub> (M + H)<sup>+</sup> 477.7283, found 477.7282.

**4-(1,3,3,3-Tetrabromopropyl)benzonitrile (30).** Following the general procedure: yellow oil; 80 mg, 0.13 mmol; 35% yield;  $R_{\rm f} = 0.55$  (petroleum ether : ethyl acetate = 9 : 1); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>), δ 7.67 (d, J = 8.0 Hz, 2H), 7.61 (d, J = 8.5 Hz, 2H), 5.32 (dd, J = 8.5, 3.5 Hz, 1H), 4.14–3.99 (m, 2H), <sup>13</sup>C NMR (126 MHz, CDCl<sub>3</sub>), δ 145.6, 132.7, 129.0, 118.2, 112.8, 65.9, 47.9, 34.1. MS (ESI): m/z (% relative intensity) = 457.7 (100) (M + H)<sup>+</sup>; HRMS (ESI): m/z calcd for  $C_{13}H_{10}O_2$  (M + H)<sup>+</sup> 457.7283, found 457.7385.

#### Carboxylation of substituted arylvinyls

(*E*)-2,3-Diphenylacrylic acid (5a).<sup>30</sup> Following the general procedure using (*E*)-1,2-diphenylethene in place of styrene. The reaction was stirred at 100 °C for 6 h: white solid; 40 mg, 0.18 mmol; 36% yield; mp 170–171 °C (Lit. 171–172 °C);  $R_f = 0.41$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01); <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.72 (brs, 1H), 7.76 (s, 1H), 7.39–7.34 (m, 3H), 7.24–7.13 (m, 5H), 7.04 (d, *J* = 7.5 Hz, 2H); <sup>13</sup>C NMR (125 MHz, DMSO- $d_6$ )  $\delta$  168.8, 139.4, 136.8, 134.9, 133.8, 130.6, 129.9, 129.5, 128.9, 128.7, 128.0; MS (ESI): *m/z* (% relative intensity) = 223.0 (100) (M – H)<sup>-</sup>.

(*E*)-2,3-Diphenylacrylic acid (5a'). Following the general procedure using (*Z*)-1,2-diphenylethene in place of styrene. The reaction was stirred at 100 °C for 6 h: white solid; 17 mg, 0.08 mmol; 15% yield; mp 170–171 °C (Lit. 171–172 °C);  $R_f = 0.41$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01); <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  12.71 (brs, 1H), 7.75 (s, 1H), 7.40–7.34 (m, 3H), 7.24–7.14 (m, 5H), 7.04 (d, *J* = 7.5 Hz, 2H); <sup>13</sup>C NMR (125 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  168.8, 139.4, 136.8, 134.9, 133.8, 130.6, 129.9, 129.5, 128.9, 128.7, 128.0; MS (ESI): *m/z* (% relative intensity) = 223.0 (100) (M – H)<sup>-</sup>.

(*E*)-3-Phenyl-2-(4-(trifluoromethyl)phenyl)acrylic acid (5b). Following the general procedure using (*Z*/*E*)-1-styryl-4-(trifluoromethyl)benzene in place of styrene. The reaction was stirred at 100 °C for 6 h: white solid; 32 mg, 0.11 mmol; 22% yield; mp 192–194 °C;  $R_f = 0.45$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  8.04 (s, 1H), 7.65 (d, *J* = 8.5 Hz, 2H), 7.39 (d, *J* = 8.0 Hz, 2H), 7.29 (t, *J* = 7.5 Hz, 1H), 7.21 (t, *J* = 8.0 Hz, 2H), 7.06 (d, *J* = 8.0 Hz, 2H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  171.9, 143.6, 139.1, 133.7, 130.7, 130.4,

This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.

130.1, 129.9, 128.5, 125.6 (d, J = 3.4 Hz). <sup>19</sup>F NMR (470 MHz, CDCl<sub>3</sub>)  $\delta$  – 62.61 (s, 3F); MS (ESI): m/z (% relative intensity) = 291.0 (100) (M – H)<sup>-</sup>; HRMS (ESI): m/z calcd for C<sub>16</sub>H<sub>11</sub>F<sub>3</sub>O<sub>2</sub> (M – H)<sup>-</sup> 291.0638, found 291.0638.

(*E*)-2-(4-Nitrophenyl)-3-phenylacrylic acid (5c). Following the general procedure using (*E*)-1-nitro-4-styrylbenzene in place of styrene. The reaction was stirred at 100 °C for 6 h: white solid; 32 mg, 0.11 mmol; 22% yield; mp 220–223 °C;  $R_{\rm f} = 0.34$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  8.25 (d, *J* = 8.5 Hz, 2H), 8.08 (s, 1H), 7.45 (d, *J* = 8.5 Hz, 2H), 7.30 (t, *J* = 7.5 Hz, 1H), 7.22 (t, *J* = 8.0 Hz, 2H), 7.06 (d, *J* = 7.5 Hz, 2H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  171.3, 147.6, 144.4, 142.3, 133.3, 131.2, 130.7, 130.3, 129.4, 128.6, 123.9; MS (ESI): *m/z* (% relative intensity) = 268.0 (100); HRMS (ESI): *m/z* calcd for C<sub>15</sub>H<sub>11</sub>NO<sub>4</sub> (M – H)<sup>-</sup> 268.0615, found 268.0614.

(*E*)-2-Methyl-3-phenylacrylic acid (5d). Following the general procedure using (*Z*/*E*)-prop-1-en-1-ylbenzene in place of styrene. The reaction was stirred at 100 °C for 6 h: white solid; 20 mg, 0.13 mmol; 25% yield; mp 59–61 °C;  $R_f = 0.41$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01); <sup>1</sup>H NMR (600 MHz, DMSO- $d_6$ )  $\delta$  12.52 (brs, 1H), 7.60 (s, 1H), 7.48–7.42 (m, 4H), 7.36 (t, *J* = 5.5 Hz, 1H), 2.03 (s, 3H); <sup>13</sup>C NMR (150 MHz, DMSO- $d_6$ )  $\delta$  169.3, 137.6, 135.5, 129.6, 128.7, 128.5, 128.4, 13.9; MS (ESI): *m*/*z* (% relative intensity) = 161.0 (100) (M – H)<sup>-</sup>; HRMS (ESI): *m*/*z* calcd for C<sub>10</sub>H<sub>10</sub>O<sub>2</sub> (M – H)<sup>-</sup> 161.0680, found 161.0680.

**3,3-Diphenylacrylic acid (5e)**.<sup>31</sup> Following the general procedure using (*Z*/*E*)-prop-1-en-1-ylbenzene in place of styrene: white solid; 68 mg, 0.38 mmol; 76% yield; mp 146–148 °C (Lit. 158 °C);  $R_{\rm f} = 0.44$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01). <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  12.15 (brs, 1H), 7.40–7.33 (m, 6H), 7.27–7.22 (m, 2H), 7.15–7.12 (m, 2H), 6.35 (s, 1H); <sup>13</sup>C NMR (125 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  167.2, 154.1, 141.0, 139.3, 129.7, 129.4, 129.0, 128.3, 119.3; MS (ESI): *m*/*z* (% relative intensity) = 223.0 (100) (M – H)<sup>-</sup>.

**1***H***-Indene-2-carboxylic acid** (**5f**).<sup>32</sup> Following the general procedure using (*Z*/*E*)-prop-1-en-1-ylbenzene in place of styrene: white solid; 50 mg, 0.32 mmol; 63% yield; mp 233–235 °C;  $R_{\rm f} = 0.46$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01). <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>) 7.68 (s, 1H), 7.61–7.56 (m, 1H), 7.55–7.52 (m, 1H), 7.35–7.31 (m, 2H), 3.62 (s, 2H); <sup>13</sup>C NMR (125 MHz, DMSO-*d*<sub>6</sub>)  $\delta$  166.1, 145.1, 143.0, 140.6, 138.9, 127.7, 127.2, 124.8, 123.8, 38.6; MS (ESI): *m/z* (% relative intensity) = 159.0 (100) (M – H)<sup>-</sup>.

(*E*)-3-(Thiophen-2-yl)acrylic acid (5g).<sup>33</sup> Following the general procedure using 2-vinylthiophene in place of styrene: white solid; 45 mg, 0.29 mmol; 58% yield; mp 143–147 °C (Lit. 140–142 °C);  $R_f = 0.44$  (hexane : ethyl acetate : acetic acid = 2 : 2 : 0.01). <sup>1</sup>H NMR (500 MHz, DMSO- $d_6$ )  $\delta$  12.43 (brs, 1H), 7.73 (d, J = 16.0 Hz, 1H), 7.68 (d, J = 10.0 Hz, 1H), 7.49 (d, J = 3.0 Hz, 1H), 7.14–7.11 (m, 1H), 6.17 (d, J = 16.0 Hz, 1H). <sup>13</sup>C NMR (125 MHz, DMSO- $d_6$ )  $\delta$  167.98, 139.18, 137.35, 132.19, 129.96, 129.03, 117.66. MS (ESI): m/z (% relative intensity) = 153.0 (100) (M – H)<sup>-</sup>.

#### <sup>18</sup>O Labeling experiment

<sup>18</sup>O Labeled  $H_2O$  experiment. Styrene (26 mg, 0.25 mmol, 1.00 equiv.), CBr<sub>4</sub> (166 mg, 0.50 mmol, 2.00 equiv.), 10 mol% eosin Y disodium (15 mg, 0.03 mmol), CoI<sub>2</sub> (78 mg, 0.25 mmol,

2.00 equiv.), <sup>18</sup>O–H<sub>2</sub>O (45 mg, 5.00 mmol, 10.00 equiv.) and 0.50 mL DMSO were combined in dried flask. The solution was stirred at 50 °C for 15 h on the condition of visible light (18 W fluorescent lamp) under nitrogen. After cooling to room temperature, the reaction mixture was extracted with ethyl acetate (10.00 mL  $\times$  3) and then washed with brine (10.00 mL). The organic layer was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> followed by filtration and then condensation. The residue was purified by silica gel column chromatography to afford the corresponding product. ESI-ms analysis of this product clearly assigned the presence of both compounds **2a** (M<sup>-</sup> at 147.0) and <sup>18</sup>O-**2a** (M<sup>-</sup> at 149.0) in noticeable amount.

Method for preparation of <sup>18</sup>O labeled DMSO.<sup>34</sup> Solid dimethylsulfur dibromide (5.00 g, 22.50 mmol, 1.00 equiv.) prepared as per known procedure<sup>35</sup> was added portion wise over 15 min to a vigorously stirred solution of triethylamine (6.30 mL, 45.00 mmol, 2.00 equiv.) freshly distilled from calcium hydride and <sup>18</sup>O-labeled water (97 atom% <sup>18</sup>O) (0.20 mL, 11.00 mmol, 0.50 equiv.) in 15.00 mL of tetrahydrofuran (freshly distilled from sodium metal). The temperature of the reaction was maintained below 50 °C. The precipitate of triethylamine hydrobromide was removed by centrifugation and washed twice with ether. The combined yellow supernatant and washings were dried on high vacuum pressure pump at room temperature (15 mm) to remove the solvent and was given 0.50 g of a brownish black liquid. Without further purification the reaction was performed styrene (26 mg, 0.25 mmol, 1.00 equiv.), CBr<sub>4</sub> (166 mg, 0.50 mmol, 2.00 equiv.), 10 mol% eosin Y disodium (15 mg, 0.03 mmol), CoI<sub>2</sub> (78 mg, 0.25 mmol, 2.00 equiv.) following the general procedure giving the corresponding product 2a. ESI-ms analysis of this product clearly assigned the presence of both compounds  $2a (M^- at 147.0)$ , <sup>18</sup>O- $2a (M^- at$ 149.0) and  ${}^{18}O_2$ -2a (M<sup>-</sup> at 151.0) in noticeable amount.

# Acknowledgements

Y. T. thanks National Natural Science Foundation of China for generous founding [21172169, 21572154] and the Fundamental Research Funds for the Central Universities [201612013] for financial support.

### Notes and references

1 (a) M. S. Kharasch, E. V. Jensen and W. H. Urry, J. Am. Chem. Soc., 1947, 69, 1100; (b) T. Nguyen, J. Leroy and C. Wakselman, J. Org. Chem., 1993, 58, 3772; (c) M. Aslam, W. F. Stansbury, R. J. Reiter and D. R. Larkin, J. Org. *Chem.*, 1997, **62**, 1550; (*d*) J. N. Moorthy and K. N. Parida, J. Org. Chem., 2014, 79, 11431; (e) C. Wuensch, S. M. Glueck, J. Gross, D. Koszelewski, M. Schober and K. Faber, Org. Lett., 2012, 14, 8; (f) C. Wuensch, T. Pavkov-Keller, Steinkellner, J. Gross, M. Fuchs, A. Hromic, G. A. Lyskowski, K. Fauland, K. Gruber, S. M. Glueck and K. Faber, Adv. Synth. Catal., 2015, 357, 1909; (g) S. Beaumont, E. A. Ilardi, L. R. Monroe and A. Zakarian, J. Am. Chem. Soc., 2010, 132, 1482; (h) Z. H. Gu, A. T. Herrmann and A. Zakarian, Angew. Chem., Int. Ed.,

2011, **50**, 7136; (*i*) A. T. Herrmann, L. L. Smith and A. Zakarian, *J. Am. Chem. Soc.*, 2012, **134**, 6976; (*j*) E. A. Ilardi and A. Zakarian, *Chem. – Asian J.*, 2011, **6**, 2260.

- 2 (a) K. Huang, C. L. Sun and Z. J. Shi, Chem. Soc. Rev., 2011, 40, 2435; (b) M. Cokoja, C. Bruckmeier, B. Rieger, W. A. Herrmann and F. E. Kühn, Angew. Chem., Int. Ed., 2011, 50, 8510; (c) Q. Liu, L. Wu, R. Jackstell and M. Beller, Nat. Commun., 2015, 6, 5933; (d) Y. Tsuji and T. Fujihara, Chem. Commun., 2012, 48, 9956; (e) M. D. Greenhalgh and S. P. Thomas, J. Am. Chem. Soc., 2012, 134, 11900; (f) T. G. Ostapowicz, M. Schmitz, M. Krystof, J. Klankermayer and W. Leitner, Angew. Chem., Int. Ed., 2013, 52, 12119; (g) C. M. Williams, J. B. Johnson and T. Rovis, J. Am. Chem. Soc., 2008, 130, 14936; (h) L. Wu, Q. Liu, I. Fleischer, R. Jackstell and M. Beller, Nat. Commun., 2014, 5, 3091.
- 3 (a) L. Zhang, J. Cheng, T. Ohishi and Z. Hou, Angew. Chem., Int. Ed., 2010, 49, 8670; (b) I. I. F. Boogaerts, G. C. Fortman, M. R. L. Furst, C. S. J. Cazin and S. P. Nolan, Angew. Chem., Int. Ed., 2010, 49, 8674; (c) I. I. F. Boogaerts and S. P. Nolan, J. Am. Chem. Soc., 2010, 132, 8858; (d) H. Mizuno, J. Takaya and N. Iwasawa, J. Am. Chem. Soc., 2011, 133, 1251; (e) H. Inomata, K. Ogata, S. I. Fukuzawa and Z. Hou, Org. Lett., 2012, 14, 3986.
- 4 S. Tanaka, K. Watanabe, Y. Tanaka and T. Hattori, *Org. Lett.*, 2016, **18**, 2576.
- 5 H. Togo, Advanced Free Radical Reactions for Organic Synthesis, Elsevier, Amsterdam, 2004.
- 6 (a) D. M. Schultz and T. P. Yoon, Science, 2014, 343, 985; (b)
  J. M. R. Narayanam and C. R. J. Stephenson, Chem. Soc. Rev., 2011, 40, 102; (c) Y. Xi, H. Yi and A. W. Lei, Org. Biomol. Chem., 2013, 11, 2387; (d) C. K. Prier, D. A. Rankic and D. W. C. MacMillan, Chem. Rev., 2013, 113, 5322; (e)
  M. N. Hopkinson, B. Sahoo, J. L. Li and F. Glorius, Chem. Eur. J., 2014, 20, 3874; (f) L. Shi and W. J. Xia, Chem. Soc. Rev., 2012, 41, 7687; (g) T. P. Yoon, M. A. Ischay and J. Du, Nat. Chem., 2010, 2, 527.
- 7 M. Majek, F. Filace and A. J. von Wangelin, *Beilstein J. Org. Chem.*, 2014, **10**, 981.
- 8 (a) A. K. Yadav, V. P. Srivastava and L. D. S. Yadav, New J. Chem., 2013, 37, 4119; (b) V. P. Srivastava, A. K. Yadav and L. D. S. Yadav, Synlett, 2013, 24, 465; (c) J. Zhang, L. Wang, Q. Liu, Z. Yang and Y. Huang, Chem. Commun., 2013, 49, 11662; (d) M. Majek and A. J. von Wanglin, Chem. Commun., 2013, 49, 5507; (e) D. P. Hari, P. Schroll and B. König, J. Am. Chem. Soc., 2012, 134, 2958; (f) D. P. Hari, T. Hering and B. König, Org. Lett., 2012, 14, 5334; (g) Y. Q. Zou, J. R. Chen, X. P. Liu, L. Q. Lu, R. L. Davis, K. A. Jørgensen and W. J. Xiao, Angew. Chem., Int. Ed., 2012, 51, 784; (h) K. Fidaly, C. Ceballos, A. Falguières, M. S. I. Veitia, A. Guy and C. Ferroud, Green Chem., 2012, 14, 1293; (i) M. Neumann, S. Füldner, B. König and K. Zeitler, Angew. Chem., Int. Ed., 2011, 50, 951; (j) D. P. Hari and B. König, Org. Lett., 2011, 13, 3852.
- 9 For recent reviews of cobalt-catalyzed C-H bond functionalization, see: (a) M. Moselage, J. Li and L. Ackermann, ACS Catal., 2016, 6, 498; (b) D. Wei, X. Zhu, J. L. Niu and M. P. Song, ChemCatChem, 2016, 8, 1242; (c)

T. K. Hyster, *Catal. Lett.*, 2015, **145**, 458; (*d*) O. Daugulis, J. Roane and L. D. Tran, *Acc. Chem. Res.*, 2015, **48**, 1053.

- 10 M. Majek and A. J. von Wangelin, *Chem. Commun.*, 2013, **49**, 5507.
- 11 (a) J. D. Nguyen, J. W. Tucker, M. D. Konieczynska and C. R. J. Stephenson, J. Am. Chem. Soc., 2011, 133, 4160; (b)
  C. J. Wallentin, J. D. Nguyen, P. Finkbeiner and C. R. J. Stephenson, J. Am. Chem. Soc., 2012, 134, 8875; (c)
  C. Dai, J. M. R. Narayanam and C. R. J. Stephenson, Nat. Chem., 2011, 3, 140; (d) M. Pirtsch, S. Paria, T. Matsuno, H. Isobe and O. Reiser, Chem.-Eur. J., 2012, 18, 7336; (e)
  E. Arceo, E. Montroni and P. Melchiorre, Angew. Chem., Int. Ed., 2014, 53, 12064; Angew. Chem., Int. Ed., 2014, 126, 12260.
- 12 Q. Q. Yang, M. Marchini, W. J. Xiao, P. Ceroni and M. Bandini, *Chem.-Eur. J.*, 2015, 21, 18052.
- 13 Q. Kong, M. Wulff, J. H. Lee, S. Bratos and H. Ihee, *J. Am. Chem. Soc.*, 2007, **129**, 13584.
- 14 (a) N. Assaf-Anld, K. F. Hayes and T. M. Vogel, *Environ. Sci. Technol.*, 1994, 28, 246; (b) P. Chunchiu and M. Reinhard, *Environ. Sci. Technol.*, 1996, 30, 1882.
- 15 H. Yi, L. B. Niu, C. l. Song, Y. Y. Li, B. Dou, A. K. Singh and A. W. Lei, Angew. Chem., Int. Ed., 2016, 56, 1120.
- 16 (a) H. Shimakoshi and Y. Hisaeda, Angew. Chem., Int. Ed., 2015, 54, 15439; (b) H. Shimakoshi, Z. Luo, T. Inaba and Y. Hisaeda, Dalton Trans., 2016, 45, 10173.
- 17 (a) R. Tomita, Y. Yasu, T. Koike and M. Akita, Angew. Chem., Int. Ed., 2014, 53, 7144; (b) N. Kornblum, J. W. Powers, G. J. Anderson, W. J. Jones, H. O. Larson, O. Levand and W. M. Weaver, J. Am. Chem. Soc., 1957, 79, 6562; (c) M. A. Khuddus and D. Swern, J. Am. Chem. Soc., 1973, 95, 8393; (d) A. J. Mancuso and D. Swern, Synthesis, 1981, 165; (e) Y. Ashikari, T. Nokami and J.-I. Yoshida, J. Am. Chem. Soc., 2011, 133, 11840; (f) Y. Ashikari, A. Shimizu, T. Nokami and J. I. Yoshida, J. Am. Chem. Soc., 2013, 135, 16070.
- 18 J. Z. Zhang and Y. Tang, Adv. Synth. Catal., 2016, 358, 752.
- 19 D. F. Taber and C. G. Nelson, J. Org. Chem., 2006, 71, 8973.
- 20 R. Kumar, A. Sharma, N. Sharma, V. Kumar and A. K. Sinha, *Eur. J. Org. Chem.*, 2008, **2008**, 5577.
- 21 J. Waser, B. Gaspar, H. Nambu and E. M. Carreira, J. Am. Chem. Soc., 2006, **128**, 11693.
- 22 C. Gioia, A. Hauville, L. Bernardi, F. Fini and A. Ricci, *Angew. Chem., Int. Ed.*, 2008, **120**, 9376.
- 23 M. L. Salum, C. J. Robles and R. Erra-Balsells, *Org. Lett.*, 2010, **12**, 4808.
- 24 T. Fukuyama, M. Arai, H. Matsubara and I. Ryu, *J. Org. Chem.*, 2004, **69**, 8105.
- 25 C. Pardin, J. N. Pelletier, W. D. Lubell and J. W. Keillor, *J. Org. Chem.*, 2008, **73**, 5766.
- 26 W. Szymanski, B. Wu, B. Weiner, S. de Wildeman,
  B. L. Feringa and D. B. Janssen, *J. Org. Chem.*, 2009, 74, 9152.
- 27 T. Thiemann, M. W. Elshorbagy, M. H. F. A. Salem, S. A. N. Ahmadani, Y. Al-Jasem, M. A. Azani, M. A. M. Al-Sulaibi and B. Al-Hindawi, *Int. J. Org. Chem.*, 2016, 6, 126.
- 28 J. Takaya, S. Tadami, K. Ukai and N. Iwasawa, Org. Lett., 2008, 10, 2697.

#### Paper

- 29 M. S. Kharasch, E. V. Jensen and W. H. Urry, J. Am. Chem. Soc., 1946, 68, 154.
- 30 S. Li, W. Yuan and S. Ma, Angew. Chem., Int. Ed., 2011, 50, 2578.
- 31 M. Bellassoued, S. Mouelhi, P. Fromentin and A. Gonzalez, *J. Org. Chem.*, 2005, **690**, 2172.
- 32 T. Mita, K. Suga, K. Sato and Y. Sato, Org. Lett., 2015, 17, 5276.
- 33 G. F. Xua, X. L. Yang, B. B. Jiang, P. Lei, X. L. Liu, Q. M. Wang, X. B. Zhang and Y. Ling, *Bioorg. Med. Chem. Lett.*, 2016, 26, 1849.
- 34 N. Mupparapu, S. Khan, S. Battula, M. Kushwaha, A. P. Gupta, Q. N. Ahmed and R. A. Vishwakarma, *Org. Lett.*, 2014, 16, 1152.
- 35 A. H. Fenselau and J. G. Moffatt, *J. Am. Chem. Soc.*, 1966, **88**, 1762.