# **RSC Advances**



## PAPER

Check for updates

Cite this: RSC Adv., 2017, 7, 27981

Received 21st February 2017 Accepted 11th May 2017

DOI: 10.1039/c7ra02137b

rsc.li/rsc-advances

## Introduction

Protein folding is a very complex physiochemical process whose underlying mechanisms in different environments remain obscure.<sup>1-17</sup> The native structure of a protein is the result of a critical interplay of intramolecular interactions, *i.e.* hydrophobic effects, hydrogen bonding, salt bridges, and disulfide bonds, as well as interactions of the protein with its solvent environment.<sup>18–25</sup> Water is a canonical solvent for biological systems, and its role has been widely explored in protein folding.<sup>26–31</sup> While hydrophobic effects are the major driving force behind protein folding in aqueous solvent, electrostatic interactions are important in folding, stability, and molecular recognition. The strength of electrostatic interactions is

# Scrutiny of electrostatic-driven conformational ordering of polypeptide chains in DMSO: a study with a model oligopeptide<sup>†</sup>

Kinshuk Raj Srivastava, 📴 🛊 Bhupesh Goyal, 回 § Anil Kumar¶ and Susheel Durani

The physicochemical effects of the solvent DMSO on protein confirmation remain enigmatic despite its diverse applications in the proteomics field. Many attempts to understand the effects of DMSO have focused on the unfolding of  $\alpha$ -helical-rich proteins; however, the cause of the profound stability of  $\beta$ sheets in DMSO remains to be elucidated. Therefore, we designed an octapeptide as a  $\beta$ -hairpin fold to serve as a model  $\beta$ -sheet; we then performed combined experimental and simulation studies to investigate the effects of DMSO on the structure and stability of the  $\beta$ -hairpin. We compared the results of the designed octapeptide with its cognate polyalanine model to directly analyze the side chain interactions responsible for ordering the octapeptide in a specific conformation. NMR and simulation results established the ordering of the octapeptide as a  $\beta$ -hairpin fold, while simulations manifested the unfolded conformation of the cognate polyalanine in DMSO. It appears that owing to their weaker dielectric and strong dipolar strengths, DMSO abolishes the  $\alpha$ -conformation as well as the solvated backbone amidic NH groups through hydrogen bonds; therefore, it destabilizes the intramolecular backbone hydrogen bonds, which leads to the unfolding of polyalanine peptides. Furthermore, our results conform to the possibility that DMSO stabilizes electrostatic and quadrupolar interactions among polar side chain atoms due to its low dielectric strength. Accordingly, we propose that the molecular mechanism of DMSO-induced stabilization of  $\beta$ -sheets is a combination of polar electrostatic interactions among the side chains and backbone desolvation through bulky side chains, which promotes backbone hydrogen bonding.

inversely proportional to the dielectric constant of the environment. Therefore, electrostatic interactions are stronger in the buried solvent-excluded core of a protein where the dielectric constant is low ( $\varepsilon = 4$  to 10) compared to the solvent-exposed surface, which experiences dielectric constants of  $\varepsilon \sim$  78.5 (that of water).<sup>32</sup> Furthermore, the dielectric constants of non-aqueous solvents are lower than that of water, such as  $\varepsilon \sim$  33 for methanol and  $\varepsilon \sim$  46 for DMSO; therefore, it is believed that electrostatic interactions drive the folding and stability of proteins in non-aqueous environments.<sup>33</sup> The role that non-aqueous solvents play in the folding and stability of proteins is not fully understood; therefore, further investigations along this line will facilitate the use of non-aqueous solvents in diverse applications in the biotech and pharma industries.

Non-aqueous solvents have been extensively used in proteomic research; among these, dimethyl sulfoxide (DMSO) is unique due to its capacity to solubilize peptides and act as a denaturant, activator, cryoprotectant, inhibitor, and molecular chaperone.<sup>34-41</sup> Furthermore, DMSO has drawn major attention in the study of transmembrane peptides/proteins due to its amphipathic nature and good membrane-mimicking properties, which are attributed to its lower dielectric constant ( $\varepsilon = 46$ ) and high dipolar strength (dipole moment =

Department of Chemistry, Indian Institute of Technology Bombay, Mumbai-400076, India. E-mail: kinshuks@umich.edu; kinshukraj1@gmail.com; Tel: +91-734-834-8017 † Electronic supplementary information (ESI) available: Fig. S1–S8 and Tables ST1 and ST2. See DOI: 10.1039/c7ra02137b

<sup>‡</sup> Life Science Institute, University of Michigan, Ann Arbor, MI, USA 48105.

<sup>§</sup> Department of Chemistry, School of Basic and Applied Sciences, Sri Guru Granth Sahib World University, Fatehgarh Sahib-140406, Punjab, India.

<sup>¶</sup> Department of Chemistry, University of Toronto, Toronto, ON, M5S 3H6.

#### **RSC Advances**

4.0 D).<sup>25,33,42-57</sup> Although DMSO is the most commonly used cosolvent, the understanding of its physicochemical effects on protein folding and structure remains elusive. A high concentration of DMSO in aqueous solvent has been reported to unfold peptides or proteins by disrupting intra-peptide hydrogen bonds. With a dipole strength of 3.96 D, compared to 1.85 D for water and 1.70 D for methanol, against 3.50 D of the peptide, DMSO has a stronger dipole than the peptide, while water has a much weaker dipole.33 DMSO is a much stronger acceptor of hydrogen bonds from peptide-NH groups than water, according to quantum mechanical calculations;58 therefore, proteins unfold by establishing hydrogen bonds with the oxygen atoms of DMSO and their backbone amides.25 Recently, DMSOinduced unfolding of α-helical rich proteins has been extensively studied; this is primarily governed by preferential solvation of exposed hydrophobics through the methyl groups of DMSO, followed by solvation of exposed backbone NHs via the oxygen atoms of DMSO.44,47-50 Contrastingly, short β-hairpin peptides and  $\beta$ -sheet rich proteins were found to be stable in DMSO.<sup>56,59</sup> Furthermore, β-barrel motifs, *e.g.* ion channels, have been found to be very stable in membranes.60-71 The exceptional stability of β-sheets in the low dielectric environment of DMSO and in membranes motivated us to design a β-hairpin peptide as a model of  $\beta$ -sheets and to investigate its conformational stability in DMSO. We believe that the results will provide a better molecular-level understanding of the interactions responsible for ordering octapeptides as  $\beta$ -hairpins in DMSO. We designed the octapeptide by grafting hydrophobic and polar residues onto two different faces of a β-hairpin to achieve amphipathicity. The octapeptide was subjected to combined experimental and simulation studies in DMSO. Additionally, a simulation study of a cognate polyalanine peptide was performed, and the results were compared with that of the octapeptide to address the role of side chain interactions in conformational ordering. NMR and simulation results have established that the octapeptide folds as a well-ordered  $\beta$ hairpin in DMSO, whereas the simulation results affirm the unfolded conformation of the polyalanine model. Further analysis of the simulation results was conducted to elucidate the physicochemical factors responsible for the ordering of the octapeptide in DMSO.

## Results

We designed an octapeptide with a full complement of side chains as a  $\beta$ -hairpin and performed combined experimental and simulation studies in DMSO to investigate the physiochemical factors responsible for the folding of the polypeptide chain in the low dielectric environment of DMSO. The peptide design was accomplished by harnessing noncovalent interactions between strands, such as salt bridges and hydrogen bonding, as well as weakly polar interactions, *i.e.* aromaticaromatic, cation– $\pi$ , and CH– $\pi$  interactions. The octapeptide (Ac–Glu<sub>1</sub>–Arg<sub>2</sub>–Tyr<sub>3</sub>–<sup>D</sup>Asn<sub>4</sub>–<sup>D</sup>Phe<sub>5</sub>–Gln<sub>6</sub>–Trp<sub>7</sub>–Lys<sub>8</sub>–NH<sub>2</sub>) was designed as a type I'  $\beta$ -hairpin, as presented in Fig. 1. There are several conformational promoting effects; for example, Trp<sub>7</sub>, Phe<sub>5</sub>, and Arg<sub>2</sub> were grafted on one face of the  $\beta$ -hairpin plane to



Fig. 1 Cartoon representations of the designed octapeptide (Ac– $Glu_1$ -Arg<sub>2</sub>-Tyr<sub>3</sub>-<sup>D</sup>Asn<sub>4</sub>-<sup>D</sup>Phe<sub>5</sub>-Gln<sub>6</sub>-Trp<sub>7</sub>-Lys<sub>8</sub>-NH<sub>2</sub>) and cognate polyalanine.

maximize the possible combination of  $\pi$ - $\pi$  and cation- $\pi$ interactions required for stabilizing the  $\beta$ -hairpin conformation. Moreover, on the other face of the  $\beta$ -hairpin, Lys<sub>8</sub> and Glu<sub>1</sub> were planned to stabilize the hairpin through salt bridge interactions. Additionally, Tyr<sub>3</sub> and Gln<sub>6</sub> were grafted to further stabilize the  $\beta$ -hairpin through a network of hydrogen bonding interactions involving Glu<sub>1</sub>. D-Amino acid residues (<sup>D</sup>Asn<sub>4</sub>--<sup>D</sup>Phe<sub>5</sub>) were used in the middle to nucleate the type I' turn in the targeted  $\beta$ -hairpin design. The peptide was synthesized by solid phase Fmoc chemistry and was found to be pure by analytical HPLC, eluting as a sharp peak. In the MALDI-MS peak shown in Fig. S1 (ESI<sup>†</sup>), the peptide displays a molecular ion peak at 1212, corresponding to M<sub>H<sup>+</sup></sub> against M<sub>calc</sub> 1211.

#### **Experimental NMR characterization**

The <sup>1</sup>H NMR spectrum of the octapeptide was recorded at 2.5 mM; the trace is shown in Fig. S2 (ESI<sup>†</sup>). We observed no noticeable concentration-dependent variations in the chemical shifts or line widths in spectra recorded at 2.5 mM and 0.25 mM concentrations in DMSO (results not shown). Presumably, the peptides are free of aggregation under the NMR concentration regime. Assignment of chemical shifts, as presented in Table 1, for the  $\beta$ -hairpin octapeptide was achieved by the combined use of TOCSY and NOESY.<sup>72</sup> Some illustrative results as portions of the 2D spectra are given in Fig. S3 (see ESI<sup>†</sup>) for the  $\beta$ -hairpin octapeptide with labeled cross peaks. The <sup>3</sup>*J*<sub>NH-C<sub>a</sub>H</sub> observed coupling constraints in the 1D spectra are listed in Table 1. The *J* values are >7 Hz, suggesting a general bias towards  $\beta$ -sheet type  $\phi$ -torsional angles in DMSO.

The peptide was examined for NOEs; the observed NOEs between the side chains are presented in Fig. 2, while the NOEs between main chain atoms are shown in Fig. S4 (ESI<sup>†</sup>). A full listing of the observed NOEs and relative distances between interacting protons derived from duly calibrated NOE volumes are listed in Table ST1 (see the ESI<sup>†</sup>). The octapeptide features NOEs that support its folded conformation. In addition to sequential backbone NH–NH and NH–C<sub> $\alpha$ </sub>H connectivity, the peptide also features short ( $i \rightarrow i \pm 2$ ) and medium ( $i \rightarrow i \pm 3$ )

Table 1	<sup>1</sup> H-NMR	chemical	shift	assignments	of the	octapeptide	in	DMSO-d	6
---------	--------------------	----------	-------	-------------	--------	-------------	----	--------	---

Residue	${}^{3}\!J_{\mathrm{NH-C}_{\alpha}\mathrm{H}}$	NH	$C_{\alpha}H$	$C_{\beta}H$	$C_{\gamma} H/C_{\delta} H$	Others
Glu-1	7.2	8.097	4.218	1.725, 1.872	2.232	
Arg-2	7.2	8.052	4.258	1.654	1.441, 2.996	7.456
Tyr-3	8.8	7.871	4.409	2.849, 2.689		6.620, 6.967
<sup>D</sup> Asn-4	5.6	8.261	4.569	2.512, 2.312		7.386, 6.963
<sup>D</sup> Phe-5	7.2	8.070	4.405	2.814, 2.996		7.206, 7.149, 7.073
Gln-6	_	8.167	4.182	1.650, 1.814	1.952	6.812, 7.232
Trp-7	7.2	8.088	4.498	2.996, 3.156		7.586, 7.320, 7.056, 6.974, 7.180, 10.789
Lys-8	8.0	7.912	4.165	1.663	1.258, 1.503	2.750, 7.656

Paper



Fig. 2 Schematic of selected long range NOEs observed for the octapeptide in  $\mathsf{DMSO-d}_6.$ 

range interactions, *e.g.* NH(Tyr<sub>3</sub>)–Hα(Gln<sub>6</sub>), SC(Tyr<sub>3</sub>)–SC(Gln<sub>6</sub>), NH(Tyr<sub>3</sub>)–NH(Phe<sub>5</sub>), and Hα(Tyr<sub>3</sub>)–SC(Gln<sub>6</sub>), indicating the existence of a turn conformation between Tyr<sub>3</sub>–<sup>D</sup>Asn<sub>4</sub>–<sup>D</sup>Phe<sub>5</sub>– Gln<sub>6</sub>. The NOEs that appear between the Arg<sub>2</sub>–Trp<sub>7</sub>, Arg<sub>2</sub>–Phe<sub>5</sub>, Phe<sub>5</sub>–Trp<sub>7</sub> and Arg<sub>2</sub>–Asn<sub>4</sub> side chain atoms suggest their spatial proximity, especially the NOEs observed between Arg<sub>2</sub>(H<sup>ε</sup>)– Trp<sub>7</sub>(H<sup>n</sup>/H<sup>ζ</sup>), Arg<sub>2</sub>(H<sup>ε</sup>)–Phe<sub>5</sub>(H<sup>δ</sup>), Arg<sub>2</sub>(H<sup>β</sup>)–Phe<sub>5</sub>(H<sup>δ</sup>/<sup>ζ</sup>), Phe<sub>5</sub>(H<sup>ε</sup>)– Trp<sub>7</sub>(H<sup>n</sup>, and Asn<sub>4</sub>(H<sup>δ</sup>)–Phe<sub>5</sub>(H<sup>δ</sup>/<sup>ζ</sup>), which suggest the likelihood of  $\pi$ – $\pi$  and CH– $\pi$ , NH– $\pi$  type interactions. The observed NOEs between Arg<sub>2</sub>(H<sup>β</sup>)–Trp<sub>7</sub>(H<sup>δ</sup>) and Arg<sub>2</sub>(H<sup>ε</sup>)–Phe<sub>5</sub>(H<sup>δ</sup>) suggest their possible interaction through cation– $\pi$  interactions. The observed SC–SC interactions between the residue pairs Glu<sub>1</sub>– Lys<sub>8</sub>, Arg<sub>2</sub>–Lys<sub>8</sub>, Tyr<sub>3</sub>–Lys<sub>8</sub>, and Tyr<sub>3</sub>–Gln<sub>6</sub> suggest that the octapeptide adopts a β-hairpin conformation in DMSO.

The conformation modeling was approached with CYANA 2.1 (ref. 73 and 74) using NMR-derived distance restraints

calibrated to NOE volumes and  $\phi$  constraints derived from  ${}^{3}J_{\rm NH-}$ C.H values, depending upon the residue stereochemistry. The modeling with CYANA 2.1 was implemented with total 101 distance restraints (40 intra, 31 sequential, 5 short, 11 medium, and 14 long-range) which resulted in a convergence of conformations as a  $\beta$ -hairpin fold. The ten lowest energy CYANA structures, shown in Fig. 3, displayed pairwise RMSD values for backbone atoms of 0.73  $\pm$  0.28 Å for the top ten structures and  $0.68 \pm 0.27$  Å for the mean structure. The mean structure was further energy minimized with GROMACS75,76 using the GRO-MOS force field and is presented in Fig. S5 (see the ESI<sup>†</sup>). The close proximity of the residues Arg<sub>2</sub>, Asn<sub>4</sub>, Phe<sub>5</sub>, and Trp<sub>7</sub> on one face combined with the proximity of Glu<sub>1</sub> and Lys<sub>8</sub> on another face of this energy-minimized structure further supports the hypothesis that salt bridges, hydrogen bonding, and polar interactions  $(\pi - \pi$  and cation  $-\pi$ ) drive the conformational ordering of the octapeptide into a  $\beta$ -hairpin fold.

# Computational characterization: simulation of sequence effects in folding

To investigate the sequence-dependent ordering of the octapeptide, we performed simulations of the octapeptide and its cognate polyalanine in the explicit solvent DMSO using the GROMOS 43A1 force field of GROMACS75,76 at 298 K under NVT conditions. The ensembles were prepared with Molecular Dynamics (MD) simulations over 250 ns. The peptide structures of entire ensembles were clustered at 0.15 nm RMSD over backbone atoms which results in different microstates. The time-dependent asymptotic growth of microstates suggests an approximation of equilibrium for peptides conformations. The macrostates were evaluated in the time course of microstate evolution during MD; the results, given in Fig. 4, imply that the simulations attained an asymptote in their microstate evolution. Therefore, we assume each ensemble is a reasonable model for the equilibrium of interest. According to the results given in Table 2, the octapeptide achieves equilibrium earlier and populates 8 times fewer microstates than the cognate polvalanine. Therefore, the presence of side chains appears to restrict conformational freedom, which results in shrinkage of the octapeptide conformational landscape. Conformational equilibria were analyzed in the radius-of-gyration  $(R_g)$  over the peptide main chain. The results, summarized in Table 2 and Fig. 5, established that the octapeptide order has a more compact conformational fold than the cognate polyalanine.



Fig. 3 Tube representation of the ten lowest energy structures modeled with CYANA (a) and the lowest energy models with the full complement of side chains (b).



**Fig. 4** Time evolution of the microstates of the oligopeptide structures during molecular dynamics simulations.

Bimodal  $R_g$  distribution of the octapeptide over the main chain atoms indicates the existence of two major conformations in the entire equilibria. The first maxima correspond to the first microstate (M1), comprising ~51% of the entire ensemble population, which is ordered as a  $\beta$ -hairpin fold as shown in Fig. 6. Furthermore, we calculated the  $R_g$  values of the side chain atoms, presented in Fig. 8, to investigate their close proximity and their possible mutual interactions leading to the conformational ordering of the octapeptide. The calculated  $R_g$ distributions of  $Arg_2$ -Phe<sub>5</sub> and  $Arg_2$ -Trp<sub>7</sub> pair display first maxima at 0.36 nm and 0.31, respectively, implying a similar affinity of  $Arg_2$  with both Phe<sub>5</sub> and Trp<sub>7</sub>. The central member of

M1, Fig. 6, suggests that the Arg side chain is sandwiched between the aromatic rings of the Trp and Phe side chains, in which its guanidinium group is oriented laterally with regard to Trp and diagonally with regard to the Phe aromatic rings. The  $R_{g}$ values of Arg<sub>2</sub> with both Phe<sub>5</sub> and Trp<sub>7</sub> feature bimodal distribution, which is indicative of two different types of conformations that the octapeptide can adopt in the entire canonical ensemble. The Rg values of the side chain pairs towards another face of the designed  $\beta$ -hairpin, such as Tyr<sub>3</sub>-Lys<sub>8</sub>, Glu<sub>1</sub>-Tyr<sub>3</sub>, and Glu<sub>1</sub>-Lys<sub>8</sub>, were found to be 0.33 nm, 0.31 nm, and 0.25 nm, respectively, suggesting their clustering and possible mutual interactions in the conformational ordering of the octapeptide. The broader  $R_g$  distribution of the Asn<sub>4</sub>-Phe<sub>5</sub> and Phe<sub>5</sub>-Trp<sub>7</sub> pair suggests the non-specific nature of their side chain interactions. The  $R_{g}$  and NOE results suggest the role of polar electrostatic interactions (salt bridges, hydrogen bonding, dispersions such as cation- $\pi$  and CH- $\pi$ ) among the side chains drives the conformational ordering of the octapeptide as a βhairpin fold in DMSO.

A half-ensemble lower in free energy was accessed to obtain statistics for the intra-peptide hydrogen bonds involving the main chain (MC) and side chain (SC) atoms; these are reported in Table ST2 (see the ESI<sup>†</sup>). For the octapeptide, we observed 2.55 MC–MC hydrogen bonds, 5.44 per conformer, which includes MC–MC, SC–SC, and MC–SC hydrogen bonds, whereas the cognate polyalanine is devoid of intramolecular hydrogen bonds. The total of 5.44 intramolecular hydrogen bonds per conformation provides further evidence for a folded conformation of the octapeptide. We further analyzed the patterns of the hydrogen bonds to anticipate the conformation

 Table 2
 Statistics of oligopeptide microstates comprising the equilibrium and radius of gyration over the polypeptide main chain structure

			0/ manulation	Radius of gyration (nm)		
Model system	No. of conformers	No. of microstates	in microstate 1	Macrostate	Microstate 1	
Octapeptide	25 000	21	52	$0.51\pm0.05$	$0.47\pm0.02$	
Polyalanine	25 000	167	19	$0.70\pm0.07$	$0.75\pm0.03$	



Fig. 5 Radius of gyration ( $R_g$ ) distributions over the main chain atoms of the octapeptide and the cognate polyalanine over the entire macrostate as well as the first microstate only.

characteristics of the octapeptide. The presence of both medium  $(i \rightarrow i \pm 3)$  and long-range  $(i \ge i \pm 6)$  MC–MC as well as both short-range  $(i \rightarrow i \pm 2)$  and long-range  $(i \ge i \pm 6)$  SC–SC hydrogen bonds strongly suggests that the octapeptide is a  $\beta$ -hairpin fold.

We sought to analyze the  $\phi$ ,  $\psi_s$  distributions of the equilibrium ensemble in the  $\alpha$ -,  $\beta$ -, and PPII-basins to further understand the conformational characteristics of peptides in DMSO. We calculated the percentage population in each basin for all residues along with the residue-wise distributions; the results are given in Table ST2, Fig. S6 (see ESI<sup>†</sup>). The observed occupancy of D-amino acids in the  $\alpha$ -basin and L-amino acids in the  $\beta$ -sub-basin ( $\beta$  + PPII) suggests ordering of the octapeptide in a  $\beta$ -hairpin conformation and provides support for our design. Contrastingly, the cognate polyalanine is present in only the  $\beta$ -sub-basin; this is indicative of the unfolded nature of the conformational ensemble. The residue-wise distribution of  ${}^{\rm D}Asn_4-{}^{\rm D}Phe_5$  in D-specific  $\alpha$ -basins implies their ordering in a turn conformation and further advocates the role of stereo-chemistry in the nucleation of the hairpin.

The central members of microstates which populate half of the entire equilibrium ensemble that is lower in free energy are presented in Fig. 6 for the octapeptide and Fig. 7 for the cognate polyalanine. The first microstate of the octapeptide, comprised of a ~51% population of ensembles, is folded in a type I'  $\beta$ hairpin conformation as per the sequential  $\phi$ ,  $\psi_s$  distribution and hydrogen bonding pattern. However, the cognate polyalanine top 5 microstates populate the half ensemble, and all are unfolded conformations in which the first microstate populates only 18% of the total ensemble. The polyalanine peptide features kinks or curvature in its backbone due to the presence of D-amino acid residues in the middle of the peptide sequence, which suggests that stereochemical constraints are aiding the nucleation of a type I' turn even in the absence of side chains. Without side chains, the microstates are essentially unfolded; meanwhile, with a complement of side chains, the octapeptide is well ordered due to the apparent interactions between the specific side chains.

#### Solvation shell analysis

We calculated the pairwise radial distribution function (RDF) of solvent atoms around peptide atoms to gain insight about the local concentration and orientation of DMSO molecules around the peptide; this will enable us to interpret the role of DMSO as a solvent in the conformational folding–unfolding of the



Fig. 6 Tube and stick representations of the central members of the top microstate of the octapeptide with the percentage population in parentheses (a) and the residue-by-residue meander in  $\phi$ ,  $\psi$  space (b).

Paper





Fig. 7 Backbone trace representations and residue-by-residue meanders in  $\phi$ ,  $\psi$  space of a few top microstates of the cognate polyalanine with their percentage populations given in parentheses.

polypeptide chain. The RDF data, presented in Fig. S7 and S8 for the octapeptide and in Fig. S9 for the polyalanine peptide (see ESI<sup>†</sup>), were calculated for specific atoms of the peptide over the entire conformational ensemble populated in the simulated trajectories. The results show that the density of solvent DMSO atoms varies significantly from that of bulk solvent and also between different atoms of the peptides, which is indicative of the specific solvent structure around the peptide atoms of interest. For the octapeptide, the first RDF maximum of oxygen against the NH groups of the backbone is at 0.29 nm, which corresponds to hydrogen bonding  $[NH \cdots O = S(CH_3)_2]$  distances. The peptidic carbonyl and carboxylate oxygen feature RDF maxima at 0.32 nm against the DMSO methyls, which suggests solvation of the peptidic oxygens by the DMSO methyls. However, the intensity of the first RDF maxima of the solvent DMSO atoms against the backbone amides and carbonyls are significantly smaller compared to the side chain atoms, which suggests shielding of the octapeptide backbone from DMSO. Contrastingly, the RDF maxima of the DMSO atoms for the cognate polyalanine atoms are higher than for the octapeptide, which suggests that the polyalanine backbone is more solventexposed and the octapeptide backbone atoms are preferably solvent sequestered. These data clearly indicate that the unfolding of polyalanine peptides is due to preferential solvation of their backbones, while sequestration of the octapeptide backbone from the solvent could promote intramolecular hydrogen bonding and therefore contribute to the stabilization of the folded state of the octapeptide. Furthermore, we propose that side chain-mediated shielding of the octapeptide can promote intramolecular backbone hydrogen bonding and hence its ordering in a compact conformation.

The intensities of the RDF maxima of DMSO atoms against the side chain terminal atoms are higher; this is indicative of

their preferential solvation, which aids in solubilizing the peptide in DMSO. Arg  $N^{\eta,\epsilon}$  and Trp  $N^{\epsilon}$  display first RDF maxima against the oxygen atoms of DMSO at 0.29 nm, which highlight their solvation through hydrogen bonding. Similarly, the Asn and Gln side chain amides display peaks at 0.29 nm, which is indicative of their solvation by the oxygen of DMSO. The RDF maxima of the side chain carbonyl oxygens of Asn and Gln against the DMSO methyls features first maxima at 0.31 nm; this suggests solvation by the methyls of DMSO, presumably through CH---O interactions. Furthermore, the side chain carboxylate oxygen of Glu displays a similar trend of RDF maxima at 0.30 nm, and the results suggest its solvation by the methyls of DMSO via CH…O interactions. The Tyr phenolic oxygen features first RDF maxima against the DMSO oxygens at 0.29 nm, suggesting their solvation through hydrogen bonding  $[OH \cdots O = S(CH_3)_2]$ . Interestingly, we observed a higher intensity of the first RDF maximum of the DMSO methyls against the carbonyl oxygen atoms of the ASN and Gln side chains compared to the carboxylate oxygen of Glu and the phenolic oxygen of the Tyr side chain. These data further support the observed computed R<sub>g</sub> values and suggest that the Glu and Tyr side chains are involved in side chain interactions, whereas the Asn and Gln side chains are preferably more solvated. Furthermore, we observed that the intensities of the RDF maxima for the aromatic carbons of the Trp, Tyr, and Phe residues, as well as the methylenes of the Arg, Lys, Glu, and Gln side chains, display very low magnitudes, close to the value of the bulk solvent. The lower magnitude of the RDF maxima of the DMSO atoms against the aromatic and methylene groups of the octapeptide suggests their possible involvement in mutual side chain interactions which can potentially contribute to the stabilization of the octapeptide as a  $\beta$ -hairpin fold. Our



solvation shell analysis further supports the amphipathic nature of the DMSO solvation of peptides and proteins.

## Discussion

In the present study, we investigated the effects of DMSO, a nonaqueous solvent, on the conformational ordering of an octapeptide as a  $\beta$ -hairpin conformation. To directly study the role of the side chains, we compared the equilibrium ensemble properties of the octapeptide with its cognate polyalanine peptide, studied herein. The exceptional stability of  $\beta$ -sheets in the low dielectric environment of DMSO motivated us to de novo design an octapeptide as a  $\beta$ -hairpin and investigate the physicochemical factors responsible for stabilization of the  $\beta$ hairpin fold in DMSO. The destabilizing effects of DMSO on αhelical structures have been explored thoroughly; however, the exceptional stability of  $\beta$ -sheets in DMSO remains enigmatic. DMSO is reported to denature  $\alpha$ -helical rich proteins through preferential solvation of exposed hydrophobic surfaces.<sup>42-45,47,48,50</sup> It is important to note that compared to  $\beta$ - sheets, the backbones of  $\alpha$ -helices are more exposed; additionally, its hydrophobic surface can therefore easily be attacked by the solvent DMSO.<sup>77</sup> To develop a molecular-level understanding of the physicochemical factors, we performed combined experimental and simulation studies for the octapeptide, while the cognate polyalanine was investigated by simulations only. We matched the experimentally observed and calculated (simulation) equilibrium ensemble properties to verify the simulation results, followed by analysis of the conformational ensembles obtained from the simulation, to elucidate the interactions responsible for the conformational ordering of peptides in DMSO.

The higher dipole strength of DMSO compared to the peptide groups makes it a better acceptor of backbone NH groups ( $\mu_{\text{DMSO}} = 3.96$ ;  $\mu_{\text{water}} = 1.85$ ; peptide group  $\mu_{\text{peptide}} = 3.50$ ).<sup>25,33</sup> Additionally, DMSO is a weak dielectric solvent compared to water ( $\varepsilon_{\text{DMSO}} = 46$ ;  $\varepsilon_{\text{water}} = 80$ ) and cannot screen the local repulsive interactions between peptide dipoles; therefore, it indirectly promotes unfolding of the polypeptide chain.<sup>25,78</sup> Thus, unfolding of the protein main chain appears to

DMSO, the cognate polyalanine studied herein was found to be

unfolded. However, the lower magnitudes of the RDF maxima of

the DMSO oxygen around the octapeptide backbone amides

compared to those of the polyalanine indicates steric blocking

of DMSO access to the octapeptide backbone through the side

chains. Protrusion of the side chains from both the faces away

from the  $\beta$ -hairpin plane appears to sterically block the access

of DMSO to the backbone NH groups. Furthermore, the octa-

peptide was found to populate 2.5 hydrogen bonds per

conformer between the backbone atoms, whereas no hydrogen

bond was detected for the polyalanine peptide in the entire

simulated ensemble; this suggests that shielding of the octa-

peptide backbone promotes intramolecular hydrogen bonding.

The low density of solvent atoms around the aromatic and

methylene atoms of the side chains suggests their desolvation

and possible involvement in mutual interactions to stabilize the

folded conformation of the octapeptide. The  $\phi$ ,  $\psi$  distributions

and number and pattern of backbone hydrogen bonds, along

with the observed NOEs between the backbone atoms and the

 ${}^{3}J_{\text{NH-C}_{a}\text{H}}$  values, provide evidence of ordering of the octapeptide

as a  $\beta$ -hairpin conformation. Moreover, the observed NOEs and

calculated  $R_{\rm g}$  distributions between the side chain atoms

demonstrate stable interactions, such as salt bridges (Glu<sub>1</sub>-

Lys<sub>8</sub>), hydrogen bonding (Tyr<sub>3</sub> with both Glu<sub>1</sub> and Lys<sub>8</sub>) and

cation  $-\pi$  (Arg<sub>2</sub> with Trp<sub>7</sub> and Phe<sub>5</sub>), among the side chains of

the octapeptide; this appears to provide further stabilization to

the  $\beta$ -hairpin conformation. The observation that electrostatic

interactions are major contributing factors in the folding of the

octapeptide in the low dielectric environment of DMSO further

advocates the importance of electrostatic effects on the folding

effects of DMSO as a solvent. The results of the present study

will not only aid understanding of DMSO effects in the ordering of an octapeptide as a  $\beta$ -hairpin, but will also allow us to infer

information about the stability of  $\beta$ -sheets of larger proteins in

DMSO. The stability of polar electrostatic interactions in DMSO,

observed in the present study, further explains why DMSO is

reported to enhance the self-association of tubulin with the

microtubules.87,88 The tubulin self-assembly process, which is

primarily driven by electrostatic interactions between the

charged tubulin monomers, may be further promoted by the

presence of DMSO. Interestingly, the profound stability and

amphipathic nature of the octapeptide studied herein can offer

advantages for its use in several other biotechnological appli-

cations, such as antimicrobial activity, as an inhibitor against

transmembrane protein diseases, and as an amphiphilic

reagent for protecting membrane proteins, to facilitate their

structural and functional studies.<sup>89</sup> Furthermore, similar to other reported studies,<sup>71</sup> we believe that our designed octapep-

tide can serve as a good model to investigate the folding mechanism of  $\beta$ -sheets in the form of  $\beta$ -barrel motifs inside

The present study reveals many important physicochemical

be achieved through a complex interplay of both the dipolar and dielectric effects of DMSO.<sup>25</sup> Consistent with our previous studies of octaalanine and lysine-solubilized polyalanine in We performed a complex interplay of both the dipolar and the dipolar

We performed a comparative study of an octapeptide and its cognate polyalanine to investigate the physicochemical factors responsible for conformational ordering of the octapeptide as a  $\beta$ -hairpin fold in the low dielectric environment of DMSO. To achieve this objective fold, we de novo designed the octapeptide with a full complement of side chains by maximizing interactions along the strand to stabilize the octapeptide in a  $\beta$ -hairpin conformation. The designed octapeptide was then subjected to combined experimental and simulation studies, whereas only simulation studies were performed with the cognate polyalanine in DMSO. Data from solvation shell analysis, hydrogen bond statistics, and  $\phi$ ,  $\psi$  distributions demonstrate unfolding of the polyalanine peptide, which could be due to preferential solvation of the backbone atoms by DMSO. Contrastingly, NOEs,  $R_g$  values, backbone dihedral angle distributions, hydrogen bond statistics, and solvation shell analysis demonstrate ordering of the octapeptide in a  $\beta$ -hairpin conformation. These results affirm that the combination of intramolecular hydrogen bonding, promoted by side chain-mediated steric blocking of DMSO access to the octapeptide backbone, along with electrostatic interactions between the side chain atoms, is responsible for the stability of the  $\beta$ -hairpin conformation of the octapeptide. Based on the understanding gained from the present study, we can draw inferences about the possible mechanism of DMSO-induced stabilization of the β-sheets; this can be mainly attributed to backbone desolvation, which promotes backbone hydrogen bonding. The insights gained in the present study further advance our current understanding of the physicochemical effects of DMSO and therefore will facilitate the use of DMSO in diverse pharmaceutical and biotechnological applications.

## Materials and methods

### Peptide synthesis

The synthesis was performed manually on Rink Amide AM resin using standard Fmoc chemistry and HOBt/DIC as coupling reagents. Each coupling, monitored with Kaiser and chloranil tests, typically required about 6 hours. Deprotections were carried out with 30% (v/v) piperidine–DMF. The N-terminus was acetylated (–NHCOCH<sub>3</sub>) with Ac<sub>2</sub>O : DIPEA : DMF in a 1 : 2 : 20 ratio. The cleavage of the final polypeptide and deprotection of the side chains were achieved together with reagent K (82.5% TFA/5% dry-phenol/5% thioanisole/2.5% ethanedithiol/5% water). The product was precipitated with anhydrous diethyl ether and was then lyophilized from a 1 : 4 ratio of H<sub>2</sub>O : <sup>*t*</sup>BuOH solution to obtain a white powder. Peptide purity was assessed with HPLC over RP-C18 (10 µM, 10 mm × 250 mm; Merck) eluting with CH<sub>3</sub>CN\H<sub>2</sub>O (0.1% TFA) 0% to 100% gradients.

#### **MALDI-MS** measurements

Mass spectra were recorded in MALDI-TOF mode on a duly calibrated AXIMA-CFR Kratos instrument. Positive ions were detected in linear/reflectron mode.

27988 | RSC Adv., 2017, 7, 27981-27991

membranes.

and stability of proteins.25,78-86

#### Nuclear magnetic resonance

<sup>1</sup>H NMR spectra were recorded in 100% DMSO-d<sub>6</sub> at 2.5 mM and 0.25 mM concentrations at 298 K on a 800 MHz Bruker instrument; VT spectra were recorded on a 400 MHz Varian NMR instrument using a cryoprobe. Phase-sensitive TOCSY90 and NOESY<sup>72</sup> experiments were performed for conformational characterization of the peptides. The solvent was suppressed with a pre-saturation or WATERGATE sequence, as provided by Bruker or VARIAN. The 800 MHz NMR data were processed with TOPSPIN, provided by Bruker, or CARA (Computer Aided Resonance Assignment). Typically, sine-squared window function phases shifted by 70° were applied in both dimensions with the data zero filled to 2 K  $\times$  1 K or 4 K  $\times$  4 K before the Fourier transformations. J values were extracted directly from the 1D <sup>1</sup>H NMR traces. Structure calculation was performed with CYANA 2.1.73 Dihedral angle restraints and lower limit distance restraints for hydrogen bonding were used for structure refinement. D-Amino acid residues were introduced in the CYANA library under the guidance of the developer. Representative structures were energy minimized using the GRO-MACS75,76 software package. All structural figures presented in this report were made using either Pymol or MolMol.

#### Preparation of equilibrium ensembles

Molecular dynamics simulations were performed with the GROMOS-96 43A1 force field in the GROMACS 3.3.3 package<sup>75,76</sup> in a periodic box of explicit solvent, using the DMSO model of Geerke et al.91 The simulation was performed under NVT conditions,92 viz., a fixed number of particles and constant volume and temperature. The non-bonded list cutoff was 1.4 nm with a shift at 0.8 nm. The integration step was 2 fs. The initial velocities were drawn from the Maxwellian distribution. The temperature was coupled to an external bath with a relaxation time constant of 0.1 ps. Bond lengths were constrained with SHAKE<sup>93</sup> to a geometric accuracy of  $10^{-4}$ . The electrostatics were treated in this case with Particle Mesh Ewald (PME)94,95 methods implementing a Coulomb cutoff of 1.4 nm, a Fourier spacing of 0.12 nm, and an interpolation order of 4. The oligopeptides were modeled with our in-house software, CAPM (Computer Aided Peptide Modeling), and PDBmake and cartoon representations were created with either MOLMOL or Pymol. Oligopeptides constrained to the center of a periodic cubic box with an edge length of 5 nm were soaked in DMSO, which was added to 1 atm density at 298 K. The box was rescaled to maintain a density of 1095 kg m<sup>-3</sup>, corresponding to the experimental density of DMSO.<sup>91</sup> First, the solute was energy minimized; then, the solvent was energy minimized while restraining the solute, and finally both were energy minimized after removing restraints. MD was initialized, and an 3 ns trajectory was exempted from the analysis as a pre-equilibration period. The trajectory was sampled thereafter at 10 ps intervals.

Characterization of polypeptide macrostates and microstates

The peptide conformers were clustered in Cartesian space to

≤0.15 nm RMSD cutoff over backbone atoms using an

algorithm by Daura *et al.*<sup>96</sup> This gives microstates in a diminishing population, *viz.*, diminishing thermodynamic stability. The Helmholtz free energy was calculated from the relative probabilities  $p_A$  and  $p_B$  of finding the system in microstates A and B as  $\Delta F_{A-B} = -RT \ln p_B/p_A$ , where *R* is the gas constant, *T* is the temperature, and  $p_A$  and  $p_B$  are the number of members in microstates A and B.

#### Solvation shell analysis

Radial and spatial distribution functions of specific solvent atoms were calculated over the most populous polypeptide microstate in each ensemble using the g\_rdf and g\_spatial utilities in GROMACS.

## Acknowledgements

We acknowledge DST (09DST028), Government of India, for financial support and IIT Bombay for the computing facility "Corona". KRS, BG, and AK were recipients of fellowships from the Council of Scientific and Industrial Research (CSIR). We acknowledge TIFR, Mumbai for help with the NMR experiments.

## References

- 1 C. B. Anfinsen, Science, 1973, 181, 223-230.
- 2 R. L. Baldwin, Adv. Protein Chem., 2002, 62, 361-367.
- 3 R. L. Baldwin, J. Biol. Chem., 2003, 278, 17581-17588.
- 4 R. L. Baldwin, J. Mol. Biol., 2007, 371, 283-301.
- 5 R. P. Bywater, J. Biomol. Struct. Dyn., 2013, 31, 351-362.
- 6 Y. Cote, G. G. Maisuradze, P. Delarue, H. A. Scheraga and P. Senet, *J. Phys. Chem. Lett.*, 2015, **6**, 1082–1086.
- 7 V. Daggett and A. Fersht, Nat. Rev. Mol. Cell Biol., 2003, 4, 497-502.
- 8 K. A. Dill, Biochemistry, 1990, 29, 7133-7155.
- 9 K. A. Dill and J. L. MacCallum, Science, 2012, 338, 1042-1046.
- 10 K. A. Dill, S. B. Ozkan, M. S. Shell and T. R. Weikl, *Annu. Rev. Biophys.*, 2008, 37, 289–316.
- 11 S. W. Englander and L. Mayne, *Proc. Natl. Acad. Sci. U. S. A.*, 2014, **111**, 15873–15880.
- 12 A. Onofrio, G. Parisi, G. Punzi, S. Todisco, M. A. Di Noia, F. Bossis, A. Turi, A. De Grassi and C. L. Pierri, *Phys. Chem. Chem. Phys.*, 2014, 16, 18907–18917.
- 13 J. N. Onuchic and P. G. Wolynes, Curr. Opin. Struct. Biol., 2004, 14, 70–75.
- 14 M. B. Prigozhin and M. Gruebele, *Phys. Chem. Chem. Phys.*, 2013, **15**, 3372–3388.
- 15 G. D. Rose, P. J. Fleming, J. R. Banavar and A. Maritan, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 16623–16633.
- 16 S. Sacquin-Mora, J. R. Soc., Interface, 2015, 12, 20150876.
- 17 P. G. Wolynes, *Biochimie*, 2015, **119**, 218–230.
- 18 D. R. Canchi and A. E. Garcia, Annu. Rev. Phys. Chem., 2013, 64, 273–293.
- 19 J. L. England and G. Haran, *Annu. Rev. Phys. Chem.*, 2011, **62**, 257–277.

- 20 Q. R. Johnson, R. B. Nellas and T. Shen, *Biochemistry*, 2012, 51, 6238–6245.
- 21 A. Merlino, N. Pontillo and G. Graziano, *Phys. Chem. Chem. Phys.*, 2016, **19**, 751–756.
- 22 S. Murakami and M. Kinoshita, J. Chem. Phys., 2016, 144, 125105.
- 23 H. Oshima and M. Kinoshita, J. Chem. Phys., 2015, 142, 145103.
- 24 P. Ren, J. Chun, D. G. Thomas, M. J. Schnieders, M. Marucho, J. Zhang and N. A. Baker, *Q. Rev. Biophys.*, 2012, **45**, 427–491.
- 25 K. R. Srivastava, A. Kumar, B. Goyal and S. Durani, *J. Phys. Chem. B*, 2011, **115**, 6700–6708.
- 26 R. L. Baldwin and G. D. Rose, *Proc. Natl. Acad. Sci. U. S. A.*, 2016, **113**, 12462–12466.
- 27 M. C. Bellissent-Funel, A. Hassanali, M. Havenith, R. Henchman, P. Pohl, F. Sterpone, D. van der Spoel, Y. Xu and A. E. Garcia, *Chem. Rev.*, 2016, **116**, 7673–7697.
- 28 J. Grdadolnik, F. Merzel and F. Avbelj, Proc. Natl. Acad. Sci. U. S. A., 2017, 114, 322–327.
- 29 D. J. Huggins, J. Struct. Biol., 2016, 196, 394-406.
- 30 F. Palazzesi, M. Salvalaglio, A. Barducci and M. Parrinello, *J. Chem. Phys.*, 2016, **145**, 121101.
- 31 J. B. Udgaonkar, Arch. Biochem. Biophys., 2013, 531, 24-33.
- 32 L. Li, C. Li, Z. Zhang and E. Alexov, *J. Chem. Theory Comput.*, 2013, **9**, 2126–2136.
- 33 C. Reichardt, Solvents and solvent effects in organic chemistry, WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim, 2003.
- 34 V. I. Gordeliy, M. A. Kiselev, P. Lesieur, A. V. Pole and J. Teixeira, *Biophys. J.*, 1998, 75, 2343–2351.
- 35 Q. Guo, Q. Wu, D. Bai, Y. Liu, L. Chen, S. Jin, Y. Wu and K. Duan, *Antimicrob. Agents Chemother.*, 2016, **60**, 7159–7169.
- 36 S. W. Jacob, M. Bischel and R. J. Herschler, *Curr. Ther. Res.*, 1964, **6**, 193–198.
- 37 J. E. Lovelock and M. W. Bishop, *Nature*, 1959, **183**, 1394– 1395.
- 38 H. Mi, D. Wang, Y. Xue, Z. Zhang, J. Niu, Y. Hong, K. Drlica and X. Zhao, *Antimicrob. Agents Chemother.*, 2016, **60**, 5054– 5058.
- 39 N. C. Santos, J. Figueira-Coelho, J. Martins-Silva and C. Saldanha, *Biochem. Pharmacol.*, 2003, **65**, 1035–1041.
- 40 A. C. Williams and B. W. Barry, *Adv. Drug Delivery Rev.*, 2004, 56, 603–618.
- 41 Z. W. Yu and P. J. Quinn, Biosci. Rep., 1994, 14, 259-281.
- 42 A. N. Batista, J. M. Batista Jr, L. Ashton, V. S. Bolzani, M. Furlan and E. W. Blanch, *Chirality*, 2014, 26, 497–501.
- 43 A. N. Batista, J. M. Batista Jr, V. S. Bolzani, M. Furlan and
  E. W. Blanch, *Phys. Chem. Chem. Phys.*, 2013, 15, 20147–20152.
- 44 S. Bhattacharjya and P. Balaram, Proteins: Struct., Funct., Genet., 1997, 29, 492–507.
- 45 W. Dzwolak, J. Kalinowski, C. Johannessen, V. Babenko, G. Zhang and T. A. Keiderling, *J. Phys. Chem. B*, 2012, **116**, 11863–11871.
- 46 M. Jackson and H. H. Mantsch, *Biochim. Biophys. Acta*, 1991, 1078, 231–235.
- 47 S. Roy and B. Bagchi, J. Phys. Chem. B, 2014, 118, 5691-5697.

- 48 S. Roy, B. Jana and B. Bagchi, *J. Chem. Phys.*, 2012, **136**, 115103.
- 49 T. Tretyakova, M. Shushanyan, T. Partskhaladze, M. Makharadze, R. van Eldik and D. E. Khoshtariya, *Biophys. Chem.*, 2013, **175–176**, 17–27.
- 50 I. K. Voets, W. A. Cruz, C. Moitzi, P. Lindner, E. P. Areas and P. Schurtenberger, *J. Phys. Chem. B*, 2010, **114**, 11875–11883.
- 51 T. Arakawa, Y. Kita and S. N. Timasheff, *Biophys. Chem.*, 2007, **131**, 62–70.
- 52 M. Bellanda, E. Peggion, R. Burgi, W. van Gunsteren and S. Mammi, *J. Pept. Res.*, 2001, 57, 97–106.
- 53 A. M. Duarte, C. P. van Mierlo and M. A. Hemminga, J. Phys. Chem. B, 2008, 112, 8664–8671.
- 54 A. M. Duarte, C. J. Wolfs, N. A. van Nuland, M. A. Harrison,
  J. B. Findlay, C. P. van Mierlo and M. A. Hemminga, *Biochim. Biophys. Acta*, 2007, **1768**, 218–227.
- 55 I. L. Karle, J. L. Flippen-Anderson, K. Uma and P. Balaram, *Biopolymers*, 1993, **33**, 827–837.
- 56 R. Mahalakshmi, S. Raghothama and P. Balaram, J. Am. Chem. Soc., 2006, **128**, 1125–1138.
- 57 A. Giugliarelli, M. Paolantoni, A. Morresi and P. Sassi, *J. Phys. Chem. B*, 2012, **116**, 13361–13367.
- 58 Y. J. Zheng and R. L. Ornstein, J. Am. Chem. Soc., 1996, 118, 4175–4180.
- 59 S. Rana, B. Kundu and S. Durani, *Chem. Commun.*, 2004, 2462–2463.
- 60 J. U. Bowie, Nature, 2005, 438, 581-589.
- 61 H. Hong, Arch. Biochem. Biophys., 2014, 564, 297-313.
- 62 S. H. White and W. C. Wimley, Annu. Rev. Biophys. Biomol. Struct., 1999, 28, 319-365.
- 63 T. Doherty, A. J. Waring and M. Hong, *Biochim. Biophys. Acta*, 2006, **1758**, 1285–1291.
- 64 S. J. Ganesan, H. Xu and S. Matysiak, *Phys. Chem. Chem. Phys.*, 2016, **18**, 17836–17850.
- 65 G. Manzo, M. A. Scorciapino, P. Wadhwani, J. Burck, N. P. Montaldo, M. Pintus, R. Sanna, M. Casu, A. Giuliani, G. Pirri, V. Luca, A. S. Ulrich and A. C. Rinaldi, *PLoS One*, 2015, **10**, e0116379.
- 66 M. Meier and J. Seelig, J. Mol. Biol., 2007, 369, 277-289.
- 67 M. Meier and J. Seelig, J. Am. Chem. Soc., 2008, 130, 1017– 1024.
- 68 M. A. Scorciapino, G. Pirri, A. V. Vargiu, P. Ruggerone, A. Giuliani, M. Casu, J. Buerck, P. Wadhwani, A. S. Ulrich and A. C. Rinaldi, *Biophys. J.*, 2012, **102**, 1039–1048.
- 69 M. Tang and M. Hong, Mol. BioSyst., 2009, 5, 317-322.
- 70 J. Thundimadathil, R. W. Roeske and L. Guo, *Biochem. Biophys. Res. Commun.*, 2005, 330, 585–590.
- 71 W. C. Wimley, K. Hristova, A. S. Ladokhin, L. Silvestro,
   P. H. Axelsen and S. H. White, *J. Mol. Biol.*, 1998, 277, 1091–1110.
- 72 A. Kumar, R. R. Ernst and K. Wuthrich, *Biochem. Biophys. Res. Commun.*, 1980, **95**, 1–6.
- 73 P. Guntert, C. Mumenthaler and K. Wuthrich, *J. Mol. Biol.*, 1997, **273**, 283–298.
- 74 K. Wüthrich, NMR of Proteins and Nucleic Acids, 1986.
- 75 E. Lindahl, B. Hess and D. van der Spoel, *J. Mol. Model.*, 2001,
   7, 306–317.

- 76 W. F. Van Gunsteren, S. R. Billeter, A. A. Eising, P. H. Hünenberger, P. Krüger, A. E. Mark, W. R. P. Scott and I. G. Tironi, *Biomolecular Simulation: The GROMOS96 Manual and User Guide*, 1996.
- 77 N. Juranic, S. Macura and F. G. Prendergast, *Protein Sci.*, 2003, **12**, 2633–2636.
- 78 A. Kumar, V. Ramakrishnan, R. Ranbhor, K. Patel and S. Durani, J. Phys. Chem. B, 2009, 113, 16435–16442.
- 79 F. Avbelj, Biochemistry, 1995, 34, 755-764.
- 80 F. Avbelj and R. L. Baldwin, *Proc. Natl. Acad. Sci. U. S. A.*, 2002, **99**, 1309–1313.
- 81 E. G. Baker, G. J. Bartlett, M. P. Crump, R. B. Sessions, N. Linden, C. F. Faul and D. N. Woolfson, *Nat. Chem. Biol.*, 2015, **11**, 221–228.
- 82 J. Batra, H. Tjong and H. X. Zhou, *Protein Eng., Des. Sel.*, 2016, 29, 301–308.
- 83 S. Kumar and R. Nussinov, ChemBioChem, 2002, 3, 604-617.
- 84 S. Kumar and R. Nussinov, ChemBioChem, 2004, 5, 280-290.
- 85 H. Nakamura, Q. Rev. Biophys., 1996, 29, 1-90.
- 86 M. Y. Tsai, W. Zheng, D. Balamurugan, N. P. Schafer, B. L. Kim, M. S. Cheung and P. G. Wolynes, *Protein Sci.*, 2016, 25, 255–269.

- 87 E. Hamel, C. M. Lin, S. Kenney and P. Skehan, Arch. Biochem. Biophys., 1991, 286, 57–69.
- 88 J. Robinson and Y. Engelborghs, J. Biol. Chem., 1982, 257, 5367–5371.
- 89 H. Tao, S. C. Lee, A. Moeller, R. S. Roy, F. Y. Siu, J. Zimmermann, R. C. Stevens, C. S. Potter, B. Carragher and Q. Zhang, *Nat. Methods*, 2013, **10**, 759–761.
- 90 B. Borah, J. S. Cohen and A. Bax, *Biopolymers*, 1985, 24, 747–765.
- 91 D. P. Geerke, C. Oostenbrink, N. F. A. Van Der Vegt and W. F. Van Gunsteren, *J. Phys. Chem. B*, 2004, **108**, 1436–1445.
- 92 V. Ramakrishnan, R. Ranbhor, A. Kumar and S. Durani, J. Phys. Chem. B, 2006, 110, 9314–9323.
- 93 J. Ryckaert, G. Ciccotti and H. Berendsen, *J. Comput. Phys.*, 1977, 23, 327–341.
- 94 T. Darden, D. York and L. Pedersen, *J. Chem. Phys.*, 1993, **98**, 10089–10092.
- 95 U. Essmann, L. Perera, M. L. Berkowitz, T. Darden, H. Lee and L. G. Pedersen, J. Chem. Phys., 1995, 103, 8577–8593.
- 96 X. Daura, W. F. Van Gunsteren and A. E. Mark, Proteins: Struct., Funct., Genet., 1999, 34, 269–280.