



Cite this: *RSC Adv.*, 2017, 7, 31372

Isotope selective activation: a new insight into the catalytic activity of urease

Sanchi Maithani,^{†a} Mithun Pal,^{†a} Abhijit Maity^a and Manik Pradhan ^{*ab}

Urease, a metalloenzyme, requires carbon dioxide (CO₂) for its activation. But, whether this activation is isotope-specific to ¹²CO₂ or ¹³CO₂, is not yet known and even the potential role of CO₂ in the enzymatic activity of urease is poorly understood. Here, we provide direct experimental evidence that the catalytic activity of urease exhibits a unique isotope-specific response where the ¹²CO₂ isotope is strongly preferred over the ¹³CO₂ isotope during its catalytic activation. Moreover, this isotope-selective activation depends on different isotopic fractionations (¹²C:¹³C) of the reaction-environment as well as the substrate urea (¹³C-urea and ¹²C-urea), where the ¹²CO₂ isotope in the reaction medium essentially facilitates the hydrolysis of ¹³C-enriched urea. This deepens our understanding of the isotope-specific urease activation and its potential role in hydrolytic reaction. Our findings thus may offer novel opportunities for a better fundamental understanding of isotope-specificity in chemical reactions involving metalloenzymes.

Received 15th May 2017
 Accepted 14th June 2017

DOI: 10.1039/c7ra05489k

rsc.li/rsc-advances

Introduction

Urease, the first enzyme to be isolated in crystalline form, was found to have a dinuclear nickel(II) active site.^{1,2} This Ni(II)-containing metalloenzyme is found in various soil microbes, bacteria, fungi and plants. It plays an important role in environmental nitrogen transformations.³ Urease is also a potent virulence factor in certain pathogens and is particularly known for its potential link to human diseases like peptic ulcers and stomach cancer.^{3,4} Moreover, this enzyme acts as a unique catalyst for its very specific action on urea [CO(NH₂)₂] and subsequently has found widespread applications in enzymatic assays and as urea sensors for routine clinical measurements of urea in blood, urine and different body fluids.^{5,6} Besides this, it is also widely used in waste water treatment for removal of urea⁵ and as a model enzyme for the study of enzymatic kinetics.⁵

The urease-catalyzed hydrolysis of urea has extensively been studied in the past.^{5,7} Urease catalyzes the hydrolysis of urea to produce ammonium carbamate [H₂NCO₂NH₄] which rapidly decomposes into bicarbonate [HCO₃⁻] and ammonium ions [NH₄⁺] in a non-enzymatic and buffer-mediated system. The bicarbonate finally gets converted into carbon dioxide (CO₂). Early studies focused on several aspects of this reaction such as

reaction kinetics,⁸ action of buffers⁹ and effect of temperature.¹⁰ But the mechanism underlying the activation of urease or its potential role in hydrolytic reaction still remains controversial. Moreover, it was demonstrated in the past that the metalloenzyme urease requires CO₂ for its activation.^{11,12} Therefore, the product of the urease-catalyzed hydrolysis of urea and the activator of the catalyst (*i.e.* CO₂) being the same entity makes the reaction mechanism more complex. In this context, the detailed study of isotopic signatures of CO₂ (*i.e.* ¹³CO₂ and ¹²CO₂) would provide better insight into the reaction mechanisms. However, so far there have been no studies focused on the potential role of isotope-specific CO₂ environment on the isotopic-fractionations of *in situ* CO₂ production in the urease-urea reaction. Moreover, how the variation of isotopic compositions (¹²C:¹³C) of the substrate urea (¹³C-urea and ¹²C-urea) affects the catalytic activity of urease particularly in the environment of atmospheric CO₂ concentration, has never been explored and therefore remains an open question. Consequently, unravelling the reaction mechanism involving the isotope-selective activation of urease would open new perspectives for a better understanding of the role of CO₂ for urease activation kinetics.

In this study, we report for the first time, that the enzymatic activity of urease exhibits a unique isotope-selective CO₂ affinity. Furthermore, we have showed that the catalytic activity of urease depends on the isotopic compositions of both reaction-environment and substrate. Finally, we also demonstrate that the isotope-specific activation of urease is buffer-independent. This new knowledge is of great significance for fundamental understanding of isotope-specific chemical reactions involving metalloenzymes. Fig. 1 illustrates a scheme

^aDepartment of Chemical, Biological and Macromolecular Sciences, S. N. Bose National Centre for Basic Sciences, Salt Lake, JD Block, Sector III, Kolkata-700098, India. E-mail: manik.pradhan@bose.res.in; manik.pradhan@gmail.com; Fax: +91 33 2335 3477; Tel: +91 33 2335 5706; +91 33 2335 5708

^bTechnical Research Centre (TRC), S. N. Bose National Centre for Basic Sciences, Salt Lake, JD Block, Sector III, Kolkata-700098, India

[†] Both authors contributed equally to this work and should be considered as joint first authors.






Fig. 1 A scheme showing the isotope-specific catalytic reaction of urease enzyme. (a) Isotopic $^{12}\text{CO}_2$ concentration decreases with simultaneous *in situ* generation of $^{13}\text{CO}_2$ isotope in the urease-catalyzed hydrolysis of ^{13}C -urea, (b) urease specifically requires $^{12}\text{CO}_2$, exhibiting no intake of $^{13}\text{CO}_2$ even in an enriched $^{13}\text{CO}_2$ reaction-environment.

showing the isotope-specificity of urease enzyme and specific requirement of $^{12}\text{CO}_2$ isotope for its activation.

Result and discussion

To investigate the urease ($1\ \mu\text{M}$)-catalyzed hydrolysis of isotope enriched ^{12}C and ^{13}C -urea ($4\ \text{mM}$) at different reaction environments, we first altered the headspace of the reaction with increasing concentration of $^{12}\text{CO}_2$ (with $\sim 1.1\%$ natural abundance of $^{13}\text{CO}_2$) starting from CO_2 -free pure N_2 environment. It is noteworthy to mention here that only the concentration of $^{12}\text{CO}_2$ or $^{13}\text{CO}_2$ isotopes is regulated keeping the total pressure constant at around atmospheric pressure above the reaction medium. We observed (Fig. 2a) that the headspace $^{12}\text{CO}_2$ concentration was absorbed with simultaneous *in situ*

generation of $^{13}\text{CO}_2$ in the urease-catalyzed hydrolysis of ^{13}C -urea. We also interestingly (Fig. 2a) observed that while ^{13}C -urea derived *in situ* $^{13}\text{CO}_2$ production is increased gradually with increasing initial headspace concentration of $^{12}\text{CO}_2$, at the same time the final headspace $^{12}\text{CO}_2$ concentration is decreased more in the reaction. Taken together, these observations indicate that there is a potential link between the enzymatic activity of urease and the requirement of specific isotope of CO_2 (*i.e.* $^{12}\text{CO}_2$). These observations are indistinguishable for urease-catalyzed hydrolysis of ^{12}C -urea because the product and the headspace component are the same isotopic species (*i.e.* $^{12}\text{CO}_2$).

Another striking finding revealed in our observations (Fig. 2b) is that in CO_2 -free pure N_2 environment, the *in situ* generation of $^{13}\text{CO}_2$ concentration from ^{13}C -urea was observed to be the lowest value, whereas the ^{12}C -urea derived *in situ*




Fig. 2 Urease ($1\ \mu\text{M}$)-catalysed hydrolysis of ($4\ \text{mM}$) ^{13}C -urea ($\geq 99\%$ ^{13}C) and ^{12}C -urea ($\geq 99\%$ ^{12}C) at different headspace concentrations (partial pressures) of $^{12}\text{CO}_2$. (a) $^{13}\text{CO}_2$ is generated while headspace $^{12}\text{CO}_2$ concentration has been absorbed in the hydrolysis of ^{13}C -urea. (b) ^{13}C -urea derived $^{13}\text{CO}_2$ has been increased whereas ^{12}C -urea derived $^{12}\text{CO}_2$ has been diminished with increasing headspace $^{12}\text{CO}_2$ concentration. The dotted region shows the CO_2 -free pure N_2 headspace condition.






Fig. 3 Urease (1 μM)-catalysed hydrolysis of (4 mM) ^{13}C -urea ($\geq 99\%$ ^{13}C) and ^{12}C -urea ($\geq 99\%$ ^{12}C) at enriched headspace concentrations (partial pressures) of $^{13}\text{CO}_2$ ($^{12}\text{CO}_2$: $^{13}\text{CO}_2$: 40% : 60%). (a and b) Depict the variation of $^{13}\text{CO}_2$ and $^{12}\text{CO}_2$ concentrations before and after the reaction of urease-catalysed hydrolysis of ^{13}C -urea and ^{12}C -urea, respectively.

production of $^{12}\text{CO}_2$ concentration was found to be markedly enhanced. However, in absence of any $^{12}\text{CO}_2$, the catalytic activity of urease diminishes and subsequently the hydrolysis of ^{13}C -urea becomes very small, resulting in lower amount of $^{13}\text{CO}_2$ production. In contrast, ^{12}C -urea derived *in situ* $^{12}\text{CO}_2$ itself facilitates the enzymatic activity of urease and thus urease-catalysed hydrolysis of ^{12}C -urea is not hindered in CO_2 -free pure N_2 environment. However, ^{12}C -urea derived *in situ* production of $^{12}\text{CO}_2$ decreases with increase of initial headspace concentration of $^{12}\text{CO}_2$ and this is likely to be the combined effect of partial pressure of the same species $^{12}\text{CO}_2$ above the reaction medium as well as the over-saturation of urease enzyme in presence of excess headspace $^{12}\text{CO}_2$.

We next altered the headspace environment with $\sim 60\%$ (103 ppm) isotope enriched $^{13}\text{CO}_2$ (of 174 ppm total CO_2), (instead of $\sim 1.1\%$ natural abundance as in the previous case), to ensure the specific role of $^{12}\text{CO}_2$ in the catalytic activity of urease. We found (Fig. 3a) that initial headspace $^{13}\text{CO}_2$ concentration was not absorbed at all, while the headspace $^{12}\text{CO}_2$ was markedly decreased in the usual way for urease (1 μM)-catalysed hydrolysis of (4 mM) ^{13}C -urea. Moreover, under this altered environmental conditions in headspace, no noticeable effect of enriched headspace $^{13}\text{CO}_2$ concentration was observed (Fig. 3b)

even for urease (1 μM)-catalysed hydrolysis of (4 mM) ^{12}C -urea. Our observation clearly manifests that urease preferentially links to $^{12}\text{CO}_2$ for its activation, thus unveiling a missing link between the enzymatic activity of urease and the necessity for $^{12}\text{CO}_2$ isotope.

Next, we examined the catalytic activity of urease in response to the different isotopic-compositions (^{12}C : ^{13}C) of the substrate urea to gain a better insight into the fundamental processes of isotope – selective nature of urease with a fixed headspace atmospheric CO_2 concentration (~ 360 ppm). The different isotopic-compositions (^{12}C : ^{13}C) were achieved by suitably mixing ^{12}C -urea and ^{13}C -urea, while retaining the same concentration of urea (4 mM). We observed (Fig. 4a) that the *in situ* generation of the product $^{13}\text{CO}_2$ from hydrolysis of 1 : 2 (^{12}C : ^{13}C) isotopic-composition of urea was almost identical to that of 1 : 99 (^{12}C : ^{13}C) isotopic-composition. However, ^{13}C -isotopic enrichment of the substrate urea for 1 : 2 (^{12}C : ^{13}C) isotopic-composition was significantly lower than that of 1 : 99 (^{12}C : ^{13}C) isotopic-compositions. This observation demonstrates that the *in situ* availability of $^{12}\text{CO}_2$ derived from the ^{12}C -urea, itself activates urease for higher hydrolysis of ^{13}C -fraction of the 1 : 2 (^{12}C : ^{13}C) isotopic-composition of urea, whereas the




Fig. 4 Urease (1 μM)-catalysed hydrolysis of (4 mM) different isotopic compositions (^{12}C : ^{13}C) of substrate urea at a fixed head-space concentration of CO_2 (~ 360 ppm). (a) Demonstrates the $^{13}\text{CO}_2$ evolved in the hydrolysis of (4 mM) different isotopic compositions (^{12}C : ^{13}C) of urea. (b) Depicts a comparison between % content of ^{13}C isotope in urea and $^{13}\text{CO}_2$ produced in the reaction with urease.






Fig. 5 Concentration dependent study of urease-catalysed hydrolysis of ^{12}C -urea and ^{13}C -urea. (a and b) Elucidate the effect of different concentrations of urea and urease, respectively.

^{13}C -urea ($^{12}\text{C} : ^{13}\text{C}$: 1 : 99) was not effectively hydrolysed in absence of *in situ* generation of $^{12}\text{CO}_2$ in the reaction medium.

In order to gain a detailed and in-depth understanding of the role of $^{12}\text{CO}_2$ derived from different isotopic fractionations of ^{12}C -urea, we then normalized the *in situ* generation of $^{13}\text{CO}_2$ from the different isotopic-compositions of urea (*i.e.* 2 : 1, 1 : 1 and 1 : 2) to the percentage (%) of $^{13}\text{CO}_2$ produced from 99% ^{13}C -urea (*i.e.* 1 : 99) and this is illustrated in Fig. 4b. It was found (Fig. 4b) that the percentage of *in situ* $^{13}\text{CO}_2$ production was much higher than the ^{13}C percentage actually present in all the different isotopic-compositions of urea, suggesting the inadequate hydrolysis of ^{13}C -urea (1 : 99) in absence of sufficient *in situ* $^{12}\text{CO}_2$ concentration in the reaction medium. Therefore, urease-catalyzed hydrolysis of urea was also found to be strongly affected due to the different isotopic fractionations of substrate urea in presence of CO_2 environment. Taken together, our data clearly suggest that $^{12}\text{CO}_2$ isotope plays a vital role that facilitates the enzymatic activity of urease enzyme.

The potential effect of ^{12}C -urea and ^{13}C -urea on the urease activation was further elucidated in response to different substrate (urea) and enzyme (urease) concentrations while maintaining the headspace at a typical atmospheric CO_2

concentration (~ 360 ppm). The product ($^{12}\text{CO}_2$ or $^{13}\text{CO}_2$) of (1 μM) urease-catalyzed hydrolysis of ^{12}C -urea and ^{13}C -urea was found (Fig. 5a) to be gradually increased with increasing concentration of urea (^{12}C -urea or ^{13}C -urea) from 100 μM to 5 mM. The observation was followed by a sharp decline after 5 mM urea concentration, thus exhibiting the effect of substrate-inhibition on the urease-catalyzed hydrolysis. In contrast, keeping the urea concentration fixed (4 mM), there was no significant change of the concentration of the product ($^{12}\text{CO}_2$ or $^{13}\text{CO}_2$) with increasing concentration of the product ($^{12}\text{CO}_2$ or $^{13}\text{CO}_2$) with increasing concentration of urease enzyme (Fig. 5b), thus indicating that urease still hydrolyzes urea in an efficient way and thereby indicates the steady catalytic activity of urease. However, it is noteworthy that the isotope-specific activation of urease was observed throughout the study. Our findings also indicate that the lower *in situ* generation of $^{13}\text{CO}_2$ eventually signifies the hydrolysis of lesser amount of ^{13}C -urea in absence of adequate activation of urease resulting from the headspace $^{12}\text{CO}_2$ concentration.

Finally, we investigated the isotope-specific enzymatic activity of urease in two buffer mediums (Tris-buffer and phosphate buffer) to eliminate the potential effect of CO_2 on the pH of the reaction medium. It is worth noting that the previous experiments in the present study were carried out in a buffer-free aqueous medium to avoid the effect of urease inhibition in presence of buffer ions.⁵ Fig. 6 depicts that this activation in the buffer mediums is not significantly changed compared to the buffer-free system. However, the diminished magnitude of the concentration of the products, $^{12}\text{CO}_2$ and $^{13}\text{CO}_2$ observed in the buffer mediums is exhibited to be the well-known inhibitory effect of buffer ions in the reaction.




Fig. 6 The effect of buffer medium. The hydrolysis of ^{12}C -urea and ^{13}C -urea in $^{12}\text{CO}_2$ environment is performed in Tris buffer (pH 7.4) and phosphate buffer (pH 7).

Conclusion

In conclusion, our findings provide new evidences that the catalytic activity of urease exhibits a unique isotope specific response. We have shown that the $^{12}\text{CO}_2$ isotope is strongly preferred over the $^{13}\text{CO}_2$ isotope by the urease enzyme during its catalytic activation. It was earlier reported¹¹ that CO_2 acts as a ligand in urease-urea reaction. Hence, the isotope preferential activation of urease enzyme may be attributed to the fact that $^{12}\text{CO}_2$ isotope possibly offers lower interaction energy than $^{13}\text{CO}_2$ isotope in this ligand-enzyme interaction. However,



quantum-mechanical simulations of interaction energy for both the isotopes would provide better insight and further exploration of the isotope specific activation of urease enzyme. We also provide direct experimental evidences that deepens our understanding of how the activation of urease depends on the isotopic compositions (^{12}C : ^{13}C) of reaction-environment as well as the substrate urea. Despite these new evidences, however, there still remains a great deal to be explored about the underlying mechanisms linking the specific requirement of $^{12}\text{CO}_2$ isotope for the activation of urease. Moreover, new insights into the isotope-specific responses in urease-activation are fostering exploration of a new arena to study the chemical reactions involving metalloenzyme.

Materials

Jack-bean urease enzyme (E.C. 3.5.1.5) was purchased from Sigma Aldrich and ^{13}C -enriched urea (^{13}C -urea, 99%) (CCLM-311-GMP) was obtained from Cambridge Isotopic Laboratories, Inc., USA. All other chemicals, including ACS grade urea with 99% ^{12}C -enrichment (^{12}C -urea), were acquired from Sigma Aldrich and were used without further purification. Milli-Q water was used to prepare the aqueous solutions.

Method

The aqueous solutions of ^{12}C -urea, ^{13}C -urea and Jack-bean urease were utilized, unless stated otherwise, in the entire study. The reactions were carried out in sealed round-bottomed flasks wherein different concentrations of CO_2 were maintained. To eliminate the interference due to air, the flasks were purged with CO_2 -free nitrogen before the experiments. The products ($^{12}\text{CO}_2$ or $^{13}\text{CO}_2$) of urease-catalyzed hydrolysis of urea (^{12}C or ^{13}C -urea) were considered at the equilibrium of the reaction and the termination timing of the reaction was set to be at 60 minutes which was confirmed by Berthelot's experiment. After 60 minutes of the progress of the reaction, acidification was done using H_3PO_4 to extract the dissolved CO_2 into headspace. The headspace CO_2 was then collected from each flask and measured by a high-precision laser-based Integrated Cavity Output Spectroscopy (ICOS) technique. The net isotopic concentrations of CO_2 produced from the hydrolysis reactions were calculated after subtracting the same from a blank flask containing only urease at equal initial CO_2 concentration. All reactions were conducted at room temperature.

Measurement and analysis

The measurements were made using a CO_2 isotope analyser (CCIA 36-EP, LGR, USA) exploiting the ICOS technique as described in ref. 13–15. Briefly, a high-finesse optical cavity consisting of two very high-reflectivity mirrors ($R \sim 99.98\%$) provides an optical path length of ~ 3 km thereby increasing the sensitivity considerably. The high-resolution absorption spectra of the stable isotopes $^{12}\text{C}^{16}\text{O}^{16}\text{O}$ and $^{13}\text{C}^{16}\text{O}^{16}\text{O}$ were acquired by probing the ro-vibrational lines at the wavenumbers 4874.448 cm^{-1} and 4874.086 cm^{-1} , respectively in the vibrational combination band

of CO_2 molecule and subsequently the concentrations of the isotopic species were reported in parts per million (ppm).

Author contributions

M. P. (Manik Pradhan) arranged the funding and supervised the whole study; M. P. (Manik Pradhan) and A. M. provided the conception of the study; S. M., M. P. (Mithun Pal) and A. M. designed the study; S. M., M. P. (Mithun Pal) and A. M. performed the experiments and analysed the data; all authors drafted the manuscript and critically reviewed.

Conflict of interests

The authors declare no conflict of interests.

Acknowledgements

The authors acknowledge the funding from the Technical Research Centre (TRC) (No. All1/64/SNB/2014(C)) of the S. N. Bose Centre, Kolkata for this work. S. Maithani, M. Pal and A. Maity acknowledge the Department of Science & Technology (DST, India) for Inspire Fellowships.

References

- 1 N. E. Dixon, T. C. Gazzola, R. L. Blakeley and B. Zermer, *J. Am. Chem. Soc.*, 1975, **97**(14), 4131.
- 2 J. B. Sumner, *J. Biol. Chem.*, 1926, **69**, 435.
- 3 H. L. Mobley and R. P. Hausinger, *Microbiol. Rev.*, 1989, **53**(1), 85.
- 4 A. Maity, M. Pal, S. Maithani, B. Ghosh, S. Chaudhuri and M. Pradhan, *J. Breath Res.*, 2016, **10**(3), 036007.
- 5 Y. Qin and J. M. S. Cabral, *Appl. Biochem. Biotechnol.*, 1994, **49**(3), 217.
- 6 A. Maity, M. Pal, S. Som, S. Maithani, S. Chaudhuri and M. Pradhan, *Anal. Bioanal. Chem.*, 2017, **409**, 193–200.
- 7 B. Krajewska, *J. Mol. Catal. B: Enzym.*, 2009, **59**(1), 9–21.
- 8 T. C. Huang and D. H. Chen, *J. Chem. Technol. Biotechnol.*, 1991, **52**(4), 433.
- 9 N. D. Jespersen, *J. Am. Chem. Soc.*, 1975, **97**(7), 1662.
- 10 B. R. van Eldik and M. Brindell, *J. Biol. Inorg. Chem.*, 2012, **17**(7), 1123.
- 11 D. Walther, M. Ruben and S. Rau, *Coord. Chem. Rev.*, 1999, **182**, 67–100.
- 12 I.-S. Park and R. P. Hausinger, *Science*, 1995, **267**(5201), 1156.
- 13 E. R. Crosson, K. N. Ricci, B. A. Richman, F. C. Chilese, T. G. Owano, R. A. Provencal, M. W. Todd, J. Glasser, A. A. Kachanov, B. A. Paldus, T. G. Spence and R. N. Zare, *Anal. Chem.*, 2002, **74**, 2003.
- 14 A. Maity, S. Som, C. Ghosh, G. Dutta Banik, S. B. Daschakraborty, S. Ghosh, S. Chaudhuri and M. Pradhan, *J. Anal. At. Spectrom.*, 2014, **29**, 2251.
- 15 A. Maity, G. D. Banik, C. Ghosh, S. Som, S. Chaudhuri, S. B. Daschakraborty, S. Ghosh, B. Ghosh, A. K. Raychaudhuri and M. Pradhan, *J. Breath Res.*, 2014, **8**(1), 016005.

