RSC Advances



View Article Online

View Journal | View Issue

PAPER

Check for updates

Cite this: RSC Adv., 2017, 7, 38956

Received 31st May 2017 Accepted 17th July 2017

DOI: 10.1039/c7ra06078e

rsc.li/rsc-advances

Introduction

The genus *Illigera*, one of four that make up the family Hernandiaceae, contains approximately thirty known species distributed in southern Africa and Asia.¹ The chemical constituents of this genus include aporphine alkaloids,^{2–5} phenolic acids, and steroids,⁶ which have been reported to have various bioactivities, including antiplatelet aggregation,⁶ cytotoxic activity,^{4,5} and vasorelaxant activity.²

Illigera aromatica (Hernandiaceae) is a small liana that mainly grows in the Yunnan and Guangxi provinces in China,⁷ and is frequently used by local people for promoting blood circulation and treating tuberculosis. It has been reported that the chemical composition of *I. aromatica* includes fatty acids and their esters,⁸ steroids,⁹ monoterpenoids^{8,10} and alkaloids.¹⁰

Herbal fermentation processing, which has been practised in China for more than 4000 years, frequently causes biotransformations and changes in bioactivity. For example, Kim *et al.*¹¹ reported that pre-treating fermented *Curcuma longa* L. with *Aspergillus oryzaeis* effectively prevents CCl_4 -induced hepatic damage in rats. Hsu *et al.*¹² reported that fermenting *Radix astragali* with *Bacillus subtilis* significantly stimulates the biosynthesis of type I procollagen in a dose-dependent manner

Fermentation of *Illigera aromatica* with *Clonostachys rogersoniana* producing novel cytotoxic menthane-type monoterpenoid dimers†

Jian-Wei Dong, ២ Le Cai,* Xue-Jiao Li, Rui-Feng Mei, Jia-Peng Wang, Ping Luo, Yan Shu and Zhong-Tao Ding 匝 *

Illigera aromatica, a medicinal liana, was fermented with *Clonostachys rogersoniana*. Two novel menthanetype monoterpenoid dimers, dimericilligerates E (1) and F (2) were isolated from the fermented material. Their structures were identified by 1D/2D NMR, electric circular dichroism, derivatization, and Mosher's method. A novel elimination reaction that formed a benzene ring from a menthane moiety afforded derivative 1a, which possesses more resolved NMR signals than the dimeric monoterpenoid, dimericilligerate E (1). The determination of 1a verified the estimation of 1. Compounds 1 and 2 showed potent cytotoxic activities; especially, 1 possessed significant activities against the liver hepatocellular carcinoma cell line (SMMC-7721) with an IC₅₀ value of $3.89 \pm 0.23 \mu$ M. Three novel precursors, dimericilligerates A–C (3–5), were identified by HPLC-MS and isolated from the original *I. aromatica*. This paper presents a novel class of monoterpenoid dimers and suggests that *C. rogersoniana*-fermented *I. aromatica* is effective to produce cytotoxic dimeric monoterpenoids from inactive original materials.

in both young and old human dermal fibroblast cells. Our previous studies have shown that the fermentations of *Bletilla striata*¹³ and *Asparagus filicinus*¹⁴ both facilitate the biotransformations of their components and enhance their bioactivities.

The secondary metabolites of *Clonostachys rogersoniana*, a rhizosphere fungus, are cyclic peptides¹⁵ and polyketides.¹⁶ Previous studies have shown that *C. rogersoniana* fermentation could transform the aporphine alkaloids to 4*R*-hydrox-yaporphine alkaloids, improving their water solubility and acetylcholinesterase inhibitory activity.^{17,18}

In the present study, several fungi including *Fusarium oxy-sporum*, *Fusarium avenaceum*, *Geomyces luteus*, *Alternaria compacta*, *Talaromyces purpurogenus*, *Scytalidium lignicola*, *Cladosporium cladosporioides*, *Stagonosporopsis cucurbitacea-rum*, *Penicillium vancouverense*, and *C. rogersoniana* were used to ferment the tubers of *I. aromatica*. Fermentation with *C. rogersoniana* caused obvious transformation, prompting an investigation into the chemical constituents of *C. rogersoniana*-fermented *I. aromatica*. Two novel menthane-type



Fig. 1 Structures of compounds 1-5 and 1a.

Functional Molecules Analysis and Biotransformation Key Laboratory of Universities in Yunnan Province, School of Chemical Science and Technology, Yunnan University, Kunming 650091, P. R. China. E-mail: ztding@ynu.edu.cn; caile@ynu.edu.cn † Electronic supplementary information (ESI) available: 1D/2D NMR, HRESIMS, and LC-MS spectra. See DOI: 10.1039/c7ra06078e

Paper

monoterpenoid dimers, dimericilligerates E (1) and F (2) were isolated from the fermented material (Fig. 1). Afterwards, three precursors, dimericilligerates A–C (3–5), were identified by LC-MS and isolated from the original *I. aromatica*. The biotransformations from 3–5 to 1 and 2 were elucidated. The free monoterpenoid dimers (1–2) showed stronger cytotoxic activity than the original compounds (3–5). This paper presents an effective approach to the production of cytotoxic dimeric monoterpenoids from inactive original materials.

Results and discussion

1. .

Identification of compounds isolated from fermented *I. aromatica*

After *I. aromatica* was fermented with *C. rogersoniana*, two compounds (1 and 2) were isolated by silica gel and Sephadex LH-20 column chromatography (CC).

Compound 1, isolated as a white amorphous powder, was determined to have the molecular formula $C_{21}H_{32}O_5$ based on analysis of its HRESIMS m/z 409.2235 [M + COOH]⁻ (calcd for $C_{22}H_{33}O_7$ 409.2232), ¹³C NMR and DEPT spectra, which indicated seven degrees of unsaturation. The ¹H and ¹³C NMR data (Table 1) showed four methyl carbons, one carbonyl, and seven oxygenated alkyl groups. Comparing its NMR data with the known isolated menthane-type monoterpenoid, *cis*-4-hydroxy-5-(1-hydroxy-1-methylethyl)-2-methyl-2-cyclohexen-1-one,^{19,20} **1** possesses four angular methyl groups, which is double the number found in the known monoterpenoid, and two high-field



Fig. 2 Key ¹H-¹H COSY, HMBC, and ROESY correlations of 1 and 1a.

methylenes that appear instead of the C-7 methyl group in the menthane-type monoterpenoid, suggesting **1** is a dimer of monoterpenoids, composed of the two menthane-type monoterpenoid components I and II. In component I (shown in purple, Fig. 1), an oxygenated methine at $\delta_{\rm C}$ 77.3 (d) was assigned at C-3 based on the HMBC cross peaks of H-3/C-2, C-4, and C-8 (Fig. 2); a carbonyl at $\delta_{\rm C}$ 206.3 (s) was assigned based on the HMBC networks of H-2/C-6, H-5/C-6, and H-7/C-6; an oxygenated quaternary carbon at $\delta_{\rm C}$ 79.6 (s) located at C-1 was identified by the HMBC cross-peak of H-2/C-1; and a C(2)–O–C(8)–C(4) bridge was established by the HMBC cross-peaks of H-2/C-8, and CH₃-9,10/C-8 and C-4, indicating a bicyclo[3,2,1] menthane-type monoterpenoid as shown in Fig. 1 (component

No.	1		1a		2	
	$\delta_{ m C}$	δ _H mult. (J in Hz)	$\delta_{ m C}$	δ _H mult. (J in Hz)	$\delta_{ m C}$	δ _H mult. (J in Hz)
1	79.6 C		80.0 C		79.3 C	
2	84.6 CH	4.11 s	84.2 CH	4.16 s	89.3 CH	3.73 s
3	77.3 CH	4.87 s	77.4 CH	5.00 s	76.2 CH	4.74 s
4	47.4 CH	2.31 s	47.2 CH	2.37 d (3.2)	47.5 CH	2.22 br s
5	42.6 CH ₂	2.61 d (3.6), 2.59 d (3.6)	42.6 CH ₂	2.71 d (3.2)	42.8 CH ₂	2.50 s
6	206.3 C		206.5 C		208.8 C	
7	23.6 CH ₂	2.09 m, 1.73 ddd (13.2, 11.4, 4.8)	23.1 CH ₂	2.26 ddd (10.4, 6.0, 4.4), 2.03 ddd (10.4, 6.0, 4.4)	25.5 CH ₂	1.93 m, 1.43 m
8	84.4 C		84.6 C		84.3 C	
9	26.9 CH ₃	1.18 s	27.0 CH ₃	1.22 s	26.7 CH ₃	1.13 s
10	29.7 CH ₃	1.53 s	29.7 CH ₃	1.56 s	29.6 CH ₃	1.47 s
1'	108.2 C		122.3 C		42.1 CH	2.73 m
2'	82.3 CH	3.91 s	152.7 C		75.8 CH	3.90 s
3′	78.5 CH	4.24 s	116.8 CH	6.74 d (8.0)	68.8 CH	4.19 s
4'	48.6 CH	2.23 s	127.5 CH	6.86 dd (8.0, 6.8)	44.9 CH	1.96 d (2.4)
5′	32.9 CH ₂	2.36 d (2.4), 2.15 d (2.4)	121.1 CH	7.03 dd (8.0, 6.8)	36.8 CH ₂	2.69 br d (13.2), 2.34 dd (13.2, 2.4)
6′	146.4 C		129.6 CH	7.06 dd (8.0)	211.7 C	
7′	20.2 CH ₂	2.17 m, 2.13 m	$20.8~\mathrm{CH}_2$	2.92 ddd (10.4, 7.6, 4.4), 2.75 ddd (10.4, 7.6, 4.4)	17.4 CH ₂	2.19 m, 2.03 m
8′	82.1 C				73.3 C	
9′	27.2 CH ₃	1.30 s			28.9 CH ₃	1.38 s
10'	31.1 CH ₃	1.41 s			28.4 CH ₃	1.19 s

I). Similar to the aforementioned analysis, the component II (shown in red, Fig. 1) was determined to be a bicyclo[3,2,1] menthane-type monoterpenoid based on the HMBC data in which a double bond between C-1' and C-6' was identified based on cross-peaks of H-7'/C-1', C-2', and C-6', and H-2'/C-1'; two peaks representing oxygenated carbons at $\delta_{\rm C}$ 146.4 (s) and 78.5 (d) were assigned to C-6' and C-3' based on the cross-peaks of H-3'/C-2' and C-8', as well as a C(4')-C(8')-O-C(2') bridge that was identified based on the cross-peak of H-2'/C-8'. The absences of the two methyl carbon signals from C-7 and C-7' in menthanetype monoterpenoid fragments, and the ¹H-¹H COSY correlation of H-7/H-7' confirmed the connection between components I and II via a C(7)-C(7') bond. Combined with the molecular formula C₃₈H₄₄O₈, a linkage of C(1)-O-C(6') was determined to be a part of a six-membered ring. Hence, the planar structure of 1 involves an unprecedented carbon skeleton of two menthanetype monoterpenoids.

In the above planar structure, H-2/H-4 and H-2'/H-4' are at the bridgeheads of bicyclo[3,2,1] ring systems and are splayed out from the bicycle in opposite directions, indicate α -orientations for H-2/H-4 and H-2'/H-4', respectively. In the ROESY spectrum, the cross-peaks of H-3/H-5 and H-3'/H-5', allowed the assignment of α -orientations for H-3 and H-3', respectively. Furthermore, presence of the H-2/H-7 β [$\delta_{\rm H}$ 1.73 (1H, ddd, J = 13.2, 11.4, 4.8 Hz] cross-peak and the absence of the H-3/H-7 cross-peak suggested a β-orientation for C-7 bound to C-1. The relative configuration of components I and II was assigned as shown in Fig. 2 by the ROESY cross-peak of H-2/H-9'. Hence, the relative configuration of 1 was determined to be 1S*,2R*,3R*,4R*,2'S*,3'R*,4'R*.

To assign the absolute configuration of 1, a further derivatization was carried out for obtaining a mono-hydroxyl derivative. 1 was treated with methanesulfonyl chloride (MsCl), to promote the novel elimination reaction yielding 1a (Scheme 1).

Compound 1a, a white powder, has a molecular formula of $C_{38}H_{44}O_8$ based on analysis of its HRESIMS m/z 311.1253 (calcd for C₁₇H₂₀O₄Na 311.1254). Its ¹H and ¹³C NMR data showed the presence of a benzene ring, two methyl groups, one carbonyl, and four oxygenated alkyl groups. Analysis of coupling constants in the ¹H NMR data in the low-field region (Table 1) showed an ortho-disubstituted benzene ring. The presence of this structural feature is supported by the ¹H-¹H COSY correlations of H-3'/H-4', H-4'/H-5', and H-5'/H-6', and the HMBC cross-peaks of H-3'/C-4', C-5', C-2' and C-1' (Fig. 2). Comparing the rest of the NMR data of 1a with that of 1, they shared similar signals from C-1-C-7, indicating that 1a possessed the same menthane-type monoterpenoid as component I (purple), which is supported by the HMBC correlations shown in Fig. 2. The

¹H-¹H COSY cross-peak of H-7/H-7' and the HMBC cross-peaks of H-7'/C-1', C-6', and C-2' suggested the two components are linked by a bond at C(7)–C(7'). Combing that information with the molecular formula, leads to the conclusion that C-2' and C-1 are linked by C(1)-O-C(2'). With that, the planar structure of 1a was elucidated. Based on comparison of structures of 1 and 1a, 1a is formed from 1 by an elimination of component II, and therefore, the relative configuration of 1a is same as component I of 1; this is supported by the ROESY cross-peak of H-3/H-5. This is a novel reaction for the construction of a benzene ring from a menthane unit.

In the present study, the NMR spectrum of 1a was better resolved than the spectrum of 1; having more separation between high-field and low-field peaks in the spectrum of 1a also helped to confirm the structure of 1. Besides that, this elimination could produce the mono-hydroxyl compound 1a, which indicates that Mosher's method is an option for determination the absolute configuration.

The absolute configuration of 1a was assigned using Mosher's method; the treatment of **1a** with (*R*)- and (*S*)-(α)methoxy(trifluoromethyl)phenylacetic acid (MTPA-OH) yielded (R)- and (S)-MTPA esters, respectively. The values of the ¹H NMR $\Delta \delta_{S-R}$ for the mono-MTPA esters (shown in Fig. 3) revealed an R absolute configuration at C-3 of 1a, implying the absolute configuration of 1 is 1S,2R,3R,4R,2'S,3'R,4'R. Therefore, the structure of 1 was solved as shown and named dimericilligerate E.

Compound 2, isolated as a white amorphous powder, possessed a molecular formula of C₂₀H₃₀O₇, based on HRESIMS m/z 381.1917 [M – H]⁻ (calcd for C₂₀H₂₉O₇, 381.1919), ¹³C NMR and DEPT spectra. The ¹H NMR spectrum showed four methyl groups; the ¹³C NMR and DEPT spectra showed two carbonyls and seven oxygenated alkyl carbons located downfield. Comparing the NMR data of 2 with that of 1 reveals that 2 has a similar dimeric monoterpenoid structure. Two partial structural components, I (in purple) and II (in red), were then constructed based on analysis of NMR data. Component I of 2 was found to be same component I of 1 through detailed comparison of the NMR data and the HMBC cross-peaks of H-2/C-1, C-3, C-4, C-6, and C-8, H-3/C-4 and C-8, and H-5/C-6, as well as the cross-peak of H-4/H-5 in ¹H-¹H COSY spectrum (Fig. 4). Analysing the remaining NMR data, a six-membered menthanetype monoterpenoid was identified based on the HMBC crosspeaks of CH₃-9',10'/C-8' and C-4', and H-7'/C-1' and C-2'. The HMBC correlations of H-3'/C-6, H-4'/C-6', and H-2'/C-6' allowed C-6' to be assigned as C=O, and the HMBC cross-peaks of H-3'/C-2' and H-2'/C-4' indicated that two of the oxygenated





Scheme 1 Derivative reaction of dimericilligerate E (1)



R = (R)/(S)-MTPA

Fig. 3 $\Delta \delta_{S-R}$ of ¹H-NMR for S- and R-MTPA esters of compound **1a**.

methines were located at C-3' and C-2'. The linkage of components I and II via the bonds of C(7)-C(7') and C(2')-O-C(1), forming a stable six-membered tetrahydropyran ring, was determined based on the key ¹H-¹H COSY interaction of H-7/H-7' and the HMBC correlation of H-2'/C-1. Thus, the planar structure of 2 was identified as an ester of phenylpropionic acid and novel dimeric monoterpenoids composed of a bicyclo[3,2,1] group and a six-membered menthane-type monoterpenoid.

The relative configuration of 2 was assigned based on a ROESY experiment, which showed correlations of H-3/H-5 and H-2/H-7 β [$\delta_{\rm H}$ 1.43 (1H, m)] and lacked a correlation of H-3/H-7, indicating a $\beta,\alpha,\alpha,\alpha$ -orientation of CH₂-7, H-2, H-3, and H-4, respectively. Meanwhile, H-1' β , H-3' α , and H-4' α configurations were assigned based on the existence of cross-peaks of H-1'/H-2 and CH₃-9' and the absence of a cross-peak of H-1'/H-3'; and an H-2' α configuration was deduced by the cross-peaks of H-2'/H-7 α [$\delta_{\rm H}$ 1.93 (1H, m)], and H-4'. Therefore, the relative configuration of 2 was determined to be 1S*,2R*,3R*,4R*,1'R*,2'R*,3'S*,4'S*.

Two possible configurations, (1S, 2R, 3R, 4R, 1'R, 2'R, 3'S, 4'S)and (1R,2S,3,4S,1'S,2'S,3'R,4'R), were optimized using the density functional theory (DFT) method at the B3LYP/6-31G(d,p) level.²¹ Subsequently, the ECD was calculated using time-dependent density functional theory (TDDFT) at the B3LYP/6-31G(d,p) level with the PCM solvent model²² and simulated by SpecDic 1.64.23,24 The calculated ECD curve of the (1S,2R,3R,4R,1'R,2'R,3'S,4'S)-isomer (Fig. 5A) resembled the experimental ECD curve, indicating that the absolute configuration of 2 is (1S,2R,3R,4R,1'R,2'R,3'S,4'S). The structure of 2 was thus established as depicted and named dimericilligerate F.

в



Experimental and calculated ECD spectra of 1-5. Fia. 5

Analysis of non-fermented and fermented I. aromatica

To find the precursors of dimericilligerates E(1) and F(2), HPLC-ESIMS was used to identify possibilities from the different extracted fractions of the original I. aromatica. The peaks corresponding to 1 and 2 were not detected in any fraction; and two major peaks at m/z 627.4 [M – H]⁻ and 513.2 [M – H^{-} were observed in the petroleum ether extract (Fig. S52[†]). Therefore, 1 and 2 might be produced from the components of 627.4 $[M - H]^-$ and 513.2 $[M - H]^-$. Inspired by that, three compounds, 3-5, were isolated from the petroleum ether extract and identified by 1D/2D NMR and ECD spectroscopy.

Identification of precursors

Compound 3, isolated as a colourless gummy oil, was found to have a molecular formula of C38H44O8 based on analysis of its HRESIMS m/z 651.2920 $[M + Na]^+$ (calcd for $C_{38}H_{44}O_8Na$ 651.2928), ¹³C NMR and DEPT spectra, which indicated seventeen degrees of unsaturation. The ¹H and ¹³C NMR data (Table 2) showed four methyl groups, two benzene rings, three carbonyls (one ketone and two esters), and seven oxygenated alkyl groups. Comparing the NMR data with phenylpropionic acid^{25,26} and monoterpenoid phenylpropionic acid esters¹⁰ previously isolated from *I. aromatica* indicated that 3 contains two phenylpropionic esters. Comparison of the rest of the NMR data of 3 with that of 1, suggests that the rest of the structure of 3 is same as 1, which is supported by the ${}^{1}H{}^{-1}H$ COSY and HMBC correlations shown in Fig. 6. The cross-peaks of H-3/C-9" and H-3'/C-9" revealed two phenylpropionic units attached to C-3 and C-3', respectively. Hence, the planar structure of 3 was established as shown.

The relative configurations were assigned by a ROESY experiment. The ROESY cross-peaks of H-3/H-5 and H-3'/H-5', allowed the assignment of an α -orientations for H-3 and H-3', respectively. Furthermore, the presence of a H-2/H-7 β [$\delta_{\rm H}$ 2.03 (1H, m)] cross-peak and the absence of a H-3/H-7 cross-peak suggested a β-orientation of C-7 linked at C-1. Combined with the ROESY cross-peak of H-2/H-9', the relative configuration of 3 was determined to be 1S*,2R*,3R*,4R*,2'S*,3'R*,4'R*, which is same as 1.

Compound 4, isolated as a colourless gummy oil, has a molecular formula of $C_{29}H_{36}O_7$ based on the HRESIMS m/z519.2355 $[M + Na]^+$ (calcd for C₂₉H₃₆O₇Na 519.2353), ¹³C NMR and DEPT spectra. From the ¹H and ¹³C NMR data, four methyl groups, one benzene ring, two carbonyls (one ketone and one ester), and seven oxygenated alkyl groups were identified. Comparison of the NMR data of 4 with that of dimeric menthane-type monoterpenoid 3 (Table 2) revealed that 4 possessed a same skeleton as 3, which was also supported by the 2D NMR analysis shown in Fig. 6. The main difference is that only one phenylpropionic substituent is found in 4, while 3 has two. Based on the cross-peak of H-3'/C-9" found in the HMBC spectrum, the phenylpropionic unit was attached at C-3'. The relative configuration of 4 was determined to be $1S^{*}, 2R^{*}, 3R^{*}, 4R^{*}, 2'S^{*}, 3'R^{*}, 4'R^{*}$ (which is the same as 1 and 3) based on its ROESY cross-peaks of H-3/H-5, H-3'/H-5', and H-2/ H-7 β [$\delta_{\rm H}$ 2.14 (1H, ddd, J = 13.8, 11.4, 5.4 Hz)].



2*R*,3*R*,4*R*,1'*R*,2'*R*,3'*S*,4'

	3		4		5	
No.	$\delta_{ m C}$	δ _H mult. (J in Hz)	$\delta_{ m C}$	$\delta_{\rm H}$ mult. (<i>J</i> in Hz)	$\delta_{ m C}$	δ _H mult. (J in Hz)
1	79.7 C		79.6 C		78.3 C	
2	81.0 CH	4.10 s	84.7 CH	4.11 d (1.2)	81.6 CH	4.16 s
3	78.5 CH	5.69 s	77.4 CH	4.88 s	82.1 CH	5.67 s
4	45.5 CH	2.28 s	47.5 CH	2.33 s	46.2 CH	2.39 d (7.2)
5	42.5 CH ₂	2.63 br s	42.7 CH ₂	2.62 d (3.0), 2.59 d (3.0)	$44.8~\mathrm{CH}_2$	2.79 d (2.4), 2.59 dd (7.2, 2.4)
6	205.0 C		205.9 C		210.9 C	
7	19.8 CH ₂	2.05 m, 2.03 m	23.5 CH ₂	2.11 ddd (13.8, 11.4, 5.4), 1.73 ddd (13.8, 11.4, 5.4)	19.8 CH ₂	1.90 m, 1.81 m
8	84.1 C		84.3 C		84.5 C	
9	26.4 CH ₃	1.09 (3H, s)	27.0 CH ₃	1.19 s	29.7 CH ₃	1.33 s
10	29.1 CH ₃	1.25 (3H, s)	29.8 CH ₃	1.53 s	26.7 CH ₃	1.14 s
1'	108.4 C		108.7 C		56.1 CH	
2'	79.3 CH	3.94 s	79.5 CH	4.03 s	84.1 CH	6.74 d (6.0)
3'	80.1 CH	5.02 s	80.3 CH	5.11 s	77.6 CH	4.70 dd (6.0, 3.2)
4'	46.4 CH	2.18 s	46.6 CH	2.29 d (15.0)	47.2 CH	1.93 dd (10.0, 3.2)
5'	32.7 CH ₂	2.41 m, 2.17 m	32.9 CH ₂	2.46 dd (15.0, 3.6), 2.22 d (3.6)	41.8 CH ₂	2.84 dd (10.0, 4.0) 2.53 d (4.0)
6′	146.3 C		146.0 C		210.9 C	
7′	22.6 CH ₂	2.04 m, 1.70 ddd (8.0, 5.6, 2.4)	20.2 CH ₂	2.19 ddd (13.8, 11.4, 5.4), 2.14 ddd (13.8, 11.4, 5.4)	29.7 CH ₂	2.33 m, 2.22 m
8'	82.2 C		82.2 C		83.9 C	
9′	26.7 CH ₃	1.12 s	26.9 CH ₃	1.30 s	29.7 CH ₃	1.44 s
10'	30.2 CH ₃	1.19 s	30.4 CH ₃	1.28 s	27.1 CH_3	1.24 s
1''	140.0 C		140.5 C		140.2 C	
2'', 6''	128.3 CH	7.13 d (overlapped, olp)	128.5 CH	7.20 d (olp)	128.4 CH	7.23 d (olp)
3", 5"	128.5 CH	7.19 dd (7.2, 6.0)	128.7 CH	7.28 dd (7.2, 3.6)	128.0 CH	7.28 dd (8.2, 2.4)
4''	126.3 CH	7.12 dd (olp)	126.5 CH	7.19 d (olp)	126.6 CH	7.21 dd (olp)
7″	30.8 CH ₂	2.87 t (7.2)	31.0 CH ₂	2.95 t (8.1)	30.9 CH ₂	2.99 t (8.8)
8″	36.0 CH ₂	2.59 t (7.2)	36.3 CH ₂	2.66 t (8.1)	36.2 CH ₂	2.73 t (8.8)
9″	171.9 C		172.5 C		172.4 C	
1‴	140.3 C					
2‴, 6‴	128.3 CH	7.13 d (olp)				
3‴, 5‴	128.6 CH	7.19 dd (7.2, 6.0)				
4‴	126.3 CH	7.12 dd (olp)				
7‴	30.8 CH ₂	2.87 t (7.2)				
8‴	36.2 CH ₂	2.59 t (7.2)				
9‴	172.4 C					

Because 3 and 4 shared skeletal features with 1, they possessed similar ECD spectra (Fig. 5B), indicating the absolute configurations of both 3 and 4 are 1S,2R,3R,4R,2'S,3'R,4'R. The structure of 3 and 4 were therefore identified and named dimericilligerates A and B, respectively.

Compound 5, obtained as a colourless gummy oil, has a molecular formula $C_{29}H_{38}O_8$ based on its HRESIMS m/z559.2552 [M + COOH]⁻ (calcd for $C_{30}H_{39}O_{10}$ 559.2549), ¹³C NMR and DEPT spectra, which showed eleven degrees of unsaturation. The ¹H NMR spectrum showed four methyl groups and five aromatic protons; the ¹³C NMR and DEPT spectra showed three carbonyls (two ketones and one ester), and seven oxygenated alkyl carbons located downfield. Comparing the NMR data of 5 with that of 3 and 4 implied that 5 consisted of one phenylpropionic substituent and one similar dimeric monoterpenoid. The dimeric monoterpenoid was confirmed to be 2 by comparing their NMR data and analysing their respective ¹H–¹H



Fig. 6 Key ${}^{1}H-{}^{1}H$ COSY, HMBC, and ROESY correlations of **3–5**.

		Cytotoxic activity (IC ₅₀ /µM)						
Sample	NO production (IC ₅₀ / μ M)	HL-60	A-549	SMMC-7721	MCF-7	SW480		
1	>25	18.42 ± 0.35	27.40 ± 1.11	3.89 ± 0.23	17.29 ± 2.98	19.41 ± 0.76		
2	23.61 ± 2.77	19.83 ± 1.54	27.47 ± 0.98	14.21 ± 0.87	16.74 ± 2.98	16.63 ± 0.83		
3	>25	>40	>40	>40	>40	>40		
4	>25	>40	>40	>40	>40	>40		
5	>25	>40	>40	>40	>40	>40		
MG132 ^a	0.24 ± 0.02	_	_	_	_	_		
DDP^{b}	_	1.72 ± 0.22	6.82 ± 0.30	12.19 ± 0.85	17.40 ± 0.93	16.63 ± 2.90		
$Taxol^b$		<0.008	<0.008	<0.008	<0.008	<0.008		

^{*a*} Positive control for inhibitory effect on NO production in RAW 264.7 at the concentration of 50 μM. ^{*b*} DDP (*cis*-dichlorodiammineplatinum) and taxol used as the positive control for cytotoxic activity.

COSY and HMBC correlations (Fig. 6). The relative configuration of 5 was assigned as $1S^*$, $2R^*$, $3R^*$, $4R^*$, $1'R^*$, $2'R^*$, $3'S^*$, $4'S^*$ because of its ROESY cross-peaks of H-3/H-5; H-2/H-7 β [$\delta_{\rm H}$ 1.56 (1H, dd, J = 13.8, 7.2 Hz)]; H-1'/H-2 and CH₃-9'; H-1'/H-3' and H-2' α ; and H-2'/H-7 α [$\delta_{\rm H}$ 2.03 (1H, dd, J = 13.8, 7.2 Hz)] and H-4', which is the same as **2**.

Subsequently, the theoretical ECD spectra were calculated using TDDFT at the B3LYP/6-31G(d,p) level. The results showed that the theoretical ECD curve of the (1S,2R,3R,4R,1'R,2'R,3'S,4'S)isomer (Fig. 5C) resembled the experimental ECD curve, indicating an absolute configuration of (1S,2R,3R,4R,1'R,2'R,3'S,4'S)for 5. The structure of 5 was thus established as depicted, and 5 was named dimericilligerate C.

Bioactivity

Compounds 1–5 were tested for cytotoxic activity^{27,28} and an inhibitory effect on NO production in LPS-stimulated RAW 264.7 cells.^{29,30} As shown in Table 3, compound 2 showed moderate inhibitory effects on NO production. Both 1 and 2 possessed cytotoxic activities against five cell lines. In particular, 1 possessed potent cytotoxic activity against a liver hepatocellular carcinoma cell line (SMMC-7721) with an IC₅₀ value of $3.89 \pm 0.23 \ \mu$ M, which was more potent than positive control (*cis*-dichlorodiammineplatinum). However, the phenyl-propionic esters did not show any activity.

Verification the biotransformation

To confirm the biotransformation of monoterpenoid dimer phenylpropionic esters (3-5) into free monoterpenoid dimers (1and 2), the main compound 3 was selected as the substrate for fermentation. After 30 days of solid-state fermentation, 3 was almost completely transformed to 1, indicating that *C. rogersoniana*-fermentation of *I. aromatica* was effective at producing cytotoxic free monoterpenoid dimers.

Conclusions

In the present study, five monoterpenoid dimer derivatives (1-5) were isolated from fermented and non-fermented *C. rogersoniana*. Compounds 1 and 2 isolated from fermented *I.*

aromatica showed stronger cytotoxic activity than those isolated from the original material. This paper presents a novel class of monoterpenoid dimers and suggests that *C. rogersoniana*-fermented *I. aromatica* is effective at the production of cytotoxic dimeric monoterpenoids from inactive original materials.

Experimental

General procedures

Melting points were determined on an XRC-1 melting point apparatus (Optical Instrument Factory of Sichuan University, Chengdu, China) and are uncorrected. Optical rotations were measured with a Jasco P-1020 digital polarimeter (Jasco, Tokyo, Japan). A Shimadzu UV-Vis 2550 spectrometer (Shimadzu, Tokyo, Japan) was used for the acquisition of UV-Vis spectra. Electronic circular dichroism spectra were recorded using a Chirascan circular dichroism spectrometer (Applied Photophysics Ltd., UK). NMR spectra were recorded on a Bruker Avance 400 MHz spectrometer or Bruker Avance III 600 MHz spectrometer (Bruker, Karlsruhe, Germany) using TMS as an internal reference. HRESIMS and ESIMS analyses were carried out with an Agilent G3250AA MSD TOF-MS or Agilent 6540 Q-TOF-MS (Agilent, Santa Clara, CA, USA). Silica gel (200-300 mesh, Qingdao Marine Chemical Group Co., Qingdao, China) was used for column chromatography (CC). Fractions were monitored by TLC and visualized by spraying with a 10% H₂SO₄-EtOH solution.

Plant material

The tubers of *I. aromatica* were collected from Wenshan, Yunnan, China, in December 2015 and were identified by assistant professor Shu-Da Yang from School of Pharmacy, Kunming Medical University, Kunming, China. A voucher specimen (2015ia-01) has been deposited in School of Chemical Science and Technology, Yunnan University, Kunming, China.

Microorganism and fermentation processing of I. aromatica

C. rogersoniana 828H2 was obtained from the Yunnan Institute of Microbiology, Yunnan Province, China. The nucleotide

sequence data have been deposited in GenBank (accession number KT625993).

PDA (1 L water, 200 g potato, 20 g dextrose, and 15 g agar) slant culture media were inoculated with the above fungus and incubated in a constant temperature incubator at 28 °C for 5 days. Afterwards, the mature fungus was cultivated in PDB (1 L water, 200 g potato, 20 g dextrose) media for 5 days at 28 °C as seed culture mediums.

The dried and powdered tubers (2.0 kg) were added to 350 mL glass spawn bottles (40 g per bottle) as the SSF culture media. After being treated with 8 mL water and sterilized at 121 $^{\circ}$ C for 30 min, the seed culture media were added and incubated at 28 $^{\circ}$ C for 30 days.

Extraction and isolation

Extraction and isolation of fermented I. aromatica. After 30 days of fermentation, the I. aromatica was immersed in 95% EtOH at room temperature for 24 h, then extracted at 60 °C three times (2 h each). The removal of solvents under reduced pressure afforded the crude product (250 g). The crude mixture was extracted with petroleum ether (PE), ethyl acetate (EA), and *n*-butanol successively to obtain three extracts (E_{PE} , E_{EA} , and $E_{n-butanol}$). E_{EA} (23 g) was chromatographed over Biotech Flash 65i silica gel CC using a MPLC (Biotage, Uppsala, Sweden) eluting with a CH₃Cl-MeOH gradient system (100% CH₃Cl, 1.5 CV; 2% MeOH-CH₃Cl, 1.5 CV; 5% MeOH-CH₃Cl, 1.5 CV; 10% MeOH-CH₃Cl, 1.5 CV; 20% MeOH-CH₃Cl, 1.5 CV; 35% MeOH-CH₃Cl, 1.5 CV; 50% MeOH-CH₃Cl, 1.5 CV; and 2% MeOH-CH₃Cl, 1.5 CV). Absorbance at 220 nm and TLC were used to identify fractions EA1-5. Further purification of fraction EA2 (2.0) was accomplished by repeated silica gel (PE-acetone) and Sephandex LH-20 (methanol) CC to afford 2 (10.0 mg). EA3 (5.0 g) was separated by repeated silica gel CC (PE-acetone and CH₃Cl-MeOH) and purified by Sephadex LH-20 (methanol) CC to afford 1 (105.0 mg).

Extraction and isolation of non-fermented I. aromatica. The dried and powdered tubers of I. aromatica (2.5 kg) were immersed in 95% EtOH at room temperature for 24 h and extracted at 60 $^\circ C$ three times (2 h each). The removal of the solvent under reduced pressure afforded the crude extract (500 g). Afterwards, the crude mixture was dispersed in water and extracted with PE, EA, and *n*-butanol successively to obtain three extracts (E_{PE} , E_{EA} , and $E_{n-butanol}$). E_{PE} (100 g) was further subjected to Sephadex LH-20 gel CC (chloroform-methanol, 1:1) to give fractions PE1-3. PE2 (5.0 g) was chromatographed over Sephadex LH-20 gel (chloroformmethanol, 1:1) to yield PE2-1, PE2-2, and PE2-3. PE2-2 (220 mg) was further separated by repeated silica gel CC (petroleum ether-acetone) to afford 4 (3.5 mg) and 5 (1.0 mg). Further purification of PE3 (100 mg) was accomplished by repeated silica gel (petroleum ether-acetone) and Sephadex LH-20 (chloroformmethanol, 1:1) CC to yield 3 (50 mg).

Dimericilligerate *E* (1). White amorphous powder; mp 109–110 °C; $[\alpha]_{D}^{25}$ –104.2 (*c* = 0.19, MeOH); UV (CH₃OH) λ_{max} (log ε) 212 (3.88), 262 (sh, 2.86), 310 (1.76); FT-IR: $\nu_{KBr max}$ (cm⁻¹) 3432 (O–H), 2972, 2933, 1711, 1634, 1369, 1095, 1049; ¹H and ¹³C NMR data, see Table 1; HRESIMS *m*/*z* 409.2235 [M + COOH]⁻ (calcd for C₂₂H₃₃O₇ 409.2232).

Dimericilligerate F (2). White amorphous powder; mp 115–118 °C; $[\alpha]_D^{25}$ –79.1 (c = 0.1, MeOH); UV (CH₃OH) λ_{max} (log ε) 198 (3.81), 267 (3.01), 319 (2.68); ¹H and ¹³C NMR data, see Table 1; HR-ESI-MS *m/z* 381.1917 [M – H]⁻ (calcd for C₂₀H₂₉O₇ 381.1919).

Dimericilligerate A (3). Colourless gummy oil; $[\alpha]_{D}^{20} - 29.2 (c = 0.17, CH_3CN)$; UV (CH₃CN) λ_{max} (log ε) 210 (4.07), 280 (3.09), 328 (2.06); ¹H and ¹³C NMR data, see Table 2; HRESIMS *m*/*z* 651.2920 [M + Na]⁺ (calcd for C₃₈H₄₄O₈Na 651.2928).

Dimericilligerate *B* (4). Colourless gummy oil; $[\alpha]_{D}^{20}$ -57.5 (*c* = 0.12, CH₃CN); UV (CH₃CN) λ_{max} (log ε) 210 (4.17), 281 (3.24), 329 (2.45); ¹H and ¹³C NMR data, see Table 2; HRESIMS *m/z* 519.2355 [M + Na]⁺ (calcd for C₂₉H₃₆O₇Na 519.2353).

Dimericilligerate C (5). Colourless gummy oil; $[\alpha]_{D}^{20}$ –9.0 (c = 0.10, CH₃CN); UV (CH₃CN) λ_{max} (log ε) 210 (4.13), 274 (3.39), 325 (2.44); ¹H and ¹³C NMR data, see Table 2; HRESIMS *m/z* 559.2552 [M + COOH]⁻ (calcd for C₃₀H₃₉O₁₀ 559.2549).

Preparation of compounds 1a

Triethylamine (11 µL) was added to a stirred solution of **1** (10 mg, 0.027 mmol) in CH₂Cl₂ (2 mL) at 0 °C. After 15 min, 20 µL of methanesulfonyl chloride (MsCl) was added and the mixture stirred at room temperature for 18 h, after which, the reaction was quenched by the addition of 5 µL of water. The mixture was concentrated under vacuum and dried over anhydrous Na₂SO₄. The extract was further separated by silica gel CC (petroleum ether–acetone, $8: 1 \rightarrow 5: 1$) to yield **1a** (6.0 mg, 77%).

White amorphous powder; mp 99–102 °C; $[\alpha]_{D}^{20}$ –92.2 (c = 0.18, CH₃CN); UV (CH₃CN) λ_{max} (log ε) 198 (4.43), 219 (3.99), 274 (3.22) nm; ¹H and ¹³C NMR data, see Table 1; HRESIMS m/z 311.1253 [M + Na]⁺ (calcd for C₁₇H₂₀O₄Na 311.1254).

Preparation of MTPA esters of 1a

To a stirred solution of **1a** (2.0 mg, 0.007 mmol) in CH_2Cl_2 (2 mL), either (*S*)-(–)-MTPA-OH (5 µL, 0.029 mmol) or (*R*)-(+)-MTPA-OH (5 µL, 0.029 mmol) was added. Then, 10.0 mg of *N*,*N'*-diisopropylcarbodiimide and 10.0 mg of 4-dimethylaminopyridine were added. The reaction mixture was stirred at room temperature for 30 min and concentrated under vacuum at 45 °C. The residues were purified by silica gel CC (petroleum ether–acetone) to afford (*S*)-MTPA ester (2.5 mg, 70.9%) or (*R*)-MTPA ester (2.0 mg, 56.7%), respectively.

(*R*)-MTPA ester of 1a. White amorphous powder; ¹H NMR (400 MHz, CDCl₃): $\delta_{\rm H}$ 4.28 (1H, s, H-2), 6.11 (1H, s, H-3), 2.43 (1H, d, J = 3.0 Hz, H-4), 2.35 (1H, ddd, 10.4, 6.0, 4.4 Hz, H-7a), 2.01 (1H, ddd, J = 10.4, 6.0, 4.4 Hz, H-7b), 2.96 (1H, ddd, J = 10.4, 7.2, 4.4 Hz, H-7'a), 2.72 (1H, ddd, J = 10.4, 7.2, 4.4 Hz, H-7'b), 1.18 (3H, s, H-9), 1.01 (3H, s, H-10); HRESIMS *m*/*z* 527.1664 [M + Na]⁺ (calcd for C₂₇H₂₇O₆F₃Na 527.1652).

(S)-MTPA ester of 1a. White amorphous powder; ¹H NMR (400 MHz, CDCl₃): $\delta_{\rm H}$ 4.18 (1H, s, H-2), 6.19 (1H, s, H-3), 2.54 (1H, d, J = 3.2 Hz, H-4), 2.37 (1H, ddd, J = 10.0, 6.6, 4.4 Hz, H-7a), 1.99 (1H, ddd, J = 10.0, 6.6, 4.4 Hz, H-7b), 2.95 (1H, ddd, J = 10.0, 7.4, 4.4 Hz, H-7'a), 2.75 (1H, ddd, J = 10.0, 7.4, 4.4 Hz, H-7'a), 1.29 (3H, s, H-10); HRESIMS *m*/*z* 527.1668 [M + Na]⁺ (calcd for C₂₇H₂₇O₆F₃Na 527.1652).

LC-MS analysis

The dried and powdered tubers (2.0 g) were extracted with petroleum ether using an ultrasonic cleaner three times (30 min each). Afterwards, the solvent was removed under reduced pressure at 40 °C to afford the crude extract. Then, the crude extract was dissolved in acetone and filtrated through a 0.45 µm filter for LC-MS analysis. An LC-MS system equipped with an Agilent 1100 Series HPLC system, an Agilent G3250AA MSD TOF-MS, and Agilent Zorbax SB-C18 column (250×4.6 mm, 5 μm i.d.; Agilent, Santa Clara, CA, USA) were used to identify the isolated compounds. A gradient elution system consisting of solvents A (water) and B (acetonitrile) was used for the analysis, and the gradient program was as follows: 0-5 min, 10% B; 5-20 min, 10-60% B; and 20-30 min, 60-100% B. The flow rate was 0.8 mL min⁻¹, the column temperature was set to 25 °C, and the injection volume was 10 µL. The ion source of the MS was set to ESI- to monitor quasi-molecular ions.

Cytotoxic activity

Cytotoxicity was assayed by the trypan blue method²⁷ using the human leukaemia cell line (HL-60) and by the MTT assay²⁸ using the liver hepatocellular carcinoma cell line (SMMC-7721), lung carcinoma cell line (A-549), breast carcinoma cell line (MCF-7), and colon carcinoma cell line (SW480). The cell lines were purchased from America Type Culture Collection (ATCC; Rockville, MD, USA) and cultured in RPMI-1640 medium (Gibco, New York, NY, USA) supplemented with 100 U mL⁻¹ penicillin, 100 μ g mL⁻¹ streptomycin, 1 mM glutamine and 10% heat-inactivated foetal bovine serum (Gibco). *cis*-Dichlorodiammineplatinum (DDP) and taxol (Sigma-Aldrich Co. LLC., Shanghai, China) were selected as the positive controls. Each sample was repeated in triplicate.

NO production in RAW 264.7 macrophages

NO production in RAW 264.7 Macrophages was measured according to the method reported in the literature.^{29,30} Murine monocytic RAW 264.7 macrophages (Institute of Biochemistry and Cell Biology, SIBI, CAS, Shanghai, China) were seeded in 96-well cell culture plates (2×10^5 cells per well) and treated with serial dilutions of the compounds at a maximum concentration of 25 μ M; this treatment was followed by stimulation with LPS (1 μ g mL⁻¹; Sigma-Aldrich) for 18 h. NO production in the supernatant was assessed using Griess reagents (Sigma-Aldrich). The absorbance at 570 nm was measured using a 2104 Envision multilabel plate reader (PerkinElmer Life Sciences, Inc., Boston, MA, USA). MG132 (Sigma-Aldrich) was used as a positive control. The viability of RAW 264.7 cells were simultaneously evaluated using the MTT assay to exclude the interference of the cytotoxicity of the test compounds.

ECD calculation

The calculation of theoretical ECD spectra of compounds **3** and **4** were performed using the Gaussian Program by Yunnan Electronic Computing Center. The possible geometries were previously optimized using DFT method at the B3LYP/6-

31G(d,p) level,²¹ and the excitation energies and rotational strengths were calculated using TDDFT at the B3LYP/6-31G(d,p) level in methanol with PCM model.²² Afterwards, the ECD spectra were generated using SpecDis.^{23,31}

Acknowledgements

This work was financially supported by two grants from the Natural Science Foundation of China (No. 81660719 and 81460648), a project of Yunling Scholars of Yunnan province, and a grant from New Academic Researcher Award for Doctoral Candidates of Yunnan Province.

Notes and references

- 1 Flora of China Editorial Committee, *Flora Reipublicae Popularis Sinicae*, Science Press, Beijing, 1982.
- 2 K.-S. Chen, Y.-C. Wu, C.-M. Teng, F.-N. Ko and T.-S. Wu, J. Nat. Prod., 1997, 60, 645–647.
- 3 S. A. Ross, R. D. Minard, M. Shamma, M. O. Fagbule, G. Olatunji and Z. Gbile, *J. Nat. Prod.*, 1985, **48**, 835–836.
- 4 J.-J. Chen, H.-C. Hung, P.-J. Sung, I.-S. Chen and W.-L. Kuo, *Phytochemistry*, 2011, 72, 523–532.
- 5 J.-J. Chen, T.-H. Lee, W.-L. Kuo, P.-J. Sung, I.-S. Chen, C.-W. Shu, M.-J. Cheng, T.-C. Wang, Y.-P. Lim and R. Xu, *Chem. Nat. Compd.*, 2015, **51**, 739–742.
- 6 C.-H. Huang, Y.-Y. Chan, P.-C. Kuo, Y.-F. Chen, R.-J. Chang, I.-S. Chen, S.-J. Wu and T.-S. Wu, *Int. J. Mol. Sci.*, 2014, 15, 13424–13436.
- 7 S. Z. Huang, Guihaia, 1985, 5, 17-20.
- 8 S. Mo, Z. Li, Y. Ou, S. Wei, X. Si and D. Fan, *Lishizhen Med. Mater. Med. Res.*, 2006, 17, 2512–2513.
- 9 L. Xie, P. Li, Z. Gong and Y. Ou, *Contemp. Biomed.*, 2011, 17, 31-32.
- 10 J. W. Dong, L. Cai, X. J. Li, J. P. Wang, R. F. Mei and Z. T. Ding, Arch. Pharmacal Res., 2016, DOI: 10.1007/ s12272-016-0860-3.
- 11 Y. Kim, Y. You, H. G. Yoon, Y. H. Lee, K. Kim, J. Lee, M. S. Kim, J. C. Kim and W. Jun, *Food Chem.*, 2014, 151, 148–153.
- 12 M. F. Hsu and B. H. Chiang, *Process Biochem.*, 2009, 44, 83–90.
- J. W. Dong, L. Cai, J. Xiong, X. H. Chen, W. Y. Wang, N. Shen,
 B. L. Liu and Z. T. Ding, *Process Biochem.*, 2015, 50, 8–13.
- 14 Y. Li, L. Cai, J. W. Dong, Y. Xing, W. H. Duan, H. Zhou and Z. T. Ding, J. Agric. Food Chem., 2015, 63, 6596–6602.
- 15 K. Adachi, K. Kanoh, P. Wisespongp, M. Nishijima and Y. Shizuri, *J. Antibiot.*, 2005, **58**, 145–150.
- 16 T.-P. Yin, Y. Xing, L. Cai, J. Yu, P. Luo and Z.-T. Ding, J. Asian Nat. Prod. Res., 2017, 1–6, DOI: 10.1080/ 10286020.2017.1314271.
- 17 L. Cai, J.-W. Dong, L.-X. Zhao, H. Zhou, Y. Xing, Y. Li, Z.-J. Li, W.-H. Duan, X.-J. Li and Z.-T. Ding, *Process Biochem.*, 2016, 51, 933–940.
- 18 X.-J. Li, J.-W. Dong, L. Cai, R.-F. Mei and Z.-T. Ding, *J. Biosci. Bioeng.*, DOI: 10.1016/j.jbiosc.2017.05.012.

- 19 P. Ventura, M. Schiavi and S. Serafini, *Xenobiotica*, 1983, **13**, 139–146.
- 20 K. Hashimoto, T. Katsuhara, K. Niitsu, Y. Ikeya, K. Hayashi, M. Maruno and T. Fujita, *Phytochemistry*, 1994, 35, 969–973.
- 21 J. W. Dong, L. Cai, X. J. Li, L. Peng, Y. Xing, R. F. Mei, J. P. Wang and Z. T. Ding, *Fitoterapia*, 2016, **109**, 212–216.
- 22 Y. Liao, X. Liu, J. Yang, Y. Z. Lao, X. W. Yang, X. N. Li, J. J. Zhang, Z. J. Ding, H. X. Xu and G. Xu, *Org. Lett.*, 2015, 17, 1172–1175.
- 23 T. Bruhn, A. Schaumlöffel, Y. Hemberger and G. Bringmann, *Chirality*, 2013, **25**, 243–249.
- 24 S. Lu, S. Nishimura, G. Hirai, M. Ito, T. Kawahara, M. Izumikawa, M. Sodeoka, K. Shin-ya, T. Tsuchida and H. Kakeya, *Chem. Commun.*, 2015, **51**, 8074–8077.

- 25 T. Ohishi, L. Zhang, M. Nishiura and Z. Hou, *Angew. Chem., Int. Ed.*, 2011, **50**, 8114–8117.
- 26 P. Devi, S. Wahidullah, C. Rodrigues and L. D. Souza, *Mar. Drugs*, 2010, 8, 1203–1212.
- 27 F. Wang, H. Hua, Y. Pei, D. Chen and Y. Jing, J. Nat. Prod., 2006, 69, 807–810.
- 28 K. B. Wang, Y. T. Di, Y. Bao, C. M. Yuan, G. Chen, D. H. Li, J. Bai, H. P. He, X. J. Hao, Y. H. Pei, Y. K. Jing, Z. L. Li and H. M. Hua, *Org. Lett.*, 2014, **16**, 4028–4031.
- 29 J. Yang, W. G. Wang, H. Y. Wu, X. Du, X. N. Li, Y. Li, J. X. Pu and H. D. Sun, *J. Nat. Prod.*, 2016, **79**, 132–140.
- 30 J. H. Yang, X. Du, F. He, H. B. Zhang, X. N. Li, J. Su, Y. Li, J. X. Pu and H. D. Sun, *J. Nat. Prod.*, 2013, **76**, 256–264.
- 31 P. M. Allard, M. T. Martin, M. E. T. H. Dau, P. Leyssen, F. Gueritte and M. Litaudon, *Org. Lett.*, 2012, 14, 342–345.