



Cite this: *RSC Adv.*, 2017, 7, 45075

Influence of culture conditions on extracellular polysaccharide production and the activities of enzymes involved in the polysaccharide synthesis of *Nostoc flagelliforme*

Pei-pei Han,^{ID}†* Shun-yu Yao,† Rong-jun Guo, Rong-rong Yan, Yi-kai Wu, Shi-gang Shen and Shi-ru Jia*

Culture conditions significantly influence extracellular polysaccharide (EPS) production of *Nostoc flagelliforme*, however, the key enzyme controlling EPS synthesis has not been fully explored yet. The influence of different culture conditions including light quality, carbon source and nitrogen source on EPS production of *N. flagelliforme* and activities of EPS synthesis enzymes was investigated. Three experimental groups produced higher amounts of EPS than the control group, including the carbon source group with 1.26 g L⁻¹ NaHCO₃, the nitrogen source group with 0 g L⁻¹ NaNO₃ and the light quality group with blue light. Activities of seven related enzymes phosphoglucose isomerase (PGI), fructose-1,6-bisphosphatase (FBPase), UDP-glucose pyrophosphorylase (UGPase), UDP-galactose-4-epimerase (UGE), UDP-glucose dehydrogenase (UGDH), phosphomannose isomerase (PMI), and phosphofructokinase (PFK) were significantly influenced by culture conditions. Partial least-squares analysis and correlation analysis methods were used to analyze the relationship between the activities of these enzymes and EPS production, and a correlation between the production of EPS and the activities of PGI, PMI, FBPase, UGDH, and UGPase was found under different culture conditions. Subsequent analysis of the transcription level of genes encoding the five enzymes showed genes *pgi* and *fbp1* in three experimental groups were significantly up regulated. The results revealed PGI and FBPase might be important enzymes positively influencing the biosynthesis of *N. flagelliforme* EPS. The findings would be helpful to further understand the pathway of EPS biosynthesis aimed to improve the EPS production of *N. flagelliforme*.

Received 20th July 2017
 Accepted 12th September 2017

DOI: 10.1039/c7ra07982f

rsc.li/rsc-advances

Introduction

The cyanobacteria extracellular polysaccharide can be divided into two main forms: the capsule polysaccharide (CPS) fraction bound to the cell surface and the extracellular polysaccharide (EPS) fraction released into the surrounding environment.^{1,2} The EPS shows advantages over the polysaccharides from plants or microalgae, given its characteristic easy recovery from the culture medium and its physicochemical properties.^{3,4} The future looks bright for its potential applications in the food, pharmaceutical and environmental industries, consequently, the interest toward EPS has significantly increased in recent years.⁵⁻⁷

Nostoc flagelliforme is a species of edible terrestrial filamentous cyanobacteria, distributed throughout arid and semi-arid

areas.^{8,9} The properties of *N. flagelliforme* EPS, including anti-virus, antioxidant and anti-tumor, have been intensively investigated.¹⁰⁻¹² Considerable research efforts have been devoted to find the optimum culture medium and culture condition to obtain high EPS production yields.¹³⁻¹⁷ Light quality, carbon and nitrogen sources have been reported as important factors controlling *N. flagelliforme* EPS synthesis.^{15,17-21} As a group of photoautotrophic microorganisms, several studies reported that light quality especially red-light and blue-light could promote the accumulation of *N. flagelliforme* EPS.^{17,20-25} Most of cyanobacteria have efficient carbon concentration mechanisms (CCM) which could transport and accumulate inorganic carbon actively (Ci; HCO₃⁻, and CO₂).²⁶⁻²⁸ and could utilize very different forms of nitrogen, including combined nitrogen source or atmospheric nitrogen,^{2,29-31} which was helpful for increasing EPS production.

In previous studies, various complex EPS biosynthetic mechanisms were investigated and described in many bacterial, fungal and microalgae species, but the necessary pathways and enzymes involved in typical EPS biosynthesis, such as UDP-

Key Laboratory of Industrial Fermentation Microbiology, Ministry of Education, College of Biotechnology, Tianjin University of Science and Technology, Tianjin 300457, P. R. China. E-mail: pphan@tust.edu.cn; jiashiru@tust.edu.cn; Fax: +86 22 60602298; Tel: +86 22 60601598

† These two authors made equal contributions to the study.






Fig. 1 Summarized simplified biosynthesis pathways of EPS for *N. flagelliforme* EPS. The dashed arrows and boxes indicate putative biosynthesis pathways and precursors of EPS respectively. I: fructose-1,6-bisphosphatase (FBPase); II: phosphofructokinase (PFK); III: phosphomannose isomerase (PMI); IV: phosphoglucose isomerase (PGI); V: UDP-glucose pyrophosphorylase (UGPase); VI: UDP-galactose-4-epimerase (UGE); VII: UDP-glucose dehydrogenase (UGDH).

glucose pyrophosphorylase and UDP-glucose dehydrogenase, were relatively conserved, which played important roles in EPS production.^{32–41} Based on previous reports,^{42–44} and KEGG pathway, a simplified biosynthesis pathway for biosynthesis of cyanobacterial EPS was summarized (Fig. 1). In the complex pathway, the key enzyme that control EPS production in *N. flagelliforme* remains still unclear, and little is known about the correlation between related enzymes activities in *N. flagelliforme* EPS biosynthesis and the production of EPS under these different culture conditions.

In the study, the effects of culture conditions including different light quality, NaHCO_3 concentration, nitrogen source type and nitrogen concentration on EPS production and activities of EPS biosynthesis enzymes were investigated, and the correlation between them was subsequently analyzed for key enzyme affecting EPS production. The selected conditions (including different concentrations of NaHCO_3 and NaNO_3 , nitrogen sources and light qualities) have been found to significantly influence cell growth and EPS production in our previous study. Furthermore, expression level of genes encoding key EPS synthesis enzymes were determined for improved understanding EPS synthesis process in *N. flagelliforme*.

Materials and methods

Strain and culture conditions

The *N. flagelliforme* cells (TCCC11757) utilized in liquid suspension cultures were obtained from the Tianjin Key Lab of Industrial Microbiology (Tianjin, China). The cells were inoculated into a 500 mL shake-flask containing 200 mL BG-11 medium, and cultured at 25 °C under continuous white fluorescent light at a photon flux density of 60 mol photon per (m^2 per s) for 16 days, which was set as the basic group. Differing values of three culture conditions (carbon source, nitrogen source, and light quality) were evaluated in submerged fermentation of *N. flagelliforme*. For carbon source condition, initial concentrations of NaHCO_3 were set at 0 g L^{-1} , 1.26 g L^{-1} (equivalent to 15 mmol L^{-1}) and 2.94 g L^{-1} (equivalent to

35 mmol L^{-1}), respectively, with fixed NaNO_3 concentration at 1.50 g L^{-1} . For nitrogen source condition, one group set initial concentrations of NaNO_3 at 0 g L^{-1} , 1.50 g L^{-1} and 4.50 g L^{-1} respectively, with no exogenous addition of NaHCO_3 . The other group set the initial concentrations of NH_4Cl , $\text{CO}(\text{NH}_2)_2$ and NaNO_3 at 0.95 g L^{-1} , 0.53 g L^{-1} and 1.50 g L^{-1} respectively, with the same concentration of N element. For light quality condition, the same photon flux density of monochromatic red (660 nm) and blue (460 nm) were used instead of the white fluorescent light as the treatment group. The half-band widths are 5 nm for each monochromatic light, according to manufacturer's instruction (Shenzhen federal heavy Secco electronic Co. LTD., China).

The measurement of cell growth and EPS production

The cell growth was measured by weighing the cell dry weight. The total EPS (T-EPS) production was determined *via* phenol-sulfuric acid method.⁴⁵ The unit quality EPS (U-EPS) production was calculated as follows: U-EPS (mg g^{-1}) = T-EPS per cell dry weight.

Enzyme extract

The enzyme extract method is modified by the method of Looijesteijn *et al.*³³ and Tamoi *et al.*⁴⁶ The *N. flagelliforme* cells were harvested by centrifugation at 10 000 $\times g$ for 10 min at 4 °C, washed twice with cell extract buffer pre-cooled at 4 °C, which was 20 mM phosphate buffer containing 50 mM NaCl, 10 mM MgCl_2 , and 1 mM dithiothreitol at pH 6.5. The cells pellet resuspended in the extract buffer were disrupted ultrasonically at 20 kHz for 12 min on ice using an ultrasonic processor (SCIENTZ-IIID sonicator, Ningbo Scientz Biotechnology Inc., China). Cell debris was removed by centrifuged at 12 000 $\times g$ for 15 min at 4 °C, and the resulting supernatant (cell extract) was used for the enzyme activity assay. The protein content of the cell extract was determined by the method of Bradford.⁴⁷

Enzyme assays

Activity of EPS synthesis enzymes were measured spectrophotometrically by kinetic assay in a final volume of 250 μL at 24 °C. Reactions were initiated by adding 30 μL cell extract to various reaction mixtures depending on the enzyme. The enzyme activity was determined by monitoring the formation or disappearance of NAD(P)H by measuring the absorbance at 340 nm ($\epsilon_{340} = 6220 \text{ M}^{-1} \text{ cm}^{-1}$), and specific activities were expressed in $\mu\text{mol NAD(P)H (mg protein)}^{-1} \text{ min}^{-1}$. The blank for each enzyme was the same reaction condition with the same volume of cell extract that had been heated at 100 °C for 5 min. In all of the assays, the reaction velocity was linearly proportional to the amount of cell extract. The reaction rate ($\Delta\text{absorbance}/\text{min}$) calculated from the slope of linear plots ($R^2 = 0.95\text{--}1.0$) from 0–3 min or 0–5 min that was normalized to the control. The enzyme activities were calculated using the following equations:⁴⁸

$$\text{Enzyme activity (U, } \mu\text{mol min}^{-1}\text{)} = (\Delta A_{340}/\Delta t) \times V_1 \times D / (V_2 \times \epsilon_{340} \times d)$$



Specific activities = enzyme activity/protein content

where $\Delta A_{340}/\Delta t$ is the reaction rate, V_1 is the final reaction volume, V_2 is the cell extract volume, D is the dilution ratio, ϵ_{340} is extinction coefficients of NAD(P)H and NADH, d is colorimetric cup light path.

Phosphoglucose isomerase (PGI)³⁷ reaction mixture contained 50 mM potassium phosphate buffer (pH 6.8), 5 mM MgCl₂, 4 U glucose-6-phosphate dehydrogenase, 0.4 mM NADP⁺, 10 mM fructose-6-phosphate.

Fructose-1,6-bisphosphatase (FBPase)³⁷ reaction mixture contained 100 mM Tris-HCl buffer (pH 8.0), 0.4 mM NADP⁺, 20 mM MgCl₂, 20 mM DTT, 0.5 U phosphoglucose isomerase, 0.5 U glucose-6-phosphate dehydrogenase, 2 mM fructose-1,6-bisphosphate.

UDP-glucose pyrophosphorylase (UGPase)⁴⁹ reaction mixture contained 50 mM Tris-HCl buffer (pH 7.8), 0.4 mM UDP-glucose, 14 mM MgCl₂, 4 U glucose-6-phosphate.

UDP-galactose-4-epimerase (UGE)³⁷ reaction mixture contained 50 mM Tris-HCl buffer (pH 8.5), 5 mM MgCl₂, 0.015 U UDP-glucose dehydrogenase, 0.5 mM NAD, 0.2 mM UDP-galactose.

UDP-glucose dehydrogenase (UGDH)⁵⁰ reaction mixture contained 100 mM Tris-HCl buffer (pH 7.5), 5 mM UDP-glucose, 1 mM dithiothreitol, 1 mM MgCl₂, 1 mM NADP⁺.

Phosphomannose isomerase (PMI)⁵¹ reaction mixture contained 50 mM MOPS buffer (pH 7.0), 1 mM CoCl₂, 1 mM NADP⁺, 10 mM mannose-6-phosphate, 4 U phosphoglucose isomerase, 4 U glucose-6-phosphate dehydrogenase.

Phosphofructokinase (PFK)^{33,37} reaction mixture contained 50 mM Tris-HCl buffer (pH 7.5), 5 mM MgCl₂, 50 mM KCl, 1.25 mM ATP, 0.15 mM NADH, 4.5 U aldolase, 18 U triose-phosphate isomerase, 6.2 U glycerol-3-phosphate dehydrogenase, 5 mM fructose-6-phosphate.

Total RNA extraction

Total RNA of *N. flagelliforme* cells was prepared using MiniBEST Universal RNA Extraction Kit (TaKaRa, Japan) according to the manufacturer's instruction. For cDNA synthesis of total RNA was transcribed with the PrimeScript™ RT reagent Kit (TaKaRa, Japan) according to the manufacturer's instruction. Total RNA was quantified through the absorbance spectrum at wavelengths 260 nm and 280 nm using UV-mini1246 spectrophotometer (SHIMADZU, Japan). Total RNA solution was loaded on 2% agarose gel to assess the integrity of total RNA bands. The DNA maker was DL2000 maker (TaKaRa, Japan).

Primer designing and real-time quantitative PCR analysis

Expression levels of *pgi*, *pmi*, *fbp1*, *fbp2*, *ugp* and *ugdh* (encoding enzymes PGI, PMI, FBPase, UGPase and UGDH, respectively) were analyzed by qRT-PCR using SYBR® Premix Ex Taq™ (Tli RNaseH Plus) Kit (TaKaRa, Japan) according to the manufacturer's instructions with Bio-Rad IQ5 (Bio-Rad, American). Primers used for qRT-PCR were listed in Table 1. GAPDH was used as the reference gene to normalize the expression levels of different genes. For each gene, the reference sample (the control

Table 1 Primer sequences for qRT-PCR analysis

Enzyme	Target gene	Primer sequences (5' → 3')
GAPDH	<i>GAPDH</i>	Forward: GGTTTGCGGCTCCTTCGTAG Reverse: TCATTCCTTGGTGTGCGGTG
PMI	<i>pmi</i>	Forward: GTAGTCTCAGGGACAGCTAGGGTAAAC Reverse: AACACCAAGGGAATCACGCC
PGI	<i>pgi</i>	Forward: CCACGAGGATGGGACTGT Reverse: GCTTGCTCTGGGAGGATGAATAG
FBPase	<i>Fbp1</i>	Forward: GCTTAGTGTGTGCGCTCGCTTC Reverse: TCGGTTGCCTTACCATCGCT
FBPase	<i>Fbp2</i>	Forward: CGGGAGCGGATGAATAAAAT Reverse: GCGGGAGGTGCTGCTAACTT
UGPase	<i>ugp</i>	Forward: TCTATGAAAAGCTACGGTTGAG Reverse: TCCCTCTTCTGACACAATCTATC
UGDH	<i>ugdh</i>	Forward: ATTGGTTGGGGTGGTTCTTG Reverse: GTCCGACTGTTTTGCCTTTGAG

group) was defined as 1.0, and the $2^{-\Delta\Delta C_t}$ method was applied to calculate other gene expression level which was expressed as the fold of the reference sample.^{52,53} Negative controls (reactions without cDNA template) were included to detect the presence of any DNA contamination. All the reactions of these samples were done in triplicate to determine the accuracy and repeatability of the qRT-PCR reaction.

Data analysis and statistical analysis

Experimental data in this study was obtained from three replicates for each treatment, and values were shown as mean ± standard deviation. Supervised partial least-squares (PLS) analysis was used to verify the relative enzymes which were significant response to the EPS production for different culture conditions. Supervised PLS was performed by SIMCA 11.5 software (Umetrics, Umea, Sweden). Statistical significance of correlations between enzyme activities and EPS production was based on correlation test, which were analyzed based on data obtained under various culture conditions by SPSS statistics vision 2.0. Correlation coefficient ($R^2 > 0.5$) and significant effects ($P < 0.05$) were considered statistically significant.

Results

Effects of culture conditions on EPS production

In order to optimize the culture condition with maximum production of *N. flagelliforme* EPS, the effects of carbon source, nitrogen source, and light quality on EPS production were studied. As shown in Fig. 2A, the productions of U-EPS were similar and high under both blue-light and red-light condition, which were 48.01 ± 3.70 and 45.84 ± 0.68 mg g⁻¹ DCW, respectively. The U-EPS productions under blue-light and red-light condition were 1.52 and 1.45 times compared to that of white-light. Because of the highest biomass, the T-EPS obtained the highest production (37.32 ± 1.75 mg L⁻¹) under blue-light, which was 1.78 times to that of white-light.

The carbon sources used for submerged fermentation of *N. flagelliforme* were NaHCO₃ with different initial concentrations,






Fig. 2 Effects of culture conditions on EPS production. (A) Light quality; (B) NaHCO₃ concentration; (C) nitrogen source type; (D) NaNO₃ concentration.

and the profiles of EPS production were showed in Fig. 2B. The highest T-EPS ($28.38 \pm 3.19 \text{ mg L}^{-1}$) and U-EPS ($40.93 \pm 4.70 \text{ mg g}^{-1} \text{ DCW}$) production were both observed in the medium containing $1.26 \text{ g L}^{-1} \text{ NaHCO}_3$, which were 4.27 and 3.75 times compared to that of the medium containing $0 \text{ g L}^{-1} \text{ NaHCO}_3$ respectively.

The effects of nitrogen source type and initial concentration of NaNO₃ on EPS production by *N. flagelliforme* were shown in Fig. 2C and D, respectively. Using EPS production as maker, we tested three nitrogen source types (Fig. 2C), including NaNO₃, NH₄Cl and CO(NH₂)₂. It showed that NaNO₃ was the best nitrogen source for accumulating EPS ($29.27 \pm 0.37 \text{ mg L}^{-1}$ T-EPS and $42.87 \pm 1.52 \text{ mg g}^{-1} \text{ DCW}$ U-EPS) among these three nitrogen source types. The low level of EPS production was caused by adding NH₄Cl and CO(NH₂)₂ as nitrogen source, especially in the medium containing NH₄Cl as solo nitrogen source with few EPS production ($1.32 \pm 0.27 \text{ mg L}^{-1}$ T-EPS and $3.17 \pm 0.81 \text{ mg g}^{-1} \text{ DCW}$ U-EPS) detected, although NH₄⁺ was considered to be the preferred reduced N form for cyanobacteria uptake in general.^{30,31} As shown in Fig. 2D, the changes of initial NaNO₃ concentration had different effects on EPS production. The highest EPS production ($37.95 \pm 1.87 \text{ mg L}^{-1}$ T-EPS and $67.73 \pm 3.98 \text{ mg g}^{-1} \text{ DCW}$ U-EPS) was observed in the media without exogenous addition of NaNO₃, and EPS production was reduced when initial NaNO₃ concentration was increased.

Effects of culture conditions on enzyme activities involved in *N. flagelliforme* EPS synthesis

Activities of seven enzymes were evaluated under different culture conditions (Fig. 3), which were involved in *N. flagelliforme* EPS synthesis. Fig. 3A showed the profiles of the activity levels of seven enzymes measured under light quality condition. The activities of PGI, PMI, UGDH, UGE and UGPase were significantly improved under blue-light and red-light condition. The order of activity of FBPase was red-light > white-light > blue-light condition. As can be seen from Fig. 3B, it was interesting that for the five enzymes, the lowest activity was obtained at $1.26 \text{ g L}^{-1} \text{ NaHCO}_3$ and then their activities were increased at $2.94 \text{ g L}^{-1} \text{ NaHCO}_3$, which was an opposite trend with EPS production. However, FBPase activity was increased with the increase of NaHCO₃ concentration.

For different nitrogen source types, the activities of PGI, PMI, UGDH, UGPase in Fig. 3C had similar trend with EPS production, which was declined after adding NH₄Cl and CO(NH₂)₂ as nitrogen source, except FBPase and PFK. For different initial NaNO₃ concentrations (Fig. 3D), increased concentration of NaNO₃ led to the reduction of enzyme activity, which showed similar trend with EPS production, except UGDH. Therefore, the activity of PGI, PMI, and UGPase had the same trend with the changes of EPS production regardless of nitrogen source type and the initial NaNO₃ concentration used.






Fig. 3 Effects of culture conditions on enzyme activities involved in *N. flagelliforme* EPS synthesis. (A) Light quality;⁵⁹ (B) NaHCO_3 concentration;⁵⁹ (C) nitrogen source type; (D) NaNO_3 concentration.⁵⁹ * indicates $p < 0.05$ and ** indicates $p < 0.01$.

Relationship between activities of enzymes and EPS production under differing culture conditions

To study relationship between activities of enzymes and EPS production under differing culture conditions and avoid the impact of collinearity, multivariate statistical analysis was performed by using supervised partial least-squares (PLS) analysis and correlation analysis. The PLS was used to validate the differences between the different groups. The $t1/t2$ scatter plot used to explain the applicability of the PLS Method. The score plot displayed the space formed by the two predictive components as first predictive component ($t[1]$) and second predictive component ($t[2]$), respectively. These models were all well-constructed and possessed excellent fit and satisfactory predictive capability (Table 2). The first two components

accounted for >95% of the total variance for each culture conditions. In the PLS score plots, the confidence interval is defined by Hotelling's T2 ellipse (95% confidence interval), and observations outside the confidence ellipse are considered outliers.

Variable Importance Plot (VIP) is one of the important plots used to explain or determine the fitting effect of PLS model and VIP coefficients reflect the contribution of each variables to the PLS models. In this study, VIP values ≥ 0.5 were considered as an index indicating that these were possible key enzymes which had contribution to EPS production.

Three groups clearly separated in each profiles of PLS score plots (Fig. 4A and D and Fig. 5A and D). These results showed that the activities of enzymes involved in EPS biosynthesis of *N.*

Table 2 Statistical data from PLS analysis under different culture conditions

Culture condition	PLS					
	R^2X	R^2Y	Q^2	$t[1]$	$t[2]$	$t[1] + t[2]$
Light quality	0.985	0.971	0.958	72.43%	26.03%	98.46%
NaHCO_3 concentration	0.991	0.947	0.913	85.48%	13.65%	99.13%
Nitrogen source type	0.972	0.993	0.990	86.48%	10.74%	97.22%
NaNO_3 concentration	0.993	0.998	0.997	85.48%	13.78%	99.26%






Fig. 4 PLS score plot (A), loading plot (B) and VIP (C) results of samples treated by different light quality. PLS score plot (D), loading plot (E) and VIP (F) results of samples with the addition of different NaHCO_3 concentrations.

flagelliforme differed significantly. PLS loading plots and VIP were used to analyze the contribution of each enzyme to the EPS production. The results from Fig. 4B and C showed that PGI,

PMI, UGDH and UGPase were possible key enzymes affecting EPS production under light quality condition. As shown in Fig. 4E and F, PGI, UGDH, FBpase, UGPase and PMI might be




Fig. 5 PLS score plot (A), loading plot (B) and VIP (C) results of samples with the addition of different nitrogen source types. PLS score plot (D), loading plot (E) and VIP (F) results of samples.



Table 3 Correlation analysis between activities of enzymes and EPS production under differing culture conditions^a

Culture condition	EPS type	Related enzyme						
		PGI	FBPase	PFK	PMI	UGDH	UGE	UGPase
Light quality	T-EPS	++	+/-	+/-	+/-	++	+/-	+/-
	U-EPS	++	+/-	--	++	++	++	+/-
NaHCO ₃ concentration	T-EPS	--	+/-	--	--	--	+/-	--
	U-EPS	--	+/-	--	--	--	+/-	--
Nitrogen source type	T-EPS	+/-	--	+/-	+/-	++	+/-	++
	U-EPS	+/-	--	+/-	+/-	++	+/-	++
NaNO ₃ concentration	T-EPS	++	++	++	++	+/-	+/-	++
	U-EPS	++	++	++	++	+/-	+/-	++

^a ++: significant positive, $p < 0.01$; +: positive, $p < 0.05$; -: negative, $p < 0.05$; --: significant negative, $p < 0.01$; +/-: uncorrelated with the addition of different NaNO₃ concentrations.

the potential major drivers of EPS accumulation under carbon conditions. The order of enzyme contribution was UGDH, UGPase, PGI, FBPase, PFK and PMI under various nitrogen source conditions (Fig. 5B and C), while PGI, UGPase, PFK and PMI might be the possible key enzymes in the medium containing different concentration of NaNO₃ (Fig. 5E and F).

The correlation test was performed by correlation coefficient ($R^2 > 0.5$) and the results of linear correlation analysis showed in Table 3. The result of correlation analysis between T-EPS and enzyme activities was in accord with that between U-EPS and enzyme activities under different culture conditions, except light quality condition. PGI and UGDH showed positive correlation with both T-EPS and U-EPS under light quality condition. PGI, UGDH, UGPase, PMI and PFK were negatively associated with EPS (T-EPS and U-EPS) under carbon source condition. The effect of UGDH and UGPase was positive on EPS, while that of FBPase was negative under various nitrogen source conditions. The positive correlations were observed between five enzymes (PGI, FBPase, UGPase, PFK and PMI) and EPS production when different concentrations of NaNO₃ were added.

Combing the results of PLS analysis and correlation analysis, by choosing the enzymes showing significant correlation with EPS production under all tested conditions, PGI, PMI, UGDH, UGPase and FBPase were preliminarily selected as the potential important enzymes influencing EPS production.

Transcription levels of genes encoding key EPS synthesis enzymes

To evaluate the effects of culture conditions on expression of key enzymes, qRT-PCR was used to analyze the transcription levels of genes including *pgi* (encoding PGI), *pmi* (encoding PMI), *fbp1* and *fbp2* (encoding FBPase), *ugd* (encoding UGDH) and *ugpg* (encoding UGPase) in control group and three experimental groups with high-yield production of EPS consisting of carbon source group with 1.26 g L⁻¹ NaHCO₃, nitrogen source group with 0 g L⁻¹ NaNO₃, and light quality group with blue-light.

The relative quantitative results of the above genes under four culture conditions were shown in Fig. 6. The mRNA levels were normalized against GAPDH level of control group. In three experimental groups, *pgi* and *fbp1* genes were highly expressed compared with that of control group, while the transcription levels of other four genes were decreased in three experimental groups.

Discussion

In recent two decades, various efforts have been devoted to increase *N. flagelliforme* EPS production, which mostly focus on optimization of culture condition.^{14–18,20,21} So far, little information is available about the effects of culture conditions on activities of enzymes involved in EPS production. Therefore, in this study, the possible relationship between *N. flagelliforme* EPS production and enzymes activities related to EPS biosynthesis was evaluated.

In this study, the results showed that the EPS production of *N. flagelliforme* was significantly influenced by culture conditions, which were consistent with previous reports.^{16–18,20,21} In the PLS analysis, PGI was the only enzyme with VIP value > 1 under all four culture conditions. The results of correlation analysis showed that PGI was significantly correlated with EPS production, except under the nitrogen source condition. PGI catalyzes the reversible isomerization of glucopyranose-6-phosphate and fructofuranose-6-phosphate playing a key role in glycolysis and gluconeogenesis pathway.^{37,54,55} The highest EPS production was also associated with the highest activity of




Fig. 6 Transcription levels of key enzymes in *N. flagelliforme* under four culture conditions. * indicates $p < 0.05$ and ** indicates $p < 0.01$.



PGI under four culture conditions, except under the condition of adding different concentration of NaHCO_3 . The gene *pgi* encoding PGI was highly expressed in three experimental groups verified the results of correlation analysis, referring that *pgi* might be one important gene influencing EPS production under different culture conditions.

FBPase can catalyze the conversion of fructose-1,6-diphosphate into fructose-6-phosphate, which occurs in two essential steps: one involved in gluconeogenesis and the other in Calvin cycle.^{33,46} The transcription analysis of genes encoding FBPase revealed that two genes (*fbp1* and *fbp2*) encoding two forms of FBPase (FBPase-I and FBPase-II) were detected in *N. flagelliforme*, which were mainly involved in Calvin cycle and involved in gluconeogenesis respectively.^{46,62–64} The observed significant up-regulation of *fbp1* and down-regulation of *fbp2* under experimental conditions indicated the adjustment of intracellular carbon metabolism to promote EPS synthesis and corresponded well with the enhanced EPS production, which suggested that gene *fbp1* rather than *fbp2* was the potential gene influencing EPS synthesis in *N. flagelliforme*, coinciding with the previous reports.^{46,62–64} Interestingly, the activity of FBPase was significantly enhanced under red-light, and was obviously increased with increasing initial NaHCO_3 concentration and decreasing NaNO_3 concentration, but the FBPase activity was more closely associated with EPS production under nitrogen condition, which might indicate different regulation mechanisms were involved in different culture conditions.

UGPase was involved in the production of UDP-glucose, which was a key precursor to polysaccharide synthesis.^{34,37,48} UGDH was required for the synthesis of the precursor UDP-glucuronic acid,⁵⁶ which was an important intermediate in polysaccharide synthesis pathway.^{57,58} In this study, both partial least-squares analysis and correlation analysis showed the activities of UGPase and UGDH had strong correlation with EPS production under most of culture conditions. However, the activities were not all increased in experimental groups with higher EPS production. In addition to that, the transcription levels of corresponding genes *ugdH* and *ugpg* were decreased. Therefore it might be deduced that UGPase and UGDH were not directly associated with EPS yield. In our previous study, it was found that culture conditions showed significant influences on monosaccharide composition of EPS, and UGPase and UGDH were correlated with monosaccharide ratio under different culture conditions.⁵⁹ Thus it might be speculated that changes of UGPase and UGDH activities induced by culture conditions had greater influence on EPS monosaccharide composition than yield.

In many cyanobacteria, PMI possess mannose-6-phosphate isomerase activity and nucleotidyltransferase activity which can catalyze the reversible isomerization of fructose-6-phosphate from central metabolism into mannose-6-phosphate and forming GDP-mannose, respectively.^{60,61} The correlation between the activity of PMI and the amount of EPS produced was observed, indicating that EPS production might be affected by activity of PMI. However, in contrast to the increased activity, down-regulation of gene *pmi* expression level was detected. The reason for discrepancy between encoding

gene transcription and activity levels remains unclear and needs to be further investigation.

In addition, UGE, the Leloir pathway key enzyme catalyzing the interconversion of UDP-galactose and UDP-glucose, was not significantly changed, except light quality condition, which seemed to be not closely associated with EPS synthesis. PFK was the first committed step of glycolysis catalyzing the phosphorylation of fructose-6-phosphate to fructose 1,6-bisphosphate by ATP, which seemed not to play an important role in controlling *N. flagelliforme* EPS production.

In this study, PGI and FBPase were found to be more important enzymes positively influencing the EPS biosynthesis than the other five enzymes, however, through comparing the relationship between enzyme activities and EPS production under four kinds of culture conditions, the results showed some discrepancies that the activities of enzymes had positive effects on EPS production under some culture conditions but had negative effects under other culture conditions. It might be due to the complex EPS biosynthesis process and regulation mechanism. Different influence patterns were found under different culture conditions indicating the utilization of distinct regulation mechanism for different culture conditions, which might explain the discrepancies, and further indicate the existence of coordinated regulation of several enzymes in the EPS production, which still requires further investigation.

Conclusions

The effects of culture conditions on *N. flagelliforme* EPS production and related enzymes were investigated in the study. Culture conditions showed significant influences on EPS production, and related enzyme activities. Both PLS analysis and correlation analysis were used to analyze the relationship between activities of seven enzymes and EPS production. The results showed that the activities of PGI, PMI, FBPase, UGDH and UGPase were correlated with EPS production under different culture conditions. Subsequent analysis of the transcription level of genes encoding the five enzymes showed *pgi* and *fbp1* in three experimental groups was significantly up regulated, which implied PGI and FBPase might be the important enzymes positively influencing EPS production of *N. flagelliforme*. The findings would improve the understanding of EPS synthesis process in *N. flagelliforme*.

Conflicts of interest

There are no conflicts of interest to declare.

Abbreviations

dTDP-Rha	dTDP-rhamnose
F6P	Fructose-6-phosphate
FBP	Fructose-1,6-bisphosphatase
FBPase	Fructose-1,6-biphosphate
G1P	Glucose-1-phosphate
G6P	Glucose-6-phosphate



M6P	Mannose-6-phosphate
PFK	Phosphofructokinase
PGI	Phosphoglucose isomerase
PMI	Phosphomannose isomerase
T-EPS	Total EPS
UDP-Gal	UDP-galactose
UDP-Glu	UDP-glucose
U-EPS	Unit quality EPS
UGDH	UDP-glucose dehydrogenase
UGE	UDP-galactose-4-epimerase
UGPase	UDP-glucose pyrophosphorylase

Acknowledgements

The authors are very grateful for the financial support from the National Natural Science Foundation of China (Grant No. 31671842).

References

- B. H. A. Rehm, *Nat. Rev. Microbiol.*, 2010, **8**(8), 578–592.
- S. Pereira, A. Zille, E. Micheletti, P. Moradas-Ferreira, R. De Philippis and P. Tamagnini, *FEMS Microbiol. Rev.*, 2009, **33**(5), 917–941.
- I. W. Sutherland, *Trends Biotechnol.*, 1998, **16**(1), 41–46.
- R. De Philippis, C. Sili, R. Paperi and M. Vincenzini, *J. Appl. Phycol.*, 2001, **13**(4), 293–299.
- R. De Philippis, M. C. Margheri, R. Materassi and M. Vincenzini, *Appl. Environ. Microbiol.*, 1998, **64**(3), 1130–1132.
- R. De Philippis and M. Vincenzini, *FEMS Microbiol. Rev.*, 2006, **22**(3), 151–175.
- R. De Philippis, G. Colica and E. Micheletti, *Appl. Microbiol. Biotechnol.*, 2011, **92**(4), 697–708.
- K. X. Qian, H. R. Zhu and S. G. Chen, *Acta Phytoecol. Geobot. Sin.*, 1989, **13**, 97–105.
- H. Takenaka, Y. Yamaguchi, S. Sakaki, K. Watarai, N. Tanaka and M. Hori, *Food Chem. Toxicol.*, 1998, **36**, 1073–1077.
- K. Kanekiyo, J. B. Lee, K. Hayashi, H. Takenaka, Y. Hayakawa, S. Endo and T. Hayashi, *J. Nat. Prod.*, 2005, **68**(7), 1037–1041.
- K. Kanekiyo, K. Hayashi, H. Takenaka, J. B. Lee and T. Hayashi, *Biol. Pharm. Bull.*, 2007, **30**(8), 1573–1575.
- K. Hayashi, K. Kanekiyo, Y. Ohta, J. B. Lee, H. Takenaka and T. Hayashi, *Planta Med.*, 2008, **74**(9), 946.
- J. Moreno, M. A. Vargas, H. Olivares, J. Rivas and M. G. Guerrero, *J. Biotechnol.*, 1998, **60**(3), 175–182.
- J. Su, S. R. Jia, X. Chen and H. Yu, *J. Appl. Phycol.*, 2008, **20**(3), 213–217.
- H. F. Yu, Y. S. Dong, and S. R. Jia, *International Conference on Biomedical Engineering and Informatics*, IEEE, 2010, 5, pp. 2052–2056.
- Z. Ding, S. R. Jia, P. P. Han, N. Yuan and N. Tan, *J. Appl. Phycol.*, 2013, **25**(4), 1017–1021.
- P. P. Han, S. G. Shen, H. Y. Wang, S. Y. Yao, Z. L. Tan, C. Zhong and S. R. Jia, *J. Appl. Phycol.*, 2016, 1–11.
- H. Yu, S. Jia and Y. Dai, *Appl. Biochem. Biotechnol.*, 2010b, **160**(2), 552–560.
- P. P. Han, S. G. Shen, H. Y. Wang, Y. Sun, Y. J. Dai and S. R. Jia, *Algal Res.*, 2015, **9**, 143–150.
- P. P. Han, Y. Sun, X. Y. Wu, Y. J. Yuan, Y. J. Dai and S. R. Jia, *Appl. Biochem. Biotechnol.*, 2014a, **172**(1), 36–49.
- P. P. Han, Y. Sun, S. R. Jia, C. Zhong and Z. L. Tan, *Carbohydr. Polym.*, 2014b, **105**(1), 145–151.
- W. Kowallik, *Plant Physiol.*, 1982, **33**, 51–72.
- M. Atta, A. Idris, A. Bukhari and S. Wahidin, *Bioresour. Technol.*, 2013, **148**(11), 373.
- L. Zhang, G. Ma, K. Yamawaki, Y. Ikoma, H. Matsumoto, T. Yoshioka, O. Satoshi and K. Masaya, *J. Plant Physiol.*, 2015, **188**, 58–63.
- E. Bland and L. T. Angenent, *Bioresour. Technol.*, 2016, **216**, 579–586.
- T. Ogawa, *American Society of Plant Physiologists*, Rockville, MD, 1993.
- M. R. Badger and G. D. Price, *J. Exp. Bot.*, 2003, **54**(383), 609–622.
- G. D. Price, M. R. Badger, F. J. Woodger and B. M. Long, *J. Exp. Bot.*, 2008, **59**(7), 1441–1461.
- A. Otero and M. Vincenzini, *J. Biotechnol.*, 2003, **102**(2), 143–152.
- J. M. García-Fernández and J. Diez, *Res. Microbiol.*, 2004, **155**(10), 795–802.
- P. M. Glibert, F. P. Wilkerson, R. C. Dugdale, J. A. Raven, C. L. Dupont, P. R. Leavitt, A. E. Parker, J. M. Burkholder and T. M. Kana, *Limnology & Oceanography*, 2016, **61**(1), 284–300.
- B. L. Horecker, *Annu. Rev. Microbiol.*, 1966, **20**(20), 253–290.
- P. J. Looijesteijn, I. C. Boels, M. Kleerebezem and J. Hugenholtz, *Appl. Environ. Microbiol.*, 1999, **65**(11), 5003–5008.
- B. Degeest and L. de Vuyst, *Appl. Environ. Microbiol.*, 2000, **66**(8), 3519–3527.
- C. Petit, J. P. Grill, N. Maazouzi and R. Marczak, *Appl. Microbiol. Biotechnol.*, 1991, **36**(2), 216–221.
- A. Escalante, C. Wachter-Rodarte, M. García-Garibay and A. Farrés, *J. Appl. Microbiol.*, 1998, **84**(1), 108–114.
- G. J. Grobbs, M. R. Smith, J. Sikkema and J. A. M. de Bont, *Appl. Microbiol. Biotechnol.*, 1996, **46**(3), 279–284.
- H. Li, H. Xu, H. Xu, S. Li and P. K. Ouyang, *Appl. Microbiol. Biotechnol.*, 2010, **86**(1), 295–303.
- S. Pan, D. Yao, J. Chen and S. Wu, *Carbohydr. Polym.*, 2013, **92**(1), 629.
- Z. Y. Zhu, X. C. Liu, F. Y. Dong, M. Z. Guo, X. T. Wang, Z. Wang and M. Z. Yong, *Appl. Microbiol. Biotechnol.*, 2016, **100**(9), 3909–3921.
- F. Mozzi, D. G. G. Savoy and D. V. G. Font, *J. Appl. Microbiol.*, 2003, **94**(2), 175–183.
- A. J. Smith, *Ann. Inst. Pasteur/Microbiol.*, 1983, **134**(1), 93–113.
- L. J. Stal and R. Moezelaar, *FEMS Microbiol. Rev.*, 1997, **21**(2), 179–211.
- V. L. Knowles and W. C. Plaxton, *Plant Cell Physiol.*, 2003, **44**(7), 758–763.



- 45 M. Dubois, K. A. Gilles and J. K. Hamilton, *Anal. Chem.*, 1956, **28**, 350–353.
- 46 M. Tamoi, A. Murakami, T. Takeda and S. Shigeoka, *Biochim. Biophys. Acta*, 1998, **1383**(2), 232–244.
- 47 M. M. Bradford, *Anal. Biochem.*, 1976, **72**, 248–254.
- 48 H. Bisswanger, *Practical enzymology*, Wiley VCH Verlag GmbH & Co, KGaA, New York, 2004.
- 49 R. L. Bernstein and P. W. Robbins, *J. Biol. Chem.*, 1965, **240**(1), 391–397.
- 50 R. J. Hung, H. S. Chien, R. Z. Lin, C. T. Lin, J. Vatsyayan, H. L. Peng and H. Y. Chang, *J. Biol. Chem.*, 2007, **282**(24), 17738–17748.
- 51 D. Shinabarger, A. Berry, T. B. May, R. Rothmel, A. Fialho and A. M. Chakrabarty, *J. Biol. Chem.*, 1991, **266**(4), 2080–2088.
- 52 K. J. Livak and T. D. Schmittgen, *Methods*, 2001, **25**, 402–408.
- 53 E. Szekeres, C. Sicora, N. Dragoş and B. Drugă, *FEMS Microbiol. Lett.*, 2014, **359**(1), 102–109.
- 54 L. Peng, J. Li, Y. Liu, Z. Xu, J. Y Wu, Z. Ding, Z. Gu, L. Zhang and G. Shi, *RSC Adv.*, 2016, **6**(45), 39284–39291.
- 55 L. Peng, S. Qiao, Z. Xu, F. Guan, Z. Ding, Z. Gu, L. Zhang and G. Shi, *Carbohydr. Polym.*, 2015, **133**, 104–109.
- 56 A. T. Granja, A. Popescu and A. R. Marques, *Appl. Microbiol. Biotechnol.*, 2007, **76**(6), 1319–1327.
- 57 A. Kereszt, I. Ernök, B. L. Reuhs, R. W. Carlson, I. Ádám Kondoros and P. Péter, *J. Bacteriol.*, 1998, **180**, 5426–5431.
- 58 B. Seitz, C. Klos, M. Wurm and R. Tenhaken, *Plant J.*, 2000, **21**, 537–546.
- 59 P. P. Han, S. Y. Yao, R. J. Guo, S. G. Shen, R. R. Yan, Z. L. Tan and S. R. Jia, *Carbohydr. Polym.*, 2017, **174**, 111–119.
- 60 F. Sousa, R. Joana, H. Jorge and S. A. Leitão, *Biotechnology of Biopolymers*, InTech, 2011, pp. 257–274.
- 61 S. F. Ana, N. Inês Silva, H. O. Vítor, C. Raquel and M. M. Leonilde, *Front. Cell. Infect. Microbiol.*, 2011, **1**(12), 16.
- 62 M. Tamoi, T. Takeda and S. Shigeoka, *Plant Cell Physiol.*, 1999, **40**(2), 257–261.
- 63 W. Ma, D. Shi, Q. Wang, L. Wei and H. Chen, *J. Appl. Phycol.*, 2005, **17**(3), 273–280.
- 64 J. Becker, C. Klopprogge, O. Zelder, E. Heinzle and C. Wittmann, *Appl. Environ. Microbiol.*, 2005, **71**, 8587–8596.

