



Cite this: *RSC Adv.*, 2017, 7, 41755

Received 4th August 2017
 Accepted 22nd August 2017

DOI: 10.1039/c7ra08637g

rsc.li/rsc-advances

Conversion of glycals into vicinal-1,2-diazides and 1,2-(or 2,1)-azidoacetates using hypervalent iodine reagents and Me_3SiN_3 . Application in the synthesis of *N*-glycopeptides, pseudo-trisaccharides and an iminosugar†

Ande Chennaiah, Srijita Bhowmick‡ and Yashwant D. Vankar *

Glycals were found to react with a reagent system comprising of phenyliodine bis(trifluoroacetate) (PIFA) and Me_3SiN_3 in the presence of TMSOTf as a catalyst to form the corresponding vicinal 1,2-diazides. On the other hand, they reacted with another reagent system phenyliodine diacetate (PIDA) and Me_3SiN_3 , also in the presence of TMSOTf as a catalyst, to lead to the corresponding vicinal 1,2-azidoacetates. These azido derivatives were converted into a number of 2-azido-*N*-glycopeptides, pseudotrisaccharides, and a piperidine triol derivative, an iminosugar.

Introduction

The growing importance of aminosugars,¹ *N*-glycopeptides² and sugar derived triazoles³ (obtained *via* click-chemistry) in biological systems is apparent from the recent literature. Synthesis of such compounds has led to the development of a number of synthetic methodologies involving functionalization of monosaccharides, in particular glycals. In this regard, methodologies that allow attachment of an azido moiety⁴ at the anomeric center are frequently used in the synthesis of *N*-glycopeptides, although there are some other ways^{2e,5} of introducing a 'nitrogen' unit. The azide group also allows click-chemistry, to link with another sugar unit leading to pseudo di-, tri- or oligosaccharides, to be carried out.³ Apart from this, introduction of an amino functionality (or an azido group) at the C-2 position of a sugar moiety, with an appropriate functional group at the anomeric carbon for effecting glycosylation, is one of the most important reactions known in carbohydrate chemistry.^{1a,6} This allows access to amino sugars with the amino group at the C-2 position, a structural feature that is prevalent in a number of oligosaccharides including aminoglycoside antibiotics. Toward this endeavor, conversion of glycals into sugar derived 2-azido-1-nitrates using a combination of ceric ammonium nitrate and NaN_3 (and its improvements) in CH_3CN , a protocol developed by Lemieux *et al.*,^{6a} is universally followed.

At the same time, there exist many oligosaccharides that contain *N*-glycopeptide units in which the sugar moiety carries a C-2 amino functionality, and the glycopeptide part has an Asn-linkage at the anomeric carbon.^{2b,7} Interest in the synthesis of such molecules stems from the fact that protein *N*-glycosylation occurs during post-translational modifications in biological systems. This has led to newer approaches towards functionalization of glycals (or any other sugar derivative such as glycosamines) to introduce 1,2-diamino (or equivalent) functionalities that are useful in the synthesis of *N*-glycopeptides.⁸ Thus, improved or alternate approaches toward the introduction of a nitrogen based functional group at C-1 and/or C-2 of a sugar moiety are important. In view of our continued interest^{2f,4a-c,8a,9} in functionalizing glycals *en route* to the synthesis of iminosugars, aminosugars and glycopeptides, we herein report an easy access to 1,2-diazido sugars from glycals upon treatment with phenyliodine bis(trifluoroacetate) (PIFA) and Me_3SiN_3 reagent system. Also, we have developed conditions that allow conversion of glycals into 1-azido-2-acetoxy sugars or 2-azido-1-acetoxy sugars upon reaction with phenyliodine diacetate (PIDA) and Me_3SiN_3 . Further, we have demonstrated the utility of these azido compounds in the synthesis of 2-amino-*N*-glycopeptides, pseudotrisaccharides, and an iminosugar.

Results and discussion


Hypervalent iodine reagents have gained enormous importance in recent years due to their low toxicity, ease of handling and ready availability.¹⁰ The use of hypervalent iodine reagents, most notably PIDA, in conjunction with a halide ion source has been reported on a few occasions in functionalizing olefins,

Department of Chemistry, Indian Institute of Technology, Kanpur – 208016, India.
 E-mail: vankar@iitk.ac.in

† Electronic supplementary information (ESI) available: CCDC 1554251 and 1554252. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c7ra08637g

‡ M.Sc. research project participant (2016).






Scheme 1 Exploratory reaction of galactal derivative 1 with PIFA–Me₃SiN₃.

including glycols, to form vicinal halo alkoxides in an intermolecular fashion.¹¹ Introduction of a nitrogen based functional group at C-2 of a glucal derivative has also been realized in an intramolecular fashion from glucal 3-carbamate upon its reaction with PIDA along with Rh₂(OAc)₄.^{6c} However, one of the most interesting reactions to stereoselectively convert glycols into sugar derived *trans*-1,2-diacetates involves their reaction with PIDA in presence of BF₃·Et₂O as a catalyst as reported by Gin *et al.*¹² Further, the *in situ* formed *trans*-1,2-diacetates lead to *O*-glycosylation when treated with an alcohol (ROH) and TfOH (as a catalyst) in the same pot. The proposed mechanism suggests that PIDA acts as an electrophile making a π -complex with a glycol, followed by the attack of acetate ion at the anomeric carbon as well as at C-2 leading to the observed *trans*-1,2-diacetates. In view of this, and in view of the importance of amino sugars and *N*-glycopeptides (*vide supra*) we surmised that reaction of glycols with PIDA combined with Me₃SiN₃ may lead to 1,2-diazido sugars. Our initial experiments involving reaction of 3,4,6-tri-*O*-benzyl galactal 1 (Scheme 1) with both one equivalent of PIDA and Me₃SiN₃ in the presence of BF₃·Et₂O or TMSOTf (30 mol%) as a catalyst led to a mixture of four products. These were 1-azido-2-acetoxy sugar 2, 2-azido-1-acetoxy sugar 3; 1,2-diacetoxy sugar 4 and 1,2-diazido sugar 5. Increasing the amount of Me₃SiN₃ did not lead to increased

formation of 1,2-diazides and there was always an incorporation of the acetate moiety onto galactal 1 due to competition between the azide and the acetate ions. Although, conversion of olefins into vicinal diazides is known with few reagents^{13a} such as Mn(OAc)₃–NaN₃,^{13b} azido iodine(III) reagent,^{13c} NaIO₄/NaN₃,^{13d} and Zhdankin reagent (a hypervalent iodine reagent) along with a copper catalyst,^{13e} formation of sugar derived 1,2-diazides from glycols has been reported rarely.^{13b,14} Thus, for example, 2,4,6-tri-*O*-acetyl galactal has been reported to react with Me₃SiN₃–PIDA–PhSeSePh^{14a} to give *cis*-1,2-diazido galactal albeit in low yield and as a side product. On the other hand, more recently Xu *et al.*^{14b} demonstrated that the *in situ* prepared Zhdankin reagent in presence of an iron catalyst converts a broad range of olefins, including one example of a glucal derivative, to the corresponding 1,2-diazides. In order to find an alternate route to convert glycols into sugar derived 1,2-diazides, we explored their reactivity with PIFA–Me₃SiN₃ reagent system¹⁵ in presence of an acid catalyst. Such a reagent system has been used to introduce an azide group into aromatics and some heterocycles akin to aromatic substitution,^{15a,b} for substitution of an azide moiety^{15c} at benzylic positions, and more recently to perform C–H activation.^{15d,e} Apart from this, an interesting intramolecular azidoarylation of alkenes using PIFA–Me₃SiN₃ combination has been reported by Antonchick *et al.*^{15f} In these examples, PIFA–Me₃SiN₃ reagent system was proposed to lead to the formation of PhI(N₃)₂ or PhIN₃(OCOCF₃) as intermediate species which act as a source of azide radical to effect the observed reactions. We expected that if PhI(N₃)₂ is formed as an intermediate, in presence of an acid catalyst it may convert glycols to sugar-derived 1,2-diazides in an analogous manner as sugar derived 1,2-diacetates were obtained from glycols upon reaction with PIDA as reported by Gin *et al.*¹² Towards this endeavor, initial experiments involved treatment of 3,4,6-tri-*O*-acetyl galactal 6a with PIFA (1.0 eq.) and 2 eq. of Me₃SiN₃ in presence of BF₃·Et₂O (30 mol%) at –30 °C which gave only 38% of the expected 1,2-diazide 7a as a single isomer. Further optimizations with respect to azide source and acid catalyst (Table 1) led to the use of 3 eq. of TMSN₃ and 30 mol%

Table 1 Optimization of reaction conditions to form 1,2-diazides^a



Entry	Catalyst	N ₃ -Source	Equiv.	Solvent	Temp (°C)	Time (h)	Yield (%)
1	None	NaN ₃	2	CH ₃ CN	rt	24	No reaction
2	None	TMSN ₃	2	CH ₂ Cl ₂	0	24	No reaction
3	BF ₃ ·OEt ₂	TMSN ₃	2	CH ₂ Cl ₂	–30	0.5	38 ^b
4	BF ₃ ·OEt ₂	TMSN ₃	3	CH ₂ Cl ₂	–30	0.5	44 ^b
5	BF ₃ ·OEt ₂	Bu ₄ NN ₃	3	CH ₂ Cl ₂	–30	0.5	37 ^b
6	TMSOTf	Bu ₄ NN ₃	3	CH ₂ Cl ₂	–30	0.5	33 ^b
7	TMSOTf	TMSN ₃	3	CH ₂ Cl ₂	–30	0.5	61

^a Reaction conditions: glycols (0.36 mmol), TMSN₃, PhI(OCOCF₃)₂ (0.36 mmol), TMSOTf (0.10 mmol), CH₂Cl₂, –30 °C, N₂ atmosphere, isolated yields after purification by silica gel column chromatography. ^b Yield based on recovered starting material.



of TMSOTf at $-30\text{ }^{\circ}\text{C}$ as the best condition forming **7a** in 61% yield (entry 7, Table 1). With this optimized condition we carried out the remaining studies. Thus, a variety of differently protected glycal derivatives led to the corresponding 1,2-diazides in moderate yields (Scheme 2).


It was generally found that while the galactal derivatives led exclusively to *cis*-1,2-diazides, the glucal derivatives gave a mixture of α - and β -anomers in varied ratios with the α -anomer being major in most of the cases. The structures and the α/β ratios were established based on COSY, NOE, homonuclear decoupling and DEPT experiments.^{16a} Thus, for example, in the ^1H NMR spectrum of compound **7a** the anomeric proton (H-1) appeared as a doublet at δ 5.48 with $J = 4.12$ Hz. The homonuclear decoupling of H-1 led the proton H-2 to appear as a doublet, from a doublet of doublet, with $J = 11.00$ Hz indicating that H-2 and H-3 are *trans*-diaxial. Further, in the NOE experiments no enhancement was observed for protons H-3 and H-5 when H-1 was irradiated and *vice versa* indicating that H-1 is equatorially oriented. In addition, the structure of the galactal derivative **5** was proved by single crystal X-ray analysis.^{16b} Likewise, the spectral data of 1,2-diazide **9a**, derived from glucal **8a**, also showed that the azido moieties are α -oriented.^{16a} Thus, in NOE experiments, irradiation of proton H-4 at δ 5.00 led to the enhancement of signal for H-2 at δ 3.61 and *vice versa* indicating that H-2 is axially oriented. On the other hand, no enhancement was observed for H-3 and H-5 protons when H-1 was irradiated at δ 5.44 suggesting that H-1 is equatorially oriented.

Interestingly, the arabinal derivatives **10a** and **10b** gave the corresponding 1,2-diazides **11a** and **11b** having $^1\text{C}_4$ conformations which was confirmed by spectral studies.^{16a} In case of **11a**,


irradiation of H-1 at δ 5.43 in the homonuclear decoupling experiment led the doublet of doublet for H-2 to appear as a doublet with $J = 10.6$ Hz indicating that H-2 and H-3 are diaxially oriented. Likewise, irradiation of H-4 at δ 5.27 caused the doublet of doublet for H-3 at δ 5.13 to appear as a doublet with $J = 10.6$ Hz. Further, in NOE experiments irradiation of H-3 did not lead to the enhancement of the signal for H-1, and also irradiation of H-2 did not show the enhancement of H-4. These data confirm that H-2 and H-3 are diaxially oriented and that **11a** possesses $^1\text{C}_4$ conformation. In a similar fashion, the structure of **11b** was established.

It is important to note that in the absence of an acid catalyst no reaction was found to take place between a glycal derivative and PIFA– Me_3SiN_3 reagent combination indicating that the azido radical, if formed,¹⁵ does not add on to the electron rich glycal double bond. Therefore we presume that in the present case, the reaction does not proceed *via* radical pathway,^{17a} instead it proceeds *via* ionic pathway similar to the ones proposed by Moriarty,^{17b} and Kirschning.^{17c} Thus, *in situ* generated $\text{PhI}(\text{N}_3)_2$ upon π -interaction (complex **A** in Scheme 3) with the double bond from the α -side (in case of a galactal and a glucal derivative) leads to intermediate **B** and transfer of the azide moiety to C-2 occurs in an $\text{S}_{\text{N}}1$ fashion from the α -side only. Following this (or simultaneously) the second azide preferentially attacks at the anomeric carbon of the galactal derivative from the α -side (path a) due to steric hindrance caused by the substituents at C-3, C-4 and C-5. In case of glucal derivatives, however, some product was also formed due to β -attack of the azide moiety *via* path b since the glucal derivatives show less steric bias from the β -side as compared to galactal derivatives.

Further, in case of arabinal derivatives it appears that the double bond preferentially forms π -complex from the less hindered β -side (complex **C**), followed by azide ion attachments at C-2 and C-1 from the opposite face to the two –OR groups, in a similar manner as happens in galactal cases, eventually




Scheme 2 Percentage yield and α/β ratio of the 1,2-diazides.



Scheme 3 Proposed mechanism for the formation of 1,2-diazides.






Scheme 4 Conversion of 1,2-diazides into bis-triazole **13** and pseudo-trisaccharide **15**.

leading to the observed 1,2-diazides **11a** and **11b** (Scheme 3) which assume 1C_4 conformations. It is clear that the success of the 1,2-diazide formation is mainly due to the low nucleophilicity of the trifluoroacetate ion compared to the azide ion.

In order to demonstrate the utility of these 1,2-diazides, we have utilized the click chemistry to prepare two bis-triazoles, one from a non-sugar and the other from a sugar based alkyne. Thus, 1,2-diazide **5** was reacted with 2 eq. of phenylacetylene **12** to form the bis-triazole **13** (Scheme 4) in 80% yield using standard conditions.^{3d} Likewise, a sugar derived alkyne¹⁸ **14** led to the corresponding bis-triazole **15**, a pseudo-trisaccharide,³ in 75% yield. The structures of these triazoles were confirmed from their spectral data.^{16a}

The importance of *N*-glycopeptides is apparent from the introduction part (*vide supra*) and a number of synthetic approaches have been reported to procure them, and newer synthetic routes are still being developed. In view of this, we demonstrate the utility of 1,2-diazides **7a** and **9a** to form the corresponding 2-azido-*N*-glycopeptides **18–23** (Scheme 5). Thus, selective reduction of the anomeric azide of **7a** and **9a** was carried out with ammonium tetrathiomolybdate $[(NH_4)_2MoS_4]^{2-}$ ¹⁹ to give the corresponding amino compounds **16** and **17** and, without isolating them, the crude amines were coupled with a number of amino acids and small peptides, mediated by 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC)^{20a} to lead to the corresponding 2-azido-*N*-glycopeptides **18–23**. The stereochemistry of the 2-azido-*N*-glycopeptides was confirmed to be 1,2-*trans* based on COSY, HETCOR and homonuclear decoupling experiments^{16a} of **18**. Thus, selective decoupling of C_1NH at δ 6.05 led the anomeric proton to appear as a doublet with coupling constant $J = 9.96$ Hz indicating that both H-1 and H-2 are axial. The facile mutarotation of free anomeric amines has been reported in many cases^{4b,20b,c} and their functionalization mainly leads to the β -anomers. It is therefore not surprising that in the present case, reduction of α -azides to the corresponding free amines exclusively lead to



Scheme 5 Selective reduction of 1,2-diazides and synthesis of β -*N*-glycopeptides.

β -glycopeptides *via* mutarotation. The azido moiety at C-2 of the azido-glycopeptide **19** was readily reduced²¹ with Zn/AcOH/Ac₂O to form the 2-amino-*N*-glycopeptide **24** in 85% yield, and thus the present method forms an alternate route to 2-amino-*N*-glycopeptides.

Having explored the conversion of glycals to 1,2-diazides using PIFA- Me_3SiN_3 reagent system, we studied their reactivity with PIDA- Me_3SiN_3 under analogous conditions. As discussed above, the galactal derivative **1** (Scheme 1) upon reaction with PIDA and Me_3SiN_3 (both 1 eq.) gave a mixture of four products **2–5** and thus it was not a synthetically useful reaction. We therefore examined the reaction conditions to incorporate both, the acetate (more nucleophilic compared to trifluoroacetate ion) as well as the azide moieties onto glycals. Our aim was to procure synthetically useful 1-azido-2-acetoxy sugars and/or 2-azido-1-acetoxy sugars, and to avoid the formation of 1,2-diacetates and 1,2-diazides. After exploring various combinations of the azide source and Lewis acids (Table 2), it became clear that a combination of 3 eq. of Me_3SiN_3 and 1 eq. of PIDA along with 30 mol% of TMSOTf at -30 °C was optimum to form vicinal azidoacetates. Thus, galactal derivative **1** led to the formation of an inseparable mixture of two products, 1-azido-2-acetoxy sugar **2** and 2-azido-1-acetoxy sugar **3** in 61% combined yield (Scheme 6). The two compounds were formed in 1 : 2 ratio which was ascertained after selectively reducing the mixture of **2** and **3** with ammonium tetrathiomolybdate (*vide supra*) followed by acetylation that gave chromatographically separable **25** and unreacted **3**.^{22a} Further, 2-azido-1-acetoxy sugar **3** was found to be a mixture of two anomers ($\alpha/\beta = 63/37$). The glucal derivative **8b**, on the other hand, gave a chromatographically separable mixture of **26**^{16a} and **27**^{22a} (α/β anomers = 60/40) in 1 : 2 ratio. Compound **26** was again reduced and acetylated to form **28**^{22b}



Table 2 Optimization of reaction conditions to form vicinal-azidoacetates from galactal derivative 1^a

Entry	Catalyst	N ₃ -Source	Equiv.	Solvent	Temp (°C)	Time (h)	Yield (%)		
							(2 + 3)	4	5
1	BF ₃ ·OEt ₂	TMSN ₃	1	CH ₂ Cl ₂	-30	1	30%	30%	21%
2	BF ₃ ·OEt ₂	TMSN ₃	2	CH ₂ Cl ₂	-30	1	20%	20%	40%
3	BF ₃ ·OEt ₂	TMSN ₃	3	CH ₂ Cl ₂	-30	0.5	52%	—	12%
4	BF ₃ ·OEt ₂	Bu ₄ NN ₃	3	CH ₂ Cl ₂	-30	0.5	46%	—	17%
5	TMSOTf	Bu ₄ NN ₃	3	CH ₂ Cl ₂	-30	0.5	38%	—	30%
6	TMSOTf	TMSN ₃	3	CH ₂ Cl ₂	-30	0.5	61%	—	—

^a Reaction conditions: glycols (0.36 mmol), TMSN₃ (1.10 mmol), PhI(OAc)₂ (0.36 mmol), TMSOTf (0.10 mmol), CH₂Cl₂, -30 °C, N₂ atmosphere, isolated yields after purification by silica gel column chromatography.


which was spectroscopically characterized. Interestingly, the arabinal derivative **10b** (ref. 23) led to a single product **29** in 56% yield whose structure was established based on spectral data and also by single crystal X-ray analysis.^{16b}

Based on these product distributions in the reactions with glycol derivatives **1** and **8b** with PIDA-Me₃SiN₃ reagent system we propose a tentative mechanism as shown in Scheme 7. Accordingly, we expect PIDA to react with excess of Me₃SiN₃ (3 eq.) resulting into species **X** and TMSOAc with the equilibrium being more favorable on the right side. Reaction of glycols **1** and **8b** with the species **X** in presence of TMSOTf should lead to a π -complex **A** which should generate another molecule of TMSOAc on the way to intermediate 'B'. Thus, chances of TMSOAc acting as a preferred nucleophile to form products **3** and **27** appear to be high compared to TMSN₃. Subsequently,


the intermediate **B** will undergo attack of the acetate ion either from α - or β -side (paths a and b) at the anomeric carbon accompanied by an S_Ni type reaction with the azide ion at C-2 to give compounds **3** and **27** with the loss of PhI. Likewise, formation of **2** and **26** can be rationalized by invoking intermediates **C** (a π -complex) and **D** in the reaction of glycols with PIDA. The π -complex **C** will expel a molecule of TMSOAc only upon the formation of the intermediate **D**. Thus at any given time, during this pathway, more amount of Me₃SiN₃ is available to attack at intermediate **D** which leads to the observed products **2** and **26**.

In case of arabinal derivative **10b**, a complex similar to 'C' should form from the β -face to avoid the steric repulsions from the two -OBn groups. This should be followed by azide ion attack in an intermolecular fashion from the axial orientation at the anomeric carbon. Subsequent acetate ion attack at C-2 occurs, in an intramolecular fashion, again from the axial side leading to the observed product **29**. It is however not clear at this moment why the product **29** prefers to have ⁴C₁ conformation as against compounds **11a** and **11b** which prefer ¹C₄ conformations.

We further checked the reactivity of 2,4,6-tri-O-acetylated glycols **6a** and **8a** towards the PIDA-Me₃SiN₃ reagent system. The reaction was extremely sluggish in presence of 30 mol% of TMSOTf even at room temperature for prolonged reaction




Scheme 6 Reaction of glycol derivatives **1**, **8b** and **29** with PIDA-Me₃SiN₃.



Scheme 7 Proposed mechanisms for the reaction of glycols with PIDA and Me₃SiN₃.







Scheme 8 Reaction of glycal derivatives **6a** and **8a** with PIDA- Me_3SiN_3 .

times. Gradual increase in the amount of TMSOTf ultimately led to the use of 1 eq. of it for complete consumption of the starting material at $-30\text{ }^\circ\text{C}$. However, the galactal derivative **6a**, interestingly, led to 2-hydroxy-1-azido sugar **30** as a product in 68% yield (Scheme 8) whose structure was established based on spectral data including COSY, NOE, homonuclear decoupling and DEPT experiments.^{16a} Thus, in NOE experiments, irradiation of H-3 at δ 4.89, led to the enhancement of the signals for protons H-1 and H-5 at δ 4.19 and δ 3.96 respectively. Also, when H-1 at δ 4.19 was irradiated, the signals for H-3 and H-5 were enhanced suggesting that H-1 is axially oriented. In the homonuclear decoupling experiment of **30** when H-4 at δ 5.36 was irradiated, the proton H-3 appeared as a doublet with $J = 3.68\text{ Hz}$ suggesting that H-2 is equatorially oriented. These data support the structure assigned to **30**. Compound **30** was also derivatized to **31** and **32** and their structures confirmed based on spectral analysis. On the other hand, the glucal derivative **8a** led to the Ferrier reaction to form **33**²⁴ in 76% yield as an anomeric mixture ($\alpha : \beta = 60 : 40$).

Mechanistically, in case of 2,4,6-tri-*O*-acetyl galactal **6a**, the intermediate **F** (Scheme 9), resulting from the π -complex **E**, should allow azide ion attack from the β -side leading to the intermediate **G**. It is possible that intermediate **G** then undergoes intramolecular participation by $\text{C}_4\text{-OAc}$ group to form intermediate **I** which then hydrolytically decomposes to give **30**. Since the glucal derivative **8a** does not give similar reaction, instead undergoes the Ferrier reaction, intramolecular participation of $\text{C}_4\text{-OAc}$ group in **6a** becomes crucial.




Scheme 9 Proposed mechanisms for the reaction of glycal **6** with PIDA and Me_3SiN_3 .



Scheme 10 Synthesis of a pseudotriscaccharide **38**.

In view of the importance of 2-amino-*O*-glycosides, 1-acetoxy-2-azido galactose derivative **3** was hydrolyzed²⁵ with benzyl amine to form 2-azido-3,4,6-tri-*O*-benzyl-galactopyranose **34** (Scheme 10). The corresponding trichloroacetimidate **35**, a glycosyl donor, was readily prepared by following a literature procedure²⁶ and was reacted with **36**,²⁷ an orthogonal glycosyl acceptor having a thioglycoside linkage, followed by the click reaction with sugar derived alkyne **14** to form a pseudotriscaccharide **38**.^{16a}

Further, we also show the importance of 1-azido-2-acetoxy sugar derivative **29** in the synthesis of a piperidine triol **44**, that belongs to a class of potential glycosidase inhibitors^{9h,i} (Scheme 11). Thus, the acetate moiety of azidoacetate **29** was deprotected and converted to the MOM derivative **40** via **39** under standard reaction conditions. Subsequently, the



Scheme 11 Synthesis of a piperidine derivative **44** from the azidoacetate **29**.



anomeric azide group was reduced with LiAlH_4 followed by protection of the resulting amine as *N*-nosyl group to give **41** which was cyclized to the piperidine derivative **42** under the Mitsunobu conditions. Deprotections of the nosyl, benzyl and the MOM groups led to the formation of the piperidine **44**, an iminosugar as a potential glycosidase inhibitor.²⁸

Conclusions

In conclusion, we have shown that the reagent system PIFA- Me_3SiN_3 in presence of TMSOTf as a catalyst conveniently converts glycals into the corresponding 1,2-diazido derivatives. These vicinal azido derivatives are converted to 2-azido-*N*-glycopeptides, which are convenient precursors of 2-amino-*N*-glycopeptides, and also into a pseudotrissaccharide **15**. Likewise, glycals are converted into a mixture of 1-azido-2-acetoxy sugars and 2-azido-1-acetoxy sugars using the reagent system PIDA- Me_3SiN_3 in the presence of TMSOTf as a catalyst. The -OAc group of 2-azido-1-acetates can be hydrolyzed and converted into a trichloroacetimidate group for effecting glycosylations. One of the azidoacetates **3** was converted into a pseudotrissaccharide **38** after glycosylation followed by click chemistry. Further, 1-azido-2-acetoxy sugar derivative **29**, derived from the arabinal derivative **10b**, was eventually converted into a piperidine triol **44**, a potential glycosidase inhibitor. Tentative mechanisms have been proposed to account for the formation of different azido sugars.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We thank the Department of Science and Technology, New Delhi, India, for a J. C. Bose National Fellowship (No. SR/S2/JCB-26/2010) to Y. D. V. A. C. thanks the Council of Scientific and Industrial Research, New Delhi for a Senior Research Fellowship.

Notes and references

- (a) S. Mirabella, F. Cardona and A. Goti, *Org. Biomol. Chem.*, 2016, **14**, 5186–5204 and references cited therein; (b) J. Zeng, G. Sun, W. Yao, Y. Zhu, R. Wang, L. Cai, K. Liu, Q. Zhang, X.-W. Liu and Q. Wa, *Angew. Chem., Int. Ed.*, 2017, **56**, 5227–5231.
- (a) C. Pöhner, V. Ullmann, R. Hilpert, E. Samain and C. Unverzagt, *Tetrahedron Lett.*, 2014, **55**, 2197–2200; (b) J. Kerékgyérvó, L. Kalmár, Z. Szurmai, O. Hegyi and G. K. Tóth, *Int. J. Pept. Res. Ther.*, 2012, **18**, 1–5; (c) T. Buskas, S. Ingale and G.-J. Boons, *Glycobiology*, 2006, **16**, 113R–136R; (d) H. Herzner, T. Reipen, M. Schultz and H. Kunz, *Chem. Rev.*, 2000, **100**, 4495–4537; (e) G. K. Rawal, A. Kumar, U. Tawar and Y. D. Vankar, *Org. Lett.*, 2007, **9**, 5171–5174; (f) R. Lahiri, S. Dharuman and Y. D. Vankar, *Chimia*, 2012, **66**, 905–912; (g) V. Ullmann, M. Rädisch, I. Boos, J. Freund, C. Pöhner, S. Schwarzinger and C. Unverzagt, *Angew. Chem., Int. Ed.*, 2012, **51**, 11566–11570; (h) C. M. Kaneshiro and K. Michael, *Angew. Chem., Int. Ed.*, 2006, **45**, 1007–1081; (i) R. Joseph, F. B. Dyer and P. Garner, *Org. Lett.*, 2013, **15**, 732–735.
- (a) L. Marmuse, S. A. Nepogodiev and R. A. Field, *Org. Biomol. Chem.*, 2005, **3**, 2225–2227; (b) V. K. Tiwari, B. B. Mishra, K. B. Mishra, N. Mishra, A. S. Singh and X. Chen, *Chem. Rev.*, 2016, **116**, 3086–3240 and references cited therein; (c) S. Chandrasekaran and R. Ramapanicker, *Chem. Rec.*, 2017, **17**, 63–70; (d) S. Tejera, R. L. Dorta and J. T. Vázquez, *Tetrahedron: Asymmetry*, 2016, **27**, 896–909.
- (a) Y. S. Reddy, A. P. John Pal, P. Gupta, A. A. Ansari and Y. D. Vankar, *J. Org. Chem.*, 2011, **76**, 5972–5984; (b) B. G. Reddy, K. P. Madhusudanan and Y. D. Vankar, *J. Org. Chem.*, 2004, **69**, 2630–2633; (c) G. K. Rawal, S. Rani, K. P. Madhusudanan and Y. D. Vankar, *Synthesis*, 2007, 294–298; (d) T. Cui, R. Smith and X. Zhu, *Carbohydr. Res.*, 2015, **416**, 14–20; (e) M. L. Lepage, A. Bodlenner and P. Compain, *Eur. J. Org. Chem.*, 2013, 1963–1972; (f) Z. Györgydeák, L. Szilágyi and H. Paulsen, *J. Carbohydr. Chem.*, 1993, **12**, 139–163; (g) G. Pastuch-Gawolek, K. Malarz, A. Mrozek-Wilczkiewicz, M. Musioł, M. Serda, B. Czaplinska and R. Musiol, *Eur. J. Med. Chem.*, 2016, **112**, 130–144.
- (a) R. Ahuja, N. K. Singhal, B. Ramanujam, M. Ravikumar and C. P. Rao, *J. Org. Chem.*, 2007, **72**, 3430–3442; (b) E. Honda and D. Y. Gin, *J. Am. Chem. Soc.*, 2002, **124**, 7343–7352.
- (a) R. U. Lemieux and R. M. Ratcliffe, *Can. J. Chem.*, 1979, **57**, 1244–1251; (b) R. R. Schmidt and Y. D. Vankar, *Acc. Chem. Res.*, 2008, **41**, 1059–1073 and references cited therein; (c) R. Bodner, B. K. Marcellino, A. Severino, A. L. Smenton and C. M. Rojas, *J. Org. Chem.*, 2005, **70**, 3988–3996; (d) J. Liu and D. Y. Gin, *J. Am. Chem. Soc.*, 2002, **124**, 9789–9797; (e) J. Liu, V. D. Bussolo and D. Y. Gin, *Tetrahedron Lett.*, 2003, **44**, 4015–4018; (f) D. A. Griffith and S. J. Danishefsky, *J. Am. Chem. Soc.*, 1990, **112**, 5811–5819; (g) B.-Y. Lee, P. H. Seeberger and D. V. Silva, *Chem. Commun.*, 2016, **52**, 1586–1589; (h) L. Corcilius, G. Santhakumar, R. S. Stone, C. J. Capicciotti, S. Joseph, J. M. Matthews, R. N. Ben and R. J. Payne, *Bioorg. Med. Chem.*, 2013, **21**, 3569–3581.
- (a) B. Premdjee, A. L. Adams and D. Macmillan, *Bioorg. Med. Chem. Lett.*, 2011, **21**, 4973–4975; (b) G. Arsequell and G. Valencia, *Tetrahedron: Asymmetry*, 1999, **10**, 3045–3094; (c) C. Brocke and H. Kunz, *Bioorg. Med. Chem.*, 2002, **10**, 3085–3112.
- (a) P. K. Kancharla, Y. S. Reddy, S. Dharuman and Y. D. Vankar, *J. Org. Chem.*, 2011, **76**, 5832–5837; (b) J. Thiem and T. Wiemann, *Angew. Chem., Int. Ed. Engl.*, 1990, **29**, 80–82; (c) Y.-A. Yuan, D.-F. Lu, Y.-R. Chen and H. Xu, *Angew. Chem., Int. Ed.*, 2016, **55**, 534–538; (d) R. Joseph, F. B. Dyer and P. Garner, *Org. Lett.*, 2013, **15**, 732–735; (e) V. Kumar and N. G. Ramesh, *Chem. Commun.*, 2006, 4952–4954; (f) V. Kumar and N. G. Ramesh, *Org. Biomol. Chem.*, 2007, **5**, 3847–3858; (g) S. Xiang, J. Ma, B. K. Gorityala and X.-W. Liu, *Carbohydr. Res.*, 2011, **346**,



- 2957–2959; (h) Y. V. Mironov, A. A. Sherman and N. E. Nifantiev, *Tetrahedron Lett.*, 2004, **45**, 9107–9110.
- 9 (a) A. Palanivel, S. Dharuman and Y. D. Vankar, *Tetrahedron: Asymmetry*, 2016, **27**, 1088–1100; (b) A. Mallick, N. Kumari, R. Roy, A. Palanivel and Y. D. Vankar, *Eur. J. Org. Chem.*, 2014, 5557–5563; (c) S. Dharuman and Y. D. Vankar, *Org. Lett.*, 2014, **16**, 1172–1175; (d) A. A. Ansari, P. Rajasekaran, M. M. Khan and Y. D. Vankar, *J. Org. Chem.*, 2014, **79**, 1690–1699; (e) A. A. Ansari and Y. D. Vankar, *RSC Adv.*, 2014, **4**, 12555–12567; (f) A. A. Ansari and Y. D. Vankar, *J. Org. Chem.*, 2013, **78**, 9383–9395; (g) S. Dharuman, P. Gupta, P. K. Kancharla and Y. D. Vankar, *J. Org. Chem.*, 2013, **78**, 8442–8450; (h) Y. S. Reddy, P. K. Kancharla, R. Roy and Y. D. Vankar, *Org. Biomol. Chem.*, 2012, **10**, 2760–2773; (i) P. Gupta and Y. D. Vankar, *Eur. J. Org. Chem.*, 2009, 1925–1933.
- 10 (a) A. Yoshimura and V. V. Zhdankin, *Chem. Rev.*, 2016, **116**, 3328–3435; (b) V. V. Zhdankin, *ARKIVOC*, 2009, **i**, 1–62.
- 11 (a) A. Kirschning, C. Plumeier and L. Rose, *Chem. Commun.*, 1998, 33–34; (b) P. Pandit, K. S. Gayen, S. Khamarui, N. Chatterjee and D. K. Maiti, *Chem. Commun.*, 2011, **47**, 6933–6935; (c) M. Islam, N. D. Tirukoti, S. Nandi and S. Hotha, *J. Org. Chem.*, 2014, **79**, 4470–4476; (d) T. R. Reddy, D. S. Rao, K. Babachary and S. Kashyap, *Eur. J. Org. Chem.*, 2016, 291–301.
- 12 L. Shi, Y.-J. Kim and D. Y. Gin, *J. Am. Chem. Soc.*, 2001, **123**, 6939–6940.
- 13 (a) K. Wu, Y. Liang and N. Jiao, *Molecules*, 2016, **21**, 1–21; (b) B. B. Snider and H. Lin, *Synth. Commun.*, 1998, **28**, 1913–1922; (c) M.-Z. Lu, C.-Q. Wang and T.-P. Loh, *Org. Lett.*, 2015, **17**, 6110–6113; (d) D. A. Kamble, P. U. Karabal, P. V. Chouthaiwale and A. Sudalai, *Tetrahedron Lett.*, 2012, **53**, 4195–4198; (e) G. Fumagalli, P. T. G. Rabet, S. Boyd and M. F. Greaney, *Angew. Chem., Int. Ed.*, 2015, **54**, 11481–11484.
- 14 (a) Y. V. Mironov, A. A. Sherman and N. E. Nifantiev, *Tetrahedron Lett.*, 2004, **45**, 9107–9110; (b) Y.-A. Yuan, D.-F. Lu, Y.-R. Chen and H. Xu, *Angew. Chem., Int. Ed.*, 2016, **55**, 534–538.
- 15 (a) Y. Kita, H. Tohma, K. Hatanaka, T. Takada, S. Fujita, S. Mitoh, H. Sakurai and S. Oka, *J. Am. Chem. Soc.*, 1994, **116**, 3684–3691; (b) P. Li, J. Zhao, C. Xia and F. Li, *Org. Chem. Front.*, 2015, **2**, 1313–1317; (c) Y. Kita, H. Tohma, T. Takada, S. Mitoh, S. Fujita and M. Gyoten, *Synlett*, 1994, 427–428; (d) K. Matcha and A. P. Antonchick, *Angew. Chem., Int. Ed.*, 2013, **52**, 2082–2086; (e) R. Narayan and A. P. Antonchick, *Chem.–Eur. J.*, 2014, **20**, 4568–4572; (f) K. Matcha, R. Narayan and A. P. Antonchick, *Angew. Chem., Int. Ed.*, 2013, **52**, 7985–7989.
- 16 (a) See the ESI† for spectral details; (b) CCDC 1554252 for compound 5 and CCDC 1554251 for compound 29.
- 17 (a) For the formation of $\text{PhI}(\text{N}_3)_2$ from $\text{PhIO-Me}_3\text{SiN}_3$ and its utility in presence of TEMPO via radical pathway see: (i) P. Magnus, M. B. Roe and C. Hulme, *J. Chem. Soc., Chem. Commun.*, 1995, 263–265; (ii) P. Magnus, J. Lacour, P. A. Evans, M. B. Roe and C. Hulme, *J. Am. Chem. Soc.*, 1996, **118**, 3406–3418; (b) R. M. Moriarty and J. S. Khosrowshahi, *Tetrahedron Lett.*, 1986, **27**, 2809–2812; (c) A. Kirschning, *Eur. J. Org. Chem.*, 1998, 2267–2274.
- 18 A. Mishra and V. K. Tiwari, *J. Org. Chem.*, 2015, **80**, 4869–4881.
- 19 P. R. Sridhar, K. R. Prabhu and S. Chandrasekaran, *J. Org. Chem.*, 2003, **78**, 5261–5264.
- 20 (a) J. L. Torres, I. Haro, E. Bardaji, G. Valencia, J. M. Garcia-Anton and F. Reig, *Tetrahedron*, 1988, **44**, 6131–6136; (b) A. Bianchi and A. Bernardi, *J. Org. Chem.*, 2006, **71**, 4565–4577; (c) A. D. Dorsey, J. E. Barbarow and D. Trauner, *Org. Lett.*, 2003, **5**, 3237–3239.
- 21 R. J. Payne, S. Ficht, S. Tang, A. Brik, Y.-Y. Yang, D. A. Case and C.-H. Wong, *J. Am. Chem. Soc.*, 2007, **129**, 13527–13536.
- 22 (a) N. V. Bovin, S. E. Zurabvan and A. Y. Khorlin, *Carbohydr. Res.*, 1981, **98**, 25; (b) A. A. Pavia, S. N. Ung-Chhun and J. L. Durand, *J. Org. Chem.*, 1981, **46**, 3158–3160.
- 23 K. Matsuo and M. Shindo, *Tetrahedron*, 2011, **67**, 971–975.
- 24 S. Chandrasekhar, C. R. Reddy and G. Chandrashekar, *Tetrahedron Lett.*, 2004, **45**, 6481–6484.
- 25 S. Kopitzki and J. Thiem, *Eur. J. Org. Chem.*, 2013, 4008–4016.
- 26 (a) R. R. Schmidt and J. Michel, *Angew. Chem., Int. Ed.*, 1980, **19**, 731–732; (b) J. Kalikanda and Z. Li, *J. Org. Chem.*, 2011, **76**, 5207–5218.
- 27 L. X. Wang, N. Sakairi and H. Kuzuhara, *J. Chem. Soc., Perkin Trans. 1*, 1990, 1677–1682.
- 28 X. Garrabou, J. A. Castillo, C. G. Hélaïne, T. Parella, J. Joglar, M. Lemaire and P. Clapés, *Angew. Chem., Int. Ed.*, 2009, **48**, 5521–5525.

