

Cite this: *RSC Adv.*, 2017, 7, 46932

Configuration and stability of naturally occurring all-*cis*-tetrahydrofuran lignans from *Piper solmsianum*†

Clécio S. Ramos,^a Harrauld V. Linnert,^b Marcilio M. de Moraes,^b João H. do Amaral,^b Lydia F. Yamaguchi^b and Massuo J. Kato^b ^{*,b}

The study of variability of the tetrahydrofuran lignan (–)-grandisin in leaves of *Piper solmsianum* (Piperaceae) revealed two unknown compounds, that were isolated and determined to be the all-*cis* tetrahydrofuran lignans **1a** [rel-(7*R*,8*S*,7'*S*,8'*R*)-3,4,5,3',4',5'-hexamethoxy-7,7'-epoxy] and **1b** [rel-(7*R*,8*S*,7'*S*,8'*R*)-3',4'-methylenedioxy-3,4,5,5'-tetramethoxy-7,7'-epoxy]. Their structures were determined by spectroscopic analysis including 1D and 2D-NMR while their configurations were determined by ECD associated to the density functional theory (DFT) at the B3LYP/6-31G(d,p) level. The hydrogen bonds between methoxy groups in trimethoxyphenyl rings stabilizes the all-*trans* tetrahydrofuran lignan grandisin by 6.5 kcal mol⁻¹ as compared to the corresponding all-*cis* isomer of grandisin. The occurrence of all-*cis* tetrahydrofuran lignans as natural products is a very rare event.

Received 21st August 2017
Accepted 29th September 2017

DOI: 10.1039/c7ra09262h


rsc.li/rsc-advances

Introduction

Lignans are dimers of phenylpropanoid units widely found in the plant kingdom and with a variety of biological properties such as cytotoxic, antioxidant, antimicrobial, anti-inflammatory, antiviral and immunosuppressive activities.^{1,2} The dimerization process in their formation requires an oxidative coupling mechanism forming a β–β bonding, which can be followed by further reactions yielding eight different sub-classes including tetrahydrofuran, furofuran, dibenzylbutane, dibenzylbutyrolactone, aryltetralin, aryl-naphthalene, dibenzocyclooctadiene and dibenzylbutyrolactol lignans.^{3–5} Tetrahydrofuran (THFs) lignans can be further subdivided in 7,7'-monoepoxy-, 7,9'-monoepoxy- and 9,9'-monoepoxy-lignans. The THFs lignans have been described so far from several plant species and their structural diversity results from the type of substituent in the aromatic ring that can be hydroxy, methoxy, methylenedioxy and a combination of them. The relative configuration among aryls and methyl groups can give rise to further stereochemistry diversity. In case of having the same aromatic rings, ten isomers are possible consisting of four diastereomeric pairs besides two *meso* forms.⁶ The (–)-grandisin is an example of the all-*trans* tetrahydrofuran lignan, which was formerly isolated from the Lauraceae plants, *Litsea grandis*⁷ and then from *Licaria aurea*⁸ and *Cryptocarya crassinevia*.⁹ Grandisin

has also been isolated from monocotyledonous Araceae,¹⁰ from other Angiosperms families such as Piperaceae,¹¹ Myristicaceae,¹² Magnoliaceae,¹³ Annonaceae¹⁴ but also from Rosids including Meliaceae,¹⁵ Violaceae species,¹⁶ and other plant species. The THFs lignans have displayed several biological activities such as anti-PAF,¹⁷ antiprotozoal,^{18–21} larvicidal,^{22,23} cytotoxic,^{24,25} estrogenic,²⁵ anti-inflammatory,²⁶ and NF-κB inhibitory activity.²⁷ Thus, some studies have also been addressed to generate analogues.^{28–31} To date, the natural occurrence of all-*cis* tetrahydrofuran lignan has only been described from *Anogeissus acuminata* (Combretaceae)³² and from *Illicium floridanum* (Illiciaceae).³³ Besides, a synthetic version of all-*cis* had been obtained by catalytic hydrogenation of a furan derivative.³⁴ Additionally, a series of synthetic tetrahydrofuran lignans with various stereochemistry were prepared, including the all-*cis* tetrahydrofuroguaiacin B, in order to evaluate their plant growth inhibitory activity.³¹ Herein we report on the isolation and structural determination of two novel all-*cis* tetrahydrofuran lignans **1a** and **1b** from the leaves of two specimens of *Piper solmsianum*. Besides, two known lignans **2b** and **3a** were also isolated (Fig. 1). The structures of the last two tetrahydrofuran lignans had been only reported without the assignment of their configurations or optical rotations.^{10,15,35} The determination of configuration of THFs lignans (**1a**, **1b**, **2b** and **3a**) were carried out using electronic circular dichroism (ECD) having (–)-grandisin as reference data since its configuration was previously reported based on crystallographic studies.¹⁴ Additionally, since natural all-*cis* tetrahydrofuran lignans, such as **1a** and **1b**, are of rare occurrence a study of structural stability were undertaken using density functional theory (DFT) at the B3LYP/6-31G(d,p) level. The relative energies of lignans **1a** and **3a** were compared to that of all-*trans* lignan (–)-grandisin (**2a**). Solvent effects were analysed by the self-

^aDepartment of Chemistry, Rural Federal University of Pernambuco, Rua Dom Manoel de Medeiros, s/n, 52.171-030 Recife, PE, Brazil^bInstitute of Chemistry, University of São Paulo, Av. Prof. Lineu Prestes, 748, 05508-000 São Paulo, SP, Brazil. E-mail: majokato@iq.usp.br† Electronic supplementary information (ESI) available: NMR spectra of **1a** and **1b**; Fig. S1–S11, Tables S1–S7. See DOI: 10.1039/c7ra09262h

Ar₂=3,4,5-trimethoxyphenylAr₃=3,4-methylenedioxy-5-methoxyphenylFig. 1 Tetrahydrofuran lignans from *P. solmsianum*.

consistence reaction field (SCRF) method, and the energy difference in water and cyclohexane environments were determined with the polarizable continuum model PCM(B3LYP/6-31G(d,p)) method.

Results and discussions

Analysis of crude extracts from *P. solmsianum*

The tetrahydrofuran lignan (–)-grandisin has displayed several bioactivities and thus, the initial approach was to determine its variability in specimens of *P. solmsianum* occurring in several

sites in São Paulo State. The initial survey in samples from leaves of different sites revealed common profiles based on the occurrence of phenylpropanoids apiole–dillapiole and grandisin (**2a**). Then, the analysis of chemical variability of a population of *P. solmsianum* occurring in the Campus of University of São Paulo was undertaken and several samples collected presented an unusual chromatographic profile with one major peak with shorter retention time than that of grandisin. These extracts were submitted to ¹H NMR analysis, which suggested the occurrence of a major tetrahydrofuran lignan (**1a**) with similar oxygenation pattern in the aromatic rings with six methoxy groups. Crude extracts from different *P. solmsianum* individuals collected in Alberto Löffgren State Park analysed by HPLC indicated a second tetrahydrofuran lignan, having methoxy and 3,4-methylenedioxy groups (**1b**). Thus, both compounds were isolated and had their structures determined by spectroscopic analysis.

Structural determination of lignans **1a**, **1b**, **2b** and **3a**

The lignan **1a** was isolated as a pale white powder with purity of 99.6% by HPLC. Its HR-ESIMS spectrum displayed a quasi-molecular ion [M + H]⁺ peak at *m/z* 433.2232 corresponding to

Table 1 NMR spectral data^a for lignans **1a** and **1b**

Position	1a				1b			
	δ ¹ H	COSY	δ ¹³ C	HMBC	δ ¹ H	COSY	δ ¹³ C	HMBC
1	—	—	137.0	C6, C2	—	—	135.3	C6, C2
2	6.67 (s, 1H)	H7	103.6	C6, C7, C8, C4, C3	6.65 (s, 1H)	H7	103.5	C6, C7
3	—	—	153.1	C2, OCH ₃ -3	—	—	153.0	C2, OCH ₃ -3
4	—	—	136.0	C-6, OCH ₃ -4	—	—	137.5	C6, OCH ₃ -4
5	—	—	153.1	C-6, OCH ₃ -5	—	—	153.0	C6, OCH ₃ -5
6	6.67 (s, 1H)	H7	103.6	C2, C7, C8, C4, C5	6.65 (s, 1H)	H7	103.5	C2, C7
7	5.11 (d, <i>J</i> 6.7 Hz, 1H)	H8, H9, H2, H6	82.9	C9, C8', C2, C3, C1	5.09 (d, <i>J</i> 7.0 Hz, 1H)	H8, H9 H2, H6	82.8	C9, C2, C3
8	2.70 (m, 1H)	H7, H9	41.6	C7, C7', C8', C9, C9', C1	2.67 (m, 1H)	H7, H9	41.5	C7, C7', C8', C9, C9'
9	0.64 (d, <i>J</i> 6.7 Hz, 3H)	H8, H7	11.6	C9', C8, C8', C7, C1	0.64 (d, <i>J</i> 6.9 Hz, 3H)	H8, H7	11.6	C8, C8', C7
1'	—	—	137.0	C6', C2'	—	—	136.1	C6', C2'
2'	6.67 (s, 1H)	H7'	103.6	C6', C7', C8', C4', C3'	6.63 (s, 1H)	H7'	105.8	C6', C7'
3'	—	—	153.1	O ₂ CH ₂ , C2'	—	—	143.4	O ₂ CH ₂ , C2'
4'	—	—	136.0	O ₂ CH ₂ , C6'	—	—	139.8	O ₂ CH ₂ , C6'
5'	—	—	153.1	C6', OCH ₃ -5'	—	—	148.8	C6', OCH ₃ -5'
6'	6.67 (s, 1H)	H7'	103.6	C2', C7', C8', C4', C5'	6.61 (s, 1H)	H7'	100.7	C2', C7'
7'	5.11 (d, <i>J</i> 6.7 Hz, 1H)	H8', H9', H2', H6'	82.9	C9', C8, C2', C3', C1'	5.09 (d, <i>J</i> 7.0 Hz, 1H)	H8', H9', H2', H6'	82.9	C9', C2', C3'
8'	2.70 (m, 1H)	H7', H9'	41.6	C7, C7', C8, C9, C9', C1'	2.67 (m, 1H)	H7', H9'	41.6	C7, C7', C8, C9, C9'
9'	0.64 (d, <i>J</i> 6.7 Hz, 3H)	H8', H7'	11.6	C9, C8, C8', C7', C1'	0.62 (d, <i>J</i> 6.9 Hz, 3H)	H8', H7'	11.7	C8, C8', C7'
OCH ₃ /3	3.88 (s, 3H)	—	56.1	C3	3.88 (s, 3H)	—	56.1	—
OCH ₃ /3'	3.88 (s, 3H)	—	56.1	C3'	—	—	—	—
OCH ₃ /4	3.86 (s, 3H)	—	60.9	C4	3.86 (s, 3H)	—	60.9	—
OCH ₃ /4'	3.86 (s, 3H)	—	60.9	C4'	—	—	—	—
OCH ₃ /5	3.88 (s, 3H)	—	56.1	C5	3.88 (s, 3H)	—	56.1	—
OCH ₃ /5'	3.88 (s, 3H)	—	56.1	C5'	3.91 (s, 3H)	—	56.6	—
O ₂ CH ₂	—	—	—	—	5.98 (s, 2H)	—	101.3	—

^a 500 MHz for ¹H and 125 MHz for ¹³C; compounds dissolved in CDCl₃; δ in ppm relative to TMS, *J* in Hz.



the molecular formula $C_{24}H_{33}O_7$. The IR of **1a** showed no absorptions near 3400 cm^{-1} , but an intense band at 1128 cm^{-1} suggesting C–O–C functions. The simplicity of the ^1H and ^{13}C NMR spectra (Fig. S1 and S2; ESI†) indicated that **1a** is a symmetrical compound (Table 1). The dimethyl-tetrahydrofuran moiety could be readily recognized in the ^1H NMR spectrum by the doublets at δ_{H} 0.64 (6H, $J = 6.7\text{ Hz}$, H-9/H-9') and 5.11 (2H, $J = 6.7\text{ Hz}$, H-7/H-7'), and a multiplet at δ_{H} 2.70 (2H, H-8/H-8'). The singlet at δ_{H} 6.74 (H-2/H-2', H-6/H-6') was assigned to the two equivalent *ortho*-methoxy aromatic hydrogens, while the intense singlets at δ_{H} 3.88 ($4 \times \text{OCH}_3$; 3,3',5,5') and 3.86 ($2 \times \text{OCH}_3$; 4,4') corresponded to the six methoxy groups. The structure of **1a** bearing two 3,4,5-trimethoxyphenyl groups was firmly supported by further ^{13}C NMR analysis (Table 1).^{3,35,36} All NMR assignments were confirmed by HSQC and HMBC correlations in which cross peaks were observed between the aromatic and aliphatic nuclei (Fig. S3 and S4†).

Considering the symmetry observed in the ^1H and ^{13}C NMR spectra and that **1a** is optically inactive, the compound was determined as a *meso* isomer with configuration similar to the all-*cis* (**1c**) or *trans-cis-trans* (**5**) (Table S1†). The determination of the relationship between the methyl and the aryl groups was based on the chemical shifts of methyl (H-9/H-9') and oxymethine (H-7/H-7') protons, and of the oxybenzylic carbons (C-7/C-7') (Tables 1 and S1†).^{3,8,32} The protons at H-9/H-9' and H-7/H-7' of **1a** showed signals at δ_{H} 0.64 and 5.11, respectively and the coupling constants of J 6.7 Hz for the H-7/H-8 (H-7'/H-8') hydrogens is compatible with a dihedral angle of approximately 30° , indicating all-*cis* relationship between the methyl and aryl substituents.³² The up-field shifted methyl carbons at δ_{C} 11.6 in the ^{13}C NMR is contrasting to the all-*trans* isomer grandisin (**2a**), which appeared at δ_{C} 13.9. This shielding effect has been assigned to the anisotropic effect of the pseudo-axially oriented aromatic ring on the methyl groups of **1a**. The chemical shifts of C-7/C-7' observed at δ_{C} 82.9, and of C-8/C-8' at δ_{C} 41.6, provided further evidences for the *cis* configurations between the methyl and aryl groups of **1a** (Tables 1 and S1†). All the ^1H and ^{13}C NMR data indicated that the lignan **1a** is all-*cis* isomer of **2a** (grandisin)³⁵ thus, the structure was determined as *rel*-(7*R*,8*S*,7'*S*,8'*R*)-3,4,5,3',4',5'-hexamethoxy-7,7'-epoxylignan.

Compound **1b** was isolated as pale-yellow oil by preparative TLC of a methanolic extract of the leaves of *P. solmsianum* with 99.2% purity indicated by HPLC analysis. The molecular formula of **1b** was determined as $C_{23}H_{28}O_7$ according to the quasi-molecular ion $[\text{M} + \text{H}]^+$ observed in the HR-ESIMS at m/z 417.1915.




Fig. 2 Important ^1H – ^{13}C observed in the HMBC correlations of lignan **1b**.

Analysis of the ^1H and ^{13}C NMR spectra of **1b** indicated the structure of an all-*cis* tetrahydrofuran lignan similarly to **1a** (Table 1; Fig. S5 and S6†). The major differences were associated to the replacement of two methoxy by one methylenedioxy group, which account for the difference of 16 Da between the molecular ions of **1a** and **1b**. The chemical shift of the equivalent aromatic hydrogens of the 3,4,5-trimethoxyphenyl ring (δ_{H} 6.65, s, 2H) was similar to that of **1a**, while the hydrogens of the second aromatic ring appeared as two broad singlets at δ_{H} 6.63 and 6.61 and were assigned to the two aromatic hydrogens of the 3-methoxy-4,5-methylenedioxyphenyl ring. The HMBC contour plot (Fig. S8†) displayed correlations between the oxymethine protons (H-7/H-7') to C-6/C-6' and C-2/C-2', and between the H-7/H-7' hydrogens to signals of C-9/C-9' (Fig. 2). All NMR data, including HSQC (Fig. S7†), confirmed the assignments made and thus **1b** was determined as *rel*-(7*R*,8*S*,7'*S*,8'*R*)-3',4'-methylenedioxy-3,4,5,5'-tetramethoxy-7,7'-epoxylignan.

Compound **2b** has been reported previously as *rel*-(7*R*,8*S*,7'*S*,8'*R*)-3',4'-methylenedioxy-3,4,5,5'-tetramethoxy-7,7'-epoxylignan.³⁵

The compound **3a**, determined as (–)-epigrandisin based on the spectrometric data analysis, is a lignan formerly isolated from the stem bark of *Aglaia leptantha* but characterized in a mixture with (–)-grandisin (**2a**) with an estimated $[\alpha]_{\text{D}}^{21}$ value of -15 ± 10 .¹⁵ (–)-Epigrandisin, without configuration assigned, was also reported from leaves of *Rhaphidophora decursiva*.¹⁰

Determination of relative configurations of lignans **1a**, **1b**, **2b** and **3a**

The relative configurations of the THF lignans **1a**, **1b**, **2b** and **3a** were determined by comparison of ECD spectra with that of (–)-grandisin (**2a**) (Fig. 3), in which its configuration had been previously determined by X-ray analysis.⁷ Lignan **1a** is optically inactive and no Cotton effect was observed in its ECD curve. The ECD curve of lignan **2b** was very similar to that of (–)-grandisin (**2a**) with a negative Cotton effect at 233 nm, indicating the *trans* configuration between the aromatic rings. In contrast, the lignans **1b** and **3a** showed a positive Cotton effect at 220 nm in the ECD curve, *i.e.* opposite to that of **2a**, hence the *cis* configuration between the aromatic rings were assigned (Fig. 3).

Energetics of the lignans in gas phase

Optimization of the tetrahydrofuran lignans (gas phase) structures was performed with the aid of Gaussian 03 (revision B.05) software,³⁷ using DFT at the B3LYP level with the 6-31G(d,p) basis set. Becke's three parameter adiabatic connection approach is well established to provide structural parameters for large molecules including the intramolecular and intermolecular hydrogen bondings.^{38,39} Harmonic frequencies were calculated in order to confirm the equilibrium geometries corresponding to minima energy and zero point energy (ZPE). Correction term was added to the electronic energies of the calculated structures to estimate differences in the relative energies between the lignans **1a**, **2a** and **3a** (Table S2†).

In order to predict the energy and the solvent effect on lignans stability in polar and apolar environments, a single






Fig. 3 ECD curves of tetrahydrofuran lignans from *P. solmsianum*.

point energy calculation was performed, with the PCM (B3LYP) SCRF models using the 6-31G(d,p) basis set. The tight SCF convergence criteria were specified for all jobs and the predicted energy and solvent effects in cyclohexane and water media were calculated (ESI, Table S3†).

The calculations at the B3LYP/6-31G(d,p) level indicated lignan **2a** as the most stable among the lignans **1a** and **3a** in gas phase, cyclohexane and water. An exception was observed for lignans **2a** and **3a**, which have similar electronic energy in water medium (Tables S2 and S3†). The calculations showed that inclusion of solvent effects does not change the overall relative results obtained in the gas phase. Furthermore, the stabilities increased with the solvent polarity, and only the change in the energy produced in the high polar medium (*i.e.*, water) was more effective in stabilizing the conformers as compared to the gas phase or apolar solvent (*i.e.*, cyclohexane).

Gas phase optimization geometry of the THF lignans

A remarkable feature in the analysis of the geometrical parameters of **1a**, **1b**, **2a**, **2b** and **3a** lignans is the stabilization resulting from intramolecular C–H⋯O hydrogen bonding.

Energy calculations predict that the all-*trans* THF lignan **2a** would be 0.42 kcal mol⁻¹ more stable than epigrandisin **3a**, and 6.5 kcal mol⁻¹ more stable than the all-*cis* THF lignan **1a**. Such sequence of stability is somewhat compatible with the rare natural occurrence of all-*cis* THF lignan isomer. In fact, there are only two previous report for it.^{32,33} It is also noteworthy that the methyl group (C9') attached to the THF ring lies close to the aromatic rings in the all-*cis* lignan **1a**, revealing one further possible hydrogen bonding contributing to an unfavourable gauche conformation around the C8' bond (Fig. S9†). Accordingly, the calculated angle at C8–C8'–C7' is 3° smaller than the corresponding C7–C8–C8' angle at the THF ring, a feature also observed in **1b**. Additionally, the phenyl rings in **1a** lie closer to the THF ring compared with the all-*trans* isomer, resulting in smaller dihedral angles, which is less favourable.

The results concerning structural changes associated to the phenyl rings suggested that the geometrical equilibrium structures of **1b** and **2b** are dependent on intramolecular hydrogen bonding. On the other hand, the geometrical energy minimization of **2b** taking into account both hydrogens but not hydrogen bonding (a difference of ~1 kcal mol⁻¹), showed that these additional hydrogen bonds are not energetically significant. It is also worth to note that in these geometry optimizations, the methoxy groups attached to each of the phenyl ring do not necessary lie close to the ring plane, but assumed the most favourable hydrogen bonding with conformers differing from those of crystallographic data.¹⁴

The calculated equilibrium geometries for lignans **1a–2a** are shown in Fig. S9 and S10 (ESI†), and the main geometrical parameters, obtained after minimized energies using the B3LYP method, are presented in Tables S4–S7 (ESI†).

Relative energies of the THF isomers without methoxy groups in the phenyl rings

In order to compare the behavior of the THF lignan isomers, a singular qualitative calculation was made for the ten possible tetrahydrofuran isomers (four diastereomeric pairs and two *meso* forms) (Table S8†) with a lower level of theory as B3LYP/3-21G*. The methoxy groups were not included in the calculation in order to minimize the effect of the hydrogen bondings in the stabilities and thus to determine only the effect of configuration in the stabilities of THF rings. The electronic energy of the optimized structures indicates that the all *trans* and the all *cis* structures are the most and the least stable among the ten stereoisomers, respectively (Table S8†). With this finding, we can predict that the configuration and the degree of methoxylation contribute to the stabilization of the THFs lignans.

Conclusions

The structures and absolute configurations of two novel THF lignans (**1a**, **1b**), isolated from leaves of the species *P. solmsianum*, and of two known lignans (**2b**, **3a**),^{10,15,35} were determined. In addition, the theoretical electronic calculations provided a new data set for the structures of the THF lignans in gas phase showing that the configuration of the tetrahydrofuran



ring and intramolecular hydrogen bonding are related to the geometrical equilibrium energy profile of the conformers. The calculated energies in the gas phase and in solvents (water and cyclohexane) indicated that lignan **2a** is the most stable among the isomeric lignans **1a–3a**. Moreover, the calculated data for **1b** and **2b** could explain the theoretical equilibrium geometries of these lignans since the configuration of the THF ring and intramolecular hydrogen bonding originated from the methoxy groups in the phenyl ring, compared to the methylenedioxy moiety, contributed to the stabilization of the THF lignan structures.

Experimental

Plant material

Leaves of *Piper solmsianum* C. DC were collected at the Instituto de Biociências (IB), Universidade de São Paulo (SP, Brazil) and at the Horto Florestal (HF), Instituto Florestal do Estado de São Paulo (SP, Brazil) in January 2003 (Permit ICMBio-15780-1). Plants were identified by Dr Elsie Franklin Guimarães (Instituto de Pesquisas Jardim Botânico do Rio de Janeiro, RJ, Brazil), and voucher specimens were deposited at the herbarium of the Instituto de Pesquisas Jardim Botânico do Rio de Janeiro (IB specimen – Kato-0367; HF specimen – Kato-0175).

Extraction and isolation of lignan **1a**

Dried leaves (3.2 g) of *P. solmsianum* (IB specimens) were milled and the powder extracted with CH_2Cl_2 for 48 h. The extract was concentrated under vacuum to yield 300 mg of a residue that was submitted to CC over C18 column (7 g, 15×2.5 cm i.d.). Elution with mixtures of $\text{H}_2\text{O} : \text{MeOH}$ containing increasing amounts of MeOH (from 50 to 100%) gave 10 fractions. Fractions 3 and 4 (eluted with 70% MeOH) were pooled, extracted with CH_2Cl_2 (3×100 mL), the organic fractions dried over anhydrous Na_2SO_4 and concentrated under vacuum to yield **1a** (6 mg). Data for compound **1a**: pale white powder; $[\alpha]_{\text{D}}^{21} 0$ ($c 1 \times 10^{-4}$ mol L^{-1} , CH_2Cl_2); UV λ_{max} 225 nm ($c 3.0 \times 10^{-2}$ mol L^{-1} MeOH); CD $\theta_{220 \text{ nm}} + 6.6$ ($c 2.5 \times 10^{-3}$ mol L^{-1} MeOH); IR $\nu_{\text{max}}^{\text{film}}$ cm^{-1} : 2965, 2930, 2837, 1593, 1127, 1095, 843; ^1H and ^{13}C NMR: see Table 1; EI/MS m/z (rel. int): 432 $[\text{M}]^+$ (31), 236 (100), 221 (70), 205 (83); HR-ESIMS $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{24}\text{H}_{33}\text{O}_7$ 433.2221, found 433.2232. Anal. calcd for $\text{C}_{24}\text{H}_{32}\text{O}_7$: C, 66.66; H, 7.40; O, 25.92. Found C, 66.57; H, 7.25; O 25.88.

Extraction and isolation of lignans **1b**, **2b** and **3a**

Dried leaves (6 g) of *P. solmsianum* (HF specimens) were milled and the powder extracted with methanol for 48 h. The extract was concentrated under vacuum to yield 400 mg of a residue that was suspended in 150 mL of $\text{MeOH} : \text{H}_2\text{O}$ (4 : 1), filtered over a bed of Celite, and extracted with CH_2Cl_2 (3×100 mL). The CH_2Cl_2 fraction was dried over anhydrous Na_2SO_4 , concentrated under vacuum to yield 70 mg of residue that was submitted to preparative TLC developed twice with hexanes : EtOAc (85 : 15) to afford **1b** (10 mg), **2b** (3 mg) and **3a** (5 mg). Data for compound **1b**: pale yellow oil; $[\alpha]_{\text{D}}^{21} + 80.7$ ($c 1 \times 10^{-4}$ mol L^{-1} , CH_2Cl_2); UV λ_{max} 225 nm ($c 3.0 \times 10^{-2}$ mol L^{-1}

MeOH); CD $\theta_{220 \text{ nm}} + 6.6$ ($c 2.5 \times 10^{-3}$ mol L^{-1} MeOH); IR $\nu_{\text{max}}^{\text{film}}$ cm^{-1} : 2965, 2935, 1634, 1507, 1233, 1228, 1095, 843; ^1H and ^{13}C NMR: see Table 1; EI/MS m/z (rel. int): 416 $[\text{M}]^+$ (14), 220 (100), 205 (85), 236 (43), 208 (21); HR-ESIMS $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{23}\text{H}_{28}\text{O}_7$ 417.1907, found 417.1915. Data for compound **2b**: pale white powder; $[\alpha]_{\text{D}}^{21} - 62.2$ ($c 1 \times 10^{-4}$ mol L^{-1} , CH_2Cl_2); UV λ_{max} 240 nm ($c 5.0 \times 10^{-2}$ mol L^{-1} MeOH); CD $\theta_{233 \text{ nm}} - 10.5$ ($c 3.0 \times 10^{-3}$ mol L^{-1} MeOH); IR $\nu_{\text{max}}^{\text{film}}$ cm^{-1} : 2965, 2935, 1634, 1507, 1233, 1228, 1095, 843; ^1H and ^{13}C NMR;³⁶ EI/MS m/z (rel. int): 432 $[\text{M}]^+$ (12), 205 (100), 2221 (85), 236 (90), 208 (30); HR-ESIMS $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{24}\text{H}_{33}\text{O}_7$ 433.2221 found 433.2234. Data for compound **3a**: pale yellow oil; $[\alpha]_{\text{D}}^{21} - 71.4$ ($c 1 \times 10^{-4}$ mol L^{-1} , CH_2Cl_2); ^1H and ^{13}C NMR.¹⁵

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was funded by grants provided by FAPESP (09/51850-9, 14/50316-7), CNPq, and PRP-USP (Pró-Reitoria de Pesquisas da USP). MJK and HVL are grateful to CNPq for research fellowships, and CSR acknowledges FAPESP for a scholarship. The authors are indebted to the Laboratório de Computação Científica Avançada (LCCA), Universidade de São Paulo (SP, Brazil) for computer facilities.

References

- R. S. Ward, *Nat. Prod. Rep.*, 1999, **16**, 75–96.
- J. Y. Pan, S. L. Chen, M. H. Yang, J. Wu, J. Sinkkonen and K. Zou, *Nat. Prod. Rep.*, 2009, **26**, 1251–1292.
- P. K. Agrawal and R. S. Thakur, *Magn. Reson. Chem.*, 1985, **23**, 389–418.
- D. A. Whiting, *Nat. Prod. Rep.*, 1985, **2**, 191–211.
- L. B. Davin, M. Jourdes, A. M. Patten, K. W. Kim, D. G. Vassão and N. G. Lewis, *Nat. Prod. Rep.*, 2008, **25**, 1015–1090.
- T. Biftu and R. Stevenson, *Phytother. Res.*, 1987, **1**, 97–106.
- D. Holloway and F. Scheinmann, *Phytochemistry*, 1974, **13**, 1233–1236.
- J. M. Barbosa, M. S. da Silva, M. Yoshida and O. R. Gottlieb, *Phytochemistry*, 1989, **28**, 2209–2211.
- J. M. Saad, E. Soepadamo, X. P. Fang, J. L. McLaughlin and P. E. Fanwick, *J. Nat. Prod.*, 1991, **54**, 1681–1683.
- H. J. Zhang, P. A. Tamez, V. D. Hoang, G. T. Tan, N. Van Hung, L. T. Xuan, L. M. Huong, N. M. Cuong, D. T. Thao, D. D. Soejarto, H. H. S. Fong and J. M. Pezzuto, *J. Nat. Prod.*, 2001, **64**, 772–777.
- C. S. Ramos, S. A. Vanin and M. J. Kato, *Phytochemistry*, 2008, **69**, 2157–2161.
- N. P. Lopes, M. J. Kato and M. Yoshida, *Phytochemistry*, 1999, **51**, 29–33.
- M. Kuroyanagi, K. Yoshida, A. Yamamoto and M. Miwa, *Chem. Pharm. Bull.*, 2000, **48**, 832–837.



- 14 H. K. Fun, K. Sivakumar, B. C. Yip, A. H. Othman and I. M. Said, *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.*, 1996, **52**, 414–416.
- 15 H. Greger, T. Pacher, S. Vajrodaya, M. Bacher and O. Hofer, *J. Nat. Prod.*, 2000, **63**, 616–620.
- 16 Y. Qin, C. L. Yin and Z. H. Cheng, *Molecules*, 2013, **18**, 13636–13644.
- 17 T. Biftu, J. C. Chabala, J. Acton, T. Beattie, D. Brooker, R. Bugianesi, M. N. Chang, P. C. Chiang, N. Gamble, N. Girotra, D. Graham, C. H. Kuo, M. Ponpipom, S. Sahoo, T. Y. Shen, K. L. Thompson, S. S. Yang, S. B. Hwang and T. Doebber, *Prostaglandins*, 1988, **35**, 846.
- 18 R. C. C. Martins, L. R. Latorre, P. Sartorelli and M. J. Kato, *Phytochemistry*, 2000, **55**, 843–846.
- 19 L. D. Ferreira, D. R. Callejon, A. Engemann, B. Cramer, H. U. Humpf, V. P. de Barros, M. D. Assis, D. B. da Silva, S. de Albuquerque, L. T. Okano, M. J. Kato and N. P. Lopes, *Planta Med.*, 2012, **78**, 1939–1941.
- 20 L. G. Felipe, D. C. Baldoqui, M. J. Kato, V. D. Bolzani, E. F. Guimarães, R. M. B. Cicarelli and M. Furlan, *Phytochemistry*, 2008, **69**, 445–450.
- 21 T. J. Schmidt, *Phytochem. Lett.*, 2012, **5**, 632.
- 22 M. M. O. Cabral, J. A. Alencar, A. E. Guimarães and M. J. Kato, *J. Am. Mosq. Control Assoc.*, 2009, **25**, 103–105.
- 23 A. Leite, M. J. Kato, R. O. A. Soares, A. E. Guimarães, J. R. Santos-Mallet and M. M. O. Cabral, *Rev. Bras. Farmacogn.*, 2012, **22**, 517–521.
- 24 J. J. Chen, E. T. Chou, C. Y. Duh, S. Z. Yang and I. S. Chen, *Planta Med.*, 2006, **72**, 351–357.
- 25 H.-W. Liu, X.-Z. Yu, D. Padula, G. Pescitelli, Z.-W. Lin, F. Wang, K. Ding, M. Lei and J.-M. Gao, *Eur. J. Med. Chem.*, 2013, **59**, 265–273.
- 26 J. L. Wu, N. Li, T. Hasegawa, J. Sakai, S. Kakuta, W. X. Tang, S. Oka, M. Kiuchi, H. Ogura, T. Kataoka, A. Tomida, T. Tsuruo and M. Ando, *J. Nat. Prod.*, 2005, **68**, 1656–1660.
- 27 M.-M. Bai, W. Shi, J.-M. Tian, M. Lei, Y. H. Kim, J. H. Kim, Y. N. Sun and J.-M. Gao, *J. Agric. Food Chem.*, 2015, **63**, 2198–2205.
- 28 H. S. Kim, C. M. Wooten, Y. Park and J. Y. Hong, *Org. Lett.*, 2007, **9**, 3965–3968.
- 29 K. Nihei, K. Konno, L. S. C. Bernardes, N. P. Lopes, S. Albuquerque, I. de Carvalho, M. T. Pupo, R. C. C. Martins and M. J. Kato, *ARKIVOC*, 2004, 112–126.
- 30 L. S. C. Bernardes, M. J. Kato, S. Albuquerque and I. Carvalho, *Bioorg. Med. Chem.*, 2006, **14**, 7075–7082.
- 31 H. Nishiwaki, K. Nakayama, Y. Shuto and S. Yamauchi, *J. Agric. Food Chem.*, 2014, **62**, 651–659.
- 32 A. M. Rimando, J. M. Pezzuto, N. R. Farnsworth, T. Santisuk, V. Reutrakul and K. Kawanishi, *J. Nat. Prod.*, 1994, **57**, 896–904.
- 33 T. J. Schmidt and J. Heilmann, *Planta Med.*, 2000, **66**, 749–751.
- 34 J. G. Blears and R. D. Haworth, *J. Chem. Soc.*, 1958, 1985–1987.
- 35 R. C. C. Martins, J. H. G. Lago, S. Albuquerque and M. J. Kato, *Phytochemistry*, 2003, **64**, 667–670.
- 36 S. F. Fonseca, L. E. S. Barata, E. A. Ruveda and P. M. Baker, *Can. J. Chem.*, 1979, **57**, 441–443.
- 37 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, J. A. Montgomery Jr, T. Vreven, K. N. Kudin, J. C. Burant, J. M. Millam, S. S. Iyengar, J. Tomasi, V. Barone, B. Mennucci, M. Cossi, G. Scalmani, N. Rega, G. A. Petersson, H. Nakatsuji, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, M. Klene, X. Li, J. E. Knox, H. P. Hratchian, J. B. Cross, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, P. Y. Ayala, K. Morokuma, G. A. Voth, P. Salvador, J. J. Dannenberg, V. G. Zakrzewski, S. Dapprich, A. D. Daniels, M. C. Strain, O. Farkas, D. K. Malick, A. D. Rabuck, K. Raghavachari, J. B. Foresman, J. V. Ortiz, Q. Cui, A. G. Baboul, S. Clifford, J. Cioslowski, B. B. Stefanov, G. Liu, A. Liashenko, P. Piskorz, I. Komaromi, R. L. Martin, D. J. Fox, T. Keith, M. A. Al-Laham, C. Y. Peng, A. Nanayakkara, M. Challacombe, P. M. W. Gill, B. Johnson, W. Chen, M. W. Wong, C. Gonzalez and J. A. Pople, *Gaussian, 03-Revision B.05*, Gaussian Inc., Pittsburgh, 2003.
- 38 H. Matsuura, H. Yoshida, M. Hieda, S. Yamanaka, T. Harada, K. Shin-ya and K. Ohno, *J. Am. Chem. Soc.*, 2003, **125**, 13910–13911.
- 39 P. Gilli, V. Bertolasi, L. Pretto, L. Antonov and G. Gilli, *J. Am. Chem. Soc.*, 2005, **127**, 4943–4953.
- 40 S. Hanessian, G. J. Reddy and N. Chahal, *Org. Lett.*, 2006, **8**, 5477–5480.

