



Cite this: *RSC Adv.*, 2017, 7, 49858

# Investigation of membrane condensation induced by CaCO<sub>3</sub> nanoparticles and its effect on membrane protein function†

Ke Luo,<sup>‡a</sup> Ki-Baek Jeong,<sup>‡a</sup> Jae-Min Oh,<sup>b</sup> Soo-Jin Choi,<sup>c</sup> Tae-Joon Jeon<sup>d</sup> and Young-Rok Kim<sup>id</sup>\*<sup>a</sup>

The membrane dynamics both across the lipid bilayer and in the lateral lipid domains play an important role in a number of key cellular processes including the normal functions of membrane proteins. Herein, we report an electrical method to monitor the changes in membrane dynamics with respect to the interactions between the lipid bilayer and CaCO<sub>3</sub> nanoparticles (nano CaCO<sub>3</sub>). Through transmembrane capacitance measurements, the electrostatic interaction between lipid molecules and Ca<sup>2+</sup> ions dissolved from nano CaCO<sub>3</sub> turned out to be responsible for the membrane condensation, whereas nano CaCO<sub>3</sub> itself didn't show any notable interaction with the lipid bilayer. Moreover, the activities of gramicidin represented by the channel forming rate and lifetime of the channel were shown to be greatly influenced by nano CaCO<sub>3</sub>. These findings demonstrate that a potential cytotoxicity of nano CaCO<sub>3</sub> could be derived from the dissolved Ca<sup>2+</sup> ions, but not from nano CaCO<sub>3</sub> itself, affecting cellular membrane dynamics by lipid condensation.

Received 1st September 2017

Accepted 17th October 2017

DOI: 10.1039/c7ra09722k

[rsc.li/rsc-advances](http://rsc.li/rsc-advances)

## Introduction

Calcium is one of the essential minerals that play an important role in bodily functions, including the formation of bones and teeth, muscle contractions, signal transductions, acting as a cofactor for a range of enzymatic reactions, *etc.* Calcium carbonate (CaCO<sub>3</sub>) is a naturally occurring calcium salt that is commonly used as a source of calcium supplement.<sup>1</sup> Since CaCO<sub>3</sub> has low solubility under physiological conditions, many efforts, for example, on the production of CaCO<sub>3</sub> in nanoparticulate form (nano CaCO<sub>3</sub>) are carried out to increase its bioavailability when consumed as a calcium supplement. Due to the great biostability and non-toxic nature of CaCO<sub>3</sub>, nano CaCO<sub>3</sub> has also emerged as promising drug carrier targeting tumour tissues and cells.<sup>2</sup> On the other hand, there has been growing concern toward potential health effect of nanoparticles over the past decades,<sup>3</sup> and most nanoparticles with the size in the range of 1 to 100 nm have been found to alter signalling

processes essential for normal cell functions and, in some cases, lead to cell death.<sup>4</sup> Moreover, recent observation in cytotoxicity assay demonstrated that Ca<sup>2+</sup> ions from nano CaCO<sub>3</sub> could cause membrane damage through oxidative stress.<sup>5</sup> However, the mechanism of the interaction between nano CaCO<sub>3</sub> and lipid membrane is not fully understood yet.

Planar lipid bilayer (PLB) is a two-dimensional structure presenting a great deal of advantages as a model system mimicking the cell membrane. Upon integration with a proper electrical setup, PLB can be used as a platform for investigating a number of membrane associated ion channels and nanopore based DNA sequencing technique.<sup>6,7</sup> In principle, lipid bilayers are a thin insulating membrane that acts as a capacitor when applying a potential bias across the membrane,<sup>8</sup> so that the changes of lipid bilayer structure in molecular level can be directly monitored by measuring the membrane capacitance.<sup>9</sup> Recently, the interaction between surface-modified gold nanoparticles and PLB was successfully investigated based on this principle, in which the adsorption of the nanoparticles to lipid membrane was quantitatively monitored by monitoring the changes in capacitance of the membranes.<sup>10</sup> In addition, gramicidin channel activity can be employed as an excellent reporter to study the effect of membrane-incorporated compound on the lipid bilayer membrane together with prediction of how such a given compound on the lipid bilayer dynamics as well as the membrane protein functions under physiological conditions.<sup>11–13</sup> Gramicidin is a cation-selective membrane channel protein whose activities, such as channel

<sup>a</sup>Graduate School of Biotechnology, Department of Food Science and Biotechnology, College of Life Sciences, Kyung Hee University, Yongin 17104, Korea. E-mail: [youngkim@khu.ac.kr](mailto:youngkim@khu.ac.kr)

<sup>b</sup>Department of Chemistry and Medical Chemistry, College of Science and Technology, Yonsei University, Wonju 26493, Korea

<sup>c</sup>Department of Applied Food System, Major of Food Science & Technology, Seoul Women's University, Seoul 01797, Korea

<sup>d</sup>Department of Biological Engineering, Inha University, Incheon 22212, Korea

† Electronic supplementary information (ESI) available: Calcium ions analysis, effect of CaCl<sub>2</sub> on membrane capacitance details. See DOI: 10.1039/c7ra09722k

‡ These authors contributed equally to this work.



forming rate and its lifetime, are closely associated with the status of membrane dynamics.

In this study, we investigated the effect of nano  $\text{CaCO}_3$  on the dynamics of lipid bilayer by monitoring the modulated capacitance of the lipid membrane. Subsequently, we employed gramicidin as a molecular force transducer to directly probe the mechanical properties of the bilayer membrane in the presence of nano  $\text{CaCO}_3$ , assuming that there would be a correlation between the changes in membrane capacitance and gramicidin-mediated ions transport activity. Finally, we used this platform to evaluate the effect of nano  $\text{CaCO}_3$  on the membrane dynamics, which would be of importance to elucidate the mechanisms of potential toxicity of nano  $\text{CaCO}_3$ .

## Experimental

### Chemicals and reagents

Phosphate buffered saline (PBS) was purchased from Noblebio Co. Ltd. (Suwon, Korea). 1,2-Diphytanoyl-*sn*-glycero-3-phosphocholine (DPhPC) was purchased from Avanti polar lipid (Alabaster, USA). *n*-Decane was purchased from Fisher Scientific. Bulk calcium carbonate ( $\text{CaCO}_3$ ) and nano  $\text{CaCO}_3$  were provided from Apexel Ltd. (Daejeon, Korea). Calcium chlorides ( $\text{CaCl}_2$ ) was purchased from Duchefa biochemie (Haarlem, Netherlands) and sodium hydroxide (NaOH) was purchased from Daejung (Siheung, Korea). Gramicidin, potassium chloride (KCl) and Quantofix-calcium test kit were purchased from Sigma-Aldrich (St. Louis, MO, U.S.A.).

### Bilayer preparation and electrical measurement

The painting method was employed to prepare a planar lipid bilayer membrane.<sup>14</sup> An aperture of diameter around 150  $\mu\text{m}$  was made on 50  $\mu\text{m}$ -thick Teflon film (HANAROTR Corp, Suwon, Korea) using a spark generator (Daedalon, Salem, MA, U.S.A.). For lipid stock, DPhPC was dissolved in chloroform and dried in a gentle stream of nitrogen gas to remove the solvent. The lipids were redissolved in *n*-decane to a final concentration of 3% (wt/v) and stored at  $-80^\circ\text{C}$  until use. The lipid/decane stock was repainted on both sides of the aperture using a 200  $\mu\text{l}$  plastic pipette tip and the Teflon film was placed horizontally between two custom-built Teflon chambers. The setup was then dried in vacuum chamber for 15 minutes to remove the excess solvent before introducing 2 ml of  $1\times$  PBS (pH 7.4) solution to both *cis* and *trans* chamber. Ag/AgCl electrodes were immersed in the *trans* and *cis* chambers to measure the capacitance and current flow across the membrane. The electrical data were collected using the Axon Digidata 1440A data digitizer and Axopatch 200b patch clamp (Molecular Devices, Sunnyvale, CA, U.S.A) at a sampling rate of 100 kHz, and then analyzed using the Clampfit 10.2 software. The output signal was filtered with a low pass Bessel filter at 300 Hz for membrane capacitance and a Gaussian filter at 10 kHz for gramicidin-mediated transmembrane current measurement.

In order to measure the changes in membrane capacitance, we applied a triangle-waveform voltage that was cyclically ramped between  $-10$  and  $10$  mV. The formation of a lipid

bilayer was confirmed by the characteristic signal of a capacitive square waveform. In our system, the membrane with a capacitance value greater than 100 pF was regarded as a bilayer.<sup>10</sup> After the capacitance of the lipid bilayer remained constant for 10 min, two types of  $\text{CaCO}_3$ , bulk  $\text{CaCO}_3$  and nano  $\text{CaCO}_3$ , were introduced to the chamber. The concentrations of  $\text{CaCO}_3$  stock solution were increased from 0.2 to 5  $\text{mg ml}^{-1}$  with a definitive time interval of 10 min. The transmembrane capacitances were measured three times per every 10 min.

For gramicidin experiment, 1 M KCl (pH 7.4) was used as a supporting electrolyte to measure the current flow through the gramicidin channels formed in lipid bilayer. After the capacitance value of membrane remained constant for 10 min at bias voltage of 200 mV, gramicidin (final conc. 2.7 nM) was introduced to the chamber. Once the first gramicidin-mediated current flow was observed, the solutions in both chambers were replaced with a fresh electrolyte to cease the incorporation of excess gramicidin into the bilayer membrane. To prevent disruption of the bilayer, buffer exchanges should be performed with an extra care by using micropipette. All experiments were performed in a Faraday cage (Warner Instruments) with a floating air table.

### Quantification of $\text{Ca}^{2+}$ ions

The amount of  $\text{Ca}^{2+}$  ions dissolved from varying concentrations of  $\text{CaCl}_2$ , bulk  $\text{CaCO}_3$ , and nano  $\text{CaCO}_3$  in  $1\times$  PBS solution were measured by using Quantofix-calcium test kit (Sigma) that can quantify the concentration of  $\text{Ca}^{2+}$  ions by using the indicator strips. After reaction, the indicator strips were dried under a gentle stream of nitrogen gas. In order to minimize the effect of the undesirable oxidation, the dried indicator strips were immediately transferred to HP scanner device (Officejet Pro 8660) to obtain the image of the strips. ImageJ software was used to convert the color of indicator strips into grayscale value to quantify the concentration of  $\text{Ca}^{2+}$  ions.

## Results and discussion

The schematic illustration of electrical measurement system is shown in Fig. 1a. In this system, PLB that separated *cis* and *trans* chamber was considered as a planar capacitor exhibiting a resistance ( $R$ ) of approximately  $10^8$  ohm  $\text{cm}^2$ .<sup>15</sup> The PLB we used to insulate these two chamber is made by diphytanoyl phosphatidylcholine (DPhPC) that has been historically known for its high mechanical stability, and the capacitance of DPhPC-based PLB is about  $0.412 \mu\text{F cm}^{-2}$ .<sup>16</sup> When applying a triangular voltage wave, the capacitive current across the bilayer is given by:

$$I_c = \frac{dQ}{dt} = \frac{d(C_m V)}{dt} = C_m \frac{dV}{dt} \quad (1)$$

where  $I_c$  is the current across the capacitor,  $Q$  is the charge on the surface of this capacitor,  $V$  is the applied potential,  $C_m$  is the membrane capacitance (F). If the specific  $C_m$  and thickness of PLB is known, the surface area of the bilayer can be calculated with






Fig. 1 Schematic illustration showing electrical setup for the measurement of membrane capacitance. The insert shows the diameter of planar lipid bilayer ( $\phi_{\text{bilayer}}$ ) prepared on the Teflon aperture. (a) A lipid bilayer separating *cis* and *trans* chambers filled with 1x PBS (pH 7.4) was constructed on Teflon film. The magnitude of the square-wave current represents the membrane capacitance at an applied triangular wave voltage of  $\pm 10$  mV (dotted red line). (b) Electrostatic interaction between lipid head and  $\text{Ca}^{2+}$  ions after addition of nano or bulk  $\text{CaCO}_3$  to the chamber and the subsequent changes in membrane capacitance.

$$C_m = \frac{\epsilon_0 \epsilon_m A}{d} \quad (2)$$

where  $\epsilon_0$  is the absolute dielectric constant,  $8.854 \times 10^{-12}$  ( $\text{F m}^{-1}$ ), and  $\epsilon_m$  is the relative dielectric constant of the membrane, which is taken to be close to 2.2 for DPhPC.<sup>17</sup> Here the medium is the mixture of the hydrophobic core of PLB and the solvent, decane.  $A$  is the surface area of the PLB ( $\text{m}^2$ ), and  $d$  is the thickness of the dielectric hydrophobic core of the PLB, which is 4.5 nm for DPhPC.<sup>16</sup> When applying a triangular voltage wave ( $\pm 10$  mV) across the PLB, we observed the square-wave form current which is a measure of bilayer capacity (Fig. 1a). As each lipid molecule has an ability to hold a certain amount of charge, the capacitance can be considered as a parameter showing structural changes of lipid molecule assuming that the area of PLB are constant.<sup>10</sup> Accordingly, we speculate that the changes in capacitance of the lipid membrane can be utilized as a measure to predict the effect of nano  $\text{CaCO}_3$  on the density of lipid molecules in the lipid membrane (Fig. 1b).

In this study, we investigated the effect of two types of calcium carbonate particles, nano  $\text{CaCO}_3$  in a size of about 100 nm and bulk  $\text{CaCO}_3$  of size about 1–3  $\mu\text{m}$  (Fig. 2a), on the




Fig. 2 (a) Scanning electron microscope image of nano  $\text{CaCO}_3$  and bulk  $\text{CaCO}_3$ . Histogram of the size distribution corresponding to nano  $\text{CaCO}_3$  and bulk  $\text{CaCO}_3$  were shown below. (b) Relative increase of membrane capacitance in response to the varying concentration of nano  $\text{CaCO}_3$  and bulk  $\text{CaCO}_3$ .

dynamics of PLB. Here the bulk  $\text{CaCO}_3$  is used as a control to determine the size effect of nano  $\text{CaCO}_3$  in terms of its effect on membrane capacitance. In  $\text{CaCO}_3$ -free condition, the PLB exhibited a structural stability with a constant capacity more than 10 min at the given potential. On the other hand, nano  $\text{CaCO}_3$  caused notable changes in membrane capacitance, which is much greater than those of bulk  $\text{CaCO}_3$  (Fig. 2b), indicating the increased lipid density of the bilayer in the presence of nano  $\text{CaCO}_3$ . Nano  $\text{CaCO}_3$  did not show pore forming activity or adsorption to the membrane according to the very stable current trace over the course of tests. Thus, we propose that changes of membrane capacitance were due primarily to the interactions between divalent calcium ions dissolved from nano  $\text{CaCO}_3$  and the phosphate group in lipid head group (Fig. 1b). Calcium ions have been known to cluster phospholipid molecules *via* ion-bridge,<sup>18</sup> so that lipid molecules are attracted to each other, resulting in the increase of lipid density of the membrane.<sup>19,20</sup> Moreover, there was a gradual increase in the membrane capacitance upon increasing the concentration of nano  $\text{CaCO}_3$  ranging from  $0.2 \text{ mg ml}^{-1}$  to  $5 \text{ mg ml}^{-1}$ , while bulk  $\text{CaCO}_3$  showed a negligible change in membrane capacitance regardless of its concentration. This might be due to the higher solubility of nano  $\text{CaCO}_3$ , which agrees well with grayscale value analysis for the dissolved calcium ions from nano  $\text{CaCO}_3$  and bulk  $\text{CaCO}_3$  (ESI Fig. S1†). On the other hand, we found that the addition of nano  $\text{CaCO}_3$  caused a slight increase of pH compared with that of the bulk  $\text{CaCO}_3$  (Fig. 3a). The effects of pH on the membrane capacitance were therefore taken into account in this study. We examined the effect of pH on membrane capacitance, and found that the membrane capacitances increased slightly in respond to the increase of pH values (Fig. 3b). This might be due to the increase in the concentration of free  $\text{OH}^-$ , some of which can enhance the negativity of polar head groups of lipid molecule, leading to the increases in the membrane capacitance.<sup>21</sup> However, the change of capacitance was insignificant ( $< 0.5\%$ ) upon increasing the concentration of NaOH up to  $0.4 \mu\text{M}$ , which brought about the pH value corresponding to that obtained from the highest concentration of nano  $\text{CaCO}_3$  we tested (Fig. 3a). In addition, we observed a concentration-dependent




Fig. 3 pH effect on membrane dynamics. (a) pH of the electrolyte solution in response to the addition of varying concentration of nano  $\text{CaCO}_3$  and bulk  $\text{CaCO}_3$ . (b) Relative increase of membrane capacitance as a function of NaOH concentration. The pH of electrolyte solution with a given concentration of NaOH was plotted in solid line. The capacitance change induced by the pH corresponding to that obtained from the highest concentration ( $5 \text{ mg ml}^{-1}$ ) of nano  $\text{CaCO}_3$  was indicated by dotted box.



increase in the capacitance when the lipid bilayer was exposed to  $\text{CaCl}_2$  that even lowered the pH value of the solution (ESI Fig. S2†). These results demonstrate that the cause of the capacitance change is more likely due to the increased lipid density of the membrane by the cation–lipid interactions rather than the increased pH in the electrolyte solution.

Subsequently, the gramicidin-mediated ions transport experiments were carried out to evaluate the effect of nano  $\text{CaCO}_3$  on mechanical properties of the lipid bilayer of which its experimental setup is shown in Fig. 4a. Considering that gramicidin can spontaneously incorporate into the membrane, a given concentration of gramicidin (2.7 nM) was added to the electrolyte solutions in both *cis* and *trans* chambers. Upon observation of the occurrence of the first leakage conductance, indicating the formation of gramicidin channel in lipid bilayer, the electrolyte solutions were replaced with fresh one without gramicidin to cease excessive incorporation of gramicidin to the membrane. In this study, the channel forming rates of gramicidin, however, was not considered as a parameter representing the effect of nano  $\text{CaCO}_3$  on the channel function due to the uncontrollable movement of gramicidin from the solvent annulus to the bilayer.<sup>12</sup> Instead, the lifetime of gramicidin channels was only taken into account to analyze the changes of the channel behavior in the presence of nano  $\text{CaCO}_3$ . In control experiment, without nano  $\text{CaCO}_3$ , gramicidin channels showed a wide range of lifetime distribution (Fig. 4b). Interestingly, the channel lifetimes were notably decreased as a function of the concentration of nano  $\text{CaCO}_3$ , indicating that the membrane condensation provided a driving force for the channel dissociation. These results indicate that membrane condensation

induced by  $\text{Ca}^{2+}$  ions that were dissolved from nano  $\text{CaCO}_3$  could alter the dynamics of membrane protein, gramicidin, by shortening the lifetime of the channels, in agreement with that observed from the DOPS bilayers.<sup>22</sup> It might be due to  $\text{Ca}^{2+}$ -mediated changes in the electrostatic interaction among lipid head groups,<sup>23</sup> which caused lipid packing and also affected the bending elasticity of bilayer.<sup>24</sup> From the results, we assume that the changes in mechanical and electrical properties of the membranes could influence the monolayer equilibrium curvature, leading to a change in the curvature stress in the bilayer,<sup>25</sup> consequently increasing the disjoining force ( $F_{\text{dis}}$ ) that influence the functions of gramicidin channels (Fig. 4c).<sup>26</sup>

## Conclusions

Although nano  $\text{CaCO}_3$  did not exhibit any notable physical effect on model cell membrane, the divalent calcium ions dissolved from nano  $\text{CaCO}_3$  were shown to induce the condensation of lipid membrane, which was reflected by the increased membrane capacitance in concentration dependent manner. Comparing to bulk  $\text{CaCO}_3$ , nano  $\text{CaCO}_3$  showed significantly higher condensation effect on lipid membrane mainly due to its higher solubility in aqueous environment. The changes in the density and bending elasticity of lipid membrane by nano  $\text{CaCO}_3$  also altered the channel function of gramicidin embedded in lipid bilayer, implying that the excess divalent calcium ions dissolved from nano  $\text{CaCO}_3$  may pose cytotoxic effect on cell membrane through altering the nature of membrane dynamics. We believe that the developed platform will be of powerful tool to study membrane dynamics in association with potential cytotoxic components in molecular level.

## Conflicts of interest

The author declares no conflicts of interest.

## Acknowledgements

This research was supported by the Pioneer Research Center Program through the National Research Foundation of Korea funded by the Ministry of Science, ICT & Future Planning (2012-0009575 & 2012-0009563), and the National Research Foundation of Korea (NRF-2012M3A7B4049864).

## Notes and references

- I. R. Reid, R. W. Ames, M. C. Evans, G. D. Gamble and S. J. Sharpe, *N. Engl. J. Med.*, 1993, **328**, 460–464.
- S. Maleki Dizaj, M. Barzegar-Jalali, M. H. Zarrintan, K. Adibkia and F. Lotfipour, *Expert Opin. Drug Delivery*, 2015, **12**, 1649–1660.
- A. Elsaesser and C. V. Howard, *Adv. Drug Delivery Rev.*, 2012, **64**, 129–137.
- S. R. Datta, A. Brunet and M. E. Greenberg, *Genes Dev.*, 1999, **13**, 2905–2927.




Fig. 4 Channel forming activities of gramicidin upon varying concentration of nano  $\text{CaCO}_3$ . (a) Experimental setup. Gramicidin was introduced to both *trans* and *cis* chambers. (b) Histogram of lifetime distribution of gramicidin channel depending on the concentration of nano  $\text{CaCO}_3$ . The average lifetime ( $T_{\text{aver}}$ ) of gramicidin channel at different concentration of nano  $\text{CaCO}_3$  were indicated in the top-right corner. (c) Effect of  $\text{Ca}^{2+}$  on curvature stress of lipid bilayer, altering the channel function of gramicidin. Single-channel conductance traces in the absence (left) and presence (right) of nano  $\text{CaCO}_3$ .



- 5 M.-K. Kim, J.-A. Lee, M.-R. Jo, M.-K. Kim, H.-M. Kim, J.-M. Oh, N. W. Song and S.-J. Choi, *Nanomaterials*, 2015, **5**, 1938–1954.
- 6 G. Maglia, M. R. Restrepo, E. Mikhailova and H. Bayley, *Proc. Natl. Acad. Sci. U. S. A.*, 2008, **105**, 19720–19725.
- 7 S. Ladha, A. Mackie, L. Harvey, D. Clark, E. Lea, M. Brullemans and H. Duclouhier, *Biophys. J.*, 1996, **71**, 1364.
- 8 W. Hanke and W. Schulue, *Planar Lipid Bilayers: Methods and Applications*, Academic Press, 2012.
- 9 L. C. Gross, A. J. Heron, S. C. Baca and M. I. Wallace, *Langmuir*, 2011, **27**, 14335–14342.
- 10 R. P. Carney, Y. Astier, T. M. Carney, K. Voitchovsky, P. H. Jacob Silva and F. Stellacci, *ACS Nano*, 2013, **7**, 932–942.
- 11 O. S. Andersen and R. E. Koeppe, *Annu. Rev. Biophys. Biomol. Struct.*, 2007, **36**, 107–130.
- 12 H. Ryu, H. Lee, S. Iwata, S. Choi, M. K. Kim, Y.-R. Kim, S. Maruta, S. M. Kim and T.-J. Jeon, *Sci. Rep.*, 2015, **5**, 11935.
- 13 J. A. Lundbaek and O. S. Andersen, *J. Gen. Physiol.*, 1994, **104**, 645–673.
- 14 P. Mueller, D. O. Rudin, H. Ti Tien and W. C. Wescott, *Nature*, 1962, **194**, 979–980.
- 15 G. Wiegand, N. Arribas-Layton, H. Hillebrandt, E. Sackmann and P. Wagner, *J. Phys. Chem. B*, 2002, **106**, 4245–4254.
- 16 J. Stern, H.-J. Freisleben, S. Janku and K. Ring, *Biochim. Biophys. Acta, Lipids Lipid Metab.*, 1992, **1128**, 227–236.
- 17 G. Valincius, F. Heinrich, R. Budvytyte, D. J. Vanderah, D. J. McGillivray, Y. Sokolov, J. E. Hall and M. Lösche, *Biophys. J.*, 2008, **95**, 4845–4861.
- 18 J. A. Lundbaek, S. A. Collingwood, H. I. Ingolfsson, R. Kapoor and O. S. Andersen, *J. R. Soc., Interface*, 2010, **7**, 373–395.
- 19 A. A. Gurtovenko and I. Vattulainen, *J. Phys. Chem. B*, 2008, **112**, 1953–1962.
- 20 R. Kagawa, Y. Hirano, M. Tajji, K. Yasuoka and M. Yasui, *J. Membr. Sci.*, 2013, **435**, 130–136.
- 21 S. Ohki, *Biophys. J.*, 1969, **9**, 1195.
- 22 J. A. Lundbæk, A. M. Maer and O. S. Andersen, *Biochemistry*, 1997, **36**, 5695–5701.
- 23 H. I. Petrache, T. Zemb, L. Belloni and V. A. Parsegian, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 7982–7987.
- 24 Y. Zhou and R. M. Raphael, *Biophys. J.*, 2007, **92**, 2451–2462.
- 25 G. L. Kirk, S. M. Gruner and D. Stein, *Biochemistry*, 1984, **23**, 1093–1102.
- 26 J. A. Lundbæk, S. A. Collingwood, H. I. Ingólfsson, R. Kapoor and O. S. Andersen, *J. R. Soc., Interface*, 2010, **7**, 373–395.

