RSC Advances

PAPER

Cite this: RSC Adv., 2017, 7, 50459

Received 21st September 2017 Accepted 24th October 2017

DOI: 10.1039/c7ra10497a

rsc.li/rsc-advances

Introduction

Lignin is a highly abundant and irregular biopolymer which constitutes lignocellulosic biomass together with cellulose and hemicellulose. During the last decades, the scientific community has deeply studied the usability of these polymers as sources of energy and high-added value compounds. On the opposite, the complex structure of lignin has overshadowed its value as a renewable alternative to the fuel industry, since high amounts of energy could be involved to break down its crosslinked structure. Nevertheless, the use of lignin as feedstock has become an important field of research in recent years, finding multiple applications in the production of bio-based polymers such as adhesives, bioplastics or resins due to its polyphenolic composition.¹ The lignin valorization is also moving to cleavage processes focused on degradation into

Synthesis of nitrogenated lignin-derived compounds and reactivity with laccases. Study of their application in mild chemoenzymatic oxidative processes†

Jesús Albarrán-Velo[,](http://orcid.org/0000-0003-0052-4771) <mark>©</mark> María López-Iglesias, Vicente Gotor, Vicente Gotor-Fernández \mathbb{D}^* \mathbb{D}^* \mathbb{D}^* and Iván Lavandera \mathbb{D}^*

The chemical synthesis of a series of lignin-derived nitrogenated compounds was performed in high yields (73–99%) through simple conventional chemical transformations starting from natural monomers such as vanillin, syringaldehyde or 3,4-dihydroxybenzaldehyde. The study of the vanillin-derived compounds as substrates for commercially available laccases from Trametes versicolor and Myceliophthora thermophila in oxidative transformations, generally led to the isolation of several dimeric species in high to excellent conversions (>70%), while for hydrazone derivatives a more rapidly oxidative coupling was evidenced by the formation of oligomers and/or polymers. Remarkably, vanillin was obtained due to the hydrolysis of some of the nitrogenated functional groups, such as the hydrazone or the hydrazono tetrazole. The three families of lignin-derived compounds can provide a great source of new laccase-mediator systems (LMS), the possibility of employing them for lignin modification being particularly attractive. Preliminary experiments showed promising levels of activity towards the oxidation of a monomer (veratryl alcohol, up to 70% conversion) and a dimer (adlerol, up to 22% conversion) lignin models, higher than those achieved with the natural vanillin and syringaldehyde (up to 7% conversion with veratryl alcohol and almost negligible conversion with adlerol), these processes being also highly influenced by the pH of the reaction medium. PAPER

(a) Check for updates

Synthesis of nitrogenated lignin-derived

Crestin fisic Adv. 2017 7.60499

Crestin fisic Adv. 2017 7.60499

Check for updates

Crestin fisic Adv. 2017 7.60499

Crestin fisic Adv. 2017 7.60499

smaller compounds,² biocatalytic methodologies being currently booming.³

Thus, it has been found that some bacterial and fungal microorganisms containing potent oxidative enzymes, have the capability to degrade the structure of lignin by promoting the cleavage of certain connections as the β -O-4-aryl ether bonds or the oxidation at C_{α} positions,^{3c} these predominant linkages being illustrated in the dimeric species of Fig. 1 (left). Particularly, some peroxidases, peroxygenases and laccases have been applied in enzyme-catalyzed delignification processes.⁴

Fig. 1 Lignin-derived model compounds, monomeric and dimeric species.

Organic and Inorganic Chemistry Department, Biotechnology Institute of Asturias (IUBA), University of Oviedo, Avenida Julián Clavería 8, 33006 Oviedo, Spain. E-mail: vicgotfer@uniovi.es; lavanderaivan@uniovi.es

[†] Electronic supplementary information (ESI) available: This file includes extensive enzymatic optimization studies, analytical data for the oxidation of adlerol and copy of ${}^{1}H$, ${}^{13}C$ and DEPT NMR spectra for novel synthesized compounds. See DOI: 10.1039/c7ra10497a

Furthermore, lignin has attracted an increasing interest not only as source of energy but also because of the valuable lowmolecular weight aromatic compounds contained in its structure, which can also serve as building blocks for synthetic processes.⁵ Within this area, many efforts have been intended to investigate the behavior of lignin-related compounds towards cleavage or oxidation processes as representative and handier models in which the most characteristic linkages are present. Among them, some dimeric species such as veratrylglycerol β guaiacyl ether (adlerol, 2, Fig. 1) or similar β -1-type lignin model compounds have found wide application, for instance, for studying the mechanism of lignin C–C linkage cleavage by pyrolytic strategies⁶ or metal catalysis.^{2a} Also Biocatalysis has been applied in lignin valorization by studying the cleavage or oxidation of β-O-4-linkages in dimeric species.⁷

Laccases (EC 1.10.3.2) are multi-copper oxidative enzymes that consume molecular oxygen as electron acceptor during the oxidation of a substrate producing water as the only by-product, which makes them an interesting and suitable tool working under mild reaction conditions.⁸ These enzymes have been employed in lignin degradation,^{4,9} and also in the opposite repolymerization processes or related lignin-surface modifications, in order to functionalize it obtaining valuable biomaterials, for example by single molecule grafting or copolymerization processes.¹⁰ Despite these enzymes present a suitable redox potential to abstract electrons from phenolic compounds, this is not high enough to transform other types of substrates such as aliphatic alcohols, so the use of laccase-mediator systems (LMS) is necessary to allow their oxidation. $s_{c,11}$ Many synthetic compounds are commonly employed in these systems, such as TEMPO, HOBt or ABTS, but also some natural compounds obtained from lignin can be used. Among them, vanillin (3a, Fig. 1), syringaldehyde (3b, Fig. 1) and their derivatives are some of the most common lignin-derived monomers that have been extensively employed in different ways, for example as natural LMS.^{9*a*,12} In fact, laccases have proven to accept a large number of lignin-**EXALM SCHE CONFIRENT CONSULTS CO**

Herein, we aim to combine the use of compounds derived from renewable resources $(e.g.$ lignocellulose) and the benefits that biocatalysis offers due to the capability of enzymes to selectively work under environmentally friendly conditions. Thus, we firstly report the chemical modification of vanillin and syringaldehyde towards the synthesis of nitrogenated derivatives whose broad structure is depicted in Fig. 1 (right). Later these compounds will be tested as suitable substrates in laccase-catalyzed processes. Overall, the synthetic strategies will provide novel systems that will be applied in the oxidation of lignin model compounds, which is the underlying purpose of the present research.

Results and discussion

Chemical modification of natural LMS

Vanillin (3a) and syringaldehyde (3b) were selected as starting materials for further modifications due to the presence of these compounds in high quantities when degrading lignin. As both substrates have a similar structure but a different number of methoxy groups in the aromatic ring, the choice of these two lignin-type monomers will allow us to compare their capacities in LMS. Moreover, anticipating the importance of both the methoxy and the hydroxyl groups in the substrate–laccase interaction, and in order to complete the study, we decided to include also 3,4-dihydroxybenzaldehyde (3c), which was synthesized starting from vanillin via demethylation in 85% yield as described in the literature (Scheme S1, see the ESI†).¹⁴

In the interest of practical designs for the potential mediators, we focused on simple and easy synthetic procedures. Moreover, the rationale for the synthesis of the proposed compounds is based on the possibility of stabilizing the oxidized radical intermediates by aromatic systems via electron

Scheme 1 An oxime derivative presenting an increasing number of reacting points.

Scheme 2 Synthesis of 4-hydroxy-3-methoxybenzonitrile (4a) and oximes 5a–c.

Table 1 Synthesis of new nitrogenated compounds derived from vanillin and syringaldehyde

^a Yield obtained after isolation of the products by solvent evaporation (for $6a-c$ and $7a-c$) or product filtration (for $8a-c$ and $9a-c$).

Table 2 Laccase-catalyzed oxidation of the new synthesized vanillin-derived compounds⁴

^a Reaction conditions: substrate 3-9a, cosolvent (10% v/v), acetate buffer pH 4.7 (total volume: 3 mL) saturated with O_2 , laccase from Trametes versicolor (LTv, 25 U) or Novozym 51003 (31.4 U). b Conversion values were measured by ¹H-NMR analysis of the reaction crudes. ^c Vanillin and a mixture of dimeric species were obtained. ^d Vanillin and a mixture of unidentified products were attained.

delocalization or by increasing the number of additional reactive points as in the case of an oxime derivative (Scheme 1), resembling in this particular case N-oxyl oxidants such as TEMPO.

In a first set of experiments, we evaluated the use of hydroxyl ammonium reagents to introduce both oxime and nitrile functions. Thus, oximes 5a–c were prepared in good to excellent yields following a described methodology¹⁵ in the presence of hydroxylammonium sulfate and sodium acetate in ethanol (Scheme 2). By varying the solvent to acetonitrile and using hydroxylamine hydrochloride, dehydration of the aldoxime intermediate took place yielding the nitrile derivative 4a from 3a after 4 h.

On the other hand, a panel of nitrogenated substituents was introduced through the aldehyde function by tuning the strength of the acid catalyst and adjusting the reaction time (Table 1).

Firstly, the reaction between hydrazine monohydrate (10) and 3a was performed under acidic conditions, observing full disappearance of the starting material by TLC analysis after 50 min. Hydrazone 6a was isolated in 90% yield after removing the solvent by distillation (entry 1). The procedure was extended to the dimethoxylated and dihydroxylated substrates 3b and 3c, leading to 6b and 6c also in good yields (entries 2 and 3). The reaction time was increased in the synthesis of semicarbazones 7a–c in the presence of semicarbazide hydrochloride (11), obtaining the desired products in good to excellent yields after solvent evaporation (entries 4–6).

To prepare the triazole derivatives 8a–c, longer reaction times and harsher acidic conditions were needed, so concentrated H_2SO_4 was employed as catalyst.¹⁶ The reaction between 3b and 3c with 4H-1,2,4-triazol-4-amine (12) was completed after 7.5 and 6.5 h, respectively (entries 8 and 9), while starting from 3a only 3 h were necessary (entry 7), recovering 8a–c with excellent yields. Concentrated HCl was used in the reactions with tetrazole 13, obtaining 9a–c in 80–97% yield (entries 10– 12). In these cases, the isolation was performed by filtration from the reaction crude, since both families of polynitrogenated compounds 8a–c and 9a–c were insoluble in the reaction medium. In the latter reactions, it was necessary to synthesize the reagent 13 through a two-step process (see Experimental section), as it was not commercially available.

Laccase-catalyzed oxidation reactions of the synthesized derivatives

To develop an efficient LMS with these compounds, first we investigated if they could react with laccases. For this, we chose commercially available laccases from Trametes versicolor (LTv)¹⁷ and Myceliophthora thermophila (Novozym 51003),¹⁸ and the enzymatic experiments were conducted with the family of vanillin-modified derivatives, in order to illustrate the oxidative processes that could take place. This study would verify the validity of the new synthesized compounds as potential laccase mediators. Following previous described protocols with these biocatalysts, enzymatic reactions were carried out in acetate buffer at pH 4.7, varying the substrate concentration from 10 to 50 mM. Due to solubility issues, 10% (v/v) of EtOH or DMSO¹⁹ were added as water-miscible organic solvents in some cases. The control reactions in the absence of enzyme led to no conversion in all cases, proving the direct role of the laccase as the only catalyst.

When the laccases and the vanillin derivatives were mixed, three processes were mainly observed. In most cases the C–C dimerization reaction of species-type A (Scheme 1) was favored, but also the formation of vanillin (via hydrolysis) and the polymerization/oligomerization of these substrates were observed (Table 2). From now on, no distinction between polymerization and oligomerization will be made, since this was not the main goal of this research, so the polymeric structures were not analyzed such depth to make that differentiation. Firstly, the enzymatic experiments were conducted with the

vanillin itself (3a), observing it was susceptible to dimerization in a very high extent (entries 1 and 2). Similarly, dimeric species 15–18 were isolated and properly characterized from the reaction crudes when starting from nitrile 4a, oxime 5a, semicarbazone 7a, and triazole 8a (entries 3–6, 9, 11 and 12). Particularly, 4a reacted at 10 mM concentration, achieving a quantitative conversion with LTv and Novozym 51003 (entries 3 and 4). Oxime 5a also led to full conversions at 50 mM (entries 5 and 6), observing with both biocatalysts the formation of a minimum amount of 3a (3%) which was no detected in the absence of the enzyme. Semicarbazone 7a was converted at 20 mM concentration in the presence of LTv conducting to dimer 17 (72% conversion, entry 9). However, the product formation differed in the case of Novozym 51003 (entry 10), in which the presence of a polymeric species, even at short reaction time, was detected due to the broad signals observed in the ¹H-NMR experiments. The generation of polymeric products in laccase-catalyzed reactions has already been reported in the literature for this type of lignin monomer.^{13b,20} Finally, dimer 18 was obtained as a precipitate from the reaction crude when starting from triazole 8a (20 mM, entries 11 and 12) in almost full conversion with both enzymes. These results can be related with the stability of the radical species-type A (Scheme 1). If this intermediate is stable enough, the dimerization process will be favored. Nitrogenated electron-withdrawing groups enhanced the stability of this radical as the C–C dimerization reaction was prominent. Paper

SEA Above: SEA Above: Secaration intervention intervention

The hydrazone bond of 6a was hydrolyzed during the enzymatic process, so vanillin was detected in significant amounts in the ¹H-NMR of the reaction crude (entries 7 and 8), a 4% of the hydrolysis product 3a being also formed in the absence of enzyme. Moreover, after aldehyde formation also dimerization took place, so a mixture of starting material, 3a and several dimeric species were detected in the reaction crude in a proportion varying depending on the biocatalyst used. Despite of the presence of unreacted hydrazone in the crude, it was noticed a fast reaction as the mixture acquired a dark purple color just after the addition of the enzyme. When the N-tetrazole substituted hydrazone derivative 9a (10 mM) was submitted to oxidation, the hydrolysis of the $C=N$ bond was also observed. Thus, a mixture of products was extracted from the reaction crude when LTv or Novozym 51003 were used, in which the starting material, vanillin and other unidentified products were already present at 2 h. Nevertheless, polymerization might also be occurring, as a significant loss of weight in the crude recovered by extraction was observed.

Since laccases are not expected to hydrolyze hydrazones, we evaluated if unspecific catalysis of this protein or other impurities that could be present in the commercial laccase preparation might be responsible of this transformation. So the LTv was denatured by a heat treatment (for details, see the ESI†). The conversion towards 3a formation was just 4%, demonstrating the role of the biocatalytic system in the hydrolysis of 6a. While it has been described the hydrolysis of hydrazones under oxidative conditions, 21 there is only one precedent in the literature showing a similar behavior of LMS, the authors observing bio-deoximation processes in the presence of TEMPO Table 3 Selected results in the laccase-catalyzed veratryl alcohol oxidation in the presence of different natural or nitrogenated lignin-derived $compounds^a$

 a Reaction conditions: natural or nitrogenated derivative (16 µmol), buffer (2 mL), veratryl alcohol (0.9 µL, 6 µmol), laccase from Trametes versicolor (LTv, 25 U, 30 C) or Novozym 51003 (31.4 U, 50 C), 72 h. Buffers used: acetate buffer 100 mM pH 4.7; citrate buffer 50 mM pH 5; citrate buffer 50 mM pH 5.5; citrate buffer 50 mM pH 6; phosphate buffer 100 mM pH 7. b Conversion values were measured by GC analysis of the reaction crudes.

as mediator.^{11a} In our experiments, we assume that the laccase is able to catalyze the synthesis of unstable radical intermediates which can undergo hydrolysis releasing vanillin.

New LMS applied to the oxidation of lignin-model compounds

After checking that these compounds were suitable substrates for the studied laccases, they were tested as potential mediators to oxidize two different lignin model compounds. The syringyl and 3,4-dihydroxybenzaldehyde families were also included in this study. Thus, veratryl alcohol (1) and adlerol (2) were employed as representative monomeric and dimeric structures of lignin units,²² respectively, the alcohol moiety being in both cases at benzylic position (Fig. 1). Control experiments in the absence of laccase and/or nitrogenated compound revealed that both must be present for an efficient catalysis.

Beginning with the veratryl alcohol oxidation, we selected 3 mM as starting substrate concentration according to some reports found in the literature, $22d$ evaluating both commercial laccases. A deep optimization was performed in order to determine the best reaction conditions, including the reaction

medium, temperature, amount of catalyst and concentration of the nitrogenated derivative, and the use of different cosolvents (see Tables S1–S18 in the ESI†). Among them, the pH was identified as key parameter, and depending on the phenolic compound employed, the optimum value varied. Similarly, changes in the amount of the co-oxidant implied important variations in the conversion degree, detecting the higher values at 8 mM concentration. The first experiments were stopped at 48 h but, after observing that the laccase was still active, the reactions were followed until 72 h. The best results were obtained in the acidic pH range of 4.7–7, as described for these laccases,17,18 and are summarized in Table 3. LTv works better under acidic conditions while Novozym 51003 accepts higher pHs. Paper

Note that the control of starbs and concentration of sytingy coinces are allows and 3b, the process being position

the control of the starb

The process was also carried out in the presence of vanillin and syringaldehyde (entries 1–4), to compare the behavior of these natural derivatives with the chemically modified ones, finding an improvement in the conversion values for several of the newly nitrogenated synthesized compounds. Still some of them displayed very low activity even at these long reaction times, as was the case of the nitrile derivative 4a, which led to less than 10% conversion with both enzymes LTv and Novozym 51003 (entries 5 and 6). On the contrary, oximes 5a–c exhibited remarkable activities, 5c providing with LTv the highest value of all derivatives tested with 70% conversion (entry 9). The reaction exceeded 30% conversion when employing vanillin and

syringyl oximes 5a and 5b, the process being promoted to 37% conversion at pH 6 with 5a and LTv, and 34% when Novozym 51003 was employed under the same pH conditions at 50 $^{\circ}$ C (entries 7 and 10). In an attempt to increase those values, the use of different cosolvents was tested but resulted ineffective (see Table S4 in the ESI†), and longer reaction times led to the disappearance of the product signal in the GC chromatogram. We hypothesized that the product was degraded, partially to form the veratryl acid, which was identified in the reaction analysis.

The presence of an additional methoxy group in oxime 5b did not affect significantly the performance of Novozym 51003, obtaining a conversion of 35% at pH 7 (entry 11), while the reaction was disfavored with LTv (entry 8), especially at higher pHs (see Table S9 in the ESI†).

When the vanillin hydrazone 6a was studied, good values of conversion were observed (entries 13 and 16), obtaining 54% in combination with Novozym 51003 at pH 6 (entry 16). On the opposite, 6b led to low or null conversions with both enzymes (entries 14 and 17). The results were slightly better but still poor with the dihydroxylated derivative 6c (entries 15 and 18), not improving in the presence of an organic cosolvent (see Table S15 in the ESI†). Conversely to the hydrazones, the semicarbazone derived from syringaldehyde 7b (entries 20 and 23) showed higher activities than the corresponding analogue from vanillin 7a (entries 19 and 22). As for the hydrazone family, the

Fig. 2 Relative activity representing micromoles of alcohol 1 converted into aldehyde 18 at different substrate concentrations, using the new synthesized compounds as co-oxidants, and Novozym 51003 (up) or LTv (down) as biocatalyst. 100% relative activity corresponds to 0.42 µmol 18 per μ L mg⁻¹ of enzyme (entry 9, Table 3). Reactions with relative activities higher than 200% were performed by duplicate, showing similar results.

dihydroxylated semicarbazone 7c displayed the lowest conversions (entries 21 and 24).

In general, the triazole family was poorly active, affording the dimethoxylated derivative 8b the highest conversion value in the reaction with Novozym 51003 at pH 6 (entry 29). Interestingly, tetrazole derivatives 9a and 9b showed enhanced conversions (entries 31–32 and 34–35). As for the case of semicarbazones 7a–b and triazoles 8a–b, Novozym 51003 exhibited higher acceptance to the syringyl-derived substrate than to the corresponding vanillin-derived one (entries 34 and 35), behaving positively the additional methoxy group. On the contrary, when the reaction was carried out in the presence of the dihydroxylated derivative 9c, the conversions decayed significantly, especially for Novozym 51003 acting as catalyst (entry 36). BSC Advances

dihydrog-lated scenarios, are the published on 30 October 2017. The common process are the common process are the common process are the common process and the steam of the common action and the common state

As the next step, we decided to increase the veratryl alcohol concentration for those cases where higher conversions were found. This would allow us to study the effect of this parameter in the reaction outcome and also to obtain a higher productivity of the aldehyde 18 in the process. Remarkably, at 10 mM substrate concentration, conversions suffered just a slight decrease in some cases, maintaining the previous values in

Fig. 3 Oxidation of adlerol (2) using natural or nitrogenated ligninderived compounds with: (a) Novozym 51003; or (b) LTv. All the experiments were performed by triplicate.

other cases, resulting in an enhancement of the relative activity for all derivatives tested (Fig. 2).

Noteworthy, it was possible to rise the substrate concentration until 15 mM and 20 mM, maintaining good levels of catalytic activity and, therefore, improving the productivities. For instance, relative activity values of 186–295% were found with oxime 5b, hydrazone 6a and tetrazole 9b within the Novozym 51003-catalyzed reactions (Fig. 2, top). In general, LTv showed less alcohol concentration tolerance, but even so oxime 5c displayed an excellent activity in the presence of this enzyme (286%, Fig. 2, bottom).

Finally, the methodology was extended to the oxidation of the benzylic position of the dimeric lignin model compound adlerol 2 using the derivatives that displayed the higher activities with the laccases for the oxidation of 1 (Fig. 3). The reaction conditions were taken in analogy to the previously shown for veratryl alcohol.

Despite the conversions were lower than for the case of 1, significant improvements were accomplished with some derivatives, obtaining the best results (22% conversion) for the combination of oxime 5b with Novozym 51003, which is a remarkable value when considering the examples reported in the literature using vanillin or syringyl-type natural mediators in this kind of process.²² In order to verify the improvement over the natural compounds, 3a and 3b were tested, observing no formation of hydroxy ketone 19.

Oxime 5c showed signicant conversions in the LTvcatalyzed process, as well as tetrazoles 9b and 9c (Fig. 3b). On the contrary, the use of Novozym 51003 as catalyst resulted in high conversion values with derivatives 6a, 9a and 9b (Fig. 3a). We figure out that the lower conversion values may be caused by a fast reaction of the phenolic compound with the laccase, leading to the formation of inactive species and disabling the oxidative capacity in the adlerol transformation. As a final remark, it must be mentioned that coupling between the synthesized nitrogenated derivatives and the substrates (veratryl alcohol and adlerol) was never observed.

Conclusions

The chemical modification of some lignin-type monomers such as vanillin and syringaldehyde has been carried out through simple methodologies toward the synthesis of several families of nitrogenated compounds. These derivatives were envisaged as suitable candidates for LMS due to the stabilization of the phenoxy radicals formed. Their combination with two commercially available laccases, the one from Trametes versicolor and the one from Myceliophthora thermophila, allowed their dimerization or oligomerization, showing that they could act as substrates for laccases.

The application of these compounds in oxidative processes of lignin model units in laccase-mediator systems has been investigated employing these commercial laccases, showing much better conversions than the natural compounds vanillin and syringaldehyde. Remarkably, both the substitution pattern of the aromatic ring and the pH have shown a dramatic influence in the reactivity behavior, conducting to higher

conversions according to the presence of additional methoxy and/or hydroxyl groups. On the one hand, veratryl alcohol was transformed into veratraldehyde in high conversions at 3 mM, and the increase in the substrate concentration up to 20 mM led to higher relative activity values (up to 295%). On the other hand, lower conversions were obtained in the oxidation of the lignin model dimer adlerol, but these results are a good starting point for future studies in the enzymatic modification of lignin surfaces.

Experimental section

General methods

Chemical reagents employed in the synthesis of starting and reference materials were purchased from Sigma-Aldrich. Solvents for chemical reactions were used without further treatment.

Laccase from *Trametes versicolor* (2.5 U mg $^{-1})$ was purchased from Sigma, while the laccase from Myceliophthora thermophila (Novozym 51003, 3.14 U μ L⁻¹) was acquired from Novozymes. Enzymatic activities were measured using the ABTS assay.²³

Thin-layer chromatography (TLC) was conducted using Merck Silica Gel 60 F254 precoated plates and visualized with UV and potassium permanganate stain. Column chromatographies were performed using silica gel 60 (230-400 mesh). ¹H-NMR, 13C-NMR, and DEPT experiments were registered using different Bruker spectrometers (${}^{1}H$, 300.13 MHz and ${}^{13}C$, 75.5 MHz). The chemical shifts (δ) are given in ppm and the coupling constants (J) in Hertz (Hz). High resolution mass spectra (HRMS) experiments were carried out by $ESI⁺$ or $ESI⁻$ using an Impact II ESI-Q-Tof spectrometer. Paper

Conversions according to the restorate of additional method some free play of the restoration and the figure of the restoration and the internet of the Crick is licensed in the Crick is article in the common of the

Synthesis of 4-hydroxy-3-methoxybenzonitrile $(4a)^{24}$

Hydroxylamine hydrochloride (82 mg, 1.18 mmol) was added to a solution of vanillin (3a, 150 mg, 0.986 mmol) in acetonitrile (1 mL) and the mixture was stirred under reflux for 4 h. After cooling the mixture at room temperature, H_2O was added and the reaction was extracted with Et₂O (3×10 mL). The organic layers were combined, washed with an aqueous saturated NaHCO₃ solution (15 mL) and dried over Na₂SO₄. The solvent was removed by distillation under reduced pressure and the reaction crude was purified by column chromatography on silica gel (eluent: 30% EtOAc/hexane) yielding the nitrile derivative 4a in 85% isolated yield. White solid. R_f (50% EtOAc/ hexane): 0.80. Mp: 86–88 °C. 1 H NMR (300.13 MHz, MeOD- d_{4}): δ 3.87 (s, 3H), 4.91 (br s, 1H), 6.87 (d, 3 J_{HH} = 8.1 Hz, 1H), 7.18 $(dd, {}^{3}J_{\text{HH}} = 8.1, {}^{4}J_{\text{HH}} = 1.9 \text{ Hz}, 1\text{H}$), 7.21 $(d, {}^{3}J_{\text{HH}} = 1.9 \text{ Hz}, 1\text{ Hz})$ ¹³C NMR (75.5 MHz, MeOD- d_4): δ 56.6 (CH₃), 103.1 (C), 115.7 (CH), 117.0 (CH), 120.5 (C), 127.7 (CH), 149.3 (C), 152.7 (C). HRMS (ESI⁺, *m*/z): calcd for $(C_8H_8NO_2)^+$ $(M + H)^+$ 150.0550, found: 150.0548.

General procedure for the synthesis of oximes 5a–c

Hydroxylamine sulfate (650 mg, 3.95 mmol) and sodium acetate (447 mg, 3.29 mmol) were successively added to a solution of 3a–c (3.29 mmol) in ethanol (3.3 mL). The mixture was stirred

under reflux for 4 h, cooled at room temperature, resuspended in CH₂Cl₂ (10 mL), filtered and washed with CH₂Cl₂ (2×2 mL). The solvent of the filtrate was evaporated under reduced pressure obtaining pure oxime 5a. Oxime 5b was submitted to column chromatography (eluent: 30% EtOAc/hexane) for further purification.

For the isolation of oxime 5c, the solvents were evaporated and the residue was redissolved in $H₂O$. The mixture was extracted with EtOAc $(3 \times 5 \text{ mL})$, the organic layers were combined, dried over $Na₂SO₄$ and concentrated in vacuo, affording pure 5c.

4-Hydroxy-3-methoxybenzaldehyde oxime (5a). ¹⁵ Light brown solid (550 mg, >99% yield). R_f (50% EtOAc/hexane): 0.67. Mp: 110–112 °C. ¹H NMR (300.13 MHz, MeOD- d_4): δ 3.83 (s, 3H), 6.79 $(d, {}^{3}J_{\text{HH}} = 8.1 \text{ Hz}, 1\text{H}), 6.95 (dd, {}^{3}J_{\text{HH}} = 8.1 \text{ Hz}, {}^{4}J_{\text{HH}} = 1.7 \text{ Hz},$ 1H), 7.23 (d, $^{4}J_{\text{HH}} = 1.7$ Hz, 1H), 8.00 (s, 1H). ¹³C NMR (75.5) MHz, MeOD- d_4): δ 56.2 (CH₃), 109.6 (CH), 116.0 (CH), 122.4 (CH) , 126.0 (C) , 149.1 (C) , 149.3 (C) , 150.6 (CH) . HRMS $(ESI⁺, m)$ z): calcd for $\left(C_8{\rm H_{10}}{\rm NO_3}\right)^t} \left({\rm M} + {\rm H}\right)^+$ 168.0655, found: 168.0656.

4-Hydroxy-3,5-dimethoxybenzaldehyde oxime (5b). Light brown solid (590 mg, 91% yield). R_f (50% EtOAc/hexane): 0.46. Mp: 126–128 °C. 1 H NMR (300.13 MHz, MeOD- d_{4}): δ 3.82 (s, 6H), 6.87 (s, 2H), 7.99 (s, 1H). ¹³C NMR (75.5 MHz, MeOD- d_4): δ 56.6 (2CH3), 105.1 (2CH), 124.9 (C), 138.2 (C), 149.2 (2C), 150.7 (CH). HRMS (ESI⁺, m/z): calcd for $(C_9H_{11}NNaO_4)^+ (M + Na)^+$ 220.0580, found: 220.0582.

3,4-Dihydroxybenzaldehyde oxime (5c). Light brown solid (448 mg, 89% yield). R_f (50% EtOAc/hexane): 0.42. Mp: decomposes at 160–162 °C. ¹H NMR (300.13 MHz, DMSO- d_6): δ 6.72 (d, δ _L, $-$ 8.1 Hz, 1H) 6.81 (dd, δ _{L, $-$} 8.1 Hz, δ _{L, $-$} 2.0 Hz, 1H) $J_{\rm HH} = 8.1$ Hz, 1H), 6.81 (dd, $^3J_{\rm HH} = 8.1$ Hz, $^4J_{\rm HH} = 2.0$ Hz, 1H), 7.04 (d, $^{4}J_{\text{HH}} = 2.0$ Hz, 1H), 7.91 (s, 1H), 9.18 (br s, 2H), 10.79 (br s 1H). ¹³C NMR (75.5 MHz, DMSO- d_6): δ 112.6 (CH), 115.5 (CH), 119.3 (CH), 124.5 (C), 145.5 (C), 146.9 (C), 148.2 (CH). HRMS (ESI⁺, *m*/z): calcd for $(C_7H_8NO_3)^+ (M + H)^+$ 154.0499, found: 154.0498.

General procedure for the synthesis of hydrazones 6a–c

Hydrazine monohydrate $(10, 207 \mu L, 6.57 \mu m)$ and glacial acetic acid (10 drops) were successively added to a solution of 3a–c (6.57 mmol) in ethanol (10 mL). The mixture was stirred under reflux for 50 min or 1 h (see Table 1), and the solvent was removed by distillation under reduced pressure isolating 6a–c in 78–90% yield.

4-(Hydrazonomethyl)-2-methoxyphenol (6a). Yellow solid (983 mg, 90% yield). R_f (50% EtOAc/hexane): 0.55. Mp: 177-179 °C. ¹H NMR (300.13 MHz, MeOD-d₄): δ 3.93 (s, 3H), 6.88 (d, ³L = 8.1 Hz, 1H) 7.52 $J_{\text{HH}} = 8.1 \text{ Hz}, 1 \text{H}$), 7.24 (dd, 3 J_{HH} = 8.3, 4 J_{HH} = 1.9 Hz, 1H), 7.52 $(d, {}^{4}J_{\text{HH}} = 1.9, 1\text{H}), 8.53 \text{ (s, 1H)}.$ ${}^{13}C$ NMR (75.5 MHz, MeOD- d_4): 56.4 (CH3), 110.8 (CH), 116.3 (CH), 125.3 (CH), 127.2 (C), 149.5 (C), 151.4 (C), 162.7 (CH). HRMS $(ESI⁺, m/z):$ calcd for $(C_8H_{11}N_2O_2)^+ (M + H)^+$ 167.0815, found: 167.0810.

4-(Hydrazonomethyl)-2,6-dimethoxyphenol (6b). Light yellow solid (1.03 g, 80% yield). R_f (50% EtOAc/hexane): 0.24. Mp: 212–214 °C. 1 H NMR (300.13 MHz, DMSO- d_{6}): δ 3.82 (s, 6H), 7.16 (s, 2H), 8.59 (s, 1H), 9.08 (br s, 1H). 13C NMR (75.5 MHz, DMSO-d₆): 56.0 (2CH₃), 105.8 (2CH), 124.3 (C), 139.9 (C), 148.1

(2C), 162.9 (CH). HRMS (ESI⁺, *m*/z): calcd for $(C_9H_{13}N_2O_3)^+$ (M + H)⁺ 197.0920, found: 197.0915.

4-(Hydrazonomethyl)benzene-1,2-diol (6c). Light orange solid (780 mg, 78% yield). R_f (50% EtOAc/hexane): 0.16. Mp: decomposes at 200 °C. 1 H NMR (300.13 MHz, DMSO- d_{6}): δ 6.81 $(d, {}^{3}J_{\text{HH}} = 8.1 \text{ Hz}, 1\text{H}), 7.10 (dd, {}^{3}J_{\text{HH}} = 8.3, {}^{4}J_{\text{HH}} = 2.0 \text{ Hz}, 1\text{H}),$ 7.32 (d, 4 J_{HH} = 2.0, 1H), 8.45 (s, 1H), 9.42 (br s, 2H). ¹³C NMR $(75.5 \text{ MHz}, \text{DMSO-}d_6)$: 113.9 (CH), 115.6 (CH), 121.9 (CH), 125.6 (C), 145.7 (C), 148.9 (C), 160.5 (CH). HRMS (ESI⁺, m/z): calcd for $(C_7H_9N_2O_2)^+$ $(M + H)^+$ 153.0659, found: 153.0658.

General procedure for the synthesis of semicarbazones 7a–c

Semicarbazide hydrochloride (11, 733 mg, 6.57 mmol) and glacial acetic acid (10 drops) were successively added to a solution of 3a–c (6.57 mmol) in ethanol (10 mL). The mixture was stirred under reflux for $1.5-2.7$ h (see Table 1) and the solvent was removed by distillation under reduced pressure, isolating semicarbazones 7a–c in good to excellent yields (73–98%).

2-(4-Hydroxy-3-methoxybenzylidene)hydrazine-1-carboxamide (7a).²⁵ Yellow solid (1.3 g, 98% yield). *R_f* (5% MeOH/EtOAc): 0.51. Mp: 228–230 °C. ¹H NMR (300.13 MHz, DMSO- d_6): δ 3.81 (s, 3H), 6.45 (br s, 2H), 6.75 (d, ${}^{3}J_{\text{HH}} = 8.1$ Hz, 1H), 6.96 (dd, ${}^{3}J_{\text{HH}} = 8.1$, ${}^{4}J_{\text{H}} = 4.1$ $J_{\text{HH}} = 1.9 \text{ Hz}, 1 \text{H}$), 7.97 (d, 4 J_{HH} = 1.9, 1H), 7.71 (s, 1H), 9.31 (br s, 1H), 10.03 (br s, 1H). ¹³C NMR (75.5 MHz, DMSO- d_6): δ 55.7 (CH3), 109.0 (CH), 115.2 (CH), 121.1 (CH), 126.4 (C), 139.9 (CH), 147.9 (C), 148.0 (C), 157.0 (C). HRMS (ESI⁺, m/z): calcd for $(C_9H_{12}N_3O_3)^+ (M + H)^+$ 210.0873, found: 210.0872. BSC Advances Wewerkets Articles. Proposed on 40 October 2017. This article is licensed on 4/22/2021. The set of the set of

2-(4-Hydroxy-3,5-dimethoxybenzylidene)hydrazine-1-carboxamide (7b). Light orange solid (1.4 g, 89% yield). R_f (5% MeOH/ EtOAc): 0.42. Mp: 148-150 °C. ¹H NMR (300.13 MHz, DMSO d_6): δ 3.79 (s, 6H), 6.49 (br s, 2H), 6.97 (s, 2H), 7.70 (s, 1H), 8.65 (br s, 1H), 10.09 (s, 1H). ¹³C NMR (75.5 MHz, DMSO- d_6): δ 56.1 (2CH3), 104.2 (2CH), 125.2 (C), 136.9 (C), 139.9 (CH), 148.1 (2C), 157.0 (C). HRMS (ESI⁺, *m*/z): calcd for $(C_{10}H_{14}N_3O_4)^+$ $(M + H)^+$ 240.0979, found: 240.0975.

2-(3,4-Dihydroxybenzylidene)hydrazine-1-carboxamide (7c). Orange solid (936 mg, 73% yield). R_f (5% MeOH/EtOAc): 0.49. Mp: decomposes at 210 °C. 1 H NMR (300.13 MHz, DMSO- d_{6}): δ 6.30 (br s, 2H), 6.72 (d, 3 J_{HH} = 8.1 Hz, 1H), 6.92 (dd, 3 J_{HH} = 8.1,
 4 I = 2.0 Hz, 1H), 7.09 (d, 4 I = 2.0, 1H), 7.66 (s, 1H), 8.98 $J_{\rm HH} = 2.0$ Hz, 1H), 7.09 (d, $^{4}J_{\rm HH} = 2.0$, 1H), 7.66 (s, 1H), 8.98 (br s, 1H), 9.31 (br s, 1H), 9.98 (s, 1H). 13C NMR (75.5 MHz, DMSO-d6): d 113.2 (CH), 115.5 (CH), 119.1 (CH), 126.3 (C), 140.2 (CH), 145.5 (C), 146.9 (C), 156.9 (C). HRMS (ESI⁺, m/z): calcd for $(C_8H_{10}N_3O_3)^+ (M + H)^+$ 196.0717, found: 196.0709.

General procedure for the synthesis of triazoles 8a–c

4-Aminotriazole (12, 55 mg, 0.657 mmol) and an aqueous concentrated sulfuric acid solution (5 drops) were successively added to a solution of $3a-c$ (0.657 mmol) in ethanol (2 mL). The mixture was stirred under reflux for $3-7.5$ h (see Table 1) and the resulting precipitate was filtrated from the reaction media, yielding 8a–c in 96–99% yield.

4-(((4H-1,2,4-Triazol-4-yl)imino)methyl)-2-methoxyphenol (8a). 16 White solid (142 mg, 99% yield). R_f (5% MeOH/EtOAc): 0.40. Mp: 198–200 °C. ¹H NMR (300.13 MHz, DMSO- d_6): δ 3.84 (s, 3H), 6.97 $(d, {}^{3}J_{\text{HH}} = 8.2 \text{ Hz}, 1\text{H}), 7.33 \text{ (dd, } {}^{3}J_{\text{HH}} = 8.2, {}^{4}J_{\text{HH}} = 1.9 \text{ Hz}, 1\text{H}), 7.44$

 $(d, {}^{4}J_{HH} = 1.9, 1H)$, 8.03 (br s, 1H), 9.00 (s, 1H), 9.66 (s, 2H). ¹³C NMR (75.5 MHz, DMSO- d_6): δ 55.7 (CH₃), 110.2 (CH), 115.9 (CH), 122.4 (C), 125.5 (CH), 139.4 (2CH), 148.3 (C), 152.1 (C), 162.5 (CH). HRMS (ESI⁺, *m*/z): calcd for $(C_{10}H_{11}N_4O_2)^+$ (M + H)⁺ 219.0877, found: 219.0878.

4-(((4H-1,2,4-Triazol-4-yl)imino)methyl)-2,6-dimethoxyphenol (8b). White solid (161 mg, 99% yield). R_f (10% MeOH/EtOAc): 0.25. Mp: decomposes at 160 °C. 1 H NMR (300.13 MHz, DMSO d_6): δ 3.84 (s, 6H), 6.44 (br s, 1H), 7.16 (s, 2H), 8.99 (s, 1H), 9.48 (s, 2H). ¹³C NMR (75.5 MHz, DMSO- d_6): δ 56.1 (2CH₃), 106.3 (2CH), 121.6 (C), 139.1 (2CH), 140.6 (C), 148.2 (2C), 160.8 (CH). HRMS (ESI⁺, *m*/z): calcd for $(C_{11}H_{13}N_4O_3)^+ (M + H)^+$ 249.0982, found: 249.0982.

4-(((4H-1,2,4-Triazol-4-yl)imino)methyl)benzene-1,2-diol (8c). White solid (129 mg, 96% yield). R_f (5% MeOH/EtOAc): 0.37. Mp: decomposes at 224-226 °C. ¹H NMR (300.13 MHz, DMSO- d_6): δ 4.29 (br s, 2H), 6.89 (d, 3 *H_H* = 8.1 Hz, 1H), 7.16 (dd, 3 *H_H* = 8.2, 4
 4 *I* = 2.0 Hz, 1H), 7.32 (d, 4 *I* = 2.0, 1H), 8.88 (c, 1H), 0.38 (c, $J_{\text{HH}} = 2.0 \text{ Hz}, 1\text{H}$), 7.32 (d, 4 $J_{\text{HH}} = 2.0, 1\text{H}$), 8.88 (s, 1H), 9.38 (s, 2H). ¹³C NMR (75.5 MHz, DMSO- d_6): δ 114.1 (CH), 116.0 (CH), 122.9 (C), 123.4 (CH), 139.2 (2CH), 146.1 (C), 150.8 (C), 161.0 (CH). HRMS (ESI⁺, *m*/z): calcd for $(C_9H_9N_4O_2)^+ (M + H)^+$ 205.0720, found: 205.0721.

General procedure for the synthesis of hydrazono tetrazoles 9a–c

5-Hydrazinyl-1H-tetrazole (13, 227 mg, 2.27 mmol) and an aqueous concentrated HCl solution (5 drops) were successively added to a solution of 3a–c (2.27 mmol) in ethanol (4 mL). The mixture was stirred under reflux for $1.75-5$ h (see Table 1) and the resulting precipitate was filtrated from the reaction media, yielding the corresponding tetrazoles 9a–c in 80–97% yield.

4-((2-(1H-Tetrazol-5-yl)hydrazono)methyl)-2-methoxyphenol (9a). White solid (500 mg, 94% yield). R_f (10% MeOH/EtOAc): 0.72. Mp: 197–198 °C. ¹H NMR (300.13 MHz, DMSO- d_6): δ 3.84 (s, 3H), 6.75 (br s, 2H), overlapped with 6.81 (d, $^{3}J_{\text{HH}} = 8.1$ Hz, 1H) and 7.04 (dd, $^3J_{\text{HH}} = 8.1, \frac{4J_{\text{HH}}}{J_{\text{HH}}} = 1.9 \text{ Hz}, 1 \text{ H}$), 7.48 (d, $^4J_{\text{HH}} =$ 1.9, 1H), 7.93 (s, 1H), 11.61 (br s, 1H). ¹³C NMR (75.5 MHz, DMSO-d₆): δ 55.9 (CH₃), 109.2 (CH), 115.4 (CH), 121.6 (CH), 125.9 (C), 143.8 (CH), 148.2 (C), 148.5 (C), 156.6 (C). HRMS (ESI⁺, *m*/z): calcd for $(C_9H_{11}N_6O_2)^+ (M + H)^+$ 235.0938, found: 235.0936.

4-((2-(1H-Tetrazol-5-yl)hydrazono)methyl)-2,6-dimethoxyphenol (9b). White solid (582 mg, 97% yield). R_f (10% MeOH/EtOAc): 0.66. Mp: decomposes at 197–199 °C. 1 H NMR (300.13 MHz, DMSO- d_{6}): d 3.82 (s, 6H), 7.07 (s, 2H), 7.93 (s, 1H), 8.09 (br s, 2H), 11.71 (br s, 1H). ¹³C NMR (75.5 MHz, DMSO- d_6): δ 56.2 (2CH₃), 104.5 (2CH), 124.8 (C), 137.3 (C), 143.7 (CH), 148.2 (2C), 156.5 (C). HRMS (ESI⁺, *m*/z): calcd for $(C_{10}H_{13}N_6O_3)^+ (M + H)^+$ 265.1044, found: 265.1038.

4-((2-(1H-Tetrazol-5-yl)hydrazono)methyl)benzene-1,2-diol (9c). White solid (400 mg, 80% yield). R_f (10% MeOH/EtOAc): 0.70. Mp: decomposes at 250 °C. ¹H NMR (300.13 MHz, DMSO-d₆): δ 6.77 (d, $\frac{3I}{I}$ = 8.1 Hz, 1H) 6.00 (dd, $\frac{3I}{I}$ = 8.1 $\frac{4I}{I}$ = 2.0 Hz, 1H) 7.20 (d) $\frac{3J_{\text{HH}}}{4}$ = 8.1 Hz, 1H), 6.99 (dd, $\frac{3J_{\text{HH}}}{4}$ = 8.1, $\frac{4J_{\text{HH}}}{2}$ = 2.0 Hz, 1H), 7.20 (d, $\frac{4J_{\text{HH}}}{4}$ = 2.0 Hz, 1H), 7.20 (d, 4 *J*_{HH} = 2.0, 1H), 7.86 (s, 1H), 8.99 (br s, 1H), 9.44 (br s, 1H), 11.45 (s, 1H), 15.23 (br s, 1H). ¹³C NMR (75.5 MHz, DMSO- d_6): δ 113.5 (CH), 115.5 (CH), 119.6 (CH), 125.9 (C), 144.1 (CH), 145.6 (C), 147.4 (C), 156.5 (C). HRMS (ESI⁺, m/z): calcd for $(C_8H_9N_6O_2)^+$ (M + H)⁺ 221.0781, found: 221.0783.

Synthesis of 5-hydrazinyl-1H-tetrazole 13

5-Aminotetrazole monohydrate (5 g, 48.51 mmol) was dissolved in a 15% aqueous NaOH solution (50 mL). The mixture was heated at 100 °C and KMD_4 (7 g) was slowly added by portions until the colour of the solution no longer changed. After stirring 30 min, ethanol was added to eliminate the excess of KMnO4. The warm solution was filtered and concentrated under reduced pressure to 1/3 of the initial volume. Sodium 5,5′-azotetrazolate was crystallized as a yellow solid (6.2 g, 61% yield).

To a suspension of sodium 5,5′-azotetrazolate (6.2 g, 29.51 mmol) in water (62 mL), an aqueous HCl 6 N solution (13 mL) was added. The mixture was warmed until the gas evolution finished, then evaporated to dryness from water three times to ensure the removal of HCl. The residue was dissolved in the minimum volume of hot water and a hot solution of NaOAc (6.2 g, 75.58 mmol) in water (6.2 mL) was added. The apparatus was flushed out with carbon dioxide and the solution was allowed to cool to give 5-hydrazinyl-1H-tetrazole (13) as a white solid (987 mg, 20% yield). Mp: 192–194 °C. $^1\mathrm{H}$ NMR (300.13 MHz, DMSO- d_6): δ 8.17 (s, 4H). ¹³C NMR (75.5 MHz, DMSO- d_6): δ 161.4 (C).

General procedure for the laccase-catalyzed oxidation of the new synthesized compounds

In a test tube, the corresponding derivative was dissolved in an organic cosolvent (EtOH or DMSO, 300 µL). Then, the corresponding buffer saturated with $O₂$ prior to the experiment (2.7 mL) and the commercial laccases from Trametes versicolor (25 U) or the laccase from Myceliophthora thermophila (Novozym 51003, 31.4 U), were successively added. A balloon that was filled with oxygen was attached to the test tube and the mixture was stirred at 30 \degree C or 50 \degree C for the time shown in Table 2. Then, the reaction was finished by extraction with EtOAc $(3 \times 2 \text{ mL})$, and the organic layers were combined and dried with $Na₂SO₄$. After solvent evaporation under reduced pressure, the reaction crude was analyzed by 1 H-NMR. In those cases in which an abundant precipitate was detected in the reaction medium, this was separated by decantation and analyzed by 1 H-NMR.

6,6′-Dihydroxy-5,5′-dimethoxy-[1,1′-biphenyl]-3,3′-dicarbaldehyde (14).²⁶ Brown solid. ¹H NMR (300.13 MHz, DMSO- d_6): δ 3.92 (s, 6H), 7.41 (d, 4 J_{HH} = 1.9 Hz, 2H), 7.44 (d, 4 J_{HH} = 1.9 Hz, 2H), 9.80 (s, 2H), 10.11 (br s, 2H). ¹³C NMR (75.5 MHz, DMSO d_6): δ 56.0 (2CH₃), 108.9 (2CH), 124.9 (2C), 127.4 (2C), 128.3 $(2CH), 148.4 (2C), 151.3 (2C), 191.1 (2CH). HRMS (ESI⁻, m/z):$ calcd for $(C_{16}H_{13}O_6)^{-}$ $(M - H)^{-}$ 301.0718, found: 301.0706.

6,6′-Dihydroxy-5,5′-dimethoxy-[1,1′-biphenyl]-3,3′-dicarbonitrile (15).²⁷ White solid. ¹H NMR (300.13 MHz, DMSO- d_6): δ 3.89 $(s, 6H)$, 7.23 $(d, {}^{4}J_{HH} = 1.9$ Hz, 2H), 7.41 $(d, {}^{4}J_{HH} = 1.9$ Hz, 2H), 9.85 (br s, 2H). ¹³C NMR (75.5 MHz, DMSO-d₆): δ 56.4 (2CH₃), 100.3 (2C), 114.1 (2CH), 119.5 (2C), 124.6 (2C) 128.1 (2CH), 147.9 (2C), 148.9 (2C). HRMS (ESI⁺, *m*/z): calcd for $(C_{16}H_{13}N_2O_4)^+$ (M + H)⁺ 297.0870, found: 297.0870.

6,6'-Dihydroxy-5,5'-dimethoxy-[1,1'-biphenyl]-3,3'-dicarbaldehyde dioxime (16). 28 Light brown solid. 1 H NMR (300.13 MHz, MeOD- d_4): δ 3.92 (s, 6H), 6.99 (d, $^4J_{HH}$ = 1.6 Hz, 2H), 7.27 (d, $^4J_{HH}$ $=$ 1.5 Hz, 2H), 8.01 (s, 2H). ¹³C NMR (75.5 MHz, MeOD- d_4): δ 56.5 (2CH₃), 108.4 (2CH), 124.9 (2CH), 125.4 (2C), 126.3 (2C), 146.7 (2C), 149.5 (2C), 150.6 (2CH). HRMS (ESI⁺, m/z): calcd for $(C_{16}H_{17}N_2O_6)^+$ $(M + H)^+$ 333.1081, found: 333.1086.

2,2′-((6,6′-Dihydroxy-5,5′-dimethoxy-[1,1′-biphenyl]-3,3′-diyl)bis(methanylylidene))bis(hydrazine-1-carboxamide) (17). ²⁹ Dark brown solid. ¹H NMR (300.13 MHz, DMSO- d_6): δ 3.89 (s, 6H), 6.47 (br s, 4H), 6.96 (s, 2H), 7.39 (s, 2H), 7.75 (s, 2H), 8.76 (br s, 2H), 10.04 (s, 2H). HRMS (ESI⁺, m/z): calcd for $\left(C_{18}H_{20}N_6NaO_6\right)^+$ $(M + Na)^+$ 439.1337, found: 439.1339.

 $5,5'-Bis((E)-((4H-1,2,4-triazol-4-yl)imino)$ methyl $]-3,3'-dime$ thoxy-[1,1'-biphenyl]-2,2'-diol (18). White solid. 1 H NMR $(300.13 \text{ MHz}, \text{DMSO-}d_6)$: δ 3.88 (s, 6H), 7.33 (d, $^4J_{HH} = 1.9 \text{ Hz}$, 2H), 7.39 (d, 4 *J_{HH}* = 1.9 Hz, 2H), 8.94 (s, 2H), 9.08 (s, 4H). ¹³C NMR (75.5 MHz, DMSO- d_6): δ 55.9 (2CH₃), 108.1 (2CH), 115.6 (2C), 121.5 (2C), 125.6 (2C), 126.4 (2CH), 138.9 (4CH), 148.8 (2C), 158.4 (2CH). HRMS (ESI⁺, m/z): calcd for $(C_{20}H_{19}N_8O_4)^+$ $(M + H)^+$ 435.1524, found: 435.1527.

General procedure for the laccase-catalyzed oxidation of lignin model compounds in the presence of natural or nitrogenated compounds

The corresponding derivative $(16 \mu \text{mol})$ was placed in a test tube. The buffer (2 mL) saturated with $O₂$ prior to the experiment (pH 4.7: acetate buffer 100 mM; pH 5–6: citric buffer 50 mM; pH 7: KPi buffer 100 mM) and the veratryl alcohol $(1, 6 \mu \text{mol})$ or adlerol $(2, 6 \mu \text{mol})$ were added. After stirring, the commercial laccases from Trametes versicolor (25 U) or the laccase from Myceliophthora thermophila (Novozym 51003, 31.4 U), were added and the reaction stirred at 30 or 50 \degree C, respectively, during 72 h. Then, the reaction was acidified with an aqueous HCl 2 N solution (4 drops) and extracted with EtOAc (2×1 mL). The organic layers were combined and dried with $Na₂SO₄$. For the oxidations of veratryl alcohol, an aliquot was taken and analyzed by GC. In the case of the oxidation of adlerol, the solvent of the organic layers was evaporated and the reaction crudes were analyzed by ¹H-NMR. Thus, the conversions were calculated. Paper

Synchromatographs of substitute (8, 415.1 amail) was dissolved $\frac{1}{2}$ and $\frac{1}{2}$ 5.6.1 (CH), 1,6.84 (CH), 1,4.84 (CH)

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by the Spanish Ministry of Economy and Competitiveness and the European Union (Project OXY-POL, PCIN-2014-134 from an ERA-NET ERA-IB 2 Call).

Notes and references

1 (a) O. Y. Abdelaziz, D. P. Brink, J. Prothmann, K. Ravi, M. Sun, J. García-Hidalgo, M. Sandahl, C. P. Hulteberg, C. Turner, G. Lidén and M. F. Gorwa-Grauslund, Biotechnol. Adv., 2016, 34, 1318–1346; (b) S. Sen, S. Patil and D. S. Argyropoulos, Green Chem., 2015, 17, 4862–4887;

(c) K. Wang, H. Yang, S. Guo, X. Yao and R.-C. Sun, J. Appl. Polym. Sci., 2014, 131, 39673.

- $2(a)$ X. Huang, O. M. M. Gonzalez, J. Zhu, T. I. Korányi, M. D. Boot and E. J. M. Hensen, Green Chem., 2017, 19, 175–187; (b) H. Yang, J. Li, J. Xu and L. Mo, BioResources, 2017, 12, 344–361; (c) H. Zhu, J. P. Max, C. L. Marcum, H. Luo, M. M. Abu-Omar and H. I. Kenttämaa, J. Am. Soc. Mass Spectrom., 2016, 27, 1813-1823; (d) M. D. Kärkäs, B. S. Matsuura, T. M. Monos, G. Magallanes and C. R. J. Stephenson, Org. Biomol. Chem., 2016, 14, 1853– 1914; (e) S. Huang, N. Mahmood, M. Tymchyshyn, Z. Yuan and C. Xu, Bioresour. Technol., 2014, 171, 95–102.
- 3 (a) G. de Gonzalo, D. I. Colpa, M. H. M. Habib and M. W. Fraaije, J. Biotechnol., 2016, 236, 110–119; (b) F. Asina, I. Brzonova, K. Voeller, E. Kozliak, A. Kubátová, B. Yao and Y. Ji, Bioresour. Technol., 2016, 220, 414–424; (c) L. Zhang, T. You, T. Zhou, L. Zhang and F. Xu, $Biokesources, 2016, 11, 3972-3986; (d) D. Salvachúa,$ R. Katahira, E. N. S. Cleveland, P. Khanna, M. G. Resch, B. A. Black, S. O. Purvine, M. Zink, A. Prieto, M. J. Martínez, A. T. Martínez, B. A. Simmons, J. M. Gladden and G. T. Beckham, Green Chem., 2016, 18, 6046–6062; (e) T. D. H. Bugg, M. Ahmad, E. M. Hardiman and R. Rahmanpour, Nat. Prod. Rep., 2011, 28, 1883–1896. Open Access Articles. Articles. Published on 2017. Downloaded on 30 October 2017. Downloaded on 4. A Notice is licensed under a common and the same of the same o
	- 4 (a) Z. Ahmad, M. Asgher, F. Hussain and M. A. Randhawa, BioResources, 2016, 11, 10511–10527; (b) C. Zhao, S. Xie, Y. Pu, R. Zhang, F. Huang, A. J. Ragauskas and J. S. Yuan, Green Chem., 2016, 18, 1306–1312; (c) P. Picart, L. Wiermans, M. Pérez-Sánchez, P. M. Grande, A. Schallmey and P. Domínguez de María, ACS Sustainable Chem. Eng., 2016, 4, 651–655.
	- 5 (a) Y. Ohta, R. Hasegawa, K. Kurosawa, A. H. Maeda, T. Koizumi, H. Nishimura, H. Okada, C. Qu, K. Saito, T. Watanabe and Y. Hatada, ChemSusChem, 2017, 10, 425– 433; (b) L. Yang, K. Seshan and Y. Li, Catal. Today, 2017, 298, 276–297; (c) T. D. Bugg and R. Rahmanpour, Curr. Opin. Chem. Biol., 2015, 29, 10–17.
	- 6 (a) X.-Y. Jiang, Q. Lu, X.-N. Ye, B. Hu and C.-Q. Dong, BioResources, 2016, 11, 6232–6243; (b) M. Akazawa, Y. Kato and Y. Kojima, J. Anal. Appl. Pyrolysis, 2016, 122, 355–364.
	- 7 (a) L. T. M. Pham, S. J. Kim and Y. H. Kim, Biotechnol. Biofuels, 2016, 9, 247; (b) E. Rosini, C. Allegretti, R. Melis, L. Cerioli, G. Conti, L. Pollegioni and P. D'Arrigo, Catal. Sci. Technol., 2016, 6, 2195–2205.
	- 8 (a) M. D. Cannatelli and A. J. Ragauskas, Chem. Rec., 2017, 17, 122–140; (b) T. Kudanga, B. Nemadziva and M. L. Roes-Hill, Appl. Microbiol. Biotechnol., 2017, 101, 13–33; (c) M. Mogharabi and M. A. Faramarzi, Adv. Synth. Catal., 2014, 356, 897–927; (d) S. Witayakran and A. J. Ragauskas, Adv. Synth. Catal., 2009, 351, 1187–1209; (e) S. Riva, Trends Biotechnol., 2005, 24, 219–226.
	- 9 (a) B. Yao, P. Kolla, R. Koodali, Y. Ding, S. Balaranjan, S. Shrestha and A. Smirnova, New J. Chem., 2017, 41, 958– 964; (b) S. Xie, Q. Sun, Y. Pu, F. Lin, S. Sun, X. Wang, A. J. Ragauskas and J. S. Yuan, ACS Sustainable Chem. Eng., 2017, 5, 2215–2223; (c) Q. Wang, S. Liu, G. Yang and J. Chen, Bioresour. Technol., 2015, 175, 638–641.
- 10 (a) L. Munk, A. M. Punt, M. A. Kabel and A. S. Meyer, RSC Adv., 2017, 7, 3358-3368; (b) M. D. Cannatelli and A. J. Ragauskas, Appl. Microbiol. Biotechnol., 2016, 100, 8685–8691; (c) D. Huber, A. Pellis, A. Daxbacher, G. S. Nyanhongo and G. M. Guebitz, Polymers, 2016, 8, 280/ 1–280/10; (d) H. Wu, J. Noro, Q. Wang, X. Fan, C. Silva and A. Cavaco-Paulo, RSC Adv., 2016, 6, 90427–90434; (e) I. F. Fițigău, C. G. Boeriu and F. Peter, Macromol. Symp., 2015, 352, 78–86.
- 11 (a) J. González-Sabín, N. Ríos-Lombardía, I. García, N. M. Vior, A. F. Braña, C. Méndez, J. A. Salas and F. Morís, Green Chem., 2016, 18, 989-994; (b) A. Díaz-Rodríguez, L. Martínez-Montero, I. Lavandera, V. Gotor and V. Gotor-Fernández, Adv. Synth. Catal., 2014, 356, $2321-2329$; (c) K. Kedziora, A. Díaz-Rodríguez, I. Lavandera, V. Gotor-Fernández and V. Gotor, Green Chem., 2014, 16, 2448–2453.
- 12 (a) A. Kirsch, K. Ostendorf, A. Kharazipour and M. Euring, BioResources, 2016, 11, 7091–7101; (b) A. Rico, J. Rencoret, J. C. del Río, A. T. Martínez and A. Gutiérrez, Biotechnol. Biofuels, 2014, 7, 6; (c) P. Könst, S. Kara, S. Kochius, D. Holtmann, I. W. C. E. Arends, R. Ludwig and F. Hollmann, ChemCatChem, 2013, 5, 3027–3032; (d) T. Rosado, P. Bernardo, K. Koci, A. V. Coelho, M. P. Robalo and L. O. Martins, Bioresour. Technol., 2012, 124, 371–378; (e) A. I. Cañas and S. Camarero, *Biotechnol. Adv.*, 2010, 28, 694–705.
- 13 (a) T. Saravanakumar, H.-S. Park, A.-Y. Mo, M.-S. Choi, D.-H. Kim and S.-M. Park, J. Mol. Catal. B: Enzym., 2016, 134, 196–205; (b) S. Slagman, J. Escorihuela, H. Zuilhof and M. C. R. Franssen, RSC Adv., 2016, 6, 99367–99375; (c) R. Reiss, J. Ihssen, M. Richter, E. Eichhorn, B. Schilling and L. Thöny-Meyer, *PLoS One*, 2013, 8, e65633.
- 14 J. Ravindran, G. V. Subbaraju, M. V. Ramani, B. Sung and B. B. Aggarwal, Biochem. Pharmacol., 2010, 79, 1658–1666.
- 15 H. E. Master, S. I. Khan and K. A. Poojari, Bioorg. Med. Chem., 2005, 13, 4891–4899.
- 16 H. S. Patel and K. B. Patel, Phosphorus, Sulfur Silicon Relat. Elem., 2009, 184, 2443–2452.
- 17 (a) M.-J. Han, H.-T. Choi and H.-G. Song, J. Microbiol., 2006, 43, 555-560; (b) P. Cassland and L. J. Jönsson, Appl. Microbiol. Biotechnol., 1999, 52, 393–400.
- 18 R. M. Berka, P. Schneider, E. J. Golightly, S. H. Brown, M. Madden, K. M. Brown, T. Halkier, K. Mondorf and F. Xu, Appl. Environ. Microbiol., 1997, 63, 3151–3157.
- 19 Dimethylsulfoxide is known as radical scavenger, but it has been described as a useful organic co-solvent in laccasecatalyzed reactions. See, for instance ref. 11c.
- 20 (a) B. Ramalingam, B. Sana, J. Seayad, F. J. Ghadessy and M. B. Sullivan, RSC Adv., 2017, 7, 11951–11958; (b) N. H. Pham, F. Hollmann, D. Kracher, M. Preims, D. Haltrich and R. Ludwig, J. Mol. Catal. B: Enzym., 2015, 120, 38–46; (c) K. Rittstieg, A. Suurnakki, T. Suortti, K. Kruus, G. Guebitz and J. Buchert, Enzyme Microb. Technol., 2002, 31, 403–410.
- 21 See, for instance: (a) J. Safaei-Ghomi and A. R. Hajipour, Org. Prep. Proced. Int., 2011, 43, 372–376; (b) D. Enders,

L. Wortmann and R. Peters, Acc. Chem. Res., 2000, 33, 157–

- 169. 22 (a) P. Nousiainen, J. Kontro, H. Manner, A. Hatakka and J. Sipilä, Fungal Genet. Biol., 2014, 72, 137-149; (b) T. M. Larson, A. M. Anderson and J. O. Rich, Biotechnol. Lett., 2013, 35, 225-231; (c) M. Díaz-González, T. Vidal and T. Tzanov, Appl. Microbiol. Biotechnol., 2011, 89, 1693–1700; (d) R. Bourbonnais and M. G. Paice, FEBS Lett., 1990, 267, 99–102. Paper

L. Wortmann and R. Peters, Art. Chon, Res., 2004, 33, 137-25 R. B. de Oliveira, E. M. Sonare-Pagnades, R. P. P. Sonare

21 Open Payer (2017), Northern A. Hensieke article. The Machine A. A. Sonardie, A. U. Service i
	- 23 M. L. Niku-Paavola, L. Raaska and M. Itävaara, Mycol. Res., 1990, 94, 27–31.
	- 24 V. R. Kanetkar, D. D. Zope and Y. V. V. Subrahmanyam, Indian J. Chem., Sect. B: Org. Chem. Incl. Med. Chem., 2005, 44, 1317–1318.
- 25 R. B. de Oliveira, E. M. de Souza-Fagundes, R. P. P. Soares, A. A. Andrade, A. U. Krettli and C. L. Zani, Eur. J. Med. Chem., 2008, 43, 1983–1988.
- 26 M. Delomenède, F. Bedos-Belval, H. Duran, C. Vindis, M. Baldas and A. Nègre-Salvayre, J. Med. Chem., 2008, 51, 3171–3181.
- 27 S. Grelier, H. Cramail, A. Llevot, S. Carlotti and E. Grau, Patent WO 2016/050988 A1, 2016.
- 28 D. R. Kelly, S. C. Baker, D. S. King, D. S. de Silva, G. Lord and J. P. Taylor, Org. Biomol. Chem., 2008, 6, 787–796.
- 29 A. M. Costero, S. Gil, M. Parra, P. M. E. Mancini, M. N. Kneeteman and M. I. Quindt, Tetrahedron Lett., 2015, 56, 3988–3991.