RSC Advances



View Article Online

View Journal | View Issue

COMMENT

Check for updates

Cite this: RSC Adv., 2017, 7, 53617

Received 30th September 2017 Accepted 27th October 2017

DOI: 10.1039/c7ra10812e

rsc.li/rsc-advances

Introduction

Anomeric Based Oxidation (ABO) has opened up a new synthetic and theoretical branch of oxidation to make organic molecules where some of them have profound medicinal and biological activities.^{1–5} In this response to the comments of Dr Salehzadeh on the two recently published papers by Zolfigol *et al.*,^{1–3} we report on a study including a response to the main basis of the "Comments"⁶ problems to provide clarification for readers and to avoid confusion on the basic concepts. The items have been responded to one by one and some more advantages were added to the subject of our studies. Fortunately, the subject of the two papers and the other related published papers has interesting

^bFaculty of Chemistry, Bu-Ali Sina University, Hamedan, Iran. E-mail: zolfi@basu.ac. ir; mzolfigol@yahoo.com

Reply to the 'Comments on "Experimental and theoretical studies of the nanostructured $\{Fe_3O_4@SiO_2@(CH_2)_3Im\}C(CN)_3$ catalyst for 2-amino-3-cyanopyridine preparation *via* an anomeric based oxidation", *RSC Adv.*, 2016, 6, 50100–50111, and "The first computational study for the oxidative aromatization of pyrazolines and 1,4-dihydropyridines using 1,2,4-triazolinediones: an anomeric-based oxidation", *RSC Adv.*, 2016, 6, 102280–102291' by S. Salehzadeh, *RSC Adv.*, 2017, 7, 39704–39707, DOI: 10.1039/c6ra27033f⁺

Avat (Arman) Taherpour*ac and Mohammad Ali Zolfigol ^b

This response to Dr Salehzadeh's comments on the papers mentioned in the title contains the comments where they have mentioned disagreement with basic chemistry concepts. The response to the comments include: (i) the experimental (X-ray) and theoretical reported results of the epimerism and the differences in the stereoisomer properties, as they are definitely not pair of enantiomers, and (ii) a discussion regarding internal molecular orbital (MO) electron transfer.

advantages relating to ABO phenomena, the MBO method and stereochemistry topics. These advantages were developed here in the response to the mentioned comment.

1. The experimental (X-ray) and theoretical reported results of the epimerism and the differences in the stereoisomer properties

Amine molecules with different substituent groups on the pyramidal N atom ($R_1R_2R_3N$; $R_1 \neq R_2 \neq R_3$) show chiral properties.⁷ The barrier energy of the two enantiomers of this type of amine could be between low and high energies depending on the R_1 , R_2 and R_3 properties and the character of the molecular structures. The barrier energy of the two enantiomers would be high if there was a main steric restraint and strong hindering effect on the N atom due to the structure becoming planar with sp² hybridization.⁷ The interconversion between the enantiomers would stop if the lone pair of the N atom became protonated and/or the sp² N could not become planar because of the above mentioned reasons. See Fig. 1. The two forms will be diastereomers if another chiral centre (stereogenic centre) appears in the molecule.

In this case, as there are two chiral centres, the two stereoisomers are *epimeric* diastereomers (and they are not

^aDepartment of Organic Chemistry, Faculty of Chemistry, Razi University, P.O.Box: 67149-67346, Kermanshah, Iran. E-mail: avatarman.taherpour@gmail.com

^eMedical Biology Research Center, Kermanshah University of Medical Sciences, Kermanshah, Iran

[†] Electronic supplementary information (ESI) available: The X-ray crystals relating to the 1,4-dihydro-4-phenylpyridine derivatives and also some educational reported results and related figures are shown in the ESI (S1–S3). The standard definition of the main arrows, which were applied, are shown in the ESI (S4). See DOI: 10.1039/c7ra10812e



Fig. 1 Interconversion of N-pyramidal atoms in enantiomeric amine derivatives and the restriction effects on the interconversion process.

enantiomers). This type of diastereomer has different chemical, energetic and structural properties such as differences in thermodynamic energy levels, kinetic properties, HOMO & LUMO energy levels, $\Delta E_{\rm HOMO-LUMO}$, dipole moments and optical activities.⁸

The experimental (X-ray) and theoretical computational calculations have confirmed that the N atoms in the structures of 1,4-dihydro-4-phenylpyridine derivatives have pyramidal geometries. See the X-ray results for some 1,4-dihydro-4-phenylpyridine derivatives reported in 2016 by Prasad and Begum in the ESI. $\dagger^{9,10a,b}$ In the X-ray results the N atoms are not planar (pyramidal form) and just one of the stereoisomers (diastereomer) was represented.

The two compounds that the author⁶ has pointed out are not enantiomers and they are essentially diastereomers (epimers). The main point is that in the structures of the epimer isomers the N atom shows pyramidality. So, the two structures are not mirror pictures of each other. As can be seen in Fig. 3 of the paper "RSC Adv., 2016, 6, 50100-50111",² there are different orientations of the N atoms in the two R and S isomers of intermediate 7. So, for "7" there are two epimers and not two enantiomers. Because of the different chemical properties of the "epimers" they have different kinetic and thermodynamic properties and this has been completely explained in the paper "RSC Adv., 2016, 6, 50100-50111".² Fig. 2 demonstrates the possible diastereomer structures ($C_R N_R$, $C_R N_S$, $C_S N_S$ and $C_S N_R$). The pyramidality of the structures has been calculated in the optimized structure using the DFT-B3LYP/6-31G** method. The pyramidality differences between the sp² N and sp³ N (and NH₃ and *NHMeEt; * = chiral amine) were obtained to be about 57° and 40°, respectively. This topic, *i.e.* the conversion of the epimers to each other, is one of the new aspects in these studies.

Due to the existence of a nitrogen atom within the structure that possesses four different substituents, the molecule has two identified stereogenic centres so the existence of diastereomers



Fig. 2 The stereochemistry relationships between the structures.

is likely. It is clear that the diastereomers are quite different in terms of their chemical and physical properties. So it is not surprising that the investigated intermediates S and R are different from a chemical stability standpoint. Due to pyramidal inversion at the N chiral centres, we have considered only the C chiral centre. See Fig. 2.

Fig. 3 demonstrates the effect that a Lewis acid has on the chiral N atoms (and also $-NH_2$ as an achiral functional group). It can be seen that the relationship between structures A'-D' (after the addition of the Lewis acid on the N atoms) is the same as the relationship between structures A-D. Upon addition of a Lewis acid the chirality of the N1 atom would be rigid and the chirality of this atom would be clearer. In this case, the C4 carbon atoms retain their chirality states. The structures of A-D and A'-D' are shown in Fig. 3 and the stereochemical relationships between the structures are shown in the box of Fig. 3.

Fig. 3 also shows the pyramidalities and the angle deviations of the chiral N atoms in the possible diastereomer structures $(C_RN_R; C_RN_S; C_SN_S; C_SN_R)$. The pyramidalities for C_SN_S , C_RN_R , C_RN_S and C_SN_R were obtained as 7.8, 10.1, 10.1 and 7.9°, respectively. The angle deviations of the chiral N atoms from sp² hybridization for C_SN_S , C_RN_R , C_RN_S and C_SN_R were obtained as: 30, 33, 33 and 30°, respectively. The van der Waals repulsions and Pitzer strains¹¹ between N–H and C–H and/or C–Ph bonds are the main components, along with other substituent group effects, for the construction of the A–D forms of the epimers.

The calculated relative energy between $C_S N_S$ and $C_S N_R$ (A and D; as the most stable epimers) was zero. The calculated relative energies between $C_S N_S$ and/or $C_S N_R$ (A and D) with $C_R N_R$ (B) and $C_R N_S$ (C) were 0.31 and 3.61 kcal mol⁻¹, respectively. The domain of the barrier energies for the N-inversion process are about 8–12 kcal mol⁻¹. The N-inversion process in the interconversion process of the epimers were calculated about 10.5 kcal mol⁻¹, respectively, by DFT-B3LYP/6-31G** method. The energy level differences, barrier to the N-inversion in the



Fig. 3 The structures of A–D (precursors) and A'–D' (after Lewis acid addition). The stereochemical relationships between the structures A–D and A'–D' are shown in the box. The possible epimer structures (C_RN_R ; C_RN_S ; C_SN_S ; C_SN_R). The van der Waals repulsion and Pitzer strain effects.² The pyramidality and angle deviations of the N chiral atoms from sp² hybridization.

interconversion process of the epimers, pyramidality and other structural and thermodynamic properties will change by changing the calculation method. However, the concepts of the discussed properties will stay the same. So, overall, in contradiction to Dr Salehzadeh's comments, the structures are not enantiomers and they are inherently structural diastereomers (epimers). The zero energy differences between the two $C_S N_S$ and $C_{S}N_{R}$ (A and D) epimers do not mean that they are structurally the same. They are different and they are diastereomers. In the 1,4-dihydro-4-phenylpyridine derivatives the N-H group achieves a planar and sp^2 geometry (zwitterionic radical) after hv irradiation (as a useful dyad).^{10b} To achieve the epimers in an experimental attempt one could apply a simple Lewis acid (for example D⁺ and not H⁺) to make the rigid structures and extract them. The X-ray results^{8,9} (as discussed) have shown just a pyramidal geometry for the N atom in the obtained crystals.

In the first paragraph of part 1 of the "Comments"⁶ the author has mentioned that: "Indeed, the energies of enantiomers can be different, in the femto-joule to pico-joule per mole

range, only because of a parity violation.⁴ However, recently Zolfigol et al. during the study on the intermediate molecule shown in Fig. 1 have reported that the S isomer is, about 0.29 kcal mol⁻¹ (\approx 1200 J mol⁻¹), more stable than the *R* one".⁶ In 2017, Juaristi et al. have reported an interesting study about stereoelectronic interactions as a probe for the existence of the intramolecular α-effect.¹² In Table S4 (ESI)† of their investigation the calculated differences in ΔE and ΔG for the axial/ equatorial conformational equilibria in some compounds with and without an anomeric effect were reported.12 In some cases the differences in ΔE and ΔG for the axial/equatorial conformations were trivial, so that in some conformers the reported calculated energy differences using the MP2/6-311+G(d,p) level of theory were between 0.1 and 0.3 kcal mol⁻¹.¹² In our study, the reported difference between the two diastereomers was 0.29 (≈ 0.3) kcal mol⁻¹. We have introduced the data with two decimal numbers in kcal mol⁻¹ (obviously, these data in our study were in kcal mol^{-1} , and not in femto-joule to pico-joule per mole), so, the mentioned phrases do not have any relationship with our study and, secondly, as it was completely discussed the zero energy difference between the two diastereomers does not mean that they are same. The 0.29 kcal mol⁻¹ value is another confirmation for their differences and by using another QM method (which calculates the steric restraint, van der Waals repulsions, Pitzer strains and other important effects with higher accuracy) we could see larger energy differences between these two structures (see the interpretations). It is obvious that the two diastereomers have different TS energies (different $\Delta G^{\#}$) in reactions with chiral reagents. In the introduction section the author of the "Comments"6 also mentioned that "the authors have proved that the transition state (TS) for R isomer is, about 4.5 kcal mol⁻¹ (\approx 19 000 J mol⁻¹), more stable than S one while we remember that these chiral molecules have reacted with an achiral ion."6 The author of the "Comments"6 should note to this point that the difference energy (4.5 kcal mol^{-1}) is related to the two different TS forms of the ABO reactions of the two diastereomers (with the special stereospecific structure of ${Fe_3O_4@SiO_2@(CH_2)_3Im}$. In the next step of the reaction the intermediate reacts with the achiral $^{-}C(CN)_{3}$ anion. In addition to the other discussed reasons related to the small differences in the energy levels, there are some studies in the literature that show that the small amounts of energy (up to about 0.2- $0.5 \text{ kcal mol}^{-1}$) are determinative of the mechanism patterns, TS and/or the results of the studies.13

In the paper "*RSC Adv.*, 2016, **6**, 50100–50111"² the *R* and *S* signs were applied to introduce just the chiral position of the anomeric chiral centre (to identify the structures) and we have noticed to this point that they are definitely diastereomers (epimers) which is in agreement with the discussion. In Fig. 1 of the "Comments" the author of the "Comments" showed the "schematic representation of chiral carbon in the compound studied by Zolfigol *et al.* (left)" and the molecular structures of the predicted *R* and *S* stereoisomers⁶ (right). The figures on the right side obviously have pyramidal and chiral N^{*} atoms and they agree with the above interpretations (in contradiction to Dr



Fig. 4 Anionic tautomers interconversion is the correct phrase for this phenomenon and it is not an "equilibrium between different structural isomers".

Salehzadeh's comments) that they are epimer diastereomers and not enantiomers.

All of the concepts and signs applied agree with standard texts and the IUPAC Gold book and are shown in Table S4 in the ESI† section.¹⁴⁻¹⁸ As shown in Table S4 in the ESI† section, arrows 1 and 2 represent electron transfer (in contradiction to Dr Salehzadeh's comments). Fig. 3 in the text of the "Comments"⁶ was not in the main papers.¹⁻⁴ Fig. 3 of Dr Salehzadeh's comments has incorrectly presented tautomers interconversion as an "equilibrium between different structural isomers".⁶ To the best of our knowledge, most of the structural isomers are detectable and/or separable from each other, whereas here the anionic tautomers are inseparable from each other (Fig. 4).

2. Discussion regarding internal molecular orbital (MO) electron transfer

Our explanation is obviously related to the ET process between the MO of the discussed molecules (such as $n_{N2} \rightarrow \pi^*_{C3C4}$ in the ET from n_{N2} and π^*_{C3C4} and $n_{N2} \rightarrow \pi^*_{C3C4}$ in the ET from n_{N2} to π^*_{C3C4}) and not between one atom/molecule/ion to another one. The final result of such an internal MO-ET process is the production of $H^{\delta-}$ and the separation of H^- from the discussed molecules in the ABO reactions. So, in addition to the above explanation, as one can see, in agreement with Marcus theory,¹⁹ there is an outer sphere ET process to produce $H^{\delta-}$ and then the separation of H^- from the discussed molecules occurs in the ABO reactions. Furthermore, an electron transfer phrase for the ET between MO orbitals was applied in the important primary and secondary sources. These concepts and our aims are too obvious so we do not need to do more interpretation about them.

Conclusion

The "Comments"⁶ on the papers "*RSC Adv.*, 2016, **6**, 50100– 50111" and "*RSC Adv.*, 2016, **6**, 102280–102291" were investigated. On the basis of the experimental (X-ray) and theoretical reported results, the epimerism and effect on the stereoisomer differences were discussed in part 1. The results have confirmed that the structures of the discussed 1,4-dihydro-4phenylpyridine derivatives are epimer diastereomers (C and N pyramidal atoms; C_RN_R , C_RN_S , C_SN_S and C_SN_R). In part 2, we explained the internal molecular orbital (MO) electron transfer on the basis of Marcus theory and its classification for ET processes. In agreement with Marcus theory, there is an outer sphere MO-ET process to produce $H^{\delta-}$ and then the separation of H^- from the discussed molecules occurs in the ABO reactions.²⁰

Conflicts of interest

There are no conflicts to declare.

Notes and references

- 1 M. A. Zolfigol, M. Kiafar, M. Yarie, A. Taherpour and M. Saeidi-Rad, *RSC Adv.*, 2016, **6**, 50100–50111.
- 2 M. Kiafar, M. A. Zolfigol, M. Yarie and A. Taherpour, *RSC Adv.*, 2016, **6**, 102280–102291.
- 3 M. Kiafar, M. A. Zolfigol, M. Yarie and A. Taherpour, *RSC Adv.*, 2016, **6**, 107295.
- 4 M. A. Zolfigol, M. Kiafar, M. Yarie, A. A. Taherpour, T. Fellowes, A. N. Hancok and A. Yari, *J. Mol. Struct.*, 2017, 1137, 674–680.
- 5 M. Yari, Iran. J. Catal., 2017, 7, 85-88.
- 6 S. Salehzadeh, RSC Adv., 2017, 7, 39704-39707.
- 7 (a) Nitrogen Inversion in regards to Stereochemistry, http:// web.chem.ucla.edu/~harding/ec_tutorials/tutorial05.pdf; (b) J. M. Lehn, Nitrogen inversion; Experiment and theory, Dynamic Stereochemistry, Part of the Fortschritte der Chemischen Forschung book series (TOPCURRCHEM, volume 15/3) Conference paper, 2006, pp. 311-377; (c) K. P. C. Vollhardt and N. E. Schore, Organic Chemistry, W. H. Freeman and Company, New York, USA, 6th edn, 2009; (d) Illustrated Glossary of Organic Chemistry, Nitrogen Inversion part, http://web.chem.ucla.edu/%7Eharding/ IGOC/N/nitrogen_inversion.html; (e) W. H. Brown, C. S. Foote, B. L. Iverson, E. V. Anslyn and B. M. Novak, Organic Chemistry, Brooks/Cole, Cengage Learning, Belmont, CA, USA, 6th edn, 2012, p. 119.
- 8 F. A. Carey and R. J. Sundberg, *Advanced Organic Chemistry-Part A*, Kluwer Academic/Plenum Publishers, New York, USA, 4th edn, 2000, pp. 103, and 131 (lines 2 and 3).
- 9 (a) E. V. Anslyn and D. A. Dougherty, *Modern Physical Organic Chemistry*, University Science Books, Sausalito, California, USA, 2006; (b) M. Abdoli-Senejani, A. A. Taherpour, H. R. Memarian and M. Khosravani, *Struct. Chem.*, 2013, 24, 191–200.
- 10 (a) N. L. Prasad and N. S. Begum, *IUCrData*, 2016, 1, x160722;
 (b) N. A. Al-Awadi, M. R. Ibrahim, M. H. Elnagdi, E. John and Y. A. Ibrahim, *Beilstein J. Org. Chem.*, 2012, 8, 441–447.
- 11 (a) IUPAC, Gold Book, Bond opposition strain, Pitzer strain, Torsional strain, https://goldbook.iupac.org/html/E/ E01886.html; (b) G. Quinkert, E. Egert and C. Greisinger, Aspects of Organic Chemistry; Structure, VCH, Verlag, New York, USA, 1996, p. 95.
- 12 E. Juaristi, G. P. Gomes, A. O. Terent'ev, R. Notario and I. V. Alabugin, *J. Am. Chem. Soc.*, 2017, **139**, 10799–10813.
- 13 (a) V. I. Minkin, B. Y. Simkin and R. M. Minyaev, Quantum Chemistry of Organic Compounds, Mechanisms of Reactions, Chapter, Low-Energy Barrier Reactions. Structural Modeling, Springer, Berlin, Heidelberg, 1990, pp. 181–189; (b)

Comment

I. Yavari, A. A. Taherpour and M. Dadgar, *J. Mol. Struct.: THEOCHEM*, 1998, **422**, 213–218; (*c*) I. Yavari and A. A. Taherpour, *J. Mol. Struct.: THEOCHEM*, 1999, **488**, 141–149.

- 14 IUPAC, Gold Book, mesomerism, https://goldbook.iupac.org/html/M/M03845.html.
- 15 IUPAC, Gold Book, resonance, https://goldbook.iupac.org/ html/R/R05326.html.
- 16 IUPAC, Gold Book, canonical forms, https://goldbook.iupac.org/html/C/C01309.html.
- 17 IUPAC, Gold Book, contributing structures, https://goldbook.iupac.org/plain/C01309-plain.html.
- 18 P. J. Taylor, G. van der Zwan and L. Antonov, *Tautomerism:* Methods and Theories, Edition: First, Chapter: Tautomerism: Introduction, History, and Recent Developments in

Experimental and Theoretical Methods, ed. L. Antonov, Wiley-VCH, 2014, pp. 1–24.

- 19 (a) R. A. Marcus, Rev. Mod. Phys., 1993, 65, 599-610; (b) M. Andrea, Marcus Theory for Electron Transfer a short introduction, MPIP-Journal Club-Mainz, January 2008, p. 29; (c) P. F. Barbara, J. Phys. Chem., 1996, 100, 13148-13161; (d) M. D. Newton, Chem. Rev., 1991, 91, 350; (e) J. Jortner and K. F. Freed, J. Chem. Phys., 1970, 52, 6272-6291; (f) R. A. Marcus, J. Chem. Phys., 1965, 43, 679; (g) R. A. Marcus and N. Sutin, Biochim. Biophys. Acta, 1985, 811, 265; (h) M. G. Kuzmin, XVIIth IUPAC Symposium on Photochemistry, Book of Abstracts, Dresden, German, 22-27 July 2000, p. 372.
- 20 Note: The calculated data in ref. 2–4 were performed by licensed Spartan '10, build 1.0.1, Wavefunction Inc., Irvine, 2011.