


Cite this: *RSC Adv.*, 2017, 7, 54196

Purification and identification of antioxidative peptides of palm kernel expeller glutelin-1 hydrolysates

Yajun Zheng,  [†]*^a Yan Li^{†a} and Yufeng Zhang^b

To obtain hydrolysates with high antioxidant activity and hydrolysis degree, palm kernel expeller glutelin-1 was hydrolyzed by pepsin assisted with high pressure pretreatment. The results of orthogonal experiment revealed that the optimum enzymatic hydrolysis conditions were as follows: enzymes concentration of 2 g/100 g, hydrolysis time of 4 h, temperature 37 °C and pH 2.0. Palm kernel expeller glutelin-1 hydrolysate was separated by ultrafiltration, Sephadex G-15 gel chromatography and reversed-phase high performance liquid chromatography. Finally, four peptides Thr-Val-Phe-Asp-Gly-Glu-Leu-Arg (935.5 Da), Ala-Asp-Val-Phe-Asn-Pro-Arg (818.7 Da), Cys-Ala-Gly-Val-Ser-Ala-Ile-Arg (832.4 Da) and Leu-Val-Tyr-Ile-Ile-Gln-Gly-Arg (819.4 Da) were identified and their IC₅₀ values on hydroxyl radical scavenging activities were 38.22 ± 2.22, 22.16 ± 1.22, 31.19 ± 1.67 and 12.85 ± 0.23 μg mL⁻¹, respectively. Furthermore, these peptides were chemically synthesized and the peptides ADVFNPR and CAGVSAIR showed good stability against simulated gastrointestinal protease digestion.

Received 22nd October 2017
Accepted 12th November 2017

DOI: 10.1039/c7ra11657h

rsc.li/rsc-advances

1. Introduction

Recently, increasing epidemiological evidence have indicated that excessive free radicals and associated oxidative damage are mediators in some chronic diseases, such as diabetes mellitus, neurodegenerative diseases and cardiovascular diseases.¹ Antioxidants play a vital role in both food systems as well as in the human body to reduce oxidative stress. Since the use of synthetic antioxidants has been suspected to threaten human health, antioxidants from natural sources have attracted increasing attention. Without marked adverse effects on the human body, peptides derived from various food proteins have been accepted as a new source of natural antioxidants. Some antioxidant peptides have been identified from varieties of food proteins such as soy protein, walnut protein, rice bran protein, potato protein and fish proteins.^{2–6} Except for the isolation and characterization, the *in vivo* physiological effects, structure–activity relationship and mechanism of action of some antioxidant peptides as well as their stability to processing and storage conditions used in food industry have been studied.⁷

Palm kernel expeller (PKE) is the main byproduct of palm milling industry, which contains about 20% protein.⁸ Palm kernel expeller protein (PKP) is a good potential edible protein due to its relatively well-balanced amino acid profile, good

solubility and emulsifying capacity.⁹ Globulin, glutelin-1 and glutelin-2 accounted for 40.10%, 24.01% and 33.32% of PKP, respectively.¹⁰ It was found that enzymatic hydrolysates of the globulin showed anticancer, antihypertensive and antibacterial activities; also several peptides obtained from the glutelin-2 could effectively lower the systolic blood pressure of spontaneously hypertensive rats.^{11–13} Dilute acid-soluble seed storage proteins are commonly known as glutelin-1 proteins.¹⁴ A previous study has demonstrated that palm kernel expeller glutelin-1 (PKEG-1) exhibited high scavenging activity of DPPH radical (77.69%), ABTS radical (64.19%), superoxide radical (49.54%) and hydroxyl radical (34.41%); also it exhibited high reducing power (0.211 ± 0.009),¹⁰ indicating that antioxidant peptides could be isolated from it. However, no study on the purification and identification of antioxidant peptides from PKEG-1 is available. In addition, nearly 2.5 million tons of PKE are produced per year in Malaysia alone,¹⁵ prompting efforts to explore it for value-addition products such as bioactive peptides.

In the current study, the optimal enzymatic hydrolysis condition of PKEG-1 was investigated and antioxidant peptide fractions were isolated by ultrafiltration membranes, Sephadex G-15 chromatography and reverse phase high performance liquid chromatography (RP-HPLC). The amino acid sequences of the individual antioxidant peptides were subsequently identified using capillary liquid chromatography/electrospray ionization/mass spectrometry/mass spectrometry (LC-MS/MS); in addition, the stability of antioxidant peptides against gastro-intestinal digestion was studied.

^aInstitute of Food Science of Shanxi Normal University, Linfen, 041004, China. E-mail: zyj_coconut@163.com; Fax: +86-0357-2051482; Tel: +86-13976563642

^bCoconut Research Institute of Chinese Academy of Tropical Agriculture Sciences, Wenchang, 571339, China

[†] The two authors contribute equally to this work and they are co-first author.



2. Materials and methods

2.1 Materials and reagents

Oil palm kernel expeller (PKE) was obtained from the Meiyue Food Processing Co. Ltd., China. Pepsin (5×10^4 U g⁻¹) was purchased from Shanghai Biotech. Co., Ltd. (China). Sephadex G-15 was purchased from Pharmacia (Uppsala, Sweden). ABTS (2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid)) and other reagents were of analytical grade.

2.2 Preparation of palm kernel expeller glutelin-1 (PKEG-1)

PKE was defatted three times with *n*-hexane (1 : 10, g mL⁻¹), and then dried, ground and passed through a sieve of 0.2 mm mesh. Then, 30 g of defatted PKE was suspended in 300 mL of 0.4 M NaCl, stirred at 4 °C for 1 h and filtered. The residue was collected and again suspended in 0.4 M NaCl and stirred at 4 °C for 1 h. This step was repeated in triplicate to remove PKE albumin and globulin. Then, the residue was suspended in 50% (v/v) glacial acetic acid (1 : 10, g mL⁻¹). After being stirred at 4 °C for 1 h, the mixture was centrifuged at 10 000g for 30 min. The supernatant was dialyzed against distilled water at 4 °C for 48 h, and then lyophilized. Subsequently, PKEG-1 was obtained and stored at -20 °C.

2.3 Optimization of the enzymatic hydrolysis

Initially, 2 g of PKEG-1 was rehydrated in 100 mL of acetate buffer (0.1 M, pH 2.0) and subjected to high pressure (HP) treatment following the method outlined by Zheng *et al.*¹³ HP treatment was performed by a HPP-L3 high-pressure apparatus (Tianjin HuaTaiSenMiao Biotechnology Co. Ltd., China) at 300 MPa and 35 °C, for 20 min. As PKEG-1 is an acid-soluble protein, pepsin was chosen to hydrolyze it. An orthogonal L₉ (4³) test design with hydroxyl radical scavenging activity (SA) as response was used to investigate the optimal hydrolysis conditions of pepsin. Four controllable variables with three levels, including enzyme-to-substrate ratio (E/S), hydrolysis time, temperature and pH were selected for optimization (Table 1).

Further, PKEG-1, treated with HP, was hydrolyzed by pepsin under the obtained optimum conditions. After digestion, the solutions were deactivated in boiling water for 10 min and centrifuged at 10 000g for 20 min. The supernatant was collected, and lyophilized to obtain PKEG-1 hydrolysates (PKEG-1H). The hydrolysis degree (DH) and scavenging activity on hydroxyl radical

([•]OH) were determined by the Adler-Nissen method¹⁶ and the 2-deoxyribose oxidation method,¹⁷ respectively.

2.4 Separation by ultrafiltration and Sephadex G-15 gel chromatography

The PKEG-1 hydrolysates (PKEG-1H) prepared, by pepsin assisted with HP, was filtered sequentially using an ultrafiltration unit (Pellicon XL, Millipore, USA) through two ultrafiltration membranes with molecular weight (MW) cut-offs of 5 and 3 kDa, respectively. Three fractions were obtained: PKEG-1H-I with MW > 5 kDa, PKEG-1H-II with MW 3–5 kDa and PKEG-1H-III with MW < 3 kDa. The fraction with the highest antioxidant activity was further separated by a Sephadex G-15 gel filtration column (Φ 1.6 cm × 100 cm), eluted with distilled water at a flow rate of 0.6 mL min⁻¹ and monitored at 220 nm. Fractions were collected, lyophilized and stored at -20 °C.

2.5 Antioxidant activity of PKEG-1H fractions

2.5.1 Superoxide radical-scavenging activity. Superoxide radical-scavenging activity was measured using the pyrogallol assay as described by Qiao *et al.*¹⁷ Briefly, 0.1 mL of sample solution was mixed with 3 mL of pyrogallol solution (3 mM); the absorbance at 320 nm was recorded at 30 s intervals. The scavenging activity was calculated from the absorbance at 320 nm in the presence or absence of samples. Glutathione (GSH) was used for comparison.

2.5.2 Scavenging activity of hydroxyl radical. Following the method reported by Qiao *et al.*,¹⁷ the reaction mixture contained sample solution (100 µg mL⁻¹), FeSO₄-EDTA (10 mM), 2-deoxyribose (10 mM), phosphate buffer (pH 7.4, 0.1 M) and H₂O₂ (10 mM). After being incubated at 37 °C for 1 h, the reaction solution was mixed with trichloroacetic acid (28 mg mL⁻¹) and thiobarbituric acid (10 mg mL⁻¹), and then incubated at 100 °C for 20 min. The absorbance was measured at 532 nm. GSH was used as the comparison. The activity was determined as follows:

$$\text{OH scavenging activity (\%)} = [1 - (A_S - A_B)/A_C] \times 100 \quad (1)$$

where *A_B* is the absorbance of the blank (distilled water instead of samples), *A_C* is the absorbance of control (without the addition of 2-deoxyribose oxidation) and *A_S* is the absorbance of mixture contained samples.

2.5.3 ABTS radical scavenging ability. According to the TEAC assay,¹⁸ 20 µL of protein solution (0.1–0.5 mg mL⁻¹) was mixed with 2 mL of ABTS solution, and then kept in the dark at 30 °C for 6 min. Then, absorbance at 734 nm was recorded. GSH was used for comparison. The activity was calculated as follows:

$$\text{Scavenging activity (\%)} = [1 - (A_S - A_B)/A_C] \times 100 \quad (2)$$

where *A_C* is the absorbance of control (distilled water instead of ABTS^{•+} solution).

2.5.4 Ferrous ion chelating capacity. Following the method reported by Joshi *et al.*,¹⁸ 450 µL sample solutions (0.1–0.5 mg mL⁻¹) were mixed with 45 µL FeCl₂ (2 mM) and 1815 µL of distilled water. Then, 90 µL ferrozine (5 mM) was

Table 1 Experimental values and levels of the independent variables for the L₉ (4³) orthogonal experiment

Independent variable	Levels		
	1	2	3
A E/S (g/100 g protein)	1	1.5	2.0
B hydrolysis time (h)	2.0	3.0	4.0
C temperature (°C)	32.0	37.0	42.0
D pH	2.0	3.0	4.0



added to start the reaction and the absorbance at 562 nm was read after 30 min. Chelating activity was calculated as follows:

$$\text{Chelating activity (\%)} = [1 - (A_S - A_B)/A_C] \times 100 \quad (3)$$

where A_C is the absorbance of control (without the addition of ferrozine).

2.6 Purification of the antioxidant peptides

The PKEG-1H fraction showing the highest antioxidant activity after the Sephadex G-15 chromatography was separated by RP-HPLC on a Zorbax semi-preparative C_{18} column (Φ 9.4 mm \times 250 mm, Agilent Technologies, USA), using a linear gradient of acetonitrile containing 0.1% TFA (5–30%, 30 min) at a flow rate of 2.5 mL min⁻¹. The fraction showing high antioxidant activity was further isolated on a Zorbax analysis C_{18} column (Φ 4.6 mm \times 250 mm, Agilent Technologies, USA) with a linear gradient of acetonitrile containing 0.1% TFA (5–25%, in 20 min) at a flow rate of 1.0 mL min⁻¹. Elution peaks were detected at 220 nm, concentrated, and lyophilized.

2.7 Molecular mass and amino acid sequence of the purified peptides

To identify molecular mass and accurate amino acid sequence of the purified peptides from RP-HPLC, the peptides were determined by LC-MS/MS coupled with a Eksigent Nano LC (Eksigent Technologies, Dublin, CA, USA) and Thermo LTQ linear ion trap mass spectrometer (Thermo Fisher, San Jose, CA, USA). The peptide sequences were matched to the published sequences of *Elaeis guineensis* proteins from the National Center for Biotechnology Information (NCBI, Bethesda, MD, USA) database.

2.8 Peptide synthesis

Following the solid phase procedure described by Zhang *et al.*,⁵ antioxidant peptides identified from PKEG-1H were synthesized with a Kromasil 100-5 C_{18} column (Φ 4.6 \times 250 mm, particle size 5 μ m). The synthetic peptides were confirmed by Liquid Chromatography coupled to Mass Spectrometry (LC-MS/ESI), and their purity was more than 95%.

2.9 Stability of the synthetic peptides

The synthetic peptides (1 mg mL⁻¹) were successively digested by pepsin (E/S = 1 : 35) and pancreatin (E/S = 1 : 25) as described by Tavares *et al.*¹⁹ The 'OH scavenging of the treated synthetic peptides was determined, and the untreated synthetic peptides were used for comparison.

2.10 Statistical analysis

All the results were the means of triplicates. Data were subjected to analysis of variance and Duncan test with a confidence interval of 95%.

3. Results and discussion

3.1 Optimization of hydrolysis conditions

The results of the orthogonal test are summarized in Table 2. It was revealed that the influential extent of the four factors to 'OH scavenging activity of PKEG-1 was: A (E/S) > C (temperature) > B (hydrolysis time) > D (pH); both E/S and temperature had significant effect on the 'OH scavenging activity ($P < 0.05$). The best level of these factors was A3B3C2D1, suggesting that the highest 'OH scavenging activity (89.14% \pm 4.84%) of PKEG-1 hydrolysates (1 mg mL⁻¹) could be obtained when the

Table 2 The result and variance analysis of the L_9 (4^3) orthogonal experiment

No.	A E/S (%)	B Hydrolysis time (h)	C Temperature (°C)	D pH	SA ^a (%)	DH ^b (%)
1	1	1	1	1	59.66 \pm 4.69	15.92 \pm 0.68
2	1	2	2	2	68.19 \pm 2.97	17.61 \pm 1.93
3	1	3	3	3	68.00 \pm 4.91	18.02 \pm 0.89
4	2	1	2	3	75.94 \pm 8.14	23.14 \pm 2.22
5	2	2	3	1	80.90 \pm 6.41	25.77 \pm 1.28
6	2	3	1	2	74.55 \pm 5.22	23.29 \pm 1.72
7	3	1	3	2	83.81 \pm 4.19	27.11 \pm 2.37
8	3	2	1	3	77.01 \pm 6.37	25.56 \pm 0.70
9	3	3	2	1	89.14 \pm 4.84	31.90 \pm 2.72
K_1	195.85	219.41	211.22	229.70		
K_2	231.39	226.11	233.26	226.55		
K_3	249.97	231.68	232.71	220.95		
Best level	A ₃	B ₃	C ₂	D ₁		
R^c	54.11	12.28	22.04	8.75		
R Order	A > C > B > D					
d_f	2	2	2	2		
Sum of squares	503.90	25.19	105.30	6.54		
F Value	38.51 ^d	1.92	8.05 ^d	1	$F_{0.05} = 6.94$	

^a Hydroxyl-radical scavenging. ^b Hydrolysis degree. ^c Refers to result of extreme analysis. ^d Significant variables.






Fig. 1 Sephadex G-15 gel chromatography profile of PKEG-1H (a) and hydroxyl radical ($\cdot\text{OH}$) scavenging activity of each fraction (b); chromatography of fraction *D* separated by semi-preparing RP-HPLC (c); $\cdot\text{OH}$ scavenging activity of fractions D1–D12 (d); fraction *D3* was separated by analytical RP-HPLC (e); $\cdot\text{OH}$ scavenging activity of fractions D3a to D3e (f); small letters (g–j) on the bars meant significant difference ($P < 0.05$).

concentration of pepsin was 2 g/100 g protein, pH was set at 2.0, and the hydrolysis time and temperature were 4.0 h and 37 °C, respectively. Consequently, the DH was $31.90\% \pm 2.72\%$, which was lower than that of palm kernel globulin (59%) digested by pepsin and trypsin.¹² PKE glutelin is an acid-soluble protein and difficult to dissolve at neutral pH, which is suitable for trypsin. Thus, trypsin was not chosen in this study.

3.2 Gel chromatography of PKEG-1H and antioxidant activity of the fractions

3.2.1 Separation by ultrafiltration and Sephadex G-15 gel chromatography. Hydroxyl radical is accepted as the indicator for purification of antioxidant peptides because it is one of the most reactive radicals and can attack almost every living cell.⁷ Among the three fractions obtained after ultrafiltration, PKEG-1H-III with the lowest molecular weight (M_w) exhibited the highest $\cdot\text{OH}$ scavenging activity (IC_{50} : $88.82 \pm 5.15 \mu\text{g mL}^{-1}$), hence it was chosen for gel chromatography analysis. As shown in Fig. 1a, four fractions (fraction *A*, *B*, *C* and *D*) were collected

separately after Sephadex G-15 gel chromatography. In gel chromatography, the separation is based on the M_w , that is, peptides with large M_w are eluted earlier, while those with lower M_w are eluted later. Thus, the order of the M_w of the four peaks was $A > B > C > D$. Fraction *D* showed the highest $\cdot\text{OH}$ scavenging activity (IC_{50} : $61.43 \pm 4.97 \mu\text{g mL}^{-1}$, Fig. 1b) among the fractions, which was in accordance with the result shown in Fig. 2a. The results confirmed the report that short peptides exhibited stronger antioxidant activity than larger polypeptides.¹⁹

3.2.2 Antioxidant activity of fractions from Sephadex G-15 gel chromatography. Superoxide anion radical is a potential precursor of stronger reactive oxidative species such as hydroxyl radical.⁶ As shown in Fig. 2b, the superoxide radical scavenging activities of PKEG-1H fractions (*A*–*D*) were all dependent on the concentration used. The fraction *D* showed a higher activity (IC_{50} : 0.486 mg mL^{-1}) than fractions *A* and *B* ($P < 0.05$), possibly due to the relatively lower M_w .⁵ The activity of fraction *D* was also higher than that of loach protein hydrolysate (IC_{50} : 0.74 mg mL^{-1}),⁶ indicating that fraction *D* could contain peptides that are more






Fig. 2 Antioxidant activity of fractions A–D including scavenging activity on hydroxyl radical (a), superoxide radical (b), ABTS radical (c) and chelating ability (d); GSH (glutathione) was used for comparison; small letters (g–k) on the bars or lines match results that are significantly different ($P < 0.05$).

easily accessible to the superoxide radical and allows these peptides to trap the radicals more easily. The ABTS radical scavenging activity was widely employed to estimate the total antioxidant activity of potential antioxidants.¹⁸ The results in Fig. 2c showed that all of the fractions exhibited considerable ABTS radical-scavenging activity at a concentration of 0.1–0.5 mg mL^{-1} , indicating their considerable total antioxidant activity. Fraction D also showed remarkably higher activity than fractions A and B ($P < 0.05$). Moreover, as shown in Fig. 2d, all the fractions except fraction B exhibited significantly higher chelating capacity

than GSH ($P < 0.05$). The chelating ability of fraction D could reach up to $93.27\% \pm 4.52\%$, which was much higher than that of sweet potato protein hydrolysates (17.62%)² and sliver carp hydrolysates (18.20%).⁵ A previous study had found that PKE glutelin-1 was rich in Glu and Asp,¹⁰ which could interact with metal ions through the charged groups and inactivate the prooxidant activity of metal ions.²⁰ In general, fraction D was selected to further purification and characterization for its high antioxidant activity.




Fig. 3 Mass spectra of the RP-HPLC chromatograms fraction D3c and D3e by LC-MS/MS.



Table 3 Antioxidant peptides identified in palm kernel expeller glutelin-1 by LC-MS/MS and their stability against the simulated gastrointestinal digestion^c

Fractions	Peptide	MW (Da)	Matched sequence in <i>Elaeis guineensis</i> ^a	Hydrophobic residues content	IC ₅₀ (μg mL ⁻¹)	IC ₅₀ ^b (μg mL ⁻¹)
D3c	TVFDGELR	935.47	K.TVFDGELR.Q glutelin-1	37.50%	38.22 ± 2.22 ^c	20.44 ± 1.35 ^d
	Thr-Val-Phe-Asp-Gly-Glu-Leu-Arg					
	ADVFNPR	817.41	R.ADVFNPR.G glutelin	57.14%	22.16 ± 1.22 ^c	21.65 ± 2.04 ^c
D3e	Ala-Asp-Val-Phe-Asn-Pro-Arg					
	CAGVSAIR	832.42	R.CAGVSAIR.R glutelin	50.00%	31.19 ± 1.67 ^c	32.04 ± 0.72 ^c
	Cys-Ala-Gly-Val-Ser-Ala-Ile-Arg					
	LVYIIQGR	819.37	R.LVYIIQGR.G glutelin-1	62.50%	12.85 ± 0.23 ^c	9.76 ± 0.88 ^d
	Leu-Val-Tyr-Ile-Ile-Gln-Gly-Arg					

^a From National Center for Biotechnology Information (NCBI). ^b Peptides were treated with pepsin and pancreatin. ^c ^d Different small letters in the same row indicates significant difference ($P < 0.05$).

3.3 Purification of the antioxidant peptides

Fraction D from Sephadex G-15 gel chromatography was further isolated by RP-HPLC on the semi-preparative C₁₈ column, and 12 major peaks named as D1–D12 were presented and collected (Fig. 1c). D3 with the lowest IC₅₀ ($51.57 \pm 1.26 \mu\text{g mL}^{-1}$, Fig. 1d) was further purified by RP-HPLC in an analytical C₁₈ column (Φ 4.6 mm \times 250 mm) and five major fractions (D3a–D3e) were obtained (Fig. 1e). The fraction D3c and D3e demonstrated significantly lower IC₅₀ values (16.22 ± 0.64 and $15.00 \pm 0.03 \mu\text{g mL}^{-1}$, respectively, Fig. 1f) than other fractions ($P < 0.05$), and they were subjected to ESI-MS/MS analysis. In RP-HPLC, hydrophilic peptides were eluted earlier while the non-polar or hydrophobic peptides eluted much later.⁴ Moreover, it was evident that high content of hydrophobic amino acids could increase the antioxidant activity of peptides.²¹ The fraction D3e with high hydrophobicity exhibited the highest antioxidant activity in this study, which is in accordance with the previous reports.

3.4 Characterization of purified peptides

As shown in Fig. 3, four peptides were identified by LC-MS/MS, of which TVFDGELR (935.5 Da) and ADVFNPR (818.7 Da) was from fraction D3c, while CAGVSAIR (832.4 Da) and LVYIIQGR (819.4 Da) were from fraction D3e. The four peptides were all short peptides with 7–8 amino acids and exhibited relatively high antioxidant activity (IC₅₀: 38.22 – $12.85 \mu\text{g mL}^{-1}$, Table 3), which was in accordance to the report that antioxidant activity of peptides was inversely correlated with M_w .¹⁹ However, both ADVFNPR and LVYIIQGR exhibited higher antioxidant activity than the smaller peptides from rice residue (464.2 Da, IC₅₀: $43.55 \mu\text{g mL}^{-1}$),²² tilapia protein (317.3 Da, IC₅₀: $26.4 \mu\text{g mL}^{-1}$)⁶ and walnut protein (423.23 Da, IC₅₀: $310.0 \mu\text{g mL}^{-1}$),³ indicating that antioxidant activity of peptides was not dependent only on M_w . Moreover, Tapal *et al.*¹² also obtained the peptide ADVFNPR from palm kernel globulin and found that it had antihypertensive and anticancer activity.

It was demonstrated that antioxidant activity of peptides is positively correlated with the hydrophobic property.²¹ The order of hydrophobic amino acid content for the four peptides was LVYIIQGR > ADVFNPR > CAGVSAIR > TVFDGELR (Table 3). With higher hydrophobicity, LVYIIQGR showed a higher

antioxidant activity (IC₅₀: $12.85 \mu\text{g mL}^{-1}$) than TVFDGELR and YPGSIDWR (IC₅₀: 31.19 – $38.22 \mu\text{g mL}^{-1}$).

Moreover, it has been reported that the amino acid constituents and sequences play an important role in the antioxidant properties of peptides.²² Some amino acids such as Leu, Cys, His, Tyr, Met and Trp contributed significantly to the antioxidant activities of peptides. Leu could increase the presence of the peptides at the water–lipid interface and therefore facilitate access to scavenge free radicals generated at the lipid phase.²³ Tyr, Trp and Phe possess phenolic groups that have special capability of to serve as hydrogen donors; Met tends to undergo oxidation to form Met sulphoxide and Cys donates the sulphur hydrogen; the imidazole group in His also exhibits proton-donation ability.^{5,24} Hence, the high antioxidant activity of the four peptides identified in the current study was attributed to the presence of Leu, Cys, Phe, Tyr, Phe and Trp (Table 3). In addition, the branched amino acids (Leu, Val and Ile) in LVYIIQGR also contribute to the antioxidant activity. Other researchers have also found that peptides containing branched amino acid residues (HPLL, LLHH and HHPLL) had good antioxidant activity.^{3,4}

3.5 Stability of synthetic peptides

As shown in Table 3, after digestion with pepsin and pancreatin, the antioxidant activity of synthetic peptides, CAGVSAIR and ADVFNPR, was not significantly decreased, suggesting that they could effectively retain the activity in the gastrointestinal digestion system. In contrast, the IC₅₀ values of TVFDGELR and LVYIIQGR decreased remarkably ($P < 0.05$). Previous investigations reported that pepsin preferentially cleaves the C-terminal to Phe, Leu and Glu, while pancreatin hydrolyses peptide bonds with aromatic side chains (Tyr, Trp and Phe) at N-terminal or C-terminal.²¹ The peptide LVYIIQGR can be split into LVY and IIQGR by pancreatin, while TVFDGELR can be split into TVF and DGELR. The antioxidant activity of split peptide TVF and LVY that contained F or Y at C-terminal was expected to increase after digestion.¹⁹

4. Conclusions

The optimum enzymatic hydrolysis conditions of palm kernel expeller glutelin-1 were as follows: using pepsin at



a concentration of 2 g/100 g, hydrolysis time of 4 h, temperature 37 °C and pH 2.0. After ultrafiltration, Sephadex G-15 gel chromatography and RP-HPLC, four peptides, TVFDGELR (935.5 Da), ADVFNPR (818.7 Da), CAGVSAIR (832.4 Da) and LVYIIQGR (819.4 Da), were identified from the PKEG-1H. LVYIIQGR showed the highest antioxidant activity (12.85 $\mu\text{g mL}^{-1}$) and CAGVSAIR and ADVFNPR exhibited good stability in gastrointestinal digestion system, indicating their potential usage in health-foods.

Conflicts of interest

The authors declare that they have no competing interests.

Acknowledgements

This work was supported by the key research project of Hainan province (No. ZDYF2017061) and the project supported by the National Natural Science Foundation of China (31771858).

References

- 1 B. J. Bhuyan and G. Mughesh, Synthesis, characterization and antioxidant activity of angiotensin I converting enzyme inhibitors, *Org. Biomol. Chem.*, 2011, **9**, 1356–1365.
- 2 S. Y. Dong, M. Y. Zeng, D. F. Wang, Z. Y. Liu, Y. H. Zhao and H. C. Yang, Antioxidant and biochemical properties of protein hydrolysates prepared from Silver carp (*Hypophthalmichthys molitrix*), *Food Chem.*, 2008, **107**, 1483–1493.
- 3 N. Chen, H. M. Yang, Y. Sun, J. Niu and S. Y. Liu, Purification and identification of antioxidant peptides from walnut (*Juglans regia* L.) protein hydrolysates, *Peptides*, 2012, **38**, 344–349.
- 4 L. Wattanasiritham, C. Theerakulkait, S. Wickramasekara, C. S. Maier and J. F. Stevens, Isolation and identification of antioxidant peptides from enzymatically hydrolyzed rice bran protein, *Food Chem.*, 2016, **192**, 156–162.
- 5 M. Zhang, T. H. Mu and M. J. Sun, Purification and identification of antioxidant peptides from sweet potato protein hydrolysates by Alcalase, *J. Funct. Foods*, 2014, **7**, 191–200.
- 6 Y. F. Zhang, X. Duan and Y. L. Zhuang, Purification and characterization of novel antioxidant peptides from enzymatic hydrolysates of tilapia (*Oreochromis niloticus*) skin gelatin, *Peptides*, 2012, **38**, 13–21.
- 7 A. G. P. Samaranayaka and E. C. Y. Li-Chan, Food-derived peptidic antioxidants: a review of their production, assessment, and potential applications, *J. Funct. Foods*, 2011, **3**, 229–254.
- 8 A. R. Alimon, The nutritive value of palm kernel cake for animal feed, *Palm Oil Development*, 2010, **40**, 12–14.
- 9 K. L. Chee, H. K. Ling and M. K. Ayob, Optimization of trypsin-assisted extraction, physicochemical characterization nutritional qualities and functionalities of palm kernel cake protein, *LWT-Food Sci. Technol.*, 2012, **46**, 419–427.
- 10 Y. J. Zheng, Y. Li, Y. L. Zhang, R. G. Zhang, Q. A. Zhang, Y. F. Zhang and S. L. Zhao, Fractionation, Physicochemical Properties, Nutritional Value, Antioxidant Activity and ACE Inhibition of Palm Kernel Expeller Protein, *RSC Adv.*, 2015, **5**, 12613–12623.
- 11 Y. N. Tan, M. K. Ayob and W. A. W. Yaacob, Purification and characterisation of antibacterial peptide-containing compound derived from palm kernel cake, *Food Chem.*, 2013, **136**, 279–284.
- 12 A. Tapal, G. E. Vegarud, A. Sreedhara, P. Hegde, S. Inamdar and P. K. Tikku, In vitro human gastro-intestinal enzyme digestibility of globulin isolate from oil palm (*Elaeis guineensis* var. *tenera*) kernel meal and the bioactivity of the digest, *RSC Adv.*, 2016, **6**, 20219–20229.
- 13 Y. J. Zheng, Y. Li, Y. L. Zhang, X. H. Ruan and R. G. Zhang, Purification, characterization, synthesis, in vitro ACE inhibition and in vivo antihypertensive activity of bioactive peptides derived from oil palm kernel glutelin-2 hydrolysates, *J. Funct. Foods*, 2017, **28**, 48–58.
- 14 K. Kwon, K. H. Park and K. C. Rhee, Fractionation and characterization of proteins from coconut (*Cocos nucifera* L.), *J. Agric. Food Chem.*, 1996, **44**(7), 1741–1745.
- 15 MPOB TOPOMPOB, Production of Palm Kernel Cake 2013, The Official Portal of Malaysian Palm Oil Board Malaysia.
- 16 J. Adler-Nissen, Determination of the degree of hydrolysis of food protein hydrolysates by trinitrobenzenesulfonic acid, *J. Agric. Food Chem.*, 1979, **27**, 1256–1262.
- 17 J. Y. Qiao, H. H. Lin, S. Qian, Q. J. Zheng and W. Xia, Isolation, identification and synthesis of four novel antioxidant peptides from rice residue protein hydrolyzed by multiple proteases, *Food Chem.*, 2015, **179**, 290–295.
- 18 R. Joshi, S. Sood, P. Dogra and M. Mehendru, In vitro cytotoxicity, antimicrobial, and metal-chelating activity of triterpene saponins from tea seed grown in Kangra valley, India, *Med. Chem. Res.*, 2015, **22**, 4030–4038.
- 19 T. Tavares, M. D. Contreras, M. Amorim, M. Pintado, I. Recio and F. X. Malcata, Novel whey-derived peptides with inhibitory effect against angiotensin-converting enzyme: in vitro effect and stability to gastrointestinal enzymes, *Peptides*, 2011, **32**, 1013–1019.
- 20 F. Shahidi and Y. Zhong, Novel antioxidants in food quality preservation and health promotion, *Eur. J. Lipid Sci. Technol.*, 2010, **112**, 930–940.
- 21 B. H. Sarmadi and A. Ismail, Antioxidative peptides from food proteins: a review, *Peptides*, 2010, **31**, 1949–1956.
- 22 Q. J. Yan, L. H. Huang, Q. Sun, Z. Q. Jiang and X. Wu, Isolation, identification and synthesis of four novel antioxidant peptides from rice residue protein hydrolyzed by multiple proteases, *Food Chem.*, 2015, **179**, 290–295.
- 23 A. Alemán, B. Giménez, E. Pérez-Santin, M. C. Gómez-Guillén and P. Montero, Contribution of Leu and Hyp residues to antioxidant and ACE-inhibitory activities of peptide sequences isolated from squid gelatin hydrolysate, *Food Chem.*, 2011, **125**, 334–341.
- 24 H. C. Wu, H. M. Chen and C. Y. Shiau, Free amino acids and peptides as related to antioxidant properties in protein hydrolysates of mackerel (*Scomber austriasicus*), *Food Res. Int.*, 2003, **36**, 949–957.

