

Cite this: *Chem. Sci.*, 2017, 8, 8193Ruthenium(II)-catalyzed olefination *via* carbonyl reductive cross-coupling†Wei Wei, ^{‡ab} Xi-Jie Dai, ^{‡a} Haining Wang, ^a Chenchen Li, ^a Xiaobo Yang ^a and Chao-Jun Li ^{*a}

Natural availability of carbonyl groups offers reductive carbonyl coupling tremendous synthetic potential for efficient olefin synthesis, yet the catalytic carbonyl cross-coupling remains largely elusive. We report herein such a reaction, mediated by hydrazine under ruthenium(II) catalysis. This method enables facile and selective cross-couplings of two unsymmetrical carbonyl compounds in either an intermolecular or intramolecular fashion. Moreover, this chemistry accommodates a variety of substrates, proceeds under mild reaction conditions with good functional group tolerance, and generates stoichiometric benign byproducts. Importantly, the coexistence of KO^tBu and bidentate phosphine dmpe is vital to this transformation.

Received 26th September 2017
Accepted 8th October 2017

DOI: 10.1039/c7sc04207h


rsc.li/chemical-science

Efficient construction of carbon–carbon double bonds has long been a central pursuit in the synthetic community. Recent decades have witnessed impressive accomplishments in the field of carbonyl olefination.^{1–6} Well-known milestones include, among others, the Wittig reaction,⁴ the Peterson reaction,⁵ the Julia olefination⁶ and the Tebbe–Petasis olefination.⁷ Parallel to these classical olefination methods, the McMurry reaction mediated by low-valent titanium (LVT) reagents enables direct reductive homo-couplings of carbonyl compounds for facile synthesis of olefins (Scheme 1, eqn (a)).⁸ Mechanistically, the


synergy between oxophilic titanium(III)/(IV) and strong metal reductants (*e.g.* LiAlH₄ and alkali metals) is crucial to form metal pinacولات as key intermediates. One problematic scenario pertaining to the original protocol, however, is that the cross-coupling of two unsymmetrical carbonyl compounds generally affords a statistic mixture of coupling products.⁸ Despite McMurry-type variants (*e.g.* external ligands and auxiliaries) having been developed to bypass this issue,⁹ two challenges endure: (1) stoichiometric quantities of metal wastes accompanied by the excessive usage of metal-based reagents, and (2) poor chemoselectivity and functional group tolerance stemmed from the presence of strong metal reductants. As the synthetic community calls for more sustainable and efficient chemical syntheses, carbonyl cross-coupling represents an ideal strategy to access olefins because naturally prevailing carbonyl functionalities are generally regarded as renewable feedstocks.

Very recently, we have disclosed a ruthenium-catalyzed deoxygenation reaction for the highly selective cleavage of aliphatic primary C–O bonds.¹⁰ Building on the similar ruthenium(II) catalysis, we further demonstrated its robustness in catalyzing a series of new carbon–carbon bond forming processes through addition reactions to various carbonyl compounds, imines and activated alkenes.^{11–13} Variations of these precedents notwithstanding, one of their commonalities is to use aldehydes/ketones as alkyl carbanion equivalents *via* hydrazone formation. As a continuation of our interests in utilizing such carbanion equivalents for useful synthetic transformations, we describe herein the development of a ruthenium(II)-catalyzed, hydrazine-mediated olefination reaction *via* carbonyl reductive cross-coupling (Scheme 1, eqn (b)). This catalytic method features good functional group tolerance and generates nitrogen and water as the only environmentally benign byproducts in stoichiometric quantities.^{14,15}

McMurry olefination: LVT-mediated carbonyl homo-coupling



Our approach: Ru(II)-catalyzed carbonyl cross-coupling

Scheme 1 Olefination methods *via* reductive carbonyl coupling.

^aDepartment of Chemistry, FQRNT Center for Green Chemistry and Catalysis, McGill University, 801 Sherbrooke St. W., Montreal, Quebec H3A 0B8, Canada. E-mail: cj.li@mcgill.ca

^bSchool of Chemistry and Chemical Engineering, Qufu Normal University, Qufu 273165, Shandong, China

† Electronic supplementary information (ESI) available: Experimental details. See DOI: 10.1039/c7sc04207h

‡ These authors contributed equally to this work.

Initially, propionaldehyde **1a** and benzophenone **2a** were used as carbonyl partners in our investigation. By varying the reaction conditions employed in carbonyl addition,¹¹ we detected the corresponding olefin **3a** in 51% yield from the cross-coupling between preformed hydrazone of **1a** and **2a** after 12 h (Table 1, entry 1). In line with our basicity rationale, a variety of bases were prioritized in the optimization process. In fact, improved yields were consistently observed using stronger ionic bases such as hydroxides and alkoxides, among which KO^tBu gave the best result (Table 1, entries 1–5; Table S3, ESI†). In contrast, organic base (e.g. Et₃N) and weaker inorganic base (e.g. K₂CO₃) were significantly inferior (Table 1, entries 6 and 7). Moreover, the amount of base matters significantly for this reaction. Specifically, lower loadings than sub-stoichiometric quantity (i.e. 50 mol%) were associated with yield attenuation (Table S4, ESI†). As expected, the olefination reaction did not proceed in the absence of base (Table 1, entry 8). In addition to the basicity, both ruthenium(II) pre-catalyst and bidentate phosphine ligand 1,2-bis(dimethylphosphino) ethane (dmpe) are essential for the elimination to occur. Likewise, no desired product was obtained without either one of them (Table 1, entries 9 and 10). Subsequent screenings on various ruthenium catalysts (Table S1, ESI†) and phosphine ligands (Table 1, entries 11–14; Table S2, ESI†)¹⁶ revealed that the combination of [Ru(*p*-cymene)Cl₂]₂ and dmpe provided the biggest catalytic turnover number, and thus the highest yield. Although other reaction parameters such as solvent, additive and temperature play minor roles in the current reaction, the following observations are worthy noting. For example, ether

solvents (e.g. THF, 1,4-dioxane, 1,2-dimethoxyethane) favor the olefination over other types of solvents (Table S5, ESI†). Addition of cesium fluoride as an additive slightly increases the reaction yield (Table S6, ESI†).

With the standard reaction conditions in hand, we moved on to study the scope of electrophilic carbonyl partners (Table 2). In general, this chemistry covers a broad spectrum of benzophenone and its derivatives, regardless of their electronic nature. Consequently, the corresponding olefins (**3a–h**) were obtained in moderate to good yields. For unsymmetrical benzophenones, a mixture of stereoisomers (*E/Z* isomers) was produced (**3e–h**). Similarly, acetophenones bearing a wide range of aryl

Table 2 Scope of electrophilic carbonyl coupling partners^{a,b}




Table 1 Optimization of reaction conditions^a

Entry	L	Base	Yield ^b (%)
1	dmpe	K ₃ PO ₄	51
2	dmpe	KO ^t Bu	84
3	dmpe	NaO ^t Bu	80
4	dmpe	CSOH	82
5	dmpe	KOH	78
6	dmpe	Et ₃ N	0
7	dmpe	K ₂ CO ₃	4
8	dmpe	—	0
9	—	KO ^t Bu	0
10 ^c	dmpe	KO ^t Bu	0
11	dppb	KO ^t Bu	11
12	dppp	KO ^t Bu	15
13	dppm	KO ^t Bu	4
14	P(<i>p</i> -Tolyl) ₃	KO ^t Bu	4

^a **1a** (0.28 mmol, 1.4 equiv.), N₂H₄·H₂O (0.3 mmol, 1.5 equiv.), THF (0.14 mL), rt, 30 min; **2a** (0.20 mmol, 1.0 equiv.), [Ru(*p*-cymene)Cl₂]₂ (0.003 mmol, 1.5 mol%), ligand (0.006 mmol, 3.0 mol%), base (0.1 mmol, 50 mol%), additive: CsF (0.15 mmol, 75 mol%), 45 °C, 12 h, under N₂. ^b Yields were determined by crude ¹H NMR using mesitylene as an internal standard. ^c Without [Ru(*p*-cymene)Cl₂]₂.

^a **1a** (0.28 mmol, 1.4 equiv.), N₂H₄·H₂O (0.3 mmol, 1.5 equiv.), THF (0.14 mL), rt, 30 min; **2** (0.20 mmol, 1.0 equiv.), [Ru(*p*-cymene)Cl₂]₂ (0.003 mmol, 1.5 mol%), dmpe (0.006 mmol, 3.0 mol%), KO^tBu (0.1 mmol, 50 mol%), CsF (0.15 mmol, 75 mol%), 45 °C, 12 h, under N₂. ^b Isolated yields and the ratio of *E/Z* isomers were determined by crude ¹H NMR analysis. ^c 80 °C. ^d K₃PO₄ (0.1 mmol, 50 mol%), 12 h. ^e Yields were determined by crude ¹H NMR using mesitylene as an internal standard. Yields of asymmetric azines were given in the parentheses.

substituents were surveyed, with the formation of *E/Z* isomers (*E*-isomer predominant) in moderate to good yields (**3i–ip**). Notably, functional groups that are commonly incompatible with traditional carbonyl olefination approach such as unprotected alcohol, ester and amide groups, were well tolerated in this chemistry and thus amenable to further functionalization (**3ij**, **3in** and **3ip**). Intriguingly, aromatic ketones substituted by linear alkyl chains (2–5 carbons) did not hamper the reaction leading to the desired products in good yields (**3j–l**). Aromatic aldehyde and aliphatic ketones can also serve as coupling partners, including the relatively complex oxo-steroid compound. Asymmetric azines were observed as major products with cyclic aliphatic ketones (**3o**). However, the complex reaction mixture was resulted in the case of acyclic aliphatic ketone (**3n**), whereby the corresponding asymmetric azine was obtained as one of the major side products. Low yields were therefore seen in all these substrates at this stage.


Subsequently, the scope of nucleophilic carbonyl coupling partners was tested (Table 3). Linear saturated aliphatic aldehydes, irrespective of carbon chain length, all delivered the corresponding olefins in moderate to good yields (**4a–f**). Although aromatic aldehydes were poorly coupled with aromatic ketones (*i.e.* The reductive coupling of benzaldehyde

with benzophenone leading to product **4g**) due to competing alcohol formation, their couplings with other aromatic aldehydes typically proceeded well to yield various stilbenes with excellent stereoselectivity (**4h–k**). In the latter cases, using K_2CO_3 as the base at elevated temperature was necessary. Encouragingly, more sterically demanding aromatic ketones (*i.e.* acetophenone) worked as nucleophilic partner, providing **4l** in moderate yield with high stereoselectivity. Finally, the intramolecular olefination was evaluated with 3-(2-benzoyl-phenyl)propanal **1m** and 6-oxo-6-phenylhexanal **1n**. The corresponding cyclohexene derivatives **4m** and **4n** were obtained in 51% and 48% yields, respectively.

Mechanistically, two scenarios are possible to form olefins. One is metal-assisted decomposition of the corresponding asymmetric azine. The other is base-mediated elimination of the corresponding alcohol. In other words, both asymmetric azines and alcohols might have been generated prior to olefins. To exclude these possibility, two parallel control experiments were conducted with the presynthesized azine **6i** and 1,1-diphenylbutan-1-ol **5a** under standard conditions, respectively (Scheme 2). However, olefin products (**3i** and **3a**) were not detected in both cases. The above results strongly suggested that olefins were eliminated from a transient intermediate, rather than azines or alcohols, *via* a E1cB-type mechanism.


Based on all experimental data, the preliminary computational calculations done on the previous carbonyl addition chemistry¹¹ and studies from others,¹⁴ a postulated reaction pathway was proposed as shown in Scheme 3. The bidentate phosphine coordinated complex **I** is initially generated by a ligand dissociation/association between $[Ru(p\text{-cymene})Cl_2]_2$ and dmpe, followed by a ligand association with carbonyl-derived hydrazone **B** and carbonyl **C** in the presence of KO^tBu , giving rise to complex **II** and **III**, respectively. Formation of the coordinately intermediate **III** sets the stage for the intramolecular isomerization. This concerted process yields the key six-membered ring intermediate **IV** by forming a new carbon–carbon bond between **A** and **C**.¹⁴ Base-catalyzed decomposition of diimide intermediate **IV** *via* a E1cB-type mechanism produces the desired olefins along with the formation of ruthenium hydroxide species **V**. To turnover the cycle, **V** then reacts with hydrazone **B** to release water and active species **II** ($X = Cl$). Alternatively, the chloride anionic ligand on **V** could have been

Table 3 Scope of nucleophilic carbonyl coupling partners^{a,b}




^a **1** (0.28 mmol, 1.4 equiv.), $N_2H_4 \cdot H_2O$ (0.3 mmol, 1.5 equiv.), THF (0.14 mL), rt, 30 min; **2** (0.20 mmol, 1.0 equiv.), $[Ru(p\text{-cymene})Cl_2]_2$ (0.003 mmol, 1.5 mol%), dmpe (0.006 mmol, 3.0 mol%), KO^tBu (0.1 mmol, 50 mol%), CsF (0.15 mmol, 75 mol%), 45 °C, 12 h, under N_2 . ^b Isolated yields and the ratio of *E/Z* isomers were determined by crude 1H NMR analysis. ^c 60 °C, 24 h. ^d $[(C_6Me_6)RuCl_2]_2$ (0.003 mmol, 1.5 mol%) was used as catalyst. ^e K_2CO_3 (0.1 mmol, 50 mol%), 120 °C, 24 h.

Olefin production from asymmetric azine




Olefin production from alcohol



Scheme 2 Control experiments for the olefin formation.





Scheme 3 Possible reaction mechanism.

replaced by **B**, leaving hydroxide on the active species **VI** ($X = OH$) and other ruthenium complexes (**III-V**).

Conclusions

In summary, we have developed a ruthenium(II)-catalyzed, hydrazine-mediated olefination method *via* reductive carbonyl coupling reactions. This chemistry possesses a distinct mechanistic profile and highlights the use of naturally abundant carbonyl functionalities for efficient olefin synthesis. Other striking features include cross-coupling capability, mild reaction conditions, good functional group tolerance and stoichiometric benign byproducts. Taken together, our findings are expected to spur more interest in developing catalytic methods in this field. Further investigations on increasing the reaction scope, synthetic applications and mechanistic details are undergoing in our laboratory.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was financially supported by the Canada Research Chair (Tier 1) foundation, FQRNT (CCVC), NSERC, CFI and McGill University. We also thank the CSC (China Scholarship Council) for the postdoctoral fellowship (W. Wei).

References

- R. Dumeunier and I. E. Markó, in *Modern Carbonyl Olefination*, ed. T. Takeda, Wiley-VCH, Weinheim, Germany, 2004.
- 2 (a) A. H. Hoveyda and A. R. Zhugralin, *Nature*, 2007, **450**, 243–251; (b) X. Guo, J. Wang and C.-J. Li, *J. Am. Chem. Soc.*, 2009, **131**, 15092–15093; (c) S. Takemoto, E. Shibata, M. Nakajima, Y. Yumoto, M. Shimamoto and H. Matsuzaka, *J. Am. Chem. Soc.*, 2016, **138**, 14836–14839; (d) J. R. Ludwig, P. M. Zimmerman, J. B. Gianino and C. S. Schindler, *Nature*, 2016, **533**, 374–379.
- 3 (a) W. S. Wadsworth and W. D. Emmons Jr, *J. Am. Chem. Soc.*, 1961, **83**, 1733–1738; (b) C. R. Johnson, J. R. Shanklin and R. A. Kirchhoff, *J. Am. Chem. Soc.*, 1973, **95**, 6462–6463; (c) S. Sano, T. Takehisa, S. Ogawa, K. Yokoyama and Y. Nagao, *Chem. Pharm. Bull.*, 2002, **50**, 1300–1302; (d) Z.-Y. Peng, F.-F. Ma, L.-F. Zhu, X.-M. Xie and Z. Zhang, *J. Org. Chem.*, 2009, **74**, 6855–6858.
- 4 (a) G. Wittig and G. Geissler, *Liebigs Ann. Chem.*, 1953, **580**, 44–57; (b) G. Wittig and U. Schollkopf, *Ber. Dtsch. Chem. Ges.*, 1954, **87**, 1318–1330; (c) J. Boutagy and R. Thomas, *Chem. Rev.*, 1974, **74**, 87–99; (d) B. E. Maryanoff and A. B. Reitz, *Chem. Rev.*, 1989, **89**, 863–927.
- 5 (a) D. J. Peterson, *J. Org. Chem.*, 1968, **33**, 780–784; (b) J. Huang, C. Wu and W. D. Wulff, *J. Am. Chem. Soc.*, 2007, **129**, 13366–13367; (c) D. J. Ager, The Peterson Olefination Reaction, *Organic Reactions*, ed. L. A. Paquette, 1990, **38**, pp. 1–224.
- 6 (a) M. Julia and J. M. Paris, *Tetrahedron Lett.*, 1973, **14**, 4833–4836; (b) P. R. Blakemore, *J. Chem. Soc., Perkin Trans. 1*, 2002, 2563–2585; (c) K. Ando, T. Kobayashi and N. Uchida, *Org. Lett.*, 2015, **17**, 2554–2557.
- 7 (a) F. N. Tebbe, G. W. Parshall and G. S. Reddy, *J. Am. Chem. Soc.*, 1978, **100**, 3611–3613; (b) I. Beadham and J. Micklefield, *Curr. Org. Synth.*, 2005, **2**, 231–250; (c) N. A. Petasis and E. I. Bzowej, *J. Am. Chem. Soc.*, 1990, **112**, 6392–6394.
- 8 (a) J. E. McMurry and M. P. Fleming, *J. Am. Chem. Soc.*, 1974, **96**, 4708–4709; (b) J. E. McMurry and L. R. Krepski, *J. Org. Chem.*, 1976, **41**, 896–897; (c) S. Tyrlik and I. Wolochowicz, *Bull. Soc. Chim. Fr.*, 1973, 2147–2148; (d) B. E. Kahn and R. D. Rieke, *Chem. Rev.*, 1988, **88**, 733–745; (e) J. E. McMurry, *Chem. Rev.*, 1989, **89**, 1513–1524; (f) T. Takeda and A. Tsubouchi, The McMurry Coupling and Related Reactions, *Organic Reactions*, ed. S. E. Denmark, 2013, **82**, pp. 1–470.
- 9 (a) N. Balu, S. K. Nayak and A. Banerji, *J. Am. Chem. Soc.*, 1996, **118**, 5932–5937; (b) C. Villiers and M. Ephritikhine, *Chem.-Eur. J.*, 2001, **7**, 3043–3051; (c) M. H. Chisholm and J. A. Klang, *J. Am. Chem. Soc.*, 1989, **111**, 2324–2325; (d) X.-F. Duan, J. Zeng, J.-W. Lü and Z.-B. Zhang, *J. Org. Chem.*, 2006, **71**, 9873–9876; (e) H. R. Diéguez, A. López, V. Domingo, J. F. Arteaga, J. A. Dobado, M. Mar Herrador, J. F. Quilez del Moral and A. F. Barrero, *J. Am. Chem. Soc.*, 2010, **132**, 254–259; (f) L. Zhang, X. Yu, L. Zhang, X. Zhou and Y. Lin, *Org. Chem. Front.*, 2014, **1**, 929–935.
- 10 (a) X.-J. Dai and C.-J. Li, *J. Am. Chem. Soc.*, 2016, **138**, 5433–5440; for an iridium-catalyzed version, see: (b) J.-L. Huang, X.-J. Dai and C.-J. Li, *Eur. J. Org. Chem.*, 2013, 6496–6500.

- 11 H. Wang, X.-J. Dai and C.-J. Li, *Nat. Chem.*, 2016, **9**, 374–378.
- 12 N. Chen, X.-J. Dai, H. Wang and C.-J. Li, *Angew. Chem., Int. Ed.*, 2017, **56**, 6260–6263.
- 13 X.-J. Dai, H. Wang and C.-J. Li, *Angew. Chem., Int. Ed.*, 2017, **56**, 6302–6306.
- 14 A few examples for the formation of carbon-carbon double bond *via* the coupling of stoichiometric metallocenes with carbonyl compounds, see: (a) G. M. Arvanitis, J. Schwartz and D. Van Engen, *Organometallics*, 1986, **5**, 2157–2159; (b) J. A. Smegal, I. K. Meier and J. Schwartz, *J. Am. Chem. Soc.*, 1986, **108**, 1322–1323; (c) G. M. Arvanitis and J. Schwartz, *Organometallics*, 1987, **6**, 421–423; (d) G. M. Arvanitis, J. Smegal, I. Meier, A. C. C. Wong, J. Schwartz and D. Van Engen, *Organometallics*, 1989, **8**, 2717–2723; for a pioneering work on using alkylhydrazone anion for olefination with carbonyls *via* a multistep process mediated by PCl_3 , see (e) J. E. Baldwin, R. M. Adlington, J. C. Bottard, J. N. Kolhe, I. M. Newington and M. W. D. Perry, *Tetrahedron*, 1986, **42**, 4235–4246.
- 15 A recent review on ruthenium-catalyzed transfer hydrogenation for C–C bond formation, see: F. Perez, S. Oda, L. M. Geary and M. J. Krische, *Top. Curr. Chem.*, 2016, **374**, 365–387.
- 16 For the reactions of $[\text{RuCl}_2(\eta^6\text{-arene})]_2$ with tertiary phosphines, please see. (a) R. A. Zelonka and M. C. Baird, *Can. J. Chem.*, 1972, **50**, 3063–3072; (b) M. A. Bennett and A. K. Smith, *J. Chem. Soc., Dalton Trans.*, 1974, 233–241; (c) H. Werner and R. Werner, *Chem. Ber.*, 1982, **115**, 3766–3780; (d) M. A. Bennett, T.-N. Huang and J. L. Latten, *J. Organomet. Chem.*, 1984, **272**, 189–205; (e) M. A. Bennett and J. L. Latten, *Aust. J. Chem.*, 1987, **40**, 841–849; (f) P. Pertici, S. Bertozzi, R. Lazzaroni, G. Vitulli and M. A. Bennett, *J. Organomet. Chem.*, 1988, **354**, 117–121.

