RSC Advances

PAPER

Cite this: RSC Adv., 2018, 8, 2759

One-pot three component synthesis of 5-allyl-1,2,3-triazoles using copper(I) acetylidest

Parigi Raghavendar Reddy, Lianji C[u](http://orcid.org/0000-0003-0802-6398)i and Jae-Sang Ryu ¹

One-pot three-component reactions using copper(I) acetylide, azide, allyl iodide, and NaOH have been developed. The reactions proceed smoothly at room temperature to afford 5-allyl-1,2,3-triazoles, which can be further transformed into a variety of 1,2,3-triazole-fused bi-/tricyclic scaffolds. This method offers the most efficient, convenient, and practical route towards useful polycyclic scaffolds in moderate to excellent yields.

Received 29th November 2017 Accepted 3rd January 2018

DOI: 10.1039/c7ra12889d

rsc.li/rsc-advances

Introduction

1,2,3-Triazoles¹ are very important heterocycles in chemistry and biology.² Synthetic molecules containing a 1,2,3-triazole scaffold exhibit diverse biological activities, which have drawn the attention of medicinal chemists in the drug discovery field. Currently, there are several 1,2,3-triazole-containing medicines on the market, and the number of potential pharmaceuticals based on these scaffolds keeps increasing.³ Beyond the drug market, 1,2,3-triazoles are utilized in a variety of areas including bioconjugation,⁴ polymer and materials science,⁵ and related areas⁶ including supramolecular chemistry,⁷ DNA labeling⁸ and oligonucleotide synthesis.⁹ Such wide applications of 1,2,3-triazoles are due to their facile synthesis through Cu(I)-catalyzed azide–alkyne cycloaddition (CuAAC), so-called 'click chemistry'. Since it was first introduced by Sharpless and co-workers, CuAAC has been rapidly adopted as a universal coupling process. However, in terms of substrate scope, CuAAC is restricted to terminal alkynes, leading to 1,4-disubstituted 1,2,3 triazoles, and the one-pot synthetic methods for the fully substituted 1,2,3-triazoles are still relatively few.¹⁰ PAPER
 CALCONG CONSULS CALCONG CONSULS CALCO

In one-pot three-component reactions used to obtain 5-halo 1,2,3-triazoles, an electrophile X^+ (X = Cl, Br, and I) is added into the CuAAC reaction to trap X^+ with an *in situ* generated 5copper(1) 1,2,3-triazole intermediate A (Scheme 1).¹¹ However, the result is usually the formation of a mixture of the desired 5 substituted 1,2,3-triazole 3 and the byproduct 1,4-disubstituted 1,2,3-triazole 4, which is generated from a competitive protonation of the 5-copper (i) 1,2,3-triazole intermediate A^{12} This competitive protonation is accelerated by a proton source, provided from the terminal alkyne substrate in the normal CuAAC. Therefore, this problem cannot be avoided in the presence of terminal alkyne substrates or protic polar solvents. Recently, this drawback was smartly overcome by using cop $per(i)$ acetylide instead of a terminal alkyne in the halogenation¹² and acylation, by Y. Hu's group.¹³

 $Copper(i)$ acetylides are highly crystalline polymeric complexes, $(RC\equiv CCu)_n$.¹⁴ They are stable to air, water, acid/ base and heat¹⁵ so they can be kept on the shelf for several months without the quality deteriorating. In the absence of exogenous ligands or additives, $copper(i)$ phenylacetylide, $PhC\equiv CCu$, is not effective under typical CuAAC conditions:¹³ It did not undergo a cycloaddition with azide and did not provide a 1,2,3-triazole product. However, when it was combined with both azide and acyl chloride, it quickly provided 5-acyl-1,2,3 triazoles.¹³ The presence of acyl chloride accelerated the cycloaddition of copper (i) phenylacetylide and azide.¹³ This intriguing result motivated us to investigate one-pot threecomponent reactions using copper(I) phenylacetylide. We are particularly interested in 5-allyl-1,2,3-triazoles as precursors for the development of new anticancer agents. However, efficient

Scheme 1 Cu(I)-catalyzed azide-alkyne cycloaddition methods for the synthesis of 5-substituted 1,2,3-triazoles.

College of Pharmacy & Graduate School of Pharmaceutical Sciences, Ewha Womans University, 52 Ewhayeodae-gil, Seodaemun-Gu, Seoul 03760, Republic of Korea. E-mail: ryuj@ewha.ac.kr

[†] Electronic supplementary information (ESI) available: Copies of ¹H and ¹³C NMR spectra for 3a–z and 6v–z. See DOI: 10.1039/c7ra12889d

methods for the synthesis of 5-allyl-1,2,3-triazoles are rare.^{10a,b} Interestingly, $copper(i)$ acetylide has never been used directly as a substrate for the synthesis of 5-allyl-1,2,3-triazoles, although it is known to be a key intermediate. Herein, we report one-pot three-component reactions for the synthesis of 5-allyl-1,2,3 triazoles from copper (I) acetylides.

Results and discussion

To assess the feasibility of the tandem CuAAC–allylation reaction, we started to investigate one-pot three component reactions in the presence of copper (i) phenylacetylide $(1a)$, 4cyanobenzyl azide (2a), allyl iodide, and base (Table 1). In preliminary screening of reaction conditions, we obtained the desired product 5-allyl-1,2,3-triazole 3a and a byproduct enyne 5a with 47% and 42% yield respectively after 4 h at room temperature when 1.5 equivalents of cyanobenzyl azide (2a), 3 equivalents of allyl iodide, and 2 equivalents of $Et₃N$ were employed (entry 1). The yield of the desired product 3a increased to 60% and the yield of the byproduct enyne 5a

Table 1 Optimization of reaction conditions⁴

Entry	Base	Solvent	Time (h)	Yield b (%)		
				3a	4a	5a
1^c	Et ₃ N	CH_2Cl_2	4	47	$\bf{0}$	42
2	Et ₃ N	CH ₂ Cl ₂	4	60	Ω	36
3	Ouinine	CH_2Cl_2	24	74	2	0
$\overline{4}$	Pyridine	CH_2Cl_2	24	63	Ω	$\mathbf{0}$
5	Na ₃ PO ₄	CH_2Cl_2	24	47	Ω	0
6	K_3PO_4	CH_2Cl_2	24	45	0	0
7	Na ₂ CO ₃	CH_2Cl_2	24	62	Ω	$\mathbf{0}$
8	K_2CO_3	CH_2Cl_2	21	71	Ω	$\mathbf{0}$
9	Cs_2CO_3	CH_2Cl_2	24	67	Ω	$\mathbf{0}$
10	NaOH	CH ₂ Cl ₂	24	84	Ω	$\mathbf{0}$
11	KOH	CH_2Cl_2	24	83	Ω	$\mathbf{0}$
12		CH_2Cl_2	24	52	Ω	0
13	NaOH	Dioxane	24	89	Ω	0
14	NaOH	THF	24	85	Ω	$\mathbf{0}$
15	NaOH	Toluene	24	94	$\bf{0}$	$\bf{0}$
16	NaOH	CH ₃ CN	24	68	$\bf{0}$	$\mathbf{0}$

^a Reaction conditions: 1a (65.8 mg, 400 µmol), 2a (94.9 mg, 600 µmol) base (800 µmol), allyl iodide (146 µL, 1.60 mmol), solvent (1 mL). All reactions were carried out under Ar. b Isolated yields. c 1.20 mmol of allyl iodide was used.

decreased to 36% when the amount of allyl iodide increased to 4 equivalents (entry 2). The use of Et_3N appeared to decompose the stable polymeric complex with the structure $[(PhC=CCu)_2]_n$ into a lower polymeric or a more reactive monomeric structure. As soon as $Et₃N$ was added, the heterogeneous reaction mixture became clear and rapidly yielded both an undesired byproduct as well as the desired product. However, replacing $Et₃N$ with pyridine, quinine, or inorganic bases such as $Na₃PO₄$, $K₃PO₄$, Na_2CO_3 , K_2CO_3 , Cs_2CO_3 , NaOH, or KOH did not change the heterogeneity of the reaction. The copper (i) phenylacetylide $(1a)$ remained suspended in $CH₂Cl₂$, which not only slowed reaction progress but also significantly reduced byproduct formation. The reaction using quinine increased the yield of desired product 3a up to 74%, but interestingly 2% of a protonated byproduct 4a was isolated instead of the allyl alkyne byproduct 5a (entry 3). The reaction using pyridine did not signicantly increase the yield compared to the reaction of $Et₃N$. However, the formation of byproducts 4a and 5a was not observed (entry 4). Similarly, the reactions using inorganic bases also generated 5-allyl-1,2,3-triazole 3a as a sole product (entries 5–11). Hydroxide bases (KOH, NaOH: entries 10 and 11) are better than phosphate bases ($Na₃PO₄$, $K₃PO₄$: entries 5 and 6) or carbonate bases (Na₂CO₃, K₂CO₃, Cs₂CO₃: entries 7-9) in terms of yield. Among bases used, the best result was obtained with NaOH (84%, entry 10). It appears the use of base is essential for the reaction. Without a base, 5-allyl-1,2,3-triazole 3a was isolated with only a moderate yield (52%, entry 12). The role of NaOH is not clear, and it is still under investigation. It is also noteworthy that the byproduct 1,4-disubstituted 1,2,3-triazole 4a, which is generated from a competitive protonation of 5-copper(i) 1,2,3triazole intermediate A, was not detected in reactions using $copper(i)$ phenylacetylide $(1a)$ with the exception of quinine. **EXALUATER CONFIDENTIFY ARTIFIED CONSULATER C**

Next, we investigated the solvent effect on the yield of $Cu(I)$ catalyzed azide–alkyne cycloaddition–allylation reactions of $copper(i)$ acetylides in the presence of NaOH. The reactions were effective in various solvents including CH_2Cl_2 , dioxane, THF, toluene, and $CH₃CN$ (entries 13–16). Considering yield of the product, toluene (entry 15) was the best solvent for the reaction. On the basis of Table 1, we chose NaOH as a base and the relatively nonpolar solvent, toluene, for further study.

Once we had established optimal reaction conditions, we examined the scope of one-pot three-component reactions (Table 2). We applied the copper (i) acetylide system to various azides in toluene at room temperature. All reactions using $copper(i)$ acetylides in Table 2 smoothly furnished 5-allyl-1,4disubstituted 1,2,3-triazoles 3b–u and a trace amount of enyne byproduct 5b–u at room temperature. As anticipated, the 1,4 disubstituted 1,2,3-triazoles 4, which are unavoidable in the typical CuAAC reactions, were not detected. First, we explored the reaction scope with benzyl azides. The reaction yields were independent of the electronic nature of azides. Substrates 2c–e bearing electron-donating substituents and substrates 2f–h bearing electron-withdrawing substituents underwent the onepot three-component reactions very smoothly, and provided the corresponding 5-allyl-1,4-disubstituted 1,2,3-triazoles 3c–h at room temperature, in good to excellent yields (85–97%). The reaction scope was not limited to only benzyl azides, but was

 a Reaction conditions: 1 (400 µmol), 2 (600 µmol), NaOH (32.0 mg, 800 mmol), allyl iodide (1.60 mmol), toluene (1 mL). All reactions were carried out under Ar.

also compatible with aryl azides 2i–k and aliphatic azide 2l. In addition, a variety of copper (i) arylacetylides were also tested. Regardless of the electronic influence of substituents on cop $per(i)$ arylacetylides, desired products $3m-p$ were obtained in moderate to good yields (54–75%). It is worth noting that this

Table 3 Synthesis of fused 1,2,3-triazoles using Cu(I)-catalyzed azide–alkyne cycloaddition–allylation reactions a and ring closing

metathesis b </sup>

^a Reaction conditions: 1a (65.8 mg, 400 µmol), 2 (600 µmol), NaOH (32.0 mg, 800 µmol), allyl iodide (146 µL, 1.60 mmol), toluene (1 mL).
All reactions were carried out under Ar. ^b Reaction conditions: 3 (200 m reactions were carried out under A. Reaction conditions. 5 (200 μ mol), Grubbs' 1st generation catalyst (5 mol%), CH₂Cl₂ (9.3 mL). ϵ Isolated yields. ^d 180 μ mol of 3 was used. ϵ 150 μ mol of 3 and Hoveyda-Grubbs' 2nd generation catalyst (5 mol%) were used.

method is not limited to the use of simple allyl iodide. Other various substituted allyl iodides could be used for one-pot threecomponent reactions, and produced the corresponding 5-allyl-1,4-disubstituted 1,2,3-triazoles 3q–t with yields in the range of 53–90%. Delightfully, the reaction scope was also expandable to copper(i) alkylacetylide, which afforded 4-alkyl 1,2,3-triazole 3u.

Finally, we studied further transformations of 5-allyl-1,4 disubstituted 1,2,3-triazoles into various fused polyheterocycles in order to demonstrate the synthetic utility of the 5-allyl-1,4-disubstituted 1,2,3-triazoles as versatile building blocks (Table 3). The alkene-tethered 5-allyl-1,2,3-triazoles 3v–x were synthesized based on Cu(I)-catalyzed azide-alkyne cycloaddition–allylation reactions of copper (i) acetylides in good yields (entries 1–3, 65–80%) and these were converted into fused 1,2,3 triazoles 6v–x by ring-closing metathesis reactions, which were effective for the synthesis of 7-, 8- and 9-membered fused 1,2,3 triazoles with good yields (entries 1–3, 65–87%). The transformations were also applicable to substrates bearing styrenetype alkenes 3y and 3z. The ring-closing metathesis of 3y and 3z proceeded smoothly to afford fused tricyclic 1,2,3-triazoles (entries 4 and 5, 70–92%).

On the basis of previous mechanistic studies¹⁶ and our own observation, we speculate that $Cu(1)$ acetylide 1 coordinates with the azide 2 to form the intermediate I-1 and following cyclization leads to the metallocycle intermediate I-2. The 6-membered $Cu(I)$ -intermediate contracts to the $Cu(I)$ -1,2,3-triazole intermediate I-3, which is readily trapped by allyl iodide to yield the desired 5-allyl-1,4-disubstituted 1,2,3-triazole 3 (Scheme 2).

Scheme 2 Plausible mechanism of Cu(I)-mediated one-pot three component synthesis of 5-allyl-1,2,3-triazoles.

Conclusions

In conclusion, we have developed a novel one-pot three component reaction method for the synthesis of 5-allyl-1,4 disubstituted 1,2,3-triazoles from copper(I) acetylides. The 5 allyl-1,4-disubstituted 1,2,3-triazoles were produced via 1,3 dipolar cycloaddition followed by in situ trapping of the $C(sp^2)$ -Cu intermediate. The byproduct 1,4-disubstituted 1,2,3-triazole 4, which is generated from a competitive protonation of 5 copper(I) 1,2,3-triazole intermediate A, was not isolated. The present method was successfully applied to achieve production of synthetically useful heterocycles. This domino reaction is characterized by mild conditions and no protonated byproduct formation, and allows for the efficient construction of functionalized fused polycyclic 1,2,3-triazoles.

Experimental

General

All reactions were performed in oven-dried glassware fitted with glass stoppers under positive pressure of Ar with magnetic stirring, unless otherwise noted. Air- and moisture-sensitive liquids and solutions were transferred via syringe or stainless-

steel cannula. TLC was performed on 0.25 mm E. Merck silica gel 60 F_{254} plates and visualized under UV light (254 nm) or by staining with cerium ammonium molybdenate (CAM), potassium permanganate (KMnO₄) or p-anisaldehyde. Flash chromatography was performed on E. Merck 230–400 mesh silica gel 60. Reagents were purchased from commercial suppliers, and used without further purification unless otherwise noted. Solvents were distilled from proper drying agents $(CaH₂)$ or Na wire) under Ar atmosphere at 760 mmHg. NMR spectra were recorded at 24 $^{\circ}$ C. Chemical shifts are expressed in ppm relative to TMS (¹H, 0 ppm), CDCl₃ (¹H, 7.26 ppm; ¹³C, 77.2 ppm), and C_6H_5F (¹⁹F, -113.15 ppm); coupling constants are expressed in Hz. High resolution mass spectra (HRMS) were obtained by electrospray ionization (ESI, TOF) or electron ionization (EI, magnetic sector). Infrared spectra were recorded with peaks reported in cm^{-1} . **Solution** (anticles 1-3, 65-60%) and these were converted into forest by the set of α -by the set of α -published on 4/2 January 2018. This are the set of α -published on 4/15/2025 2:59:25 AM. This article is a pla

General procedure for the $Cu(1)$ -catalyzed azide-alkyne cycloaddition–allylation reactions

Copper (i) acetylide (400 µmol) was placed in a 25 mL one-arm roundbottom flask. A solution of azide (600 μ mol) in anhydrous toluene (1 mL), allyl iodide (268 mg, 1.60 mmol) and NaOH (32.0 mg, 800 µmol) were sequentially added to the reaction mixture. The resulting suspension was stirred at room temperature for 24 h or 48 h as indicated in Table 1. The mixture was filtered and washed with CH_2Cl_2 (20 mL). To scavenge Cu, polymer-bound ethylenediaminetriacetic acid acetamide (3.0–4.0 mmol g^{-1} , 200 mg) was added to the filtrate, and stirred for 2 h. The polymer was filtered off and the filtrate was concentrated by rotary evaporation. Purification of crude residue by column chromatography yielded 3a–z. The reaction of 3y was carried out for 3 days at room temperature.

4-((5-Allyl-4-phenyl-1H-1,2,3-triazol-1-yl)methyl)benzonitrile (3a). TLC: R_f 0.25 (2 : 1 hexane/EtOAc). White solid (113 mg, 94%). Mp = 113-115 °C. ¹H NMR (400 MHz, CDCl₃): δ 7.68-7.63 $(m, 4H), 7.45$ -7.41 $(m, 2H), 7.38$ -7.34 $(m, 1H), 7.28$ $(d, J =$ 8.4 Hz, 2H), 5.83 (m, 1H), 5.57 (s, 2H), 5.15 (dd, $J = 10.4$ Hz, 1.2 Hz, 1H), 4.88 (dd, $J = 16.8$ Hz, 1.2 Hz, 1H), 3.45 (m, 2H). ¹³C NMR (100 MHz, CDCl₃): δ 146.2, 140.3, 132.9, 132.1, 131.0, 130.4, 128.9, 128.3, 128.0, 127.3, 118.3, 118.2, 112.6, 51.4, 27.1. HRMS (ESI) m/z calculated for C₁₉H₁₇N₄ [M + H]⁺ 301.1448, found 301.1451. IR (KBr film): ν 3061, 2229, 1639, 1506, 920, 820, 777, 699 cm $^{-1}$.

5-Allyl-1-benzyl-4-phenyl-1H-1,2,3-triazole (3b). TLC: R_f 0.52 $(2:1 \text{ hexane/EtOAc})$. Pale yellow liquid $(97.9 \text{ mg}, 89\%)$. $^1\text{H NMR}$ (400 MHz, CDCl3): d 7.70–7.67 (m, 2H), 7.44–7.40 (m, 2H), 7.36– 7.30 (m, 4H), 7.20–7.18 (m, 2H), 5.83 (m, 1H), 5.54 (s, 2H), 5.15 $(dm, J = 12.0 Hz, 1H), 4.92 (dm, J = 15.2 Hz, 1H), 3.45 (m, 2H).$
¹³C NMR (100 MHz, CDCl₃): δ 146.0, 135.1, 132.4, 131.5, 130.4, 129.1, 128.8, 128.5, 128.0, 127.3, 117.9, 52.1, 27.1. HRMS (ESI) m/z calculated for C₁₈H₁₈N₃ [M + H]⁺ 276.1495, found 276.1501. IR (KBr film): ν 3063, 3033, 1639, 1608, 1496, 918, 800, 764 cm⁻¹

5-Allyl-1-(4-methylbenzyl)-4-phenyl-1H-1,2,3-triazole (3c). TLC: R_f 0.55 (2 : 1 hexane/EtOAc). White solid (97.9 mg, 85%). $Mp = 42-44$ °C. ¹H NMR (400 MHz, CDCl₃): δ 7.69–7.66 (m, 2H), 7.44–7.39 (m, 2H), 7.36–7.31 (m, 1H), 7.14 (d, $J = 8.0$ Hz, 2H), 7.09 (d, $J = 8.0$ Hz, 2H), 5.84 (m, 1H), 5.49 (s, 2H), 5.17 $(dq, J = 10.4 \text{ Hz}, 2.0 \text{ Hz}, 1\text{H}), 4.93 (dq, J = 17.2 \text{ Hz}, 2.0 \text{ Hz}, 1\text{H}),$ 3.45 (m, 2H), 2.33 (s, 3H). ¹³C NMR (100 MHz, CDCl₃): δ 146.0, 138.3, 132.5, 132.1, 131.5, 130.4, 129.8, 128.8, 128.0, 127.3, 117.9, 52.0, 27.2, 21.3. HRMS (ESI) m/z calculated for $C_{19}H_{20}N_3$ [M + H]⁺ 290.1652, found 290.1657. IR (KBr film): ν 3056, 3031, 1639, 1495, 1360, 919, 792, 716 cm^{-1} .

5-Allyl-1-(4-methoxybenzyl)-4-phenyl-1H-1,2,3-triazole (3d). TLC: R_f 0.35 (2 : 1 hexane/EtOAc). Pale yellow liquid (107 mg, 88%). ¹H NMR (400 MHz, CDCl₃): δ 7.68–7.65 (m, 2H), 7.43–7.39 $(m, 2H)$, 7.35–7.31 $(m, 1H)$, 7.15 $(dm, J = 9.6 Hz, 2H)$, 6.86 $(dm, J$ $= 9.6$ Hz, 2H), 5.84 (m, 1H), 5.46 (s, 2H), 5.16 (dq, $J = 10.4$ Hz, 2.0 Hz, 1H), 4.92 (dq, $J = 16.8$ Hz, 2.0 Hz, 1H), 3.78 (s, 3H), 3.46 (m, 2H). ¹³C NMR (100 MHz, CDCl₃): δ 159.7, 146.0, 132.5, 131.5, 130.3, 128.9, 128.8, 128.0, 127.3, 127.1, 117.8, 114.5, 55.4, 51.7, 27.1. HRMS (ESI) m/z calculated for C₁₉H₂₀N₃O [M + H]⁺ 306.1601, found 306.1595. IR (KBr film): ν 3063, 3005, 1639, 1514, 1250, 1032, 921, 825, 715 cm⁻¹ Paper
 $\frac{1}{2}$ Sec. 41, $\frac{1}{2}$ S. 41, $\frac{1}{2}$ S. 41, $\frac{1}{2}$ S. 41, $\frac{1}{2}$ S. 41, $\frac{1}{2}$ A. 41, $\frac{1}{2}$

5-Allyl-1-(3,5-dimethoxybenzyl)-4-phenyl-1H-1,2,3-triazole (3e). TLC: R_f 0.37 (2 : 1 hexane/EtOAc). White solid (126 mg, 94%). Mp $= 80-82 \text{ °C}.$ ¹H NMR (400 MHz, CDCl₃): δ 7.64–7.61 (m, 2H), 7.38– 7.34 (m, 2H), 7.27 (tt, $J = 6.4$ Hz, 1.2 Hz, 1H), 6.33 (t, $J = 2.0$ Hz, 1H), 6.27 (d, $J = 2.0$ Hz, 2H), 5.79 (m, 1H), 5.39 (s, 2H), 5.10 (dq, J $= 12.0$ Hz, 2.0 Hz, 1H), 4.88 (dq, $J = 18.8$ Hz, 2.0 Hz, 1H), 3.67 (s, 6H), 3.40 (m, 2H). ¹³C NMR (100 MHz, CDCl₃): δ 161.4, 145.9, 137.3, 132.4, 131.4, 130.5, 128.7, 127.9, 127.2, 117.8, 105.3, 100.0, 55.5, 52.1, 27.1. HRMS (ESI) m/z calculated for C₂₀H₂₂N₃O₂ [M + H ⁺ 336.1707, found 336.1708. IR (KBr film): ν 3003, 2938, 1598, $1206, 1066, 921, 777$ $\mathrm{cm}^{-1}.$

5-Allyl-1-(4-fluorobenzyl)-4-phenyl-1H-1,2,3-triazole (3f). TLC: $R_{\rm f}$ 0.45 (2 : 1 hexane/EtOAc). Pale yellow liquid (114 mg, 97%). $^1\rm H$ NMR (400 MHz, CDCl₃): δ 7.68-7.66 (m, 2H), 7.44-7.40 (m, 2H), 7.34 (tt, $J = 6.4$ Hz, 1.2 Hz, 1H), 7.20 (dd, $J = 8.8$ Hz, 5.6 Hz, 2H), 7.03 (t, $J = 8.8$ Hz, 2H), 5.84 (m, 1H), 5.50 (s, 2H), 5.16 (dm, $J =$ 10.0 Hz, 1H), 4.91 (dm, $J = 16.4$ Hz, 1H), 3.45 (m, 2H). ¹³C NMR $(100 \text{ MHz}, \text{CDCl}_3): \delta 162.8 \text{ (d, } J_{\text{C-F}} = 246.0 \text{ Hz}), 146.1, 132.3, 131.3,$ 130.9 (d, $J_{\text{C-F}} = 3.1 \text{ Hz}$), 130.3, 129.3 (d, $J_{\text{C-F}} = 7.8 \text{ Hz}$), 128.9, 128.1, 127.3, 118.0, 116.1 $(d, J_{C-F} = 21.7 \text{ Hz})$, 51.4, 27.1. ¹⁹F NMR (376 MHz, CDCl₃): δ -113.1. HRMS (ESI) m/z calculated for $C_{18}H_{17}FN_3$ [M + H]⁺ 294.1401, found 294.1405. IR (KBr film): ν 3066, 3011, 1640, 1512, 1015, 920, 715, 698 cm^{-1} .

5-Allyl-4-phenyl-1-(4-(trifluoromethyl)benzyl)-1H-1,2,3-triazole (3g). TLC: R_f 0.47 (2 : 1 hexane/EtOAc). Light yellow solid (128 mg, 93%). Mp = 63–65 °C. ¹H NMR (400 MHz, CDCl₃): δ 7.69 (d, J = 8.8 Hz, 2H), 7.59 (d, $J = 8.0$ Hz, 2H), 7.42 (t, $J = 8.0$ Hz, 2H), 7.36– 7.29 (m, 3H), 5.84 (m, 1H), 5.57 (s, 2H), 5.15 (d, $J = 10.0$ Hz, 1H), 4.90 (d, $J = 17.2$ Hz, 1H), 3.45 (dm, $J = 3.2$ Hz, 2H). ¹³C NMR (100 MHz, CDCl₃): δ 146.1, 139.1 (q, $J_{\text{C-F}} = 1.5$ Hz), 132.1, 131.2, 130.8 $(q, J_{\text{C-F}} = 32.5 \text{ Hz})$, 130.5, 128.8, 128.1, 127.6, 127.3, 126.1 $(q, J_{\text{C-F}})$ $=$ 3.9 Hz), 124.0 (q, J_{C-F} = 270.9 Hz), 118.0, 51.4, 27.1. ¹⁹F NMR (376 MHz, CDCl₃): δ -63.1. HRMS (ESI) m/z calculated for $C_{19}H_{17}N_3F_3$ [M + H]⁺ 344.1369, found 344.1374. IR (KBr film): ν 3064, 1640, 1496, 1125, 824, 779, 699 $\mathrm{cm}^{-1}.$

5-Allyl-1-(4-nitrobenzyl)-4-phenyl-1H-1,2,3-triazole (3h). TLC: R_f 0.35 (2 : 1 hexane/EtOAc). Light yellow solid (110 mg, 86%). $\text{Mp} = 93\text{--}95 \text{ °C}.$ ¹H NMR (400 MHz, CDCl₃): δ 8.19 (dm, $J =$ 8.8 Hz, 2H), 7.67 (dm, $J = 8.8$ Hz, 2H), 7.44–7.41 (m, 2H), 7.37– 7.33 (m, 3H), 5.84 (m, 1H), 5.61 (s, 2H), 5.14 (dm, $J = 12.0$ Hz, 1H), 4.89 (dm, $J = 17.2$ Hz, 1H), 3.47 (m, 2H). ¹³C NMR (100 MHz, CDCl₃): δ 148.0, 146.2, 142.2, 132.1, 131.0, 130.5, 128.9, 128.3, 128.2, 127.3, 124.3, 118.2, 51.1, 27.1. HRMS (ESI) m/z calculated for $C_{18}H_{17}N_4O_2$ [M + H]⁺ 321.1346, found 321.1348. IR (KBr film): ν 3081, 1608, 1520, 1495, 1347, 805, 734 cm⁻¹.

5-Allyl-4-phenyl-1- $(p$ -tolyl)-1H-1,2,3-triazole (3i). TLC: R_f 0.72 $(2:1 \text{ hexane/EtOAC})$. White solid (98.3 mg, 89%). Mp = 106– 108 °C. ¹H NMR (400 MHz, CDCl₃): δ 7.82 (d, J = 8.4 Hz, 2H), 7.48–4.37 (m, 5H), 7.33 (d, $J = 8.4$ Hz, 2H), 5.91 (m, 1H), 5.19 (d, J $= 9.6$ Hz, 1H), 4.93 (d, $J = 17.2$ Hz, 1H), 3.57 (dm, $J = 4.0$ Hz, 2H), 2.45 (s, 3H). ¹³C NMR (100 MHz, CDCl₃): δ 145.4, 140.1, 134.1, 133.4, 131.4, 131.1, 130.1, 128.9, 128.1, 127.3, 125.5, 118.2, 27.7, 21.4. HRMS (ESI) m/z calculated for $C_{18}H_{18}N_3$ [M + H ⁺ 276.1495, found 276.1501. IR (KBr film): ν 2919, 1517, 992, 934, 710 $\rm cm^{-1}.$

5-Allyl-1-(4-methoxyphenyl)-4-phenyl-1H-1,2,3-triazole (3j). TLC: R_f 0.37 (3 : 1 hexane/EtOAc). Light brown solid (76.0 mg, 65%). Mp = 97–99 °C. ¹H NMR (400 MHz, CDCl₃): δ 7.81 (d, J = 7.2 Hz, 2H), 7.47-7.34 (m, 5H), 7.03 (d, $J = 9.2$ Hz, 2H), 5.91 (m, 1H), 5.19 (dd, $J = 10.0$ Hz, 1.2 Hz, 1H), 4.92 (dd, $J = 17.6$ Hz, 1.2 Hz, 1H), 3.88 (s, 3H), 3.56 (m, 2H). 13C NMR (100 MHz, CDCl3): d 160.6, 133.4, 131.4, 129.4, 128.9, 128.8, 128.0, 127.2, 127.1, 118.1, 114.7, 114.6, 55.8, 27.7. HRMS (ESI) m/z calculated for $C_{18}H_{18}N_3O$ $[M + H]^+$ 292.1444, found 292.1449. IR (KBr film): ν 3081, 1639, 1508, 1260, 1030, 835, 714 cm⁻¹.

5-Allyl-1-(4-fluorophenyl)-4-phenyl-1H-1,2,3-triazole (3k). TLC: R_f 0.45 (2 : 1 hexane/EtOAc). Light yellow solid (70.0 mg, 63%). $\text{Mp} = 121 - 123 \text{ °C}$. ¹H NMR (400 MHz, CDCl₃): δ 7.81 (d, J = 7.2 Hz, 2H), 7.54–7.48 (m, 2H), 7.48–7.44 (m, 2H), 7.40–7.36 (m, 1H), 7.24 (t, $J = 8.4$ Hz, 2H), 5.92 (m, 1H), 5.21 (dd, $J = 10.4$ Hz, 1.2 Hz, 1H), 4.92 (dd, $J = 17.2$ Hz, 1.2 Hz, 1H), 3.57 (m, 2H). ¹³C NMR (100 MHz, CDCl₃): δ 163.3 (d, $J_{C-F} = 249.2$ Hz), 145.6, 133.2, 132.6 (d, $J_{\text{C-F}} = 3.1 \text{ Hz}$), 131.2 (d, $J_{\text{C-F}} = 6.2 \text{ Hz}$), 128.9, 128.3, 127.7 $(d, J_{C-F} = 9.3 \text{ Hz})$, 127.3, 126.1, 118.4, 116.7 $(d, J_{C-F} = 22.5 \text{ Hz})$, 27.7. ¹⁹F NMR (376 MHz, CDCl₃): δ -110.9. HRMS (EI) m/z calculated for $C_{17}H_{14}FN_3$ [M]⁺ 279.1171, found 279.1172. IR (KBr film): ν 3082, 1514, 1244, 992, 846, 721, 701 cm⁻¹.

5-Allyl-1-octyl-4-phenyl-1H-1,2,3-triazole (3l). TLC: R_f 0.65 $(2:1$ hexane/EtOAc). Pale yellow liquid $(107$ mg, 90%). ¹H NMR (400 MHz, CDCl₃): δ 7.68 (dm, J = 8.4 Hz, 2H), 7.42 (tm, J = 8.4 Hz, 2H), 7.35-7.31 (m, 1H), 5.96 (m, 1H), 5.20 (dd, $J =$ 10.4 Hz, 1.2 Hz, 1H), 4.96 (dd, $J = 16.8$ Hz, 1.2 Hz, 1H), 4.23 (t, J $= 7.2$ Hz, 2H), 3.58 (m, 2H), 1.92 (quin, $J = 7.2$ Hz, 2H), 1.40– 1.26 (m, 10H), 0.87 (t, $J = 7.2$ Hz, 3H). ¹³C NMR (100 MHz, CDCl3): d 145.3, 132.9, 131.7, 129.9, 128.8, 127.9, 127.3, 117.8, 48.3, 31.8, 30.3, 29.3, 29.2, 27.2, 26.8, 22.7, 14.2. HRMS (ESI) m/z calculated for $C_{19}H_{28}N_3$ [M + H]⁺ 298.2278, found 298.2278. IR (KBr film): ν 2954, 2926, 1640, 1495, 1361, 765, 698 cm⁻¹.

5-Allyl-1-benzyl-4-(4-fluorophenyl)- $1H-1,2,3$ -triazole (3m). TLC: R_f 0.6 (3 : 1 hexane/EtOAc). White solid (81.6 mg, 70%). $\text{Mp} = 79\text{--}81 \text{ °C}. \text{ }^1\text{H} \text{ NMR}$ (400 MHz, CDCl₃): δ 7.65 (dd, J = 8.8 Hz, 5.6 Hz, 2H), 7.36–7.30 (m, 3H), 7.20–7.18 (m, 2H), 7.11 $(t, J = 8.8$ Hz, 2H), 5.81 (m, 1H), 5.53 (s, 2H), 5.15 (dd, $J =$ 10.4 Hz, 1.2 Hz, 1H), 4.90 $(dd, J = 17.2$ Hz, 1.2 Hz, 1H), 3.42 (m, 2H). ¹³C NMR (100 MHz, CDCl₃): δ 162.7 (d, $J_{\text{C-F}} = 245.3$ Hz), 145.2, 135.0, 132.2, 130.2, 129.2, 129.1 $(d, J_{C-F} = 8.5 \text{ Hz})$, 128.5, 127.6 (d, $J_{\text{C-F}} = 3.1 \text{ Hz}$), 127.4, 118.0, 115.8 (d, $J_{\text{C-F}} =$ 20.9 Hz), 52.2, 27.1. ¹⁹F NMR (376 MHz, CDCl₃): δ -110.9. HRMS (ESI) m/z calculated for $C_{18}H_{17}FN_3$ $[M + H]^+$ 294.1401, found 294.1404. IR (KBr film): ν 3081, 1640, 1510, 1224, 992, 841, 764, 698 $\rm cm^{-1}.$

5-Allyl-1-benzyl-4-(2-chlorophenyl)-1H-1,2,3-triazole (3n). TLC: R_f 0.37 (2 : 1 hexane/EtOAc). Pale yellow liquid (66.7 mg, 54%). ¹H NMR (400 MHz, CDCl₃): δ 7.47-7.42 (m, 2H), 7.38-7.29 (m, 5H), 7.18 (d, $J = 8.0$ Hz, 2H), 5.65 (m, 1H), 5.58 (s, 2H), 5.05 (dd, $J = 10.0$ Hz, 1.2 Hz, 1H), 4.92 (dd, $J = 16.8$ Hz, 1.2 Hz, 1H), 3.27 (m, 2H). ¹³C NMR (100 MHz, CDCl₃): δ 144.1, 135.0, 134.1, 132.6, 132.4, 132.2, 130.4, 130.1, 129.8, 129.1, 128.4, 127.2, 126.9, 117.9, 52.3, 27.5. HRMS (EI) m/z calculated for $C_{18}H_{16}CIN_3$ [M]⁺ 309.1031, found 309.1033. IR (KBr film): ν 3064, 3032, 1639, 1497, 920, 760, 736, 694 cm^{-1} .

5-Allyl-1-benzyl-4-(4-methoxyphenyl)-1H-1,2,3-triazole (3o). TLC: R_f 0.37 (3 : 1 hexane/EtOAc). Pale yellow liquid (70.0 mg, 57%). ¹H NMR (400 MHz, CDCl₃): δ 7.61 (d, $J = 8.8$ Hz, 2H), 7.36–7.30 (m, 3H), 7.20–7.18 (m, 2H), 6.96 (d, $J = 8.8$ Hz, 2H), 5.82 (m, 1H), 5.52 (s, 2H), 5.13 (dd, $J = 12.0$ Hz, 1.6 Hz, 1H), 4.91 (dd, $J = 17.2$ Hz, 1.6 Hz, 1H), 3.83 (s, 3H), 3.42 (m, 2H). ¹³C NMR (100 MHz, CDCl₃): δ 159.6, 145.9, 135.3, 132.5, 129.8, 129.1, 128.6, 128.4, 127.3, 124.1, 117.8, 114.3, 55.5, 52.2, 27.2. HRMS (ESI) m/z calculated for C₁₉H₂₀N₃O [M + H]⁺ 306.1601, found 306.1604. IR (KBr film): ν 2932, 1616, 1508, 1251, 1015, 836, 727, 696 $\rm cm^{-1}$. BSC Advances Weekstones Weekstones Article 2018. This article is licensed on 12 January 2018. Downloaded on 12 January 2018. Downloaded the Creative Commons Article is licensed under a Creative Commons Attribution-Noncomm

5-Allyl-1-benzyl-4-(6-methoxynaphthalen-2-yl)-1H-1,2,3-triazole (3p). TLC: R_f 0.37 (2 : 1 hexane/EtOAc). Light yellow solid (106 mg, 75%). Mp = 113-115 °C. ¹H NMR (400 MHz, CDCl₃): δ 8.04 (d, J = 0.8 Hz, 1H), 7.83 (dd, J = 8.4 Hz, 1.6 Hz, 1H), 7.79 (d, $J = 8.4$ Hz, 1H), 7.76 (d, $J = 7.6$ Hz, 1H), 7.38–7.31 (m, 3H), 7.23– 7.21 (m, 2H), 7.17–7.14 (m, 2H), 5.87 (m, 1H), 5.57 (s, 2H), 5.18 $(dd, J = 10.4 \text{ Hz}, 1.2 \text{ Hz}, 1H$, 4.98 $(dd, J = 17.2 \text{ Hz}, 1.2 \text{ Hz}, 1H$), 3.93 (s, 3H), 3.52 (m, 2H). ¹³C NMR (100 MHz, CDCl₃): δ 158.1, 146.2, 135.2, 134.2, 132.5, 130.4, 129.9, 129.2, 129.1, 128.5, 127.4 (2C), 126.8, 126.1, 126.0, 119.3, 118.0, 105.8, 55.5, 52.2, 27.3. HRMS (EI) m/z calculated for $C_{23}H_{21}N_3O$ 355.1687, found 355.1685. IR (KBr film): v 2936, 2840, 1634, 1600, 1263, 1029, 726, 695 cm^{-1} .

1-Benzyl-5-(2-methylallyl)-4-phenyl-1H-1,2,3-triazole (3q). TLC: R_f 0.73 (2 : 1 hexane/EtOAc). Light yellow solid (104 mg, 90%). Mp $=$ 56–58 °C. ¹H NMR (400 MHz, CDCl₃): δ 7.67 (d, J = 7.2 Hz, 2H), 7.42 (t, $J = 7.2$ Hz, 2H), 7.39-7.28 (m, 4H), 7.19 (dd, $J = 7.6$, 1.6 Hz, 2H), 5.51 (s, 2H), 4.90 (t, $J = 1.2$ Hz, 1H), 4.44 (s, 1H), 3.32 (s, 2H), 1.77 (d, $J = 0.4$ Hz, 3H). ¹³C NMR (100 MHz, CDCl₃): δ 146.1, 140.3, 135.0, 131.4, 130.6, 128.9, 128.7, 128.3, 127.8, 127.2, 127.1, 112.8, 52.0, 30.9, 22.9. HRMS (ESI) m/z calculated for $C_{19}H_{20}N_3$ [M $+ H$ ⁺ 290.1652, found 290.1654. IR (KBr film): ν 3063, 3031, 1605, 1496, 1251, 890, 770, 729 cm^{-1} .

1-Benzyl-4-phenyl-5-(2-phenylallyl)-1H-1,2,3-triazole (3r). TLC: $R_{\rm f}$ 0.54 (2 : 1 hexane/EtOAc). Colorless liquid (120 mg, 86%). $^1\rm H$ NMR (400 MHz, CDCl₃): δ 7.69 (dt, J = 7.2, 1.6 Hz, 2H), 7.41 (t, J = 7.2 Hz, 2H), 7.37–7.28 (m, 9H), 7.22–7.13 (m, 2H), 5.52 (s, 2H), 5.47 (t, $J = 1.6$ Hz, 1H), 4.65 (t, $J = 2.0$ Hz, 1H), 3.82 (t, $J = 2.0$ Hz, 2H). ¹³C NMR (100 MHz, CDCl₃): δ 146.4, 142.6, 139.7, 134.9, 131.2, 130.3, 129.0, 128.8, 128.6, 128.4, 128.3, 128.0, 127.4, 127.1, 125.7, 114.3, 52.2, 28.5. HRMS (ESI) m/z calculated for $C_{24}H_{22}N_3$

 $[M + H]^{+}$ 352.1808, found 352.1807. IR (KBr film): ν 3058, 3032, 1954, 1628 cm $^{-1}$.

1-Benzyl-5-(3-methylbut-2-en-1-yl)-4-phenyl-1H-1,2,3-triazole (3s). TLC: R_f 0.37 (2 : 1 hexane/EtOAc). White solid (89.0 mg, 73%). Mp = 86–88 °C. ¹H NMR (400 MHz, CDCl₃): δ 7.67 (d, J = 7.2 Hz, 2H), 7.42 (t, $J = 7.2$ Hz, 2H), 7.38-7.28 (m, 4H), 7.16 (d, J $= 6.4$ Hz, 2H), 5.54 (s, 2H), 4.93 (m, 1H), 3.41 (d, $J = 6.4$ Hz, 2H), 1.65 (d, $J = 1.6$ Hz, 3H), 1.62 (d, $J = 0.8$ Hz, 3H). ¹³C NMR (100 MHz, CDCl₃): δ 145.1, 135.2, 135.1, 132.6, 131.6, 128.9, 128.6, 128.1, 127.7, 127.4, 127.0, 118.5, 52.0, 25.5, 22.3, 18.1. HRMS (ESI) m/z calculated for $C_{20}H_{22}N_3$ $[M + H]^+$ 304.1808, found 304.1811. IR (KBr film): v 3025, 1654, 1602, 1494, 1249, 968, 728, 708 cm^{-1} .

1-Benzyl-5-cinnamyl-4-phenyl-1H-1,2,3-triazole (3t). TLC: R_f 0.51 (2 : 1 hexane/EtOAc). White solid (74.5 mg, 53%). Mp = 100–102 °C. ¹H NMR (400 MHz, CDCl₃): δ 7.72 (d, J = 7.2 Hz, 2H), 7.43 (t, $J = 7.2$ Hz, 2H), 7.38–7.26 (m, 6H), 7.25–7.18 (m, 5H), 6.21 (d, $J = 16.0$ Hz, 1H), 6.12 (dt, $J = 16.0$, 5.2 Hz, 1H), 5.58 $(s, 2H)$, 3.61 (dd, $J = 5.2$, 1.2 Hz, 2H). ¹³C NMR (100 MHz, CDCl3): d 145.9, 136.3, 135.0, 132.5, 131.3, 130.5, 129.0, 128.7, 128.6, 128.3, 129.9, 127.8, 127.3, 127.2, 126.2, 123.6, 52.2, 26.3. HRMS (ESI) m/z calculated for $C_{24}H_{21}N_3$ $[M + H]^+$ 352.1808, found 352.1812. IR (KBr film): v 3082, 3025, 1955, 1654, 727 cm^{-1} .

5-Allyl-1-benzyl-4-hexyl-1H-1,2,3-triazole (3u). TLC: R_f 0.54 (2 : 1 hexane/EtOAc). Colorless oil (87.2 mg, 77%). ¹H NMR (400 MHz, CDCl₃): δ 7.36-7.27 (m, 3H), 7.12 (dd, $J = 8.0$ Hz, 2.0 Hz, 2H), $5.72 - 5.58$ (m, 1H), 5.46 (s, 2H), 5.07 (dq, $J = 10.4$ Hz, 1.6 Hz, 1H), 4.89 (dq, $J = 16.8$ Hz, 1.6 Hz, 1H), 3.22 (dt, $J = 5.6$ Hz, 1.6 Hz, 2H), 2.60 (t, $J = 7.6$ Hz, 2H), 1.66 (quintet, $J = 6.4$ Hz, 2H), 1.41-1.22 (m, 6H), 0.87 (t, $J = 6.8$ Hz, 3H). ¹³C NMR (100 MHz, DMSO-d₆): δ 144.8, 136.1, 133.3, 130.4, 128.6, 127.8, 127.2, 116.7, 50.6, 30.9, 28.9, 28.2, 25.9, 24.2, 22.0, 13.8. HRMS (ESI) m/ *z* calculated for $C_{18}H_{25}N_3$ [M + H]⁺ 284.2121, found 284.2127. IR (KBr film): ν 3065, 3033, 2928, 1640, 1456, 992, 729 cm⁻¹.

5-Allyl-1-(but-3-en-1-yl)-4-phenyl-1H-1,2,3-triazole (3v). TLC: R_f 0.55 (2 : 1 hexane/EtOAc). Pale yellow liquid (76.8 mg, 80%). ¹H NMR (400 MHz, CDCl₃): δ 7.67 (d, J = 6.8 Hz, 2H), 7.42 (t, J = 6.8 Hz, 2H), 7.33 (tt, $J = 6.8$ Hz, 1.2 Hz, 1H), 5.97 (m, 1H), 5.78 $(m, 1H)$, 5.21 $(dq, J = 10.0$ Hz, 0.8 Hz, 1H), 5.14–5.08 $(m, 2H)$, 4.94 (dq, $J = 17.2$ Hz, 0.8 Hz, 1H), 4.29 (t, $J = 7.2$ Hz, 2H), 3.58 $(m, 2H), 2.70 (q, J = 7.2 Hz, 2H).$ ¹³C NMR (100 MHz, CDCl₃): d 145.3, 133.5, 132.8, 131.6, 130.1, 128.8, 127.9, 127.3, 118.2, 117.9, 47.5, 34.4, 27.2. HRMS (ESI) m/z calculated for C₁₅H₁₈N₃ $[M + H]^{+}$ 240.1495, found 240.1502. IR (KBr film): ν 3080, 1640, $1495, 918, 766, 699$ cm⁻¹.

5-Allyl-1-(pent-4-en-1-yl)-4-phenyl-1H-1,2,3-triazole (3w). TLC: $R_{\rm f}$ 0.70 (2 : 1 hexane/EtOAc). Pale yellow liquid (77.7 mg, 77%). ¹H NMR (400 MHz, CDCl₃): δ 7.67 (d, J = 7.2 Hz, 2H), 7.41 (t, J = 7.2 Hz, 2H), 7.33 (t, $J = 7.2$ Hz, 1H), 5.95 (m, 1H), 5.79 (m, 1H), 5.20 (d, $J = 10.0$ Hz, 1H), 5.07 (d, $J = 10.0$ Hz, 1H), 5.03 (d, $J =$ 17.2 Hz, 1H), 4.95 (d, $J = 17.2$ Hz, 1H), 4.23 (t, $J = 7.2$ Hz, 2H), 3.57 $(m, 2H)$, 2.14 $(q, J = 7.2$ Hz, 2H), 2.04 $(quin, J = 7.2$ Hz, 2H). ¹³C NMR (100 MHz, CDCl₃): δ 145.3, 136.8, 132.8, 131.6, 130.0, 128.7, 127.8, 127.2, 117.8, 116.1, 47.4, 30.7, 29.1, 27.1. HRMS (ESI) m/z calculated for $C_{16}H_{20}N_3$ [M + H]⁺ 254.1652, found 254.1660. IR (KBr film): ν 3079, 2979, 1640, 1495, 916, 765, 700 cm⁻¹.

5-Allyl-1-(hex-5-en-1-yl)-4-phenyl-1H-1,2,3-triazole (3x). TLC: R_f 0.65 (2 : 1 hexane/EtOAc). Pale yellow liquid (69.5 mg, 65%). ¹H NMR (400 MHz, CDCl₃): δ 7.68–7.65 (m, 2H), 7.43–7.39 (m, 2H), 7.34–7.31 (m, 1H), 5.95 (m, 1H), 5.76 (m, 1H), 5.20 (dd, $J =$ 10.0 Hz, 0.8 Hz, 1H), 5.04-4.98 (m, 2H), 4.94 (dd, $J = 8.8$ Hz, 0.8 Hz, 1H), 4.23 (t, $J = 7.6$ Hz, 2H), 3.57 (m, 2H), 2.10 (q, $J =$ 7.6 Hz, 2H), 1.94 (quin, $J = 7.6$ Hz, 2H), 1.47 (quin, $J = 7.6$ Hz, 2H). ¹³C NMR (100 MHz, CDCl₃): δ 145.3, 138.0, 132.8, 131.6, 129.9, 128.8, 127.8, 127.3, 117.8, 115.3, 48.0, 33.2, 29.5, 27.1, 25.9. HRMS (ESI) m/z calculated for $C_{17}H_{22}N_3 [M + H]^+$ 268.1808, found 268.1814. IR (KBr film): ν 3079, 2978, 1640, 1495, 914, 769, 698 $\rm cm^{-1}.$

5-Allyl-4-phenyl-1-(2-vinylphenyl)-1H-1,2,3-triazole (3y). TLC: R_f 0.6 (2 : 1 hexane/EtOAc). Light yellow solid (60.8 mg, 53%). $\text{Mp} = 81-83 \text{ °C}. \text{ }^{1}\text{H NMR}$ (400 MHz, CDCl₃): δ 7.83 (d, J = 7.6 Hz, 2H), 7.74 (d, $J = 8.0$ Hz, 1H), 7.54 (t, $J = 7.6$ Hz, 1H), 7.47 (t, $J =$ 7.6 Hz, 2H), 7.44-7.36 (m, 2H), 7.31 (d, $J = 7.6$ Hz, 1H), 6.18 (dd, 17.6 Hz, 11.2 Hz, 1H), 5.73 (d, $J = 17.6$ Hz, 1H), 5.72-5.65 (m, 1H), 5.27 (d, $J = 10.8$ Hz, 1H), 5.02 (d, $J = 10.0$ Hz, 1H), 4.81 (d, J $= 16.4$ Hz, 1H), 3.43 (d, $J = 5.2$ Hz, 2H). ¹³C NMR (100 MHz, CDCl3): d 144.9, 135.5, 133.7, 132.8, 132.3, 131.3, 130.9, 130.7, 128.9, 128.5, 128.1, 127.2 (2C), 126.4, 118.2, 117.9, 27.5. HRMS (ESI) m/z calculated for $C_{19}H_{18}N_3$ $[M + H]^+$ 288.1495, found 288.1492. IR (KBr film): v 3063, 2964, 1640, 1494, 991, 769, 698 cm^{-1} .

5-Allyl-4-phenyl-1-(2-vinylbenzyl)-1H-1,2,3-triazole (3z). TLC: R_f 0.35 (6 : 1 hexane/EtOAc). Light brown liquid (66.6 mg, 55%). ¹H NMR (400 MHz, CDCl₃): δ 7.70 (d, J = 6.8 Hz, 2H), 7.50 (d, J = 8.0 Hz, 1H), 7.43 (t, $J = 7.6$ Hz, 2H), 7.35 (t, $J = 7.6$ Hz, 1H), 7.30 $(t, J = 7.6 \text{ Hz}, 1\text{H})$, 7.20 $(t, J = 7.6 \text{ Hz}, 1\text{H})$, 7.03 $(dd, J = 16.8 \text{ Hz}$, 10.8 Hz, 1H), 6.81 (d, $J = 7.6$ Hz, 1H), 5.80 (m, 1H), 5.67 (dd, $J =$ 17.2 Hz, 0.8 Hz, 1H), 5.63 (s, 2H), 5.42 (dd, $J = 12.0$ Hz, 0.8 Hz, 1H), 5.13 (d, $J = 10.4$ Hz, 1H), 4.88 (d, $J = 17.2$ Hz, 1H), 3.41 (m, 2H). ¹³C NMR (100 MHz, CDCl₃): δ 146.0, 136.7, 133.4, 132.3, 132.1, 131.4, 130.7, 128.9, 128.6, 128.4, 128.0, 127.5, 127.3, 126.8, 118.2, 117.8, 49.8, 27.1. HRMS (ESI) m/z calculated for $C_{20}H_{20}N_3$ [M + H]⁺ 302.1652, found 302.1646. IR (KBr film): ν 3062, 3033, 1639, 1608, 1495, 919, 771, 699 $\mathrm{cm}^{-1}.$

Experimental procedure for ring closing metathesis

Synthesis of 3-phenyl-7,8-dihydro-4H-[1,2,3]triazolo[1,5- α] azepine (6v). The diene $3v$ (47.9 mg, 200 µmol) was dissolved in anhydrous CH₂Cl₂ (6 mL). A solution of Grubbs' 1st generation catalyst (8.2 mg, 10.0 µmol) in anhydrous CH_2Cl_2 (3.3 mL) was added to the reaction mixture via a cannula, and stirred at room temperature for 19 h. Upon completion of the reaction, the solvent was removed in vacuo. The residue was purified by column chromatography to afford 6v as a pale green liquid (33.3 mg, 79%). TLC: R_f 0.3 (2 : 1 hexane/EtOAc). ¹H NMR (400 MHz, CDCl₃): δ 7.60 (d, J = 7.2 Hz, 2H), 7.44 (t, J = 7.2 Hz, 2H), 7.35 (t, $J = 7.2$ Hz, 1H), 5.75–5.73 (m, 2H), 4.71 (t, $J = 6.0$ Hz, 2H), 3.70–3.69 (m, 2H), 2.55–2.52 (m, 2H). ¹³C NMR (100 MHz, CDCl3): d 144.7, 133.2, 131.5, 129.0, 128.8, 127.9, 127.8, 123.2, 47.9, 27.9, 25.5. HRMS (ESI) m/z calculated for C₁₃H₁₄N₃ [M + H ⁺ 212.1182, found 212.1185. IR (KBr film): ν 3058, 1660, 1624, $1498, 772, 699$ cm^{-1} .

Synthesis of 3-phenyl-4,7,8,9-tetrahydro-[1,2,3]triazolo[1,5-a] azocine (6w). The diene $3w(50.7 \text{ mg}, 200 \mu \text{mol})$ was dissolved in anhydrous CH_2Cl_2 (6 mL). A solution of Grubbs' 1st generation catalyst (8.2 mg, 10.0 µmol) in anhydrous CH_2Cl_2 (3.3 mL) was added to the reaction mixture via a cannula. The reaction mixture was stirred at room temperature for overnight, and was brought to reflux for 6 h. Upon completion of the reaction, the solvent was removed in vacuo. The residue was purified by column chromatography to afford 6w as pale yellow liquid (39.0 mg, 87%). TLC: R_f 0.35 (2 : 1 hexane/EtOAc). ¹H NMR (400 MHz, CDCl₃): δ 7.61 (d, J = 7.6 Hz, 2H), 7.44 (t, J = 7.6 Hz, 2H), 7.35 (t, $J = 7.6$ Hz, 1H), 5.79–5.67 (m, 2H), 4.53 (t, $J = 5.2$ Hz, 2H), 3.64 (d, $J = 4.8$ Hz, 2H), 1.90–1.82 (m, 4H). ¹³C NMR (100 MHz, CDCl3): d 144.4, 132.9, 131.7, 128.8, 128.4, 127.9, 127.8, 127.1, 47.2, 27.2, 24.4, 23.1. HRMS (ESI) m/z calculated for $C_{14}H_{16}N_3$ $[M + H]^{+}$ 226.1339, found 226.1336. IR (KBr film): ν 2929, 1665, $1626, 1447, 733, 698$ cm⁻¹.

Synthesis of (Z)-3-phenyl-7,8,9,10-tetrahydro-4H-[1,2,3]triazolo[1,5- α]azonine (6x) and (E)-1,10-bis(5-allyl-4-phenyl-1H-**1,2,3-triazol-1-yl)dec-5-ene (6x').** The diene $3x$ (53.5 mg, 200) μ mol) was dissolved in anhydrous CH₂Cl₂ (6 mL). A solution of Grubbs' 1st generation catalyst (8.2 mg, 10.0 µmol) in anhydrous CH_2Cl_2 (3.3 mL) was added to the reaction mixture via a cannula. The reaction mixture was brought to reflux for 8 h. Upon completion of the reaction, the solvent was removed in vacuo. The residue was purified by column chromatography to afford 6x as a white solid (31.0 mg, 65%) and a dimer $6x'$ as colorless liquid (8.0 mg, 8%). 6x: TLC: R_f 0.2 (2:1 hexane/ EtOAc). Mp = 201-203 °C. ¹H NMR (400 MHz, CDCl₃): δ 7.68 $(d, J = 7.6 \text{ Hz}, 2\text{H})$, 7.45 $(t, J = 7.6 \text{ Hz}, 2\text{H})$, 7.36 $(t, J = 7.6 \text{ Hz},$ 1H), 5.81 (m, 1H), 5.66 (m, 1H), 4.51 (t, $J = 6.0$ Hz, 2H), 3.61 (d, J $= 7.6$ Hz, 2H), 2.35–2.30 (m, 2H), 2.01 (quin, $J = 6.4$ Hz, 2H), 1.67–1.61 (m, 2H). ¹³C NMR (100 MHz, CDCl₃): δ 145.2, 131.7, 131.5, 131.0, 128.9, 128.0, 127.6, 126.3, 45.8, 25.9, 25.1, 23.0, 21.9. HRMS (ESI) m/z calculated for $C_{15}H_{18}N_3 [M + H]^2 240.1495$, found 240.1498. IR (KBr film): ν 2959, 2932, 1491, 733, 720, 697 cm^{-1} . Paper

SAMple these are typy element BFs alternation (a), TCC Symbols on a periodic is licensed on 12 AM access Article

Access Article 2012 Creative Commons Articles. Due to 2012 Creative Commons are common and the creat

> 6x': TLC: R_f 0.17 (2 : 1 hexane/EtOAc). ¹H NMR (400 MHz, CDCl₃): δ 7.67 (d, J = 7.2 Hz, 4H), 7.42 (t, J = 7.2 Hz, 4H), 7.34 (t, J $=$ 7.2 Hz, 2H), 6.00–5.91 (m, 2H), 5.39–5.35 (m, 2H), 5.21 (d, J = 10.4 Hz, 2H), 4.95 (d, $J = 17.2$ Hz, 2H), 4.25–4.22 (m, 4H), 3.59– 3.57 (m, 4H), 2.09–2.01 (m, 4H), 1.96–1.89 (m, 4H), 1.43 (quin, J $=$ 7.6 Hz, 4H). ¹³C NMR (100 MHz, CDCl₃): δ 145.2, 132.8, 131.4, 130.3, 130.1, 128.9, 128.0, 127.4, 117.9, 48.2, 32.0, 29.6, 27.2, 26.6. HRMS (ESI) m/z calculated for $C_{33}H_{39}N_6 [M + H]^+$ 507.3231, found 507.3232. IR (KBr film): ν 2925, 1660, 1600, 1495, 917, 770, 698 $\rm cm^{-1}$.

> Synthesis of 3-phenyl-4H-benzo[f][1,2,3]triazolo[1,5- α]azepine (6y). The diene $3y$ (52.0 mg, 180 μ mol) was dissolved in anhydrous CH_2Cl_2 (6 mL). A solution of Grubbs' 1st generation catalyst (8.2 mg, 10.0 µmol) in anhydrous CH_2Cl_2 (3.3 mL) was added to the reaction mixture via a cannula. The reaction mixture was stirred at room temperature for 8 h. Upon completion of the reaction, the solvent was removed in vacuo. The residue was purified by column chromatography to afford 6y as a white solid (43.0 mg, 92%). TLC: R_f 0.37 (6:1 hexane/

EtOAc). Mp = 96–98 °C. ¹H NMR (400 MHz, CDCl₃): δ 8.15 (dd, *J* $= 7.6$ Hz, 1.2 Hz, 1H), 7.71 (d, $J = 7.6$ Hz, 2H), 7.50–7.41 (m, 4H), 7.41–7.36 (m, 2H), 6.65 (d, $J = 10.4$ Hz, 1H), 6.18 (m, 1H), 3.51 (d, $J = 6.4$ Hz, 2H). ¹³C NMR (100 MHz, CDCl₃): δ 142.2, 135.0, 134.9, 130.9, 130.8, 130.2, 129.0, 128.8, 128.7, 128.2, 128.1, 127.9, 127.6, 124.0, 21.2. HRMS (ESI) m/z calculated for $C_{17}H_{14}N_3$ [M + H]⁺ 260.1182, found 260.1179. IR (KBr film): ν 2919, 1700, 1600, 1491, 992, 772, 699 $\mathrm{cm}^{-1}.$

Synthesis of (Z)-3-phenyl-4,11-dihydrobenzo[f][1,2,3]triazolo [$1,5$ - α]azocine (6z). The diene 3z (47.0 mg, 150 µmol) was dissolved in anhydrous CH_2Cl_2 (6 mL). A solution of Hoveyda-Grubbs' $2nd$ generation catalyst (5.0 mg, 7.9 µmol) in anhydrous CH_2Cl_2 (3.3 mL) was added to the reaction mixture via a cannula. The reaction mixture was stirred at room temperature for 4 h. Upon completion of the reaction, the solvent was removed in vacuo. The residue was purified by column chromatography to afford 6z as a pale green solid (29.5 mg, 72%). TLC: R_f 0.27 (6 : 1 hexane/EtOAc). Mp = 159-161 °C. ¹H NMR (400 MHz, CDCl3): d 7.58–7.54 (m, 3H), 7.45–7.38 (m, 3H), 7.37– 7.35 (m, 2H), 7.30 (d, $J = 8.4$ Hz, 1H), 7.08 (d, $J = 10.4$ Hz, 1H), 6.03 (dt, $J = 10.4$ Hz, 7.6 Hz, 1H), 5.49 (s, 2H), 3.30 (d, $J = 8.0$ Hz, 2H). ¹³C NMR (100 MHz, CDCl₃): δ 144.8, 139.2, 134.0, 131.5, 131.3, 130.9, 129.3, 128.8, 128.7, 128.6, 128.2, 128.0, 127.7, 126.2, 53.4, 22.6. HRMS (ESI) m/z calculated for C₁₈H₁₆N₃ [M + H ⁺ 274.1339, found 274.1336. IR (KBr film): ν 3025, 2360, 1494, 995, 754, 735, 699 $\rm cm^{-1}$. BSC Advances Worwide Compaction (11 January 2018) (12 January 2022) (12 January 2022) (14 Janua

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Science, ICT and Future Planning (NRF-2015R1A2A2A01004511).

Notes and references

- 1 S. G. Agalave, S. R. Maujan and V. S. Pore, Chem.–Asian J., 2011, 6, 2696.
- 2 For selected references, see: (a) P. Thirumurugan, D. Matosiuk and K. Jozwiak, Chem. Rev., 2013, 113, 4905; (b) J. Hou, X. Liu, J. Shen, G. Zhao and P. G. Wang, Expert Opin. Drug Discovery, 2012, 7, 489; (c) L. Liang and D. Astruc, Coord. Chem. Rev., 2011, 255, 2933; (d) V. Aragao-Leoneti, V. Campo, A. S. Gomes, R. A. Field and I. Carvalho, Tetrahedron, 2010, 66, 9475; (e) J. Lahann, Click Chemistry for Biotechnology and Materials Science, Wiley, Chichester, 2009.
- 3 K. M. Faraz, V. Garima, A. Wasim, M. Akranth, A. M. Mumtaz, A. Mymoona, H. Asif, H. S. Misbahul, S. Mohammad and H. S. Rashiduddin, Int. J. Drug Dev. Res., 2017, 9, 22.
- 4 (a) J. F. Lutz, Angew. Chem., Int. Ed., 2008, 47, 2182; (b) J. F. Lutz and H. G. Borner, Prog. Polym. Sci., 2008, 33, 1;

(c) R. J. Pieters, D. T. S. Rijkers and R. M. J. Liskamp, QSAR Comb. Sci., 2007, 26, 1181; (d) A. J. Dirks, J. Cornelissen, F. L. van Delft, J. C. M. van Hest, R. J. M. Nolte, A. E. Rowan and F. Rutjes, QSAR Comb. Sci., 2007, 26, 1200; (e) J. M. Baskin and C. R. Bertozzi, QSAR Comb. Sci., 2007, 26, 1211; (f) C. M. Salisbury and B. F. Cravatt, *QSAR Comb.* Sci., 2007, 26, 1229.

- 5 (a) H. Nandivada, X. W. Jiang and J. Lahann, Adv. Mater., 2007, 19, 2197; (b) D. Fournier, R. Hoogenboom and U. S. Schubert, Chem. Soc. Rev., 2007, 36, 1369; (c) P. L. Golas and K. Matyjaszewski, QSAR Comb. Sci., 2007, 26, 1116; (d) W. H. Binder and C. Kluger, Curr. Org. Chem., 2006, 10, 1791.
- 6 (a) J. F. Lutz and H. Schlaad, Polymer, 2008, 49, 817; (b) P. Wu and V. V. Fokin, Aldrichimica Acta, 2007, 40, 7; (c) K. A. Kalesh, P. Y. Yang, R. Srinivasan and S. Q. Yao, QSAR Comb. Sci., 2007, 26, 1135; (d) F. Santoyo-Gonz lez and F. Hernandez-Mateo, Top. Heterocycl. Chem., 2007, 7, 133; (e) V. D. Bock, H. Hiemstra and J. H. van Maarseveen, Eur. J. Org. Chem., 2005, 51; (f) Z. Zhou and C. J. Fahrni, J. Am. Chem. Soc., 2004, 126, 8862; (g) W. G. Lewis, F. G. Magallon, V. V. Fokin and M. G. Finn, J. Am. Chem. Soc., 2004, 126, 9152; (h) K. Sivakumar, F. Xie, B. M. Cash, S. Long, H. N. Barnhill and Q. Wang, Org. Lett., 2004, 6, 4603.
- 7 For selected reviews, see: (a) O. S. Miljanic, W. R. Dichtel, I. Aprahamian, R. D. Rohde, H. D. Agnew, J. R. Heath and J. F. Stoddart, QSAR Comb. Sci., 2007, 26, 1165; (b) I. Aprahamian, O. S. Miljanic, W. R. Dichtel, K. Isoda, T. Yasuda, T. Kato and J. F. Stoddart, Bull. Chem. Soc. Jpn., 2007, 80, 1856; (c) A. B. Braunschweig, W. R. Dichtel, O. S. Miljanic, M. A. Olson, J. M. Spruell, S. I. Khan, J. R. Heath and J. F. Stoddart, Chem.–Asian J., 2007, 2, 634; for selected papers see: (d) W. B. Zhang, Y. Tu, R. Ranjan, R. M. VanHorn, S. Leng, J. Wang, M. J. Polce, C. Wesdemiotis, R. P. Quirk, G. R. Newkome and S. Z. D. Cheng, Macromolecules, 2008, 41, 515; (e) J. S. Marois, K. Cantin, A. Desmarais and J. F. Morin, Org. Lett., 2008, 10, 33; (f) V. Aucagne, K. D. Haenni, D. A. Leigh, P. J. Lusby and D. B. Walker, J. Am. Chem. Soc., 2006, 128, 2186; (g) R. K. O'Reilly, M. J. Joralemon, C. J. Hawker and K. L. Wooley, J. Polym. Sci., Part A: Polym. Chem., 2006, 44, 5203; (h) D. D. Diaz, K. Rajagopal, E. Strable, J. Schneider and M. G. Finn, J. Am. Chem. Soc., 2006, 128, 6056.
- 8 J. Gierlich, G. A. Burley, P. M. E. Gramlich, D. M. Hammond and T. Carell, Org. Lett., 2006, 8, 3639.
- 9 A. Nuzzi, A. Massi and A. Dondoni, QSAR Comb. Sci., 2007, 26, 1191.
- 10 (a) X. Deng, J. Liang, B. B. Allison, C. Dvorak, H. McAllister, B. M. Savall and N. S. Mani, J. Org. Chem., 2015, 80, 11003; (b) W. S. Brotherton, R. J. Clark and L. Zhu, J. Org. Chem., 2012, 77, 6443; (c) F. Alonso, Y. Moglie, G. Radivoy and M. Yus, Synlett, 2012, 23, 2179; (d) V. Malnuit, M. Duca, A. Manout, K. Bougrin and R. Benhida, Synlett, 2009, 2123; (e) L. Li, G. Zhang, A. Zhu and L. Zhang, J. Org. Chem., 2008, 73, 3630; (f) D. Yang, N. Fu, Z. Liu, Y. Li and B. Chen, Synlett,

2007, 278; (g) Y.-M. Wu, J. Deng, Y. L. Li and Q.-Y. Chen, Synthesis, 2005, 1314.

- 11 (a) D. N. Barsoum, C. J. Brassard, J. H. A. Deeb, N. Okashah, K. Sreenath, J. T. Simmons and L. Zhu, Synthesis, 2013, 45, 2372; (b) L. Li, R. Li, A. Zhu, G. Zhang and L. Zhang, Synlett, 2011, 874; (c) J. C. Morris, J. Chiche, C. Grellier, M. Lopez, L. F. Bornaghi, A. Maresca, C. T. Supuran, J. Pouyssegur and S.-A. Poulsen, J. Med. Chem., 2011, 54, 6905; (d) N. W. Smith, B. P. Polenz, S. B. Johnson and S. V. Dzyuba, Tetrahedron Lett., 2010, 51, 550; (e) V. Malnuit, M. Duca, A. Manout, K. Bougrin and R. Benhida, Synlett, 2009, 2123; (f) L. Li, G. G. Zhang, A. Zhu and L. Zhang, J. Org. Chem., 2008, 73, 3630; (g) N. Joubert, R. F. Schinazib and L. A. Agrofoglio, Tetrahedron, 2005, 61, 11744; (h) Y. Wu, J. Deng, Y. Li and Q.-Y. Chen, Synthesis, 2005, 1314. Paper

2007, 278; [g] Y-M, Wi, J. Download, J. L. A. Took, N. A. Downloaded on 12 January 2018. Download, 2015, 2015

10 Solventin, J. T. Simmons and L. Chu, Syntheds, 2013, 45;

10 Solventin, J. T. Simmons and L. Chu, Syn
	- 12 B. Wang, J. Zhang, X. Wang, N. Liu, W. Chen and Y. Hu, J. Org. Chem., 2013, 78, 10519.
	- 13 B. Wang, N. Liu, C. Shao, Q. Zhang, X. Wang and Y. Hu, Adv. Synth. Catal., 2013, 355, 2564.
	- 14 (a) C. Shao, X. Wang, Q. Zhang, S. Luo, J. Zhao and Y. Hu, J. Org. Chem., 2011, 76, 6832; (b) C. Shao, X. Wang, J. Xu, J. Zhao, Q. Zhang and Y. Hu, J. Org. Chem., 2010, 75, 7002; (c) C. Shao, G. Cheng, D. Su, J. Xu, X. Wang and Y. Hu, Adv. Synth. Catal., 2010, 352, 1587; (d) S. S. Y. Chui, M. F. Y. Ng and C.-M. Che, Chem.–Eur. J., 2005, 11, 1739.
	- 15 E. C. Royer, M. C. Barral, V. Moreno and A. Santos, J. Inorg. Nucl. Chem., 1981, 43, 705.
	- 16 (a) M. S. Singh, S. Chowdhury and S. Koley, Tetrahedron, 2016, 72, 5257; (b) Y. L. Angell and K. Burgess, Chem. Soc. Rev., 2007, 36, 1674.