

Cite this: *Chem. Sci.*, 2018, 9, 181

Variations in the fuel structure control the rate of the back and forth motions of a chemically fuelled molecular switch†‡

Chiara Biagini,^a Simone Albano,^a Rachele Caruso,^a Luigi Mandolini,^a José Augusto Berrocal^b and Stefano Di Stefano^{*a}

This work deals with the use of 2-cyano-2-arylpropanoic acids as chemical fuels for an acid–base operated molecular switch that consists of a Sauvage-type catenand composed of two identical macrocycles incorporating a phenanthroline unit. When used as a base promoter of the decarboxylation of propanoic acid derivatives, the switch undergoes large amplitude motion from the neutral catenand to a protonated catenane and back again to the neutral state. The rate of back proton transfer, which determines the rate of the overall process, was markedly affected by *para*-substituents in the order Cl > H > CH₃ > OCH₃ ($\rho = +5.2$). Thus, the time required to complete a full cycle was almost two days for the OCH₃ derivative and dropped to a few minutes for the Cl derivative. These results show for the first time that the rate of operation of a molecular switch can be regulated by variations in the fuel structure.

Received 20th September 2017
Accepted 18th October 2017

DOI: 10.1039/c7sc04123c

rsc.li/chemical-science

Introduction

Great effort has been devoted in recent years to the development of molecular machines.¹ Prototypal examples of molecular machines are mechanically interlocked systems, such as rotaxanes and catenanes, characterized by various levels of structural complexity. The distinctive feature of these systems is the ability to perform back and forth motions, *i.e.* a series of two or more roto-translations during which the machine varies from an initial state A to one or more intermediate states, B', B'',..., and eventually reverts back to the initial state A.

The transition between states A and B usually requires the sequential addition of a fuel and an anti-fuel, such as an acid and a base,² a reductant and an oxidant,³ irradiation with light of different wavelengths⁴ and the like. Systems in which only one stimulus is required to guarantee the operation of a back and forth motion are rare. Early examples were reported by Leigh and Wurpel *et al.*^{5a} and Balzani, Credi and Stoddart *et al.*^{5b} who used light irradiation to induce cyclic motion in a rotaxane structure.


^aDipartimento di Chimica and Istituto CNR di Metodologie Chimiche-IMC, Sezione Meccanismi di Reazione c/o Dipartimento di Chimica, Università degli Studi di Roma "La Sapienza", P.le A. Moro 5, 00185 Rome, Italy. E-mail: stefano.distefano@uniroma1.it

^bInstitute for Complex Molecular Systems, Eindhoven University of Technology, P.O. Box 513, 5600 MB Eindhoven, The Netherlands

† This work is dedicated to Professor Carlo Galli.

‡ Electronic supplementary information (ESI) available: NMR of acids 2 (X = Cl, H, CH₃ and OCH₃), ¹H NMR monitoring of the decarboxylation of acids 2 (X = Cl, CH₃ and OCH₃) promoted by Et₃N or catenand 1 and ¹H NMR of 1H⁺CF₃CO₂⁻. See DOI: 10.1039/c7sc04123c

It is only lately that molecular machines operated by the irreversible reaction of a single chemical fuel have been reported by us⁶ and other workers.⁷ Our system is described in Scheme 1. The Sauvage-type [2]catenand **1**, composed of two identical macrocycles incorporating 1,10-phenanthroline units, performs fuelled switching between states A and B'–B''. While B' and B'' are well-defined states with the two phenanthroline



Scheme 1 Switching motions of catenane **1** fuelled by acid **2** (X = H). Fast and quantitative proton transfer (step A → B') fixes the contact segments between the interlocked macrocycles in the region of the phenanthroline nitrogen atoms. Loss of CO₂ (step B' → B''), followed by slower back proton transfer from 1H⁺ to its counteranion, restores the catenane to its original form (step B'' → A).



subunits tightly held together by interactions with the shared proton, state **A** is not co-conformationally well-defined. The fuel, 2-cyano-2-phenylpropanoic acid (2, X = H) undergoes base-promoted quantitative decarboxylation *via* a carbanion intermediate, which is rapidly formed (step **B'** → **B''**) and slowly transformed (step **B''** → **A**) to the “waste” product 2-phenylpropanonitrile (3, X = H) by back proton transfer.

Since the stability and reactivity of carbanions are strongly structure dependent,⁸ we reasoned that the rate of the back and forth motion could be conveniently controlled by structural variations of the acid fuel. Here we report an assessment of the influence of the substituents in the *para* positions of 2-cyano-2-(*p*-chlorophenyl)propanoic acid (2, X = Cl), 2-cyano-2-(*p*-methylphenyl)propanoic acid (2, X = CH₃) and 2-cyano-2-(*p*-methoxyphenyl)propanoic acid (2, X = OCH₃) on the rate of the fuelled switching of catenand **1**.

Results and discussion

Compounds **2** (X = Cl, CH₃ and OCH₃) were prepared according to the same procedure adopted for the preparation of the parent acid **2** (X = H) (Scheme 2).⁶

In a set of control experiments aimed at assessing the effect of the substituents on the rate of rupture of the R–CO₂[−] bond, the Et₃N promoted decarboxylation of the acid derivatives was monitored by ¹H NMR spectroscopy in CD₂Cl₂ at 25 °C for comparison with the corresponding reaction of the parent acid **2** (X = H). For the latter it has been established⁶ that fast and quantitative proton transfer from acid **2** (X = H) to Et₃N generates a hydrogen bonded ion pair§ RCO₂[−]⋯HNEt₃⁺. The rate-determining liberation of CO₂ leads to a carbanion intermediate R[−], presumably ion paired to Et₃NH⁺.¶ Fast proton transfer from Et₃NH⁺ to R[−] completes the reaction and restores the neutral form of Et₃N.

Decarboxylation proceeded smoothly with the other substituents as well and afforded a quantitative yield of the decarboxylated product **3** in all cases (see ESI, Fig. S1–S3†). The half-lives of 14, 36, 204 and 332 min, determined from the ¹H NMR spectra, for the sequence X = Cl, H, CH₃ and OCH₃,


respectively, are in line with expectations for a reaction in which a fractional negative charge develops on the benzylic carbon atom in the transition state. A Hammett plot of log *k*_{rel} vs. σ_p gave ρ = 2.9 (*r* = 0.98) (see ESI, Fig. S4†).

Let us now turn to the use of catenand **1** as a base promoter of decarboxylation. A marked difference from the Et₃N promoted reaction was expected, because evidence was obtained that the back proton transfer to the carbanion intermediate (Scheme 1, step **B''** → **A**) becomes the slow step in the decarboxylation of the parent acid **2** (X = H).⁶ ¹H NMR spectra of a 1 : 1 mixture of **1** and **2** (X = H) (see ESI, Fig. S6†) revealed the presence of a number of signals belonging to a reaction intermediate which was thought to be a tight ion pair composed of the protonated catenane **1H**⁺ and the carbanion produced in step **B'** → **B''**. Proton transfer from **1H**⁺ to its counteranion restores the catenane to its original form (step **B''** → **A**). The exceptional slowness of the proton removal from the tetrahedral cavity defined by the four N atoms|| was ascribed to steric hindrance.⁶

The reactions of **1** with equimolar amounts of **2** (X = Cl, CH₃ and OCH₃) in dichloromethane at 25 °C were monitored by ¹H NMR spectroscopy (see ESI, Fig. S5, S7 and S8†). As previously found for the reaction of the parent acid **2** (X = H), the spectra showed that in all cases the sole reaction product was the corresponding decarboxylated compound **3**, besides the catenane in its original neutral state. Furthermore, the ¹H NMR spectra recorded in the course of the reactions confirmed the presence of transient species, characterized by patterns of signals very similar to each other, as well as to those displayed by the reaction of the parent acid (Fig. 1). Clearly, intermediates of the same nature and with very similar structures (denoted as **B''** in Scheme 1) are involved in all cases.** Quite remarkably, the time required to achieve quantitative transformation to **3** amounts to almost two days for the OCH₃ derivative and decreases in the order OCH₃ > CH₃ > H > Cl, becoming as short as a few minutes for the Cl derivative.

The spectra related to the early stages of the sluggish reaction of **2** (X = OCH₃) (see Fig. 2 and ESI, Fig. S8†), particularly in the region of the double bond protons†† at around 5.5 ppm (Fig. 2), gave information about step **B'** → **B''** which could not be obtained from the faster reactions of **2** (X = CH₃, H and Cl). The trace at *t* = 2.5 min shows that **1** is no longer in its original, neutral state, but the signals of the ion pair **B''** are still barely perceptible. This means that the major component of the reaction mixture is the salt denoted by **B'** in Scheme 1. Since the spectrum, particularly in the region of the aromatic protons, is considerably more complex than the spectrum of the trifluoroacetate salt (see ESI, Fig. S9†), there must be a close association between **1H**⁺ and its carboxylate counterion. The concentration of this transient species is still significant at *t* = 14.5 min and becomes negligibly small in the next spectrum at 34.5 min.

Corroborating evidence came from investigation of the time-dependent UV-vis spectra of the reaction mixtures, which was prompted by the observation that the solutions became yellow in the course of the reactions, but were colourless at the end. The four sets of time-dependent UV-vis spectra (Fig. 3) are



Scheme 2 Preparation of acids **2** (X = Cl, CH₃ and OCH₃).





Fig. 1 Portions of the ^1H NMR spectra (CD_2Cl_2 , 25°C) recorded in the course of the reaction of 5 mM **1** with equimolar amounts of acid **2** ($X = \text{H}$, Cl , CH_3 and OCH_3) after 3.5, 2.5, 7.5 and 44.5 min from the start, respectively. The signals marked with an asterisk are proposed to belong to ten of the twelve protons of the two non-equivalent phenanthroline units in **B''**. The peak at 5.30 ppm is due to CHDCl_2 .

characterized by very similar absorption bands ($\lambda_{\text{max}} = 375 \text{ nm}$) belonging to transient species. The latter were identified with intermediates **B''** or, more precisely, with their carbanion components, since the absorbance of $0.30 \text{ mM } 1\text{H}^+$ is 0.30 at 350 nm, drops to 0.05 at 375 nm and becomes negligibly small at 380 nm and beyond. Indeed, the spectra related to the reaction of **2** ($X = \text{OCH}_3$) (Fig. 4) show that the absorbance reaches its maximum value after about 20 min and then decreases slowly. Clearly, this behavior agrees nicely with the behavior of intermediate **B''** as monitored by ^1H NMR spectroscopy (Fig. 2). In the reactions of acids **2** ($X = \text{CH}_3$ and H) the maximum is reached in about 3 and 1.5 min, respectively, which means that the build-up of **B''** was complete, or very nearly so, when the first ^1H NMR spectrum in Fig. S7 and S6, \ddagger respectively, was recorded. Only the fast decay of the absorption band is visible in the reaction of the Cl derivative, showing that the growing phase was over in the short time required to record the first spectrum after reactant mixing.

For the reactions of **2** ($X = \text{OCH}_3$ and CH_3), the absorbance growth and subsequent decay were treated as two separate first-order reactions with rate constants k' and k'' , respectively. Fig. 4 shows that there is a good adherence of experimental data to the lines calculated from eqn (S3) (see ESI \ddagger) and the best fit values of k' and k'' .



Fig. 2 ^1H NMR monitoring of the reaction of 5 mM **1** with 5 mM **2** ($X = \text{OCH}_3$) in CD_2Cl_2 at 25°C . The trace at time zero is the spectrum of catenand **1**. The asterisks denote the signals of the double bond protons: red = catenand **1**, blue = **B'** and green = **B''**.

A set of four kinetic runs in which the initial concentration of **1** and **2** ($X = \text{OCH}_3$) was varied over the range $0.30\text{--}1.50 \text{ mM}$ (ESI, \ddagger page S18) gave $k' = 2.5 \pm 0.5 \times 10^{-3} \text{ s}^{-1}$ and $k'' = 3.1 \pm 0.2 \times 10^{-5} \text{ s}^{-1}$, showing that k' and k'' are concentration-independent quantities. Not surprisingly, the experimental uncertainty in k' is large because of the limited number of data points in the growth phase. As for the reactions of **2** ($X = \text{H}$ and Cl), Fig. 4 shows that, except for a brief initial period, there is a strict first-order time dependence of the absorbance data in both cases.

In a last set of rate measurements the effect of added Bu_4NBr on the rate of reaction of $0.30 \text{ mM } 2$ ($X = \text{H}$) with equimolar **1** was investigated. UV-vis monitoring of the reactions showed in





Fig. 3 Time dependent UV-vis spectra of the reaction mixtures composed of equimolar amounts (0.30 mM) of catenane **1** and acid **2** (X = OCH₃, CH₃, H and Cl) in CH₂Cl₂ at 25 °C. The dotted lines represent the absorption spectrum of **1**. The spectra pictured as broken lines refer to the ascending phase of the absorbance, whereas the full lines refer to the descending phase.

all cases a rapid growth followed by a slower decay of the absorption band centered at 375 nm, the shape and maximum intensity of which were hardly affected by the added salt, even at the highest salt concentration of 5.0 mM (Fig. S10†). Again, except for a brief initial period, the absorbance was found to decrease according to first-order kinetics. The first order rate constants (k''_{obs}) increased with increasing salt concentration and showed a marked tendency to saturate. Fig. 5 shows a close adherence of the kinetic data to the line of eqn (1), which is consistent with the reaction mechanism depicted in Scheme 3, where the salt forms a 1 : 1 complex of moderate stability with **B''** ($K_{\text{ass}} = 1600 \text{ M}^{-1}$), which is significantly more reactive than uncomplexed **B''** ($k''_{\text{salt}}/k''_{\text{o}} = 20$).††.

It seems likely that the components of the ion pair **B''**, namely 1H^+ and the carbanion derived from **2** (X = H), are brought into an orientation more adapted to proton transfer when **B''** becomes a component of an ion quartet as a result of association with Bu₄NBr. However, due to the lack of structural information on the ion quartet, a detailed interpretation of the rate enhancing effect exerted by the added salt does not appear to be accessible.

$$\frac{k''_{\text{obs}}}{k''_{\text{o}}} = \frac{1 + \left(k''_{\text{salt}}/k''_{\text{o}}\right)K_{\text{ass}}[\text{salt}]}{1 + K_{\text{ass}}[\text{salt}]} \quad (1)$$

To sum up, the combination of data from ¹H NMR and UV-vis spectroscopy strongly supports the operation of the reaction mechanism outlined in Scheme 1, which involves compounds **B'** and **B''** as reaction intermediates.

Data from UV-vis spectroscopy gave pieces of information which could not be obtained from ¹H NMR spectroscopy. Firstly, the transient absorption band at 375 nm provided direct evidence of the existence and kinetic behavior of the carbanion intermediates, whose signals in the ¹H NMR spectra were hardly visible. Secondly, the kinetics based on UV-vis spectroscopy gave qualitative information about the substituent effects on the rate of step **B' → B''**. The reactivity order Cl > H > CH₃ > OCH₃ is the same as that observed in the reactions promoted by Et₃N, but the rupture of the R-CO₂⁻ bond is much faster in the reactions promoted by **1** because any stabilization of R-CO₂⁻ by hydrogen bonding is hardly conceivable when 1H^+ is the counteranion.§

Finally, kinetic treatment of the UV-vis data provided a quantitative estimate of the effect of the substituents on the rate of the back proton transfer (step **B'' → A**), whereas only a qualitative reactivity order could be obtained from the ¹H NMR spectra. The Hammett plot of log ($k''_{\text{H}}/k''_{\text{H}}$) vs. σ_{p} reported in Fig. 6 gives $\rho = 5.2$ ($r = 0.98$). Such a large ρ value is indicative of a much larger fraction of negative charge on the benzyl carbon atom in the transition state, which is equivalent to saying that the kinetic basicity of the anionic component in **B''** is enhanced by EWGs and depressed by EDGs. Unusual as it might seem, such a behavior is not unprecedented.¹⁰ Bordwell *et al.*^{10b} carried out an extensive investigation of the equilibrium and kinetic acidities of ArCHMeNO₂ (12 substituents). They found that the rates of protonation of ArCMe=NO₂⁻ were enhanced by EWGs and retarded by EDGs ($\rho > 0$) and concluded that there is a very little delocalization of the fractional negative charge to the oxygen atoms in the transition state, whereas the negative charge is mostly localized on the oxygen atoms of the nitronate anions.

Although a cyano group is less effective than a nitro group in stabilizing a negative charge on an adjacent carbon atom, we suggest that a similar interpretation applies to the protonation of our cyano-stabilized carbanions. Accordingly, it seems likely that the canonical structure **I** (Fig. 7), unlike structure **II**, contributes very little to the hybrid. This view is strongly supported by the X-ray molecular structure of the Bu₄N⁺ salt of 1-cyano-1-phenylethanide, prepared by the treatment of **2** (X = H) with Bu₄NOH.¹¹ The essential planarity of the anion, coupled with the finding that only the nitrogen of the cyano group is involved in multiple hydrogen bonding interactions with the α -methylene groups of the cations, is indicative of very little delocalization of the negative charge to the sp² benzylic carbon atom, as well as significant delocalization to the nitrogen atom of the cyano group. In our case, a high negative charge density on the nitrogen atom of the CN group would guarantee a strong electrostatic stabilization of **B''**, whose shape is supposed to be one in which the group =C=N⁻ points to the proton embedded in the two-phenanthroline core.






Fig. 4 Absorbance vs. time profiles of the decarboxylation of 0.30 mM **2** ($X = \text{OCH}_3$, CH_3 , H and Cl) promoted by 0.30 mM **1** in CH_2Cl_2 at 25 °C. $X = \text{OCH}_3$: the full line is a plot of eqn (S2) \ddagger with best fit parameters $k' = 3.1 \times 10^{-3} \text{ s}^{-1}$ and $k'' = 3.2 \times 10^{-5} \text{ s}^{-1}$. $X = \text{CH}_3$: the full line is a plot of eqn (S2) \ddagger with best fit parameters $k' = 1.7 \times 10^{-2} \text{ s}^{-1}$ and $k'' = 1.3 \times 10^{-4} \text{ s}^{-1}$. $X = \text{H}$: the full line is a plot of a first order equation with $k'' = 2.4 \times 10^{-4} \text{ s}^{-1}$. $X = \text{Cl}$: the full line is a plot of a first order equation with $k'' = 2.6 \times 10^{-2} \text{ s}^{-1}$.

Experimental section

Instruments and methods

^1H NMR spectra were recorded on a 300 MHz spectrometer. The spectra were internally referenced to the residual proton signal of the solvent at 5.30 ppm. Spectrophotometric measurements were carried out on a diode array spectrophotometer equipped with a thermostated cell compartment. Mass spectrometric measurements were carried out using an ESI-TOF spectrometer.

Materials

All reagents and solvents were purchased at the highest commercial quality and were used without further purification unless otherwise stated. CH_2Cl_2 and CD_2Cl_2 were flushed through basic alumina immediately prior to use. The same batch of solvent was used in each of the sets of kinetic measurements. Et_3N was distilled from metallic sodium before

use. Catenane **1** (a mixture of the EE, EZ and ZZ geometrical isomers) was available from a previous investigation.¹² Acid **2** ($X = \text{H}$) was available from a previous investigation⁶ while acids **2** ($X = \text{Cl}$, CH_3 and OCH_3) were prepared from the corresponding ethyl esters¹³ according to a literature procedure.¹⁴


2-Cyano-2-(4'-chlorophenyl)propionic acid (2, X = Cl). This compound had a m.p. of 81–82 °C, lit. 81–82 °C.¹⁵ ^1H NMR (300 MHz, CD_2Cl_2) δ : 7.55–7.52 (m, 2H), 7.46–7.43 (m, 2H), 3.52 (br s, 1H), 1.99 (s, 3H).

2-Cyano-2-(4'-methyl)propionic acid (2, X = CH₃). This new compound had a m.p. of 86–88 °C (dec.). ^1H NMR (300 MHz, CDCl_3) δ : 7.46–7.43 (m, 2H), 7.24–7.22 (m, 2H), 4.05 (br. s, 1H), 2.36 (s, 3H), 1.97 (s, 3H). ^{13}C NMR (75 MHz, CDCl_3) δ : 171.2, 139.1, 131.8, 129.8, 125.7, 119.0, 47.7, 24.2, 20.9. UV-vis (CH_2Cl_2): λ_{max} (ϵ) = 230 nm ($3200 \text{ cm}^{-1} \text{ M}^{-1}$), 262 nm ($400 \text{ cm}^{-1} \text{ M}^{-1}$), 285 nm ($200 \text{ cm}^{-1} \text{ M}^{-1}$). ESI-MS (negative-ion-mode): 144 ($\text{M} - \text{H}^+ - \text{CO}_2$).





Fig. 5 Effect of the added Bu_4NBr on the rate of decarboxylation of 0.30 mM **2** ($X = \text{H}$) promoted by 0.30 mM **1** in CH_2Cl_2 at 25 °C. The full line is a plot of eqn (1) with best fit parameters $K_{\text{ass}} = 1600 \text{ M}^{-1}$ and $k''_{\text{salt}} = 5.0 \times 10^{-3} \text{ s}^{-1}$.



Scheme 3 Associative mechanism involving the fast and reversible formation of a reactive 1 : 1 adduct between the added salt and the ion pair $1\text{H}^+ \cdot \text{R}^-$. The latter corresponds to **B''** in Scheme 1.




Fig. 6 Hammett plot for the decarboxylation of 2-cyano-2-arylpropionic acids **2**. The k'' values are from the caption of Fig. 4.

2-Cyano-2-(4'-methoxyphenyl)propionic acid (2, $X = \text{OCH}_3$). This new compound had a m.p. of 87–89 °C (dec.). ^1H NMR (300 MHz, CD_2Cl_2) δ : 7.51–7.46 (m, 2H), 6.99–6.93 (m, 2H), 4.95 (br. s, 1H), 3.82 (s, 3H), 1.97 (s, 3H). ^{13}C NMR (75 MHz, CD_2Cl_2) δ : 171.3, 160.0, 127.1, 126.7, 119.2, 114.3, 55.3, 47.1, 23.8. UV-vis (MeOH): λ_{max} (ϵ) = 230 nm ($13\,900 \text{ cm}^{-1} \text{ M}^{-1}$), 274 nm ($1600 \text{ cm}^{-1} \text{ M}^{-1}$), 281 nm ($1300 \text{ cm}^{-1} \text{ M}^{-1}$). ESI-MS (negative-ion-mode): 160 ($\text{M} - \text{H}^+ - \text{CO}_2$).



Fig. 7 Canonical structures of 1-cyano-1-phenyl ethanide.

Conclusions

In this work we have described the first example of a chemically operated molecular switch in which the rate of the back and forth motion could be regulated within wide limits by variations in the fuel structure, namely by variations in the nature of the *para*-substituent of the 2-cyano-2-arylpropionic acid used as a fuel. The time taken to complete a full cycle decreased in the order $\text{OCH}_3 > \text{CH}_3 > \text{H} > \text{Cl}$. It was as long as two days for the OCH_3 derivative, and dropped to a few minutes for the Cl derivative. Fine tuning of the motion rate could be further achieved by the addition of Bu_4NBr . A detailed kinetic investigation has shown the intermediacy of ion pairs **B'** and **B''**, which differ in the nature of the anionic component. The negative charge of the anion in **B''** was suggested to be mostly localized on the nitrogen atom of the CN group, and strongly shifted to the benzyl carbon atom in the transition state of the rate-determining proton transfer step.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Thanks are due to the Ministero dell'Istruzione, dell'Università e della Ricerca (MIUR, PRIN 2010CX2TLM). This work was also partially supported by the Università di Roma La Sapienza (Progetti di Ricerca 2014). The authors thank Flavia Cenesi for technical assistance. SDS thanks Professor Carlo Galli for the help and encouragement.

Notes and references

§ The idea that the ion pair is strongly hydrogen bonded is consistent, *inter alia*, with the finding that the time required to complete the decarboxylation of **2** ($X = \text{H}$) dropped from about 3 h to 10 min when Et_3N was replaced by the proton sponge 1,8-bis(dimethylamino)naphthalene (see ref. 6). The protonated proton sponge, where the proton is covalently bound to one nitrogen atom and hydrogen bonded to the other, stabilizes the carboxylate counterion much less effectively than Et_3NH^+ .

¶ Because of the low polarity of dichloromethane ($\epsilon = 9.08$) the concentration of free ions is most likely negligibly small.

|| When the counteranion of 1H^+ is such a weakly interacting anion as CF_3CO_2^- , the equivalence of the two phenanthroline units in the ^1H NMR spectrum (see ESI, Fig. S9†) points to a symmetrical species in which scrambling of the proton among the nitrogen atoms is fast on the ^1H NMR timescale.



** In our previous work (see ref. 6) the three signals that appear in the range from 4.9 to 3.7 ppm during the reaction of 2 (X = H) (see ESI, Fig. S6†) were erroneously assigned to the five aromatic protons of the 1H^+ -bound carbanion. The low precision of the integrated intensities was largely responsible for the incorrect assignment. Whereas the areas of the signals at 4.75 and 3.85 ppm were approximately in a 1 : 1 ratio, that of the signal at 4.30 ppm turned out to be approximately 2.5 times greater than those of the smallest signals. Literature ^1H NMR data (see ref. 9) showing the non-equivalence of the aromatic protons of the α -methylbenzyl carbanion prompted us to round up the relative intensity of the central peak to 3. The appearance of signals with the same patterns and almost identical chemical shifts during the reaction of 2 (X = Cl, CH_3 and OCH_3) (Fig. 1) led us to reconsider our previous assignment, because the presence of *para*-substituents should have altered their number, shape and multiplicity. Hence we concluded that these signals belong to one of the non-equivalent phenanthroline units, which is the common structural element in **B'** when X is varied, and correspond to four protons in a 1 : 2 : 1 ratio. These signals are shifted upfield by the ring current of the aryl group of the counteranion. We also concluded that the resonances of the aromatic protons of the carbanions should lie in the range from 8.2 to 7.1 ppm.

†† Analysis of the time-dependent changes of the resonances of the double bond protons of **1** is somewhat complicated by the presence of geometrical isomers in a 7 : 1 ratio (see ref. 12a). The pertinent signals are clearly visible in the ^1H NMR spectrum of the trifluoroacetate salt (see ESI, Fig. S9†). In the spectrum of **1** the signal of the less abundant isomer is hidden by the signal of CHDCl_2 .

‡‡ An alternative mechanism involves fast and reversible double exchange between ion pair partners, followed by a rate limiting second-order reaction between $1\text{H}^+\text{Br}^-$ and $\text{Bu}_4\text{N}^+\text{R}^-$.



It should be stressed that this mechanism, which is almost kinetically indistinguishable from that in Scheme 3 (see ESI,† page S20), also proceeds through an ion-quartet transition state.

- 1 (a) *Molecular Catenanes, Rotaxanes and Knots: A Journey Through the World of Molecular Topology*, ed. J.-P. Sauvage and C. O. Dietrich-Buchecker, Wiley-VCH, Weinheim, 1999; (b) F. M. Raymo and J. F. Stoddart, *Molecular Switches*, ed. B. L. Feringa, Wiley-VCH, Weinheim, 1st edn, 2001, pp. 219–248; (c) J.-P. Collin, J.-M. Kern, L. Raehm and J.-P. Sauvage, *Molecular Switches*, ed. B. L. Feringa, Wiley-VCH, Weinheim, 1st edn, 2001, pp. 249–307; (d) W. R. Browne and B. L. Feringa, *Nat. Nanotechnol.*, 2006, **1**, 25–35; (e) V. Balzani, A. Credi, S. Silvi and M. Venturi, *Chem. Soc. Rev.*, 2006, **35**, 1135–1149; (f) G. Podoprygorina and V. Böhmer, *Modern Supramolecular Chemistry: Strategies for Macrocyclic Synthesis*, ed. F. Diederich, P. J. Stang and R. R. Tykwinski, Wiley-VCH, Weinheim, 1st edn, 2008, pp. 143–184; (g) J. A. Wisner and B. A. Blight, *Modern Supramolecular Chemistry: Strategies for Macrocyclic Synthesis*, ed. F. Diederich, P. J. Stang and R. R. Tykwinski, Wiley-VCH, Weinheim, 1st edn, 2008, pp. 349–391; (h) V. Balzani, A. Credi and M. Venturi, *Molecular Devices and Machines: Concepts and Perspectives for the Nanoworld*, Wiley-VCH, Weinheim, 2nd edn, 2008; (i) J. F. Stoddart, *Chem. Soc. Rev.*, 2009, **38**, 1802–1820; (j) J. E. Beves, B. A. Blight, C. J. Cambell, D. A. Leigh and R. T. McBurney, *Angew. Chem., Int. Ed.*, 2011, **50**, 9260–9327; (k) J.-C. Chambron and J.-P. Sauvage, *New J. Chem.*, 2013, **37**,

49–57; (l) X. Liu, C.-H. Lu and I. Willner, *Acc. Chem. Res.*, 2014, **47**, 1673–1680; (m) S. Erbaş-Çakmak, D. A. Leigh, C. T. McTernan and A. L. Nussbaumer, *Chem. Rev.*, 2015, **115**, 10081–10206; (n) M. A. Watson and S. L. Cockroft, *Chem. Soc. Rev.*, 2016, **45**, 6118–6129.

- 2 For selected examples see: (a) R. A. Bissell, E. Cordova, A. E. Kaifer and J. F. Stoddart, *Nature*, 1994, **369**, 133–137; (b) J. D. Badjic, V. Balzani, A. Credi, S. Silvi and J. F. Stoddart, *Science*, 2004, **303**, 1845–1849; (c) S. Grunder, P. L. McGrier, A. C. Whalley, M. M. Boyle, C. Stern and J. F. Stoddart, *J. Am. Chem. Soc.*, 2013, **135**, 17691–17694; (d) V. Blanco, D. A. Leigh, V. Marcos, J. A. Morales-Serna and A. L. Nussbaumer, *J. Am. Chem. Soc.*, 2014, **136**, 4905–4908.
- 3 For selected examples see: (a) M. Asakawa, P. R. Ashton, V. Balzani, A. Credi, C. Hamers, G. Mattersteig, M. Montalti, A. N. Shipway, N. Spencer, J. F. Stoddart, M. S. Tolley, M. Venturi, A. J. P. White and D. J. Williams, *Angew. Chem., Int. Ed.*, 1998, **37**, 333–337; (b) N. Armaroli, V. Balzani, J.-P. Collin, P. Gavina, J.-P. Sauvage and B. Ventura, *J. Am. Chem. Soc.*, 1999, **121**, 4397–4408; (c) S. A. Vignon, T. Jarrosson, T. Iijima, H. R. Tseng, J. K. M. Sanders and J. F. Stoddart, *J. Am. Chem. Soc.*, 2004, **126**, 9884–9885; (d) C. J. Bruns, M. Frascioni, J. Iehl, K. J. Hartlieb, S. T. Schneebeli, C. Cheng, S. I. Stupp and J. F. Stoddart, *J. Am. Chem. Soc.*, 2014, **136**, 4714–4723.
- 4 For selected examples see: (a) B. M. Neilson and C. W. Bielawski, *Chem. Commun.*, 2013, **49**, 5453–5455; (b) Y. Norikane and N. Tamaoki, *Org. Lett.*, 2004, **6**, 2595–2598; (c) M. Baroncini, S. Silvi, M. Venturi and A. Credi, *Chem.–Eur. J.*, 2010, **16**, 11580–11587; (d) A. Cnossen, W. R. Browne and B. L. Feringa, *Top. Curr. Chem.*, 2014, **354**, 139–162; (e) L. Osorio-Planes, C. Rodríguez-Esrich and M. A. Pericàs, *Org. Lett.*, 2014, **16**, 1704–1707.
- 5 (a) A. M. Brouwer, C. Frochot, F. G. Gatti, D. A. Leigh, L. Mottier, F. Paolucci, S. Roffia and G. W. H. Worpel, *Science*, 2001, **291**, 2124–2128; (b) V. Balzani, M. Clemente-León, A. Credi, B. Ferrer, M. Venturi, A. H. Flood and J. F. Stoddart, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 1178–1183.
- 6 J. A. Berrocal, C. Biagini, L. Mandolini and S. Di Stefano, *Angew. Chem., Int. Ed.*, 2016, **55**, 6997–7001.
- 7 (a) M. R. Wilson, J. Solà, A. Carlone, S. M. Goldup, N. Lebrasseur and D. A. Leigh, *Nature*, 2016, **534**, 235–240; (b) M. A. Watson and S. L. Cockroft, *Angew. Chem., Int. Ed.*, 2016, **55**, 1345–1349.
- 8 E. V. Anslyn and D. A. Dougherty, *Modern Physical Organic Chemistry*, University Science Books, Sausalito (CA), 2006, p. 91.
- 9 K. Takahashi, M. Takaki and R. Asami, *Org. Magn. Reson.*, 1971, **3**, 539–543.
- 10 (a) C. Bernasconi, *Adv. Phys. Org. Chem.*, 1992, **27**, 119–238; (b) F. G. Bordwell, W. J. Boyle Jr and K. C. Yee, *J. Am. Chem. Soc.*, 1970, **92**, 5926–5932.
- 11 M. T. Reetz, S. Hütte and R. Goddard, *J. Prakt. Chem.*, 1999, **341**, 297–301.



- 12 (a) J. A. Berrocal, M. M. L. Nieuwenhuizen, L. Mandolini, E. W. Meijer and S. Di Stefano, *Org. Biomol. Chem.*, 2014, **12**, 6167–6174; (b) J. A. Berrocal, L. M. Pitet, M. M. L. Nieuwenhuizen, L. Mandolini, E. W. Meijer and S. Di Stefano, *Macromolecules*, 2015, **48**, 1358–1363.
- 13 K. Okuro, M. Furuune, M. Miura and M. Nomura, *J. Org. Chem.*, 1993, **58**, 7606–7607.
- 14 H. Brunner and P. Schmidt, *Eur. J. Org. Chem.*, 2000, 2119–2133.
- 15 F. Cramer and W. Kampe, *J. Am. Chem. Soc.*, 1965, **87**, 1115–1120.

