

Cite this: *Chem. Sci.*, 2018, 9, 3672

## The development of a novel AND logic based fluorescence probe for the detection of peroxynitrite and GSH†

Adam C. Sedgwick,<sup>‡\*a</sup> Hai-Hao Han,<sup>‡b</sup> Jordan E. Gardiner,<sup>‡a</sup> Steven D. Bull,<sup>‡\*a</sup> Xiao-Peng He<sup>‡\*b</sup> and Tony D. James<sup>‡\*a</sup>

We have developed a novel AND logic based fluorescence probe for the simultaneous detection of ONOO<sup>−</sup> and GSH (GSH-PF3). The GSH-PF3 probe was synthesised over three steps starting from commercially available fluorescein. The probe was constructed by attaching the GSH reactive motif, 2,4-dinitrobenzenesulfonyl, to the previously reported boronate fluorescence based probe, PF3. GSH-PF3 produced only a small fluorescence response towards the addition of GSH or ONOO<sup>−</sup> separately. However, when the probe was exposed to both analytes, there was a significant (40-fold) fluorescence enhancement. GSH-PF3 demonstrated an excellent selectivity towards both GSH and ONOO<sup>−</sup>. In cellular imaging experiments the probe was shown to be cell permeable with no 'turn-on' response observed for the addition of either GSH or ONOO<sup>−</sup> separately. However, in the presence of both analytes, a clear fluorescence response was observed in live cells. GSH-PF3 was further able to monitor the co-existence of metabolically produced ONOO<sup>−</sup> and GSH by exogenous stimulation.

Received 13th February 2018

Accepted 10th March 2018

DOI: 10.1039/c8sc00733k

rsc.li/chemical-science

Peroxynitrite (ONOO<sup>−</sup>), is a highly reactive nitrogen species that is formed *via* the diffusion controlled reaction between superoxide anion (O<sub>2</sub><sup>•−</sup>) and nitric oxide (NO<sup>•</sup>).<sup>1,2</sup> ONOO<sup>−</sup> acts as a signalling molecule *in vivo* for a number of pathways.<sup>1,3</sup> However, ONOO<sup>−</sup> is more commonly known for its deleterious properties, causing irreversible damage to a range of biological targets such as lipids, proteins and DNA.<sup>4</sup> Therefore, ONOO<sup>−</sup> has been implicated as a key pathogenic factor for a number of diseases, which include inflammation, cancer, ischemia-reperfusion and neurodegenerative diseases.<sup>5–7</sup> Glutathione (GSH) is a natural tripeptide (γ-L-glutamyl-L-cysteinyl-glycine), that exists in the thiol reduced form (GSH) and disulphide-oxidised (GSSG) form. GSH is the predominant form, existing at millimolar concentrations in most cells.<sup>8</sup> However, GSH can be directly oxidised by ONOO<sup>−</sup> therefore acting as a cellular defence by serving as an ONOO<sup>−</sup> scavenger. Elevated levels of GSH are common in cells under oxidative stress and the

susceptibility of a cell towards ONOO<sup>−</sup> largely depends on the concentration of intracellular GSH.<sup>1,9,10</sup>

Therefore, with this work we set out to develop a fluorescence-based probe capable of monitoring the close relationship between ONOO<sup>−</sup> and GSH. Traditionally, most fluorescence probes require a single analyte to produce a fluorescence response.<sup>11–19</sup> However, in recent years a number of fluorescence based probes for dual or multi-analyte detection have been developed.<sup>20–27</sup> These types of fluorescence based probes have been used for the construction of molecular logic gates or for medical diagnostics.<sup>28</sup> AND logic based fluorescence probes require both analytes to be simultaneously present or work in tandem in order to elicit a fluorescence response. This method has a number of advantages including: being faster than serial measurements for different analytes within the same biological sample and can provide a method for monitoring bimolecular events, which may contribute to a specific disease.<sup>20</sup>

Currently, only a few reversible fluorescence based probes for the detection of ONOO<sup>−</sup> and GSH have been developed to monitor the relationship between these analytes.<sup>29,30</sup> These include a selenium based fluorescence probe, which is oxidised by ONOO<sup>−</sup> (turn "on") and reduced by GSH (turn "off"). The fluorescence of the CyPSe probe is initially quenched by a photoinduced electron transfer (PET) process. The presence of ONOO<sup>−</sup> results in the oxidation of selenium to CyPSe=O causing the fluorescence emission to be "turned on". Then in the presence of biological thiols such as cysteine and GSH, the CyPSe=O probe is reduced back to its non-fluorescent selenide

<sup>a</sup>Department of Chemistry, University of Bath, Bath, BA2 7AY, UK. E-mail: a.c.sedgwick@bath.ac.uk; s.d.bull@bath.ac.uk; t.d.james@bath.ac.uk

<sup>b</sup>Key Laboratory for Advanced Materials and Joint International Research Laboratory of Precision Chemistry and Molecular Engineering, Feringa Nobel Prize Scientist Joint Research Center, School of Chemistry and Molecular Engineering, East China University of Science and Technology, 130 Meilong Rd., Shanghai 200237, China. E-mail: xphe@ecust.edu.cn

† Electronic supplementary information (ESI) available: Additional figures, experimental section and original spectra of new compounds. See DOI: 10.1039/c8sc00733k

‡ These authors contributed equally.




form. A similar reversible NIR Tellurium-based fluorescence probe was later developed for monitoring the redox cycles between  $\text{ONOO}^-$  and GSH. This probe was successfully applied for the visualisation of the redox cycles of  $\text{ONOO}^-$  and GSH in live cells and animals.<sup>29</sup> The probes developed by Han *et al.* are turn “on” with  $\text{ONOO}^-$  and “off” with GSH. As far as we are aware, there are currently no AND logic based fluorescence probes for GSH “and”  $\text{ONOO}^-$ .

With this research, we aimed to develop a GSH “and”  $\text{ONOO}^-$  logic based fluorescence probe (Scheme 1). We identified a suitable mono-boronate fluorescein based fluorescence probe, **PF3** that had been previously developed by Chang *et al.*, shown in Scheme 2. As previously reported the reactivity of boronate based fluorescence probes with  $\text{ONOO}^-$ ,<sup>12,14,15</sup> are significantly greater than hypochlorite ( $\text{ClO}^-$ ) and  $\text{H}_2\text{O}_2$ .<sup>31</sup> Therefore, we anticipated that the attachment of a GSH reactive motif to **PF3** would produce a selective GSH- $\text{ONOO}^-$  AND logic based fluorescence probe, **GSH-PF3** (Scheme 2). **GSH-PF3** was readily synthesised in three steps. Fluorescein was triflated using *N*-phenyl bis(trifluoromethanesulfonamide) to afford fluorescein mono-triflate in good yield. Suzuki–Miyaura conditions were then carried out to provide fluorescein mono-boronate, **PF3**. The 2,4-dinitrobenzenesulfonyl unit was then attached to **PF3** using 2,4-dinitrobenzenesulfonyl chloride,  $\text{CH}_2\text{Cl}_2$  and  $\text{NEt}_3$  at 0 °C. Using these conditions **GSH-PF3** was prepared in a reasonable yield of 52%.



**Scheme 1** Design concept for AND logic based fluorogenic probe. A fluorescence dye is masked by two functional groups, which respond to two different analytes. The fluorogenic probe requires both analytes to be present or to work in tandem in order to produce a response.




**Scheme 2** Structure of the **GSH-PF3** probe and proposed sensing mechanism for the simultaneous detection of  $\text{ONOO}^-$  and GSH.

With **GSH-PF3** in hand, fluorescence experiments for the detection of GSH and  $\text{ONOO}^-$  were performed. As shown in Fig. 1, **GSH-PF3** was initially non-fluorescent and with the addition of  $\text{ONOO}^-$  (10  $\mu\text{M}$ ), a small fluorescence increase was observed. However, incremental additions of GSH resulted in a much larger increase in fluorescence intensity (>30-fold see ESI – Fig. S1†), clearly demonstrating the need for the addition of both GSH and  $\text{ONOO}^-$  in order to achieve a full ‘turn-on’ response.

In order to demonstrate that **GSH-PF3** required both GSH and  $\text{ONOO}^-$  for a complete ‘turn-on’ response, the fluorescence experiments were performed the other way around. Therefore, an excess of GSH (200  $\mu\text{M}$ ) was added to **GSH-PF3**, and the probe was incubated for 10 min. Remarkably, this only led to a small increase in fluorescence intensity and the subsequent additions of  $\text{ONOO}^-$  resulted in a large fluorescence increase (Fig. 2 and ESI – Fig. S2†). These results confirm that the probe requires both GSH and  $\text{ONOO}^-$  for a full fluorescence ‘turn-on’ response.

The selectivity of **GSH-PF3** was then evaluated against a series of amino acids (L-cysteine, L-methionine, L-tryptophan, L-serine, L-lysine, L-leucine, L-glutamic acid, L-valine, L-arginine, L-histidine and L-aspartic acid) – see ESI Fig. S3.† Unsurprisingly, the probe responded to the other sulfhydryl containing amino acid cysteine. However, the biological concentrations of cysteine are low in cells.<sup>32</sup> More importantly, **GSH-PF3** displayed an excellent selectivity against other amino acids including serine, methionine and lysine. The probe demonstrated excellent selectivity for  $\text{ONOO}^-$  over many other ROS including  $\text{H}_2\text{O}_2$  (see ESI – Fig. S4†). The excellent selectivity for both GSH and  $\text{ONOO}^-$  allowed us to evaluate **GSH-PF3** in cell imaging experiments.

The macrophage cell line – RAW264.7 (ATCC® TIB-71™; obtained from ATCC [American Type Culture Collection]) – was used for cell imaging experiments. The cells were incubated with **GSH-PF3**, followed by either treatment of SIN-1 (an  $\text{ONOO}^-$  donor)<sup>33</sup> to produce intracellular  $\text{ONOO}^-$  or by addition of



**Fig. 1** Fluorescence spectra of **GSH-PF3** (0.5  $\mu\text{M}$ ) with addition of  $\text{ONOO}^-$  (10  $\mu\text{M}$ ) (inset) followed by the addition of GSH (0–80  $\mu\text{M}$ ), 5 min wait between addition in buffer solution [52 wt% methanol] (pH = 8.21 at 25 °C). Fluorescence intensities were measured with  $\lambda_{\text{ex}}$  = 488 nm with slit widths ex slit: 5 nm and em slit: 2.5 nm.



Fig. 2 Fluorescence spectra of **GSH-PF3** (0.5  $\mu\text{M}$ ) with addition of **GSH** (200  $\mu\text{M}$ ), 10 min wait (inset), then addition of  $\text{ONOO}^-$  (0–10  $\mu\text{M}$ ) in buffer solution [52 wt% methanol] (pH = 8.21 at 25  $^\circ\text{C}$ ). Fluorescence intensities were measured with  $\lambda_{\text{ex}}$  = 488 nm with slit widths ex slit: 5 nm and em slit: 2.5 nm.

exogenous **GSH**. As shown below in Fig. 3, addition of **GSH** separately resulted in no fluorescence response, whereas  $\text{ONOO}^-$  led to a small fluorescence response in cells. This observation is believed to be due to the presence of a low levels of endogenous biological thiols in the cells, resulting in the activation of the probe's fluorescence. As predicted, treatment of cells with both **GSH** and **SIN-1** led to a clear fluorescence increase in RAW264.7 cells with **GSH-PF3**, in clear agreement with the analytical data obtained on the fluorimeter.




Fig. 3 Fluorescence imaging (a) and quantification (b) of RAW264.7 cells with **GSH-PF3** (10  $\mu\text{M}$ ) in the absence and presence of exogenously added **GSH** (50  $\mu\text{M}$ ) and/or **SIN-1** (500  $\mu\text{M}$ ). Excitation channel = 460–490 nm, emission channel filtered = 530–590 nm. Cell nuclei were stained with Hoechst 33342. Scale bar = 100  $\mu\text{m}$ . Error bars represent SD.

We then turned our attention to evaluate the ability of **GSH-PF3** for monitoring the co-existence of metabolically produced  $\text{ONOO}^-$  by lipopolysaccharide (LPS) stimulation<sup>34</sup> and **GSH** through a drug treatment.<sup>35</sup> It has been reported that LPS can elicit  $\text{ONOO}^-$  in macrophage, which is a signature of inflammation, whereas caffeic acid (CA) is commonly used to treat inflammation through the augmentation of intracellular **GSH**. Consequently, treatment with LPS, followed by the addition of increasing CA (0–100  $\mu\text{M}$ ) led to a gradual enhancement in the fluorescence intensity of **GSH-PF3** (Fig. 4). This result is due to the co-existence of  $\text{ONOO}^-$  produced by LPS stimulation as well as **GSH** elicited by CA. Interestingly, a further increase of the CA concentration (>100  $\mu\text{M}$ ) led to the suppression of probe's fluorescence, which is believed to be due to the production of an excess of **GSH**, resulting in quenching of the  $\text{ONOO}^-$  (Fig. 4). To corroborate the presence of intracellular **GSH** in macrophage, a commercial **GSH** probe was used. The result suggested that both the exogenous addition of **GSH** and stimulation of metabolic **GSH** by treatment of CA/LPS activated the commercial probe's fluorescence (Fig. S5†). Furthermore, a cell proliferation assay indicated that **GSH-PF3** is not toxic for RAW264.7 cells (Fig. S6†).



Fig. 4 Fluorescence imaging (a) and quantification (b) of RAW264.7 cells with **GSH-PF3** (10  $\mu\text{M}$ ) in the absence and presence of LPS (1  $\mu\text{g mL}^{-1}$ ), which elicits  $\text{ONOO}^-$  and increasing caffeic acid (CA, a drug that elicits endogenous **GSH** to quench  $\text{ONOO}^-$ ). Excitation channel = 460–490 nm, emission channel filtered = 530–590 nm. Cell nuclei were stained with Hoechst 33342. Scale bar = 100  $\mu\text{m}$ . Error bars represent SD.





These results clearly demonstrate the potential of AND logic based fluorescence imaging probes for optimizing drug dosage in the treatment of inflammation and various other diseases. In addition this system could be used to help monitor the treatment of Alzheimer's disease (AD), since it is known that high levels of ONOO<sup>−</sup> correlate with oxidative stress and the progression of AD, while, high levels of GSH have been implicated in AD therapy.<sup>36–39</sup> Therefore, we believe that our **GSH-PF3** probe, which can detect GSH “and” ONOO<sup>−</sup> could potentially be used to monitor cellular resistance towards ROS and map therapeutic improvements in AD victims.

In summary, **GSH-PF3** is an easy-to-prepare fluorescence based probe providing a platform for the development of other novel AND logic based fluorescence imaging probes for use in medical diagnostics. We are currently exploring the use of 6-amino/carboxyfluorescein, which we believe could provide the opportunity to attach additional fluorophores in order to develop ratiometric fluorescence sensors<sup>40</sup> or for the attachment of targeting or therapeutic units.<sup>41–43</sup>

## Conflicts of interest

No conflicts of interest.

## Acknowledgements

We would like to thank the EPSRC and the University of Bath for funding. ACS and JEG thank the EPSRC for studentships. X.-P. He thanks the National Natural Science Foundation of China (21722801 and 21572058), the Science and Technology Commission of Shanghai Municipality (15540723800), the Fundamental Research Funds for the Central Universities (222201717003) and the Shanghai Rising-Star Program (16QA1401400) for financial support. TDJ wishes to thank the Royal Society for a Wolfson Research Merit Award and ECUST for a guest professorship. The Catalysis and Sensing for our Environment (CASE) network is thanked for research exchange opportunities. NMR characterisation facilities were provided through the Chemical Characterisation and Analysis Facility (CCAF) at the University of Bath (<http://www.bath.ac.uk/ccaf>). The EPSRC UK National Mass Spectrometry Facility at Swansea University is thanked for analyses. All data supporting this study are provided as Electronic Supplementary Information (ESI) accompanying this paper.

## Notes and references

- 1 P. Pacher, J. S. Beckman and L. Liaudet, *Physiol. Rev.*, 2007, **87**, 315.
- 2 J. S. Beckman and W. H. Koppenol, *Am. J. Physiol. Cell Physiol.*, 1996, **271**, C1424.
- 3 A. Weidinger and A. V. Kozlov, *Biomolecules*, 2015, **5**, 472.
- 4 P. Ascenzi, A. di Masi, C. Sciorati and E. Clementi, *Biofactors*, 2010, **36**, 264.
- 5 H. Ischiropoulos and J. S. Beckman, *J. Clin. Invest.*, 2003, **111**, 163.
- 6 P. Sarchielli, F. Galli, A. Floridi and V. Gallai, *Amino Acids*, 2003, **25**, 427.
- 7 D. A. Wink, Y. Vodovotz, J. Laval, F. Laval, M. W. Dewhirst and J. B. Mitchell, *Carcinogenesis*, 1998, **19**, 711.
- 8 S. C. Lu, *Mol. Aspect. Med.*, 2009, **30**, 42.
- 9 K. A. Marshall, R. Reist, P. Jenner and B. Halliwell, *Free Radic. Biol. Med.*, 1999, **27**, 515.
- 10 J. P. Bolanos, S. J. R. Heales, J. M. Land and J. B. Clark, *J. Neurochem.*, 1995, **64**, 1965.
- 11 M. Li, X. M. Wu, Y. Wang, Y. S. Li, W. H. Zhu and T. D. James, *Chem. Commun.*, 2014, **50**, 1751.
- 12 S. Palanisamy, P. Y. Wu, S. C. Wu, Y. J. Chen, S. C. Tzou, C. H. Wang, C. Y. Chen and Y. M. Wang, *Biosens. Bioelectron.*, 2017, **91**, 849.
- 13 D. Cheng, Y. Pan, L. Wang, Z. B. Zeng, L. Yuan, X. B. Zhang and Y. T. Chang, *J. Am. Chem. Soc.*, 2017, **139**, 285.
- 14 A. C. Sedgwick, X. L. Sun, G. Kim, J. Yoon, S. D. Bull and T. D. James, *Chem. Commun.*, 2016, **52**, 12350.
- 15 X. Sun, Q. Xu, G. Kim, S. E. Flower, J. P. Lowe, J. Yoon, J. S. Fossey, X. Qian, S. D. Bull and T. D. James, *Chem. Sci.*, 2014, **5**, 3368.
- 16 J. Yin, Y. Kwon, D. Kim, D. Lee, G. Kim, Y. Hu, J. H. Ryu and J. Yoon, *J. Am. Chem. Soc.*, 2014, **136**, 5351.
- 17 Q. Xu, K.-A. Lee, S. Lee, K. M. Lee, W.-J. Lee and J. Yoon, *J. Am. Chem. Soc.*, 2013, **135**, 9944.
- 18 Z. Lei, X. Li, X. Luo, H. He, J. Zheng, X. Qian and Y. Yang, *Angew. Chem., Int. Ed.*, 2017, **56**, 2979.
- 19 Y. Kuriki, M. Kamiya, H. Kubo, T. Komatsu, T. Ueno, R. Tachibana, K. Hayashi, K. Hanaoka, S. Yamashita, T. Ishizawa, N. Kokudo and Y. Urano, *J. Am. Chem. Soc.*, 2018, **140**, 1767.
- 20 A. Romieu, *Org. Biomol. Chem.*, 2015, **13**, 1294.
- 21 L. Yu, S. L. Wang, K. Z. Huang, Z. G. Liu, F. Gao and W. B. Zeng, *Tetrahedron*, 2015, **71**, 4679.
- 22 G. C. Van de Bittner, C. R. Bertozzi and C. J. Chang, *J. Am. Chem. Soc.*, 2013, **135**, 1783.
- 23 L. Yuan, W. Y. Lin, Y. N. Xie, B. Chen and S. S. Zhu, *J. Am. Chem. Soc.*, 2012, **134**, 1305.
- 24 C. G. Dai, X. L. Liu, X. J. Du, Y. Zhang and Q. H. Song, *ACS Sens.*, 2016, **1**, 888.
- 25 S. Debieu and A. Romieu, *Org. Biomol. Chem.*, 2015, **13**, 10348.
- 26 L. Yi, L. Wei, R. Y. Wang, C. Y. Zhang, J. Zhang, T. W. Tan and Z. Xi, *Chem.–Eur. J.*, 2015, **21**, 15167.
- 27 C. Y. Ang, S. Y. Tan, S. J. Wu, Q. Y. Qu, M. F. E. Wong, Z. Luo, P. Z. Li, S. T. Selvan and Y. L. Zhao, *J. Mater. Chem. C*, 2016, **4**, 2761.
- 28 S. Erbas-Cakmak, S. Kolemen, A. C. Sedgwick, T. Gunnlaugsson, T. D. James, J. Yoon and E. U. Akkaya, *Chem. Soc. Rev.*, 2018, DOI: 10.1039/c7cs00491e.
- 29 F. B. Yu, P. Li, B. S. Wang and K. L. Han, *J. Am. Chem. Soc.*, 2013, **135**, 7674.
- 30 F. B. A. Yu, P. Li, G. Y. Li, G. J. Zhao, T. S. Chu and K. L. Han, *J. Am. Chem. Soc.*, 2011, **133**, 11030.
- 31 A. Sikora, J. Zielonka, M. Lopez, J. Joseph and B. Kalyanaraman, *Free Radic. Biol. Med.*, 2009, **47**, 1401.



- 32 G. Y. Wu, Y. Z. Fang, S. Yang, J. R. Lupton and N. D. Turner, *J. Nutr.*, 2004, **134**, 489.
- 33 X. Li, R. R. Tao, L. J. Hong, J. Cheng, Q. Jiang, Y. M. Lu, M. H. Liao, W. F. Ye, N. N. Lu, F. Han, Y. Z. Hu and Y. H. Hu, *J. Am. Chem. Soc.*, 2015, **137**, 12296.
- 34 A. Vazquez-Torres, J. Jones-Carson and E. Balish, *Infect. Immun.*, 1996, **64**, 3127.
- 35 M. Y. Moridani, H. Scobie, A. Jamshidzadeh, P. Salehi and P. J. O'Brien, *Drug Metab. Dispos.*, 2001, **29**, 1432.
- 36 M. A. Smith, P. L. R. Harris, L. M. Sayre, J. S. Beckman and G. Perry, *J. Neurosci.*, 1997, **17**, 2653.
- 37 E. Rojas-Gutierrez, G. Munoz-Arenas, S. Trevino, B. Espinosa, R. Chavez, K. Rojas, G. Flores, A. Diaz and J. Guevara, *Synapse*, 2017, **71**, e21990.
- 38 C. B. Pocernich and D. A. Butterfield, *Biochim. Biophys. Acta, Mol. Basis Dis.*, 2012, **1822**, 625.
- 39 I. Cacciatore, L. Baldassarre, E. Fornasari, A. Mollica and F. Pinnen, *Oxid. Med. Cell. Longevity*, 2012, 240146.
- 40 A. E. Albers, V. S. Okreglak and C. J. Chang, *J. Am. Chem. Soc.*, 2006, **128**, 9640.
- 41 M. H. Lee, J. Y. Kim, J. H. Han, S. Bhuniya, J. L. Sessler, C. Kang and J. S. Kim, *J. Am. Chem. Soc.*, 2012, **134**, 12668.
- 42 S. Bhuniya, S. Maiti, E. J. Kim, H. Lee, J. L. Sessler, K. S. Hong and J. S. Kim, *Angew. Chem., Int. Ed.*, 2014, **53**, 4469.
- 43 X.-P. He and H. Tian, *Chem*, 2018, **4**, 246.

