Chemical Science

EDGE ARTICLE

Check for updates

Cite this: Chem. Sci., 2018, 9, 5556

Received 25th April 2018 Accepted 26th May 2018 DOI: 10.1039/c8sc01879k

rsc.li/chemical-science

Introduction

Reactive sulfur species (RSS) are a family of endogenous sulfurcontaining molecules in biological systems related to a series of signaling transduction processes.1-3 Among RSS, hydrogen sulfide (H₂S) has been studied for many years as the third most important endogenous gasotransmitter after CO and NO and has been supposed to be involved in regulating neuromodulation,4 cytoprotection,5 vascular tone regulation,6 oxygen sensing,7 inflammatory regulation,8 and cell growth.9 However, recent studies have implied that hydrogen polysulfides (H_2S_n , n > 1) might act as the real signaling mediator instead of H₂S.¹⁰⁻¹² For example, the conversion of cysteine residues (-SH) to persulfides (-S-SH) through S-sulfhydration was originally ascribed to H₂S, but H₂S itself can hardly induce S-sulfhydration.¹³⁻¹⁶ Also, studies have shown that H_2S_n can compensate for physiological functions that H₂S cannot achieve.¹⁷⁻¹⁹ There are still many unaddressed arguments associated with the biological role of H_2S_n .

A fluorescent τ -probe: quantitative imaging of ultra-trace endogenous hydrogen polysulfide in cells and *in vivo*⁺

View Article Online

View Journal | View Issue

Fan Yang, He Gao, Shan-Shan Li, Rui-Bing An, Xiao-Yang Sun, Bin Kang,* Jing-Juan Xu[®]* and Hong-Yuan Chen

Hydrogen sulfide (H₂S) has been recognized as an important endogenous gasotransmitter associated with biological signaling transduction. However, recent biological studies implied that the H₂S-related cellular signaling might actually be mediated by hydrogen polysulfides (H₂S_n, n > 1), not H₂S itself. Unraveling such a mystery strongly demanded the quantification of endogenous H₂S_n in living systems. However, endogenous H₂S_n has been undetectable thus far, due to its extremely low concentration within cells. Herein, we demonstrated a strategy to detect ultra-trace endogenous H₂S_n *via* a fluorescent τ -probe, through changes of fluorescence lifetime instead of fluorescence intensity. This τ -probe exhibited an ultrasensitive response to H₂S_n, bringing about the lowest value of the detection limit (2 nM) and a lower limit of quantification (10 nM) to date. With such merits, we quantified and mapped endogenous H₂S_n in cells and *in vivo* may have a significant implication for future research on the role of H₂S_n in biology. The methodology of the τ -probe established here might provide a general insight into the design and application of any fluorescent probes, beyond the limit of utilizing fluorescence intensity.

A good start to clarify such a conundrum might begin with the quantification of endogenous H_2S_n in living systems. Such a task seems straightforward, but actually remains a great challenge. The H₂S level in biological systems has been reported from 17 nM to 3 µM.²⁰ As a downstream product of H₂S, endogenous H_2S_n was conjectured to be <100 nM or even less. Unfortunately, it is still difficult to detect H_2S_n at such low concentration. UV-vis spectroscopy and mass spectrometry (MS) have been applied to investigate H_2S_n ^{21,22} but they were limited by poor sensitivity and insufficient operability in organisms. Fluorescent probes were feasible for H_2S_n detection in living systems, owing to their high sensitivity and compatibility with bioimaging facilities.^{23,24} In 2014, the Xian group reported the first H₂S_n-specific probes (DSP) exploiting a 2-fluoro-5nitrobenzoic ester template,²⁵ and several subsequent H₂S_n fluorescent probes were designed afterwards.26-33 The DSP-type probes showed satisfactory sensitivity and selectivity for H_2S_n ³² however, they would also react with biothiols, resulting in a slow response and low yield. To settle this issue, several other recognition groups were developed, including nucleophilic ring-opening reaction,34 a-substituted acrylate ester,35 and phenyl 2-(benzoylthio)benzoate formation.36-38 Studies have also been performed to improve the signal-to-noise ratio through the red shift of the wavelength,30,32,38,39 two-photon excitation^{31,40} or ratio detection,^{29,41,42} and also to achieve simultaneous detection of H_2S_n together with H_2S ,³⁷ glutathione,⁴³ or superoxide anions.⁴⁴ Although many efforts have

State Key Laboratory of Analytical Chemistry for Life Science, Collaborative Innovation Center of Chemistry for Life Sciences, School of Chemistry and Chemical Engineering, Nanjing University, 210023, China. E-mail: binkang@nju.edu.cn; xujj@nju.edu.cn

[†] Electronic supplementary information (ESI) available: Detailed synthesis procedure and design strategies of the τ -probe, schematic diagram of the fluorescence lifetime microscope, characterization of the τ -probe, and additional figures. See DOI: 10.1039/c8sc01879k

been done, all these probes can still detect H_2S_n only through external stimulation^{28,32,35} or H_2S_n addition,^{36,45} but not endogenous H_2S_n in native cells. In 2015, the Xian group reported a new H_2S_n -probe (PSP), showing the lowest detection limit extrapolated from theory (3 nM), and an actual working range from 0.25 μM to 20 $\mu M.^{36}$ Although the lower limit of quantification was as low as 0.25 μ M, endogenous H₂S_n in native cells still cannot be detected. It was expected that the lower limit of quantification must reach tens of nM, and the detection limit must reach a few nM, so as to detect native endogenous H_2S_n . Such a goal seemed hard to achieve by using intensity-based fluorescent probes, either from theoretical deduction or from the best result reported so far. In addition to fluorescence intensity, fluorescence lifetime is another fundamental photophysical parameter associated with any fluorescence emission. Different from fluorescence intensity that depends on emitter numbers, fluorescence lifetime mainly depends on the instinct photophysical characteristic of each emitter.46 Fluorescence lifetime imaging has been commonly applied as a powerful tool of biology studies.47 Current time-resolved techniques are able to distinguish lifetime changes as low as ps-fs levels.48

In spite of the important progress made in studying H_2S_n , the quantification of endogenous H_2S_n in native biological systems still remain unsolved. The most sensitive fluorescence intensity probes are currently unable to achieve such a low detection limit. In this work, we developed an alternative strategy of using a fluorescent τ -probe to detect H_2S_n through changes of fluorescent lifetime rather than fluorescence intensity. The detection limit of the τ -probe was as low as 2 nM, and the actual working range was from 10 nM to 5 µM, which allows us to detect ultra-trace endogenous H_2S_n in cells and in zebrafish. The concentration of endogenous H_2S_n in a HeLa cell was distributed between ~ 0 and 120 nM, with an averaged value about \sim 60–70 nM. The concentration distribution in zebrafish slightly broadened to \sim 0–160 nM, but still has a comparable averaged concentration of \sim 70-80 nM with the value of cells. To the best of our knowledge, this is the first time to detect endogenous H_2S_n in a native biological system. These results provided fundamental quantitative information about endogenous H_2S_n in cells and *in vivo* and have a significant implication for deeply understanding the role of H_2S_n in biology. Meanwhile, the methodology of the τ -probe established here might provide a general insight into the design and application of any analytical strategy exploiting fluorescent probes, to remedy the disadvantage of fluorescence intensity probes that are insensitive at extremely low concentration.

Results and discussion

Concept and design of the fluorescent τ -probe

The design of the τ -probe was inspired by Xian group's PSP-3 probe,³⁶ but its structure was modified to maximize the sensitivity. Detailed synthesis protocols and structural characterization are provided in the ESI (Scheme S1, Fig. S1–S3†). Products from the reaction between the τ -probe and H₂S_n are shown in Fig. S4.† Compared to the PSP-3 probe with two recognition groups that would consume two H₂S_n per probe, our τ -probe has

only one recognition group, which consumes one H_2S_n per probe. As shown in Fig. 1a, this probe initially showed a "dark" state with a weak fluorescence emission (Fig. 1b) and a short lifetime $\tau_{dark} = 0.75$ ns (Fig. 1d). When reacted with H_2S_n (Scheme S2†), fluorescein was released to a "bright" state with a strong fluorescence emission (Fig. 1c) and a long lifetime $\tau_{bright} = 3.47$ ns (Fig. 1e). This probe is a typical "turn-on"-type fluorescent probe. Here we presented that, in addition to fluorescence intensity, the fluorescence lifetime could also be utilized for quantitative analysis.

To prove the concept of the τ -probe, we studied the response of fluorescence intensity and lifetime to the concentration of H₂S_n. Theoretically, for an emitter with unchanged fluorescence emission characteristics, the fluorescence intensity depends on the numbers of emitters, and the fluorescence lifetime depends on the instinct photophysical properties of each emitter. However, for systems mixed with two emitters with distinct fluorescence emission characteristics, the total value of



Fig. 1 Concept and design of the τ -probe. (a) Reaction with H₂S_n switched the τ -probe from "dark" to "bright" states with distinct fluorescence characteristics. (b–e) Fluorescence intensity and lifetime of the "dark" state τ -probe (b, d) and "bright" state τ -probe (c, e), respectively.

fluorescence intensity and lifetime in such mixed systems was a result contributed by the two emitters. Particularly, for our τ -probe that has "dark" and "bright" states, the total weighted value of steady-state fluorescence intensity (I_{total}) consisted of "bright" state fluorescence intensity (I_{bright}) and "dark" state fluorescence intensity (I_{total} could be calculated by:

$$I_{\text{total}} = I_{\text{bright}} + I_{\text{dark}} \tag{1}$$

Since fluorescence intensity is proportional to the number of fluorescent molecules (*N*) and their emission cross section (σ), the fluorescence intensity (*I*) can also be expressed as:

$$I = fc V N_{\rm a} \sigma \tag{2}$$

where *f* is a proportional constant related to the incident photon flux. c_{bright} and c_{dark} are the concentrations of "bright" and "dark" state τ -probes, *V* is the volume of the solution, and N_{a} is the Avogadro constant. Combined with eqn (2), I_{total} could be expressed as:

$$I_{\text{total}} = fc_{\text{bright}} V N_a \sigma_{\text{bright}} + fc_{\text{dark}} V N_a \sigma_{\text{dark}}$$
(3)

Because the parameters f, V and N_a are constants for a given system, they could be totally represented by another constant γ . Then eqn (3) could be described as:

$$I_{\text{total}} = \gamma c_{\text{bright}} \sigma_{\text{bright}} + \gamma c_{\text{dark}} \sigma_{\text{dark}} \tag{4}$$

When the emitters were excited by a pulsed light source, for instance in fluorescence lifetime measurement, the transient fluorescence intensity at time t ($I_{total(t)}$) could be calculated by:

$$I_{\text{total}(t)} = I_{\text{bright}(t)} + I_{\text{dark}(t)} = A e^{-\frac{t}{\tau_{\text{bright}}}} + B e^{-\frac{t}{\tau_{\text{dark}}}}$$
(5)

where *A* and *B* are the compensation coefficients of fluorescence intensity at the bright state and dark state, respectively. τ_{bright} and τ_{dark} are the fluorescence lifetime of "bright" and "dark" state τ -probes. Therefore, the total weighted mean of fluorescence lifetime (τ_{total}) could be calculated by:

$$\tau_{\text{total}} = \frac{A}{A+B} \tau_{\text{bright}} + \frac{B}{A+B} \tau_{\text{dark}}$$
(6)

A and B could be described as:

$$A \propto c_{\text{bright}} \sigma_{\text{bright}}$$
 (7)

$$B \propto c_{\text{dark}} \sigma_{\text{dark}}$$
 (8)

Then the total fluorescence lifetime (τ_{total}) could be described as:

$$\tau_{\text{total}} = \frac{(c_{\text{bright}})\sigma_{\text{bright}}}{(c_{\text{bright}})\sigma_{\text{bright}} + (c_{\text{dark}})\sigma_{\text{dark}}} \tau_{\text{bright}} + \frac{(c_{\text{dark}})\sigma_{\text{dark}}}{(c_{\text{bright}})\sigma_{\text{bright}} + (c_{\text{dark}})\sigma_{\text{dark}}} \tau_{\text{dark}}$$
(9)

According to eqn (4) and (9), we find that $I_{(total)}$ and $\tau_{(total)}$ are only related to the concentration of fluorescent molecules c_{bright} and c_{dark} , owing to the fact that σ_{bright} and σ_{dark} are fixed values in a given system ($\sigma_{\text{bright}} \gg \sigma_{\text{dark}}$). Consequently, if we use c_{B0} and c_{D0} which represent the original concentration of "bright" and "dark" state τ -probes, the concentration (*c*) of H₂S_n reacted with the τ -probe could be calculated from the total fluorescence intensity ($I_{\text{total}(c)}$) or fluorescence lifetime ($\tau_{\text{total}(c)}$):

$$I_{\text{total}(c)} = \gamma(c_{B0} + c)\sigma_{\text{bright}} + \gamma(c_{D0} - c)\sigma_{\text{dark}}$$
(10)
$$\tau_{\text{total}(c)} = \frac{(c_{B0} + c)\sigma_{\text{bright}}}{(c_{B0} + c)\sigma_{\text{bright}} + (c_{D0} - c)\sigma_{\text{dark}}} \tau_{\text{bright}}$$
$$+ \frac{(c_{D0} - c)\sigma_{\text{dark}}}{(c_{B0} + c)\sigma_{\text{bright}} + (c_{D0} - c)\sigma_{\text{dark}}} \tau_{\text{dark}}$$
(11)

Fluorescence response to H_2S_n : intensity versus lifetime

Theoretical response curves of $I_{\text{total}(c)}$ and $\tau_{\text{total}(c)}$ to H_2S_n concentration (*c*) could be obtained by substituting experimental data into eqn (10) and (11): $c_{\text{B0}} = 0 \ \mu\text{M}$, $c_{\text{D0}} = 10 \ \mu\text{M}$, $\sigma_{\text{bright}} = 20\sigma_{\text{dark}}$, $\tau_{\text{bright}} = 3.47 \text{ ns}$, and $\tau_{\text{dark}} = 0.75 \text{ ns}$. Giving consideration to both accuracy and sensitivity, the concentration of the τ -probe was chosen as 10 μ M in the following experiments (Fig. S5†). Experimental response curves were obtained by measuring changes of fluorescence intensity and lifetime by the addition of 0–25 μ M Na₂S₄ (commercial H₂S_n donor), as adopted previously.^{31,35,41} Theoretical curves matched experimental data well for either fluorescence intensity (Fig. 2a)



Fig. 2 Response characteristics: intensity *versus* lifetime. (a and b) Theoretical and experimental response curves of the fluorescence intensity (a) and lifetime (b) of the τ -probe to Na₂S₄ (as a H₂S_n donor). (c) Response of fluorescence intensity and lifetime at extremely low concentration. (d) Comparison of the detection limit and working range between the current τ -probe and some intensity-based H₂S_n probes reported previously: 1-current work, 2-(ref. 36), 3-(ref. 39), 4-(ref. 37), 5-(ref. 43), 6-(ref. 25), 7-(ref. 35), and 8-(ref. 42). All experiments were repeated at least three times. Error bars represent standard deviation.

Edge Article

or lifetime (Fig. 2b). The response features of $I_{\text{total}(c)}$ and $\tau_{\text{total}(c)}$ were totally different. Itotal(c) exhibited a nearly linear response in a wide range of concentration (c). However, $\tau_{total(c)}$ showed a non-linear response, which is much more sensitive at low concentration. For instance, when the concentration of Na₂S₄ adds to 0.2 μ M, $\tau_{total(c)}$ increased 0.93 ns, and the rate of change was about 34.19% compared with the value of the "bright" state. At the same condition, $I_{total(c)}$ is almost unchanged and the variation is as low as 0.09% (Fig. 2c). According to the temporal resolution and noise level of the instrument (~ 10 ps), the detection limit of the τ -probe was about ~2 nM, and the experimental working range was from ~ 10 nM to 5 μ M. Compared to H_2S_n probes reported previously (Fig. 2d), our τ probe reached the lowest value of the detection limit and a lower limit of quantification to date. Note that even though $\tau_{\text{total(c)}}$ is sensitive at low concentration, it does not have as broad a response range as $I_{total(c)}$ does. Therefore, these two methods with high sensitivity or wide response range would complement each other, rather than replace each other.

Time-dependent response and photostability of the τ -probe to Na₂S₄ are shown in Fig. S6 and S7.† Fluorescence intensity reached almost maximum within 5 min, and thus a reaction time of 20 min was employed subsequently to ensure complete reactions. The effects of pH were investigated; the τ -probe worked effectively in the pH range of 7–8 (Fig. S8†). To obtain a working curve, varying concentrations of Na₂S₄ (0.01–25 μ M) were added to the τ -probe (10 μ M). Fluorescence intensity almost linearly correlated with Na₂S₄ in a broad range of 0.5–10 μ M (Fig. S9†), but changes at 0.5 μ M and below were hard to distinguish. However, fluorescence lifetime still has a sensitive response in the range of 10 nM to 0.2 μ M (Fig. S10†). The correlation of fluorescence lifetime (τ) with the concentration of H₂S_n (*c*) in the range of 0–0.2 μ M could be well fitted to a linear equation ($R^2 = 0.99267$):

$$\tau = 0.77 + 4.60c \ (0 < c < 0.2 \ \mu \text{M}) \tag{12}$$

Test of selectivity and specificity

To test the selectivity of the τ -probe to H_2S_n , the τ -probe was treated with a series of reactive sulfur species (RSS) including glutathione (GSH), cysteine (Cys), homocysteine (Hcy), glutathione disulfide (GSSG), H_2S , $S_2O_3^{2-}$, SO_3^{2-} , SO_4^{2-} , and CH_3 -SSSCH₃. A negligible change of fluorescence lifetime was observed for any of these compounds (Fig. 3a). Only Na_2S_4 generated a significant change (column blank). The interference of these RSS with the τ -probe upon addition of Na₂S₄ was tested and minute interference was found (column red). We also validated whether the τ -probe could sense H₂S_n formation from H_2S and reactive oxygen species (ROS) in situ. The τ -probe did not show a response (Fig. 3b) to H_2S (in the form of Na_2S) in the presence of common ROS including hydrogen peroxide (H_2O_2) , superoxide (O_2^-) , hydroxyl radical (OH), and singlet oxygen (${}^{1}O_{2}$). Only the ClO⁻/H₂S system showed an obvious response. This result was consistent with previous reports that H_2S_n can be derived from H_2S and ClO^- , but not other ROS.³⁶ All



Fig. 3 Selectivity and specificity of the τ -probe to H_2S_n . (a) Fluorescence lifetime changes caused by various reactive sulfur species, with or without Na_2S_4 (as a H_2S_n donor). (b) Fluorescence lifetime changes caused by various reactive oxygen species, with or without Na_2S (as a H_2S donor). See experimental details in the Experimental section. All experiments were repeated at least three times. Error bars represent standard deviation.

these results indicated the high sensitivity and high selectivity of the τ -probe to H₂S_n, suggesting that the τ -probe could be used to detect endogenous H₂S_n in native biological systems.



Fig. 4 Quantifying endogenous H_2S_n in a HeLa cell suspension. (a) Fluorescence lifetime change of the τ -probe in negative control cells $(H_2S_n(-))$ incubated with *N*-ethylmaleimide (NEM) to eliminate H_2S_n (left column), native cells (H_2S_n-cell) without any treatment (middle column) and positive control cells $(H_2S_n(+))$ incubated with Na_2S_4 (right column). (b) Fluorescence lifetime change of the τ -probe in native HeLa cells $(H_2S_n = 65 \text{ nM})$ and cells incubated with 0.065 μ M (1×) or 0.13 μ M (2×) additional Na_2S_4 . All experiments were repeated at least three times. Error bars represent standard deviation.

Quantifying endogenous H_2S_n within cells and zebrafish

We then demonstrated that our τ -probe could quantify H_2S_n in cells by using a suspension of HeLa cells. For positive control cells incubated with 20 μ M Na₂S₄ (as a H₂S_n donor), the fluorescence lifetime increased dramatically ($\Delta \tau = 2.70$ ns). For negative control cells incubated with 100 µM N-ethylmaleimide (NEM) to eliminate the endogenous H_2S_n , the fluorescence lifetime remained almost unchanged ($\Delta \tau = 0.05$ ns) compared to the "dark" state τ -probe (Fig. 4a). With the positive and negative controls, endogenous H_2S_n in native HeLa cells was obtained by substituting fluorescence lifetime into the working curve (eqn (3)). The averaged H_2S_n concentration in native HeLa cells is about $\sim 0.065 \ \mu M$ (65 nM). We then further confirmed this by adding 0.065 μ M (1×) and 0.130 μ M (2×) additional Na₂S₄ to cell samples, and the fluorescence lifetime changed linearly according to the working curve. Cell viability assay demonstrated that the τ -probe has nearly no cytotoxicity to cells (Fig. S11[†]).

With results in the cell suspension, we then quantified endogenous H_2S_n within single HeLa cells, utilizing both



Fig. 5 Quantitative mapping of H_2S_n in single HeLa cells. (a and b) Dark field imaging of a native HeLa cell (a) and HeLa cells treated with Na_2S_4 (b). (c and d) Fluorescence lifetime imaging of a native HeLa cell (c) and HeLa cells treated with Na_2S_4 (d). (e) Map of H_2S_n concentration in a native HeLa cell. (f) Histogram of H_2S_n concentration in the native HeLa cell showing in (e). (g) Averaged H_2S_n concentration within three different native HeLa cells. Scale bars: 10 μ m.



Fig. 6 Quantitative mapping of H_2S_n in zebrafish. (a and b) Dark field imaging of untreated zebrafish (a) and zebrafish treated with Na_2S_4 (b). (c and d) Fluorescence lifetime imaging of untreated zebrafish (c) and zebrafish treated with Na_2S_4 (d). (e) Map of H_2S_n concentration in untreated zebrafish. (f) Histogram of H_2S_n concentration in the untreated zebrafish showing in (e). Scale bars: 150 µm.

fluorescence intensity and lifetime. Fluorescence imaging was performed on a confocal microscope. As reported previously,35,36 fluorescence was observed only in positive control cells (+20 µM Na_2S_4), but not in native HeLa cells (Fig. S12⁺). Fluorescence lifetime imaging was carried out on a time-gated transient microscope (see details in the Method and Scheme S3 in the ESI[†]). Dark field imaging was utilized to locate cells (Fig. 5a and b). As expected, fluorescence lifetime changes were observed in native HeLa cells (Fig. 5c and d). Then the map and histogram of H_2S_n within a HeLa cell were obtained by applying the working curve to fluorescence lifetime imaging (Fig. 5e and f). The H_2S_n level in three different HeLa cells was determined to be about 60-70 nM (Fig. 5g), which is comparable with the value of the cell suspension (\sim 65 nM). It is interesting to note that even the intracellular concentration of H_2S_n covers a wide range of 0-120 nM (Fig. 5f), and the averaged values of different individual cells were quite close to each other (Fig. 5g). This result is sort of out of expectation since studies have shown heterogeneity between individual cells in the concentration of biomolecules.⁴⁹ Further in-depth studies are definitely needed to get a better understanding about the role of H_2S_n in cells. Afterwards, we mapped H_2S_n distribution in zebrafish (Fig. 6). The endogenous H_2S_n within zebrafish was mainly localized in its abdominal region, and the distribution covers a wide range of 0–160 nM, with an average value of about \sim 70–80 nM. It was also found that endogenous H_2S_n within zebrafish was hard to observe in the fluorescence intensity imaging (Fig. S13[†]). These results demonstrated that the τ -probe could be an ultrasensitive tool for studying H_2S_n in living systems.

Conclusions

In summary, we developed an alternative strategy to detect H_2S_n through a fluorescence lifetime probe (termed: τ -probe) rather than fluorescence intensity. The detection limit of the τ -probe

was as low as 2 nM, and the actual working range was from 10 nM to 5 μ M, which allows us to detect ultra-trace endogenous H₂S_n in cells and in zebrafish. The endogenous H₂S_n in native HeLa cells was distributed between ~0 and 120 nM, with an average value of about ~60–70 nM. The value in zebrafish slightly broadened to ~0–160 nM, but still has a comparable average level ~70–80 nM. To the best of our knowledge, this is the first time to detect endogenous H₂S_n in a native biological system. This information has a significant implication for deeply understanding the role of H₂S_n in biology. The concept of the τ -probe provided a general insight into the application of fluorescent probes, beyond the limit of fluorescence intensity.

Experimental section

Reagents

Fluorescein, methyl iodide (MeI), potassium carbonate (K₂CO₃), dimethyl formamide (DMF), sodium hydroxide (NaOH), methyl alcohol (MeOH), hydrochloric acid (HCl), dichloromethane (CH₂Cl₂), thiosalicylic acid, benzoic acid, 1-ethyl-3-(3-(dimethylamino)propyl)carbodiimide (EDC), 4-dimethylaminopyridine (DMAP), dimethyl sulfoxide (DMSO), glutathione (GSH), cysteine (Cys), homocysteine (Hcy), glutathione disulfide (GSSG), Na₂S, Na₂S₂O₃, Na₂SO₃, Na₂SO₄, CH₃SSSCH₃, hydrogen peroxide (H₂O₂), sodium hypochlorite (NaClO), KO₂, FeSO₄, and ethyl 3-aminobenzoate methanesulfonate were purchased from Sigma-Aldrich (St. Louis, MO, U.S.A.). High glucose Dulbecco's Modified Eagle Medium (DMEM), fetal bovine serum (FBS), and trypsin were purchased from Invitrogen (Carlsbad, U.S.A.). The cell cytotoxicity assay kit, human cervical cancer (HeLa) cells, and phosphate buffer solution (PBS, 10 mM, pH = 7.4) were purchased from KeyGEN Biotech. Co (Nanjing, China). Zebrafish were purchased from the Model Animal Research Center of Nanjing University. Ultra-pure water from a Millipore Milli-Q (18 M Ω cm⁻¹) was used in the experiments. All other reagents were of analytical reagent grade.

Apparatus

UV-vis absorption spectra were recorded using a UV-vis spectrophotometer (Nanodrop-2000C, Nanodrop, USA). The fluorescence emission spectra were obtained on a Shimadzu fluorescence S-3 spectrophotometer (RF-5301PC, Shimadzu Co., Japan). The fluorescence lifetime spectra were obtained on a FLS 980 spectrophotometer (Edinburgh instruments, U.K.). Confocal fluorescence images of cells were acquired with TCS SP8 confocal microscopy (Leica, Germany). The cell viability assay was performed using a Thermo Scientific Varioskan Flash (Thermo Fisher Scientific, American). The τ -Probe was purified by Ultimate 3000 high performance liquid chromatography (Thermo Fisher Scientific, American). Fluorescence lifetime imaging was achieved by using instruments built in our laboratory.

Synthesis of the τ -probe

The synthesis procedure of the τ -probe is shown in Scheme S1.† CH₂Cl₂ (20 mL) was added to a mixture of 3-O-

methylfluorescein⁵⁰ (69.2 mg, 0.2 mmol), 2-(benzoylthio)benzoic acid^{37,51} (78 mg, 0.3 mmol), EDC (58 mg, 0.3 mmol) and DMAP (12.2 mg, 0.1 mmol) at room temperature and the mixture was stirred for 5 h. Then the solvent was evaporated under reduced pressure and the resulting residue was subjected to high performance liquid chromatography for purification. The τ -probe was obtained as a yellow solid. ¹H-NMR and ¹³C-NMR spectra of τ -probe are shown in Fig. S1 and S2,[†] respectively. ¹H NMR (400 MHz, CD₃Cl) δ 8.23 (dd, J = 1.6, 8.0 Hz, 1H), 8.03-8.01 (m, 3H), 7.72-7.57 (m, 6H), 7.49-7.47 (m, 2H), 7.18 (d, *J* = 2.4 Hz, 1H), 7.16 (d, *J* = 7.6 Hz, 1H), 6.88 (dd, *J* = 2.4, 8.8 Hz, 1H), 6.80 (d, *J* = 8.8 Hz, 1H), 6.77 (d, *J* = 2.4 Hz, 1H), 6.69 (d, *J* = 8.8 Hz, 1H), 6.62 (dd, J = 2.4, 8.8 Hz, 1H), 3.82 (s, 3H). ¹³C NMR (100 MHz, CD₃Cl) δ 189.44, 169.80, 164.27, 161.51, 153.05, 152.28, 151.98, 151.92, 137.33, 136.36, 135.26, 133.90, 133.47, 132.81, 131.66, 129.92, 129.72, 129.11, 129.00, 128.93, 128.83, 127.59, 126.41, 125.16, 124.05, 117.53, 116.74, 111.99, 110.72, 110.49, 100.90, 82.97, 55.62. LC-MS (ESI⁺): m/z C₃₅H₂₃O₇S⁺, calcd 587.1168, found [M⁺] 587.1169.

Fluorescence lifetime measurement

The fluorescence lifetime measurements were performed on a FLS 980 spectrophotometer (Edinburgh instruments, U.K.). The instrument is based on time-correlated single photon counting (TCSPC). Photons are counted in a time window which sweeps across the full time range following each excitation pulse, creating a histogram of counts *versus* time. The data quality of the resulting histogram is improved by adding the data of repeated sweeps. A 488 nm picosecond laser (~3 ps, 5.5 MHz) was used to pump the fluorescence, and the collection time window is 0–50 ns in the wavelength range of 500–650 nm, with a minimum time per channel = 305 fs. The time jitter in the delay generation system is about ~10 ps.

General procedure for spectral measurements

A stock solution of the τ -probe was prepared at 1 mM in DMSO. A PBS (10 mM, pH 7.4) solution with DMSO as the cosolvent (DMSO/PBS = 1 : 9, v/v) was used for performing all spectroscopic measurements. The fluorescence signal was recorded with the excitation wavelength at 488 nm and the emission wavelength range from 500 to 650 nm. The maximum fluorescence emission was obtained at 515 nm. Glutathione (GSH, 5 mM), cysteine (Cys, 1 mM), homocysteine (Hcy, 100 μM), glutathione disulfide (GSSG, 100 µM), Na₂S (200 µM), Na₂S₂O₃ (100 µM), Na₂SO₃ (100 µM), Na₂SO₄ (100 µM), CH₃SSSCH₃ (100 µM), hydrogen peroxide (H₂O₂, 200 μ M), hypochlorite (ClO⁻, 50 μ M), superoxide (O₂⁻, 50 μ M), hydroxyl radical ('OH, 50 μ M), singlet oxygen (¹O₂, 50 μ M), H_2O_2 (200 μ M) + Na_2S (50 μ M), H_2O_2 (200 μ M) + Na_2S (100 μ M), $ClO^{-}(50 \ \mu M) + Na_2S(100 \ \mu M), O_2^{-}(50 \ \mu M) + Na_2S(100 \ \mu M), OH$ $(50 \ \mu\text{M}) + \text{Na}_2\text{S} (100 \ \mu\text{M})$, and $^1\text{O}_2 (50 \ \mu\text{M}) + \text{Na}_2\text{S} (100 \ \mu\text{M})$ were all dissolved in prepared solutions for various testing species. The preparation of reactive oxygen species was according to the literature.³⁶ Na₂S₄ (commercial H₂S_n donor) was dissolved in PBS solution, to prepare a sample solution (1 mM). A volume of 10 µL of τ -probe stock solution was added to 1 mL volumes of various analyte buffer/DMSO solutions ($H_2O/DMSO = 9: 1, v/v$), and 20

Chemical Science

 μ L of Na₂S₄ stock solution was employed in some of the experiments. After vigorous shaking, the resulting solutions were kept at room temperature for 30 min, and their fluorescence signals were collected.

Cell culture

HeLa cells were cultured in high glucose DMEM containing 10% fetal bovine serum (FBS), 100 IU mL⁻¹ penicillin and 100 IU mL⁻¹ streptomycin at 37 °C in a 5% CO₂–95% air incubator MCO-15AC (Sanyo, Tokyo, Japan).

Cell viability assay

Firstly, HeLa cells were seeded in 96-well plates (3000 cells per well) and maintained at 37 °C in a humidified atmosphere (95% air and 5% CO₂) for 24 h. After 24 h incubation, the cells were washed with PBS and then cultured in a DMSO/PBS solution (pH 7.4, 1 : 9 v/v) containing the τ -probe with various concentrations. After 30 min, the cells were washed with PBS and incubated in a fresh medium for an additional 24 h. Then, according to the manufacturer's instructions, 50 µL MTT solution was added to each well and incubated for 4 h at 37 °C. After removing the medium, 150 µL DMSO was added to solubilize the blue-colored tetrazolium, the plates were gently shaken for 5 min, and the optical density values at 550 nm were monitored using a Thermo Scientific Varioskan Flash. The cell viability was set as 100% in control cells.

Intracellular H_2S_n detection by using a fluorescence spectrometer

Cells were subjected to different treatments: the first group was incubated with 100 μ M *N*-ethylmaleimide (NEM) for 15 min, the second group was untreated, the third group was incubated with 20 μ M Na₂S₄ for 30 min, and the forth group was incubated with 0.065 μ M Na₂S₄ for 30 min, while the fifth group was incubated with 0.13 μ M Na₂S₄ for 30 min. After washing with PBS, the cells were cultured in a DMSO/PBS solution (pH 7.4, 1 : 9 v/v) containing a 10 μ M τ -probe for 30 min. Then fluorescence lifetime was acquired after washing with PBS (pH = 7.4). The cells were excited at 488 nm and the emission was collected at 515 nm.

Confocal fluorescence images of HeLa cells upon incubation

HeLa cells were digested and seeded on confocal dishes for 24 h at 37 °C. The cells were subjected to different treatments. The first group was incubated with 100 μ M *N*-ethylmaleimide (NEM) for 15 min, and the second group was untreated, while the third group was incubated with 20 μ M Na₂S₄ for 30 min. After washing with PBS, the cells were cultured in a DMSO/PBS solution (pH 7.4, 1:9 v/v) containing a 10 μ M τ -probe for 30 min. Then confocal images were acquired after washing with PBS (pH = 7.4). The cells were excited at 488 nm and the emission was collected from 505 to 550 nm.

Optical configuration of fluorescence lifetime microscopy

The homemade fluorescence lifetime microscopy system was made up of a 488 nm nanosecond pulse laser, an inverted microscope and a delay generator. The pulse laser (20 Hz, \sim 5 ns, Quanta-Ray INDI, spectra-physics, USA) acted as the excitation light source and was focused on the sample through a beam expander. The fluorescence intensity of the sample was collected by using a 20× objective and recorded with an ICCD camera (iStar CCD 334, Andor Technology Ltd., UK), with an optical gate width of \sim 2 ns. To avoid the influence from the excitation light source, a 500 nm long-pass filter was added before the ICCD camera. In addition, the recorded delay was provided by a delay generator (DG645, Stanford Research Systems, USA).

Fluorescence lifetime images of HeLa cells

HeLa cells were digested and seeded on confocal dishes for 24 h at 37 °C. The cells were washed with PBS and then cultured in a DMSO/PBS solution (pH 7.4, 1:9 v/v) containing a 10 μ M τ -probe for 30 min, and then washed with PBS and subjected to different treatments. The first group was untreated, while the second group was incubated with 20 µM Na2S4 for 30 min. Then fluorescence lifetime images were acquired after washing with PBS (pH = 7.4). The cells were excited by using a pulse laser $(\sim 5 \text{ ns})$ with a wavelength of 488 nm and the fluorescence emission was collected by using an ICCD camera through a 500 nm long-pass filter. Every group of fluorescence decaying images were captured after a delay time of 0, 2, ..., 26 ns. The map of fluorescence lifetime was obtained through fitting the emission intensities of each pixel at different delay times. The time baseline was corrected using a FLS 980 spectrophotometer by measuring the same sample, and the response curves of the laser pulse and instrument were deconvoluted from the decay curve of fluorescence emission to obtain the final lifetime. Finally, the intracellular concentration distribution of H_2S_n was obtained by substituting the fluorescence lifetime into the working equation.

Fluorescence lifetime images of zebrafish

Zebrafish were anesthetized in a PBS (pH = 7.4) solution containing 0.125% ethyl 3-aminobenzoate methanesulfonate. Firstly, the experimental group was incubated with 20 µM Na₂S₄ for 30 min, and then washed with PBS (pH = 7.4). Afterwards, the experimental group and the control group were both incubated with the τ -probe (10 μ M) for 30 min in DMSO/PBS (pH 7.4, 1:9 v/v) solution. Then fluorescence lifetime images were acquired after washing with PBS (pH = 7.4). The fluorescence lifetime of the τ -probe and intracellular concentration distribution of H_2S_n in zebrafish were obtained by the same procedure as that of the cells. The fish was too big to fit in the view field of the microscope, so the image was made up of three pieces. The experimental conditions for these three images were exactly the same. All animal work was approved by the Animal Care Committee of Nanjing University in accordance with Institutional Animal Care and Use Committee guidelines.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Edge Article

This work was supported by the National Natural Science Foundation of China (21327902, 21535003, and 21675081), State Key Laboratory of Analytical Chemistry for Life Science (5431ZZXM1715) and program B for Outstanding PhD candidate of Nanjing University (201702B052).

Notes and references

- 1 J. M. Fukuto, S. J. Carrington, D. J. Tantillo, J. G. Harrison, L. J. Ignarro, B. A. Freeman, A. Chen and D. A. Wink, *Chem. Res. Toxicol.*, 2012, 25, 769–793.
- 2 R. Wang, Physiol. Rev., 2012, 92, 791-896.
- 3 X. Wang, J. Sun, W. Zhang, X. Ma, J. Lv and B. Tang, *Chem. Sci.*, 2013, 4, 2551–2556.
- 4 J. R. Austgen, G. E. Hermann, H. A. Dantzler, R. C. Rogers and D. D. Kline, *J. Neurophysiol.*, 2011, **106**, 1822–1832.
- 5 H. Kimura, N. Shibuya and Y. Kimura, *Antioxid. Redox* Signaling, 2012, **17**, 45–57.
- 6 J. W. Calvert, S. Jha, S. Gundewar, J. W. Elrod,
 A. Ramachandran, C. B. Pattillo, C. G. Kevil and D. J. Lefer, *Circ. Res.*, 2009, **105**, 365–374.
- 7 K. R. Olson and N. L. Whitfield, *Antioxid. Redox Signaling*, 2010, **12**, 1219–1234.
- 8 L. Li, M. Bhatia, Y. Z. Zhu, Y. C. Zhu, R. D. Ramnath, Z. J. Wang, F. B. M. Anuar, M. Whiteman, M. Salto-Tellez and P. K. Moore, *FASEB J.*, 2005, **19**, 1196–1198.
- 9 R. Baskar and J. Bian, Eur. J. Pharmacol., 2011, 656, 5-9.
- 10 H. Kimura, Nitric Oxide, 2015, 47, S6.
- 11 C. Jacob, T. Burkholz, S. Griffin, L. Faulstich and E. C. Estevam, *Curr. Org. Chem.*, 2016, **20**, 211–217.
- 12 K. Ono, T. Akaike, T. Sawa, Y. Kumagai, D. A. Wink, D. J. Tantillo, A. J. Hobbs, P. Nagy, M. Xian, J. Lin and J. M. Fukuto, *Free Radical Biol. Med.*, 2014, 77, 82–94.
- 13 T. V. Mishanina, M. Libiad and R. Banerjee, *Nat. Chem. Biol.*, 2015, **11**, 457.
- 14 P. K. Yadav, M. Martinov, V. Vitvitsky, J. Seravalli, R. Wedmann, M. R. Filipovic and R. Banerjee, *J. Am. Chem. Soc.*, 2016, **138**, 289–299.
- 15 N. Krishnan, C. Fu, D. J. Pappin and N. K. Tonks, *Sci. Signaling*, 2011, 4, ra86.
- 16 H. Kimura, Proc. Jpn. Acad., Ser. B, 2015, 91, 131-159.
- 17 C. E. Paulsen and K. S. Carroll, *Chem. Rev.*, 2013, **113**, 4633–4679.
- 18 H. Kimura, Nitric Oxide, 2014, 41, 4–10.
- 19 H. Kimura, Neurochem. Int., 2013, 63, 492-497.
- 20 K. Hideo, Antioxid. Redox Signaling, 2017, 27, 619-621.
- 21 R. Greiner, Z. Pálinkás, K. Bäsell, D. Becher, H. Antelmann,
 P. Nagy and T. P. Dick, *Antioxid. Redox Signaling*, 2013, 19, 1749–1765.
- 22 Y. Kimura, Y. Mikami, K. Osumi, M. Tsugane, J.-i. Oka and H. Kimura, *Nitric Oxide*, 2014, **39**, S39.
- 23 K. P. Carter, A. M. Young and A. E. Palmer, *Chem. Rev.*, 2014, **114**, 4564–4601.
- 24 X. Li, X. Gao, W. Shi and H. Ma, Chem. Rev., 2014, 114, 590-659.

- 25 C. Liu, W. Chen, W. Shi, B. Peng, Y. Zhao, H. Ma and M. Xian, J. Am. Chem. Soc., 2014, 136, 7257–7260.
- 26 A. Basso, E. Bruno, M. Parise and M. Pepe, *Analyst*, 2015, **140**, 3766–3772.
- 27 M. Gao, F. Yu, H. Chen and L. Chen, Anal. Chem., 2015, 2015, 3631–3638.
- 28 L. Zeng, S. Chen, T. Xia, W. Hu, C. Li and Z. Liu, *Anal. Chem.*, 2015, **87**, 3004–3010.
- 29 Q. Han, Z. Mou, H. Wang, X. Tang, Z. Dong, L. Wang, X. Dong and W. Liu, *Anal. Chem.*, 2016, 88, 7206–7212.
- 30 Y. Huang, F. Yu, J. Wang and L. Chen, *Anal. Chem.*, 2016, 88, 4122–4129.
- 31 J. Zhang, X.-Y. Zhu, X.-X. Hu, H.-W. Liu, J. Li, L. L. Feng, X. Yin, X.-B. Zhang and W. Tan, *Anal. Chem.*, 2016, 88, 11892–11899.
- 32 J. Ma, J. Fan, H. Li, Q. Yao, F. Xu, J. Wang and X. Peng, J. Mater. Chem. B, 2017, 5, 2574–2579.
- 33 J. Zhang, X. Hao, W. Sang and Q. Yan, *Small*, 2017, 13, 1701601.
- 34 W. Chen, E. W. Rosser, D. Zhang, W. Shi, Y. Li, W.-J. Dong, H. Ma, D. Hu and M. Xian, Org. Lett., 2015, 17, 2776–2779.
- 35 J. Guo, S. Yang, C. Guo, Q. Zeng, Z. Qing, Z. Cao, J. Li and R. Yang, Anal. Chem., 2018, 90, 881–887.
- 36 W. Chen, E. W. Rosser, M. Tetsuro, P. Armo, A. Takaaki and M. Xian, *Angew. Chem.*, 2015, 54, 13961–13965.
- 37 W. Chen, A. Pacheco, Y. Takano, J. J. Day, K. Hanaoka and M. Xian, *Angew. Chem.*, *Int. Ed.*, 2016, 55, 9993–9996.
- 38 Y. Fang, W. Chen, W. Shi, H. Li, M. Xian and H. Ma, *Chem. Commun.*, 2017, **53**, 8759–8762.
- 39 K.-B. Li, F.-Z. Chen, Q.-H. Yin, S. Zhang, W. Shi and D.-M. Han, *Sens. Actuators, B*, 2018, **254**, 222–226.
- 40 H. Shang, H. Chen, Y. Tang, R. Guo and W. Lin, Sens. Actuators, B, 2016, 230, 773-778.
- 41 Y. Takano, K. Hanaoka, K. Shimamoto, R. Miyamoto, T. Komatsu, T. Ueno, T. Terai, H. Kimura, T. Nagano and Y. Urano, *Chem. Commun.*, 2017, 53, 1064–1067.
- 42 R. Kawagoe, I. Takashima, S. Uchinomiya and A. Ojida, *Chem. Sci.*, 2017, **8**, 1134–1140.
- 43 W. Chen, X. Yue, H. Zhang, W. Li, L. Zhang, Q. Xiao, C. Huang, J. Sheng and X. Song, *Anal. Chem.*, 2017, 89, 12984–12991.
- 44 F. Yu, M. Gao, M. Li and L. Chen, Biomaterials, 2015, 63, 93-101.
- 45 W. Chen, X. Yue, J. Sheng, W. Li, L. Zhang, W. Su, C. Huang and X. Song, *Sens. Actuators, B*, 2018, **258**, 125–132.
- 46 B. Valeur and M. N. Berberan-Santos, *Molecular Fluorescence: Principles and Applications*, Wiley, 2012.
- 47 Z. Gryczynski, I. Gryczynski and J. R. Lakowicz, in *Molecular Imaging*, ed. R. N. Day, American Physiological Society, San Diego, 2005, pp. 21–56.
- 48 R. M. C. Ammasi Periasamy, *FLIM Microscopy in Biology and Medicine*, Chapman and Hall/CRC, New York, 2009.
- 49 R. Pan, M. Xu, D. Jiang, J. D. Burgess and H.-Y. Chen, *Proc. Natl. Acad. Sci. U. S. A.*, 2016, **113**, 11436–11440.
- 50 L. Mugherli, O. N. Burchak, F. Chatelain and M. Y. Balakirev, *Bioorg. Med. Chem. Lett.*, 2006, **16**, 4488–4491.
- 51 R. J. Bahde, D. H. Appella and W. C. Trenkle, *Tetrahedron Lett.*, 2011, **52**, 4103–4105.