



Cite this: *Chem. Sci.*, 2018, 9, 7218

All publication charges for this article have been paid for by the Royal Society of Chemistry

# The hydrogen atom transfer reactivity of sulfinic acids†

Markus Griesser, Jean-Philippe R. Chauvin and Derek A. Pratt\*

Sulfinic acids ( $\text{RSO}_2\text{H}$ ) have a reputation for being difficult reagents due to their facile autoxidation. Nevertheless, they have recently been employed as key reagents in a variety of useful radical chain reactions. To account for this paradox and enable further development of radical reactions employing sulfinic acids, we have characterized the thermodynamics and kinetics of their H-atom transfer reactions for the first time. The O–H bond dissociation enthalpy (BDE) of sulfinic acids was determined by radical equilibration to be  $\sim 78 \text{ kcal mol}^{-1}$ ; roughly halfway between the RS–H BDE in thiols ( $\sim 87 \text{ kcal mol}^{-1}$ ) and RSO–H BDE in sulfenic acids ( $\sim 70 \text{ kcal mol}^{-1}$ ). Regardless, RSH, RSOH and  $\text{RSO}_2\text{H}$  have relatively similar inherent H-atom transfer reactivity to alkyl radicals ( $\sim 10^6 \text{ M}^{-1} \text{ s}^{-1}$ ). Counter-intuitively, the trend in rate constants with more reactive alkoxy radicals follows the reaction energetics:  $\sim 10^8 \text{ M}^{-1} \text{ s}^{-1}$  for  $\text{RSO}_2\text{H}$ , midway between thiols ( $\sim 10^7 \text{ M}^{-1} \text{ s}^{-1}$ ) and sulfenic acids ( $\sim 10^9 \text{ M}^{-1} \text{ s}^{-1}$ ). Importantly, since sulfinic and sulfenic acids are very strong H-bond donors ( $\alpha_2^{\text{H}} \sim 0.63$  and  $0.55$ , respectively), their reactivity is greatly attenuated in H-bond accepting solvents, whereas the reactivity of thiols is largely solvent-independent. Efforts to measure rate constants for the reactions of sulfinic acids with alkylperoxy radicals were unsuccessful. Computations predict these reactions to be surprisingly slow;  $\sim 1000$ -times slower than for thiols and  $\sim 10\,000\,000$ -times slower than for sulfenic acids. On the other hand, the reaction of sulfinic acids with sulfonylperoxy radicals – which propagate sulfinic acid autoxidation – is predicted to be almost diffusion-controlled. In fact, the rate-determining step in sulfinic acid autoxidation, and the reason they can be used for productive chemistry, is the relatively slow reaction of propagating sulfonyl radicals with  $\text{O}_2$  ( $\sim 10^6 \text{ M}^{-1} \text{ s}^{-1}$ ).

Received 31st May 2018  
Accepted 30th July 2018

DOI: 10.1039/c8sc02400f  
rsc.li/chemical-science


## Introduction

Sulfinic acids ( $\text{RSO}_2\text{H}$ ) are trivalent sulfur oxyacids isoelectronic with carboxylic acids. Although they are less prominent than the thiols from which they are derived, and the sulfonic acids to which they are oxidized, they are useful Brønsted acids and as precursors to various sulfonate derivatives. One-electron oxidation of sulfinic acids (or their conjugated bases) yields sulfonyl radicals,<sup>1,2</sup> important intermediates in industrial processes (e.g. sulfoxidation to produce surfactants)<sup>3</sup> and synthetic transformations where they often give way to sulfonyl radicals are most commonly generated from sulfonyl halides,<sup>5</sup> pseudohalides<sup>8</sup> or azides<sup>9</sup> by photolysis or atom/group transfer reactions as part of chain reactions.

Despite the fact that sulfinic acids are widely considered to be difficult to work with due to their autoxidizability, the direct

use of sulfinic acids in organic synthesis has recently gained popularity.<sup>12–21</sup> Particularly noteworthy are Lei's reports of the use of sulfinic acids for sulfonylation of alkenes<sup>12</sup> and alkynes,<sup>13</sup> and Nicewicz's use of sulfinic acids as catalytic H-atom donors in photocatalytic anti-Markovnikov alkene functionalizations

### Lei's aerobic sulfinylation of alkenes and alkynes (2013)



### Nicewicz's photocatalytic anti-Markovnikov alkene hydroacyloxylation (2013)




Fig. 1 Examples of recently reported transformations employing sulfinic acids for group transfer/radical additions.

Department of Chemistry and Biomolecular Sciences, University of Ottawa, Ottawa, Ontario, Canada K1N 6N5. E-mail: dpratt@uottawa.ca

† Electronic supplementary information (ESI) available: Synthesis and characterization data, further autoxidation data, LFP results, calculations of REqEPR and hydrogen bonding, optimized geometries and energies for computational results. See DOI: 10.1039/c8sc02400f



(*e.g.* Fig. 1).<sup>15</sup> Central to these radical chain reactions are chain-propagating steps which involve H-atom transfer (HAT) from a sulfinic acid to a peroxy radical (top, Fig. 1), to a peroxy radical and alkoxy radical (middle, Fig. 1) or alkyl radical (bottom, Fig. 1). Rate constants for these key reactions are unknown;<sup>22</sup> data which would greatly enable further development of reactions utilizing sulfinic acids for organic synthesis.

Likewise, surprisingly little is known of the stability of sulfonyl radicals relative to radicals derived from other commonly utilized reagents for addition and/or H-atom transfer in organic synthesis, such as thiols. The stability of a given radical (*e.g.* X') relative to another (*e.g.* Y') is generally estimated by comparing the X–H bond dissociation enthalpy (BDE) to that of Y–H. No definitive sulfinic acid O–H BDE has been reported. To the best of our knowledge, the first suggestion was made in 1971 (111 kcal mol<sup>-1</sup>),<sup>23</sup> but was shortly thereafter revised to be at most 104 kcal mol<sup>-1</sup>.<sup>24</sup> Benson later estimated that the phenylsulfonyl radical should have a similar stability to the benzyl radical (and therefore a BDE of *ca.* 90 kcal mol<sup>-1</sup>).<sup>25</sup> A subsequent estimate put the BDE at 73 kcal mol<sup>-1</sup>,<sup>26</sup> and a recent calculation suggested 77 kcal mol<sup>-1</sup>.<sup>27</sup> Given the central importance of the RSO<sub>2</sub> ⇌ R' + SO<sub>2</sub> equilibrium to the chemistry of sulfonyl radicals, knowledge of the inherent stability of RSO<sub>2</sub> is highly desirable.

Herein we provide the first experimental determination of the O–H BDE in sulfinic acids, and report on the kinetics of their H-atom transfer reactions with alkyl, alkoxy, and peroxy radicals. We compare these results to corresponding data for the structurally-related sulfenic acids, which have among the weakest O–H bonds<sup>28</sup> and some of the fastest HAT reactions<sup>29,30</sup> ever reported. We also study the solvent effects on these reactions, and demonstrate that sulfinic (and sulfenic) acid(s) engage in strong H-bonding interactions that slow HAT.

Our results provide key information for the development of synthetic methodology employing sulfonyl radicals and/or sulfinic acids and offer unique insights on the role of secondary orbital interactions in the reactions of peroxy radicals, which are quite significant for sulfenic acids and virtually non-existent for sulfinic acids.

## Results

Kinetic and thermochemical experiments were carried out with phenylsulfinic acid (PhSO<sub>2</sub>H) (generated from its commercially-available sodium sulfinate salt) and/or 9-triptycenesulfinic acid (TrptSO<sub>2</sub>H), which enables direct comparison with the analogous 9-triptycenesulfenic acid (TrptSOH).<sup>28,31,74</sup> The 9-triptycenesulfinic acid was prepared as described in the ESI,† in a manner similar to that we recently described for the synthesis of a fluorinated analog.<sup>32</sup>

### O–H bond dissociation enthalpies of sulfinic acids

The radical equilibration EPR (REqEPR) technique<sup>33</sup> was employed to establish a reliable O–H BDE for sulfinic acids. The approach, depicted in Fig. 2A, relies on the equilibration

of persistent radicals – one derived from a precursor whose X–H BDE is known (the reference compound) and one from the compound for which the Y–H BDE is desired. The difference in the X–H and Y–H BDEs can be estimated directly from the equilibrium constant determined by double integration of the EPR signals of the two equilibrated radicals X' and Y', assuming the entropy of the H-atom exchange is negligible.

Previous EPR studies of sulfonyl radicals<sup>24,34,35</sup> have been challenging owing to their rapid disproportionation (*e.g.*  $k = 8 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$  for *p*-tosylsulfonyl)<sup>35</sup> to sulfinyl and sulfonyloxy radicals. As such, it was hoped that the triptycene substituent in TrptSO<sub>2</sub>H would impart sufficient persistence to the sulfonyl radical to enable the REqEPR experiment. Indeed, the room temperature photolysis of a solution of 9-triptycenesulfinic acid containing *t*-BuOO*t*-Bu yielded a broad signal which could be ascribed to the 9-triptycenesulfonyl radical (Fig. 2B). Within seconds, a significantly sharper signal appeared at slightly lower field. Assuming that the latter signal corresponded to the 9-triptycenesulfinyl radical, which we had previously characterized to have  $g = 2.0114$ ,<sup>28</sup> the former signal was centered at  $g = 2.0053$  in good agreement with previously reported values for sulfonyl radicals.<sup>24,36</sup>

To identify an appropriate reference compound for the REqEPR experiment, CBS-QB3 calculations were carried out to predict the O–H BDEs of each of the phenylsulfinic acid (77.2 kcal mol<sup>-1</sup>) and 9-triptycenesulfinic acid (76.7 kcal mol<sup>-1</sup>).<sup>37</sup> We also calculated an O–H BDE of 71.0 kcal mol<sup>-1</sup> for 9-triptycenesulfinic acid using the same methodology, which is in excellent agreement with the experimental value of 71.9 kcal mol<sup>-1</sup> we previously determined using the REqEPR technique,<sup>28,38</sup> suggesting that oxidation of the sulfur atom has a relatively modest (~5 kcal mol<sup>-1</sup>) effect on the O–H BDE. *N*-Methylbenzohydroxamic acid (NMBHA) was chosen as the reference compound since its O–H BDE (78 kcal mol<sup>-1</sup>)<sup>39</sup> is similar to the predicted values for the sulfinic acids and the *N*-oxyl radical produced upon H-atom abstraction therefrom is persistent (Fig. 2C).

Photolysis of a solution of 9-triptycenesulfinic acid and NMBHA containing *t*-BuOO*t*-Bu yielded spectra consistent with the superposition of the sulfonyl and *N*-oxyl radical spectra (Fig. 2D). The equilibrium constant was readily determined from these data, but estimation of the sulfinic acid O–H BDE first required correction of the equilibrium constant for the H-bonding interactions between the sulfinic acid/NMBHA and the *t*-BuOH used to help solubilize them in benzene (10% v/v) – see ESI† for the details. The resultant value of 77.6 kcal mol<sup>-1</sup> is in excellent agreement with the CBS-QB3 calculations, and confirms that oxidation of the sulfur atom in a sulfenic acid raises the O–H BDE by 5 to 6 kcal mol<sup>-1</sup>. Analogous experiments were carried out with phenylsulfinic acid in benzene, which yielded an O–H BDE of 78.3 kcal mol<sup>-1</sup>, again consistent with the predictions by CBS-QB3. By comparison, alkylthiols and arylthiols have S–H BDEs of 87 and 79 kcal mol<sup>-1</sup>, respectively.






Fig. 2 (A) The equilibration of the sulfonyl and N-oxyl radicals derived from TrptSO<sub>2</sub>H/PhSO<sub>2</sub>H and N-methylbenzhydroxamic acid (NMBHA) enables the derivation of an O–H BDE for the sulfinic acid. (B) EPR spectrum obtained during the irradiation of a mixture of TrptSO<sub>2</sub>H and di-tert-butylperoxide in benzene at 298 K. (C) EPR spectrum of the nitroxide derived from NMBHA. (D) Representative EPR spectrum of the equilibrated radical mixture. (E and F) CBS-QB3-calculated and experimental BDEs for TrptSO<sub>2</sub>H and PhSO<sub>2</sub>H.

### Reactions with alkyl radicals

The kinetics of H-atom transfer from PhSO<sub>2</sub>H and TrptSO<sub>2</sub>H to alkyl radicals were determined using the radical clock approach employing the 1,2-aryl migration in the (naphthyl)neophyl radical as the reference reaction ( $k_r = 1.4 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$  at 25 °C).<sup>40</sup> The radicals were generated by photolysis of the corresponding Barton (PTOC) ester, and the reduced and rearranged products were quantified by GC/MS as a function of sulfinic acid concentration to yield the H-atom transfer rate constant (Fig. 3A). Initial experiments in THF yielded  $k_{\text{H}} = (5.2 \pm 0.7) \times 10^4$  and  $(8.7 \pm 1.8) \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$  for PhSO<sub>2</sub>H and TrptSO<sub>2</sub>H, respectively (Fig. 3B) – the order of reactivity being consistent with the slightly weaker O–H bond in TrptSO<sub>2</sub>H compared to PhSO<sub>2</sub>H. Given the acidity of the O–H bond in sulfinic acids, we anticipated a significant solvent effect on the H-atom transfer kinetics and therefore sought to determine rate constants in a range of H-bond accepting solvents. Unfortunately, the limited solubility of the sulfinic acids at the concentrations necessary to afford reliable kinetics by the competition method precluded such an endeavor.

Therefore, we also performed direct kinetic experiments by laser flash photolysis. Cumyl radicals were generated by the photolysis of dicumylketone with the 308 nm emission of a nanosecond-pulsed XeCl excimer laser. The rate of decay of the 330 nm absorption of the cumyl radicals was determined as


a function of sulfinic acid concentration to yield the H-atom transfer rate constant (Fig. 3C). Initial experiments in benzene yielded a rate constant of  $(1.0 \pm 0.2) \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$  for PhSO<sub>2</sub>H (Fig. 3D) – consistent with a substantial kinetic solvent effect when comparing to the data in THF. Additional experiments were carried out in chlorobenzene and anisole, which revealed a dramatic drop in the HAT rate constant with increasing hydrogen bond basicity (Fig. 3G). Plotting the data according to the Ingold–Abraham equation<sup>41</sup> that relates H-atom transfer kinetics to solvent H-bond basicity ( $\beta_2^{\text{H}}$ ),

$$\log k_{\text{inh}}^{\text{S}} = -8.32\alpha_2^{\text{H}}\beta_2^{\text{H}} + \log k_{\text{inh}}^{\text{O}}$$

suggests an H-bond acidity of the sulfinic acid of  $\alpha_2^{\text{H}} = 0.61$ . The deviation of the data points obtained by competition kinetics (empty symbols) from the correlation comprised of data points obtained by LFP is consistent with the increased reactivity of primary over tertiary alkyl radicals (*ca.* 10-fold).

Corresponding experiments carried out with the sulfinic acid (in isoctane, chlorobenzene, benzene, and anisole) yielded rate constants that were systematically *ca.* 3-fold greater than those of the sulfinic acids (*e.g.*  $(2.9 \pm 0.4) \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$  for TrptSOH in benzene). Plotting these data according to the Ingold–Abraham equation yields  $\alpha_2^{\text{H}} = 0.57$  for the sulfinic acid, in excellent agreement with our previous report ( $\alpha_2^{\text{H}} = 0.54$ ).<sup>31</sup> The modest difference in the H-atom transfer kinetics of





**Fig. 3** (A) Photolytic generation of the 2-methyl-2-(2-naphthyl)-1-propyl radical and its use as a radical clock to obtain the rate constant for HAT from sulfenic and sulfinic acids to alkyl radicals. (B) Ratio of reduced ( $R'H$ ) to rearranged and reduced ( $R''H$ ) products as a function of the concentration of  $\text{TrpSO}_2\text{H}$  (red) or  $\text{PhSO}_2\text{H}$  (black) during photocleavage of the radical precursor in THF at 25 °C. (C) Photolytic generation of cumyl radicals from dicumylketone. (D) Dependence of the pseudo-first order rate on the concentration of  $\text{PhSO}_2\text{H}$  (red) and  $\text{TrpSOH}$  (black) upon photolysis of dicumylketone in  $\text{PhCl}$  at 25 °C; inset: example decay of the cumyl radical. (E and F) Calculated (CBS-QB3) transition state structures and associated free energy barriers, estimated rate constants and reaction free energies for HAT between a model alkyl radical ( $t\text{-Bu}^\bullet$ ) and either  $\text{PhSO}_2\text{H}$  or  $t\text{-BuSOH}$  (model for  $\text{TrpSOH}$ ). (G) The HAT rate constants for  $\text{PhSO}_2\text{H}$  (red) and  $\text{TrpSOH}$  (black), determined via LFP (full symbol) and radical clock (empty symbol) plotted as a function of medium  $\beta_2^H$ .

sulfinic and sulfenic acids is surprising given the *ca.* 5–6 kcal mol<sup>-1</sup> difference in the experimental O–H BDEs, and may reflect a more favorable polarization of the H-atom transfer TS in the case of the sulfinic acid compared to the sulfenic acid (consistent with their  $\text{p}K_{\text{a}}$ s of ~1.9 and 12.5, respectively).<sup>32</sup>

The HAT transition states (TS) for the reactions of representative sulfinic and sulfenic acids are shown in Fig. 3E and F. The results are fully consistent with the experimental trends. That is, CBS-QB3 predicts a barrier of  $\Delta G^\ddagger = 10.3$  kcal mol<sup>-1</sup> for  $\text{PhSO}_2\text{H}$  (corresponding to  $k = 4.1 \times 10^6$  M<sup>-1</sup> s<sup>-1</sup>) and  $\Delta G^\ddagger = 8.7$  kcal mol<sup>-1</sup> for  $t\text{-BuSOH}$  (corresponding to  $k = 6.0 \times 10^7$  M<sup>-1</sup> s<sup>-1</sup>). Again, the difference in the kinetics (*ca.* 15-fold) is relatively modest given the significant difference in the driving force ( $\Delta\Delta G^\circ = 9.1$  kcal mol<sup>-1</sup>).


### Reactions with alkoxy radicals

The reaction of sulfinic acids with *t*-butoxy radicals has been used to generate sulfonyl radicals for low-temperature EPR studies<sup>24</sup> (as above for our REqEPR experiments), but the kinetics of this reaction have never been reported.<sup>42</sup> To provide insight to this reactivity, we chose cumyloxy radicals as model species since they possess an absorption at 485 nm. The cumyloxy radicals were generated from dicumylperoxide *via*

laser flash photolysis, and their decay monitored as a function of sulfinic acid concentration to obtain the desired rate constants. In addition, kinetics for the reactions of the sulfinic acids with the triplet state of benzophenone (monitored at 530 nm) were also determined (Fig. 4A). The benzophenone triplet state has a much higher extinction coefficient than the cumyloxy radical, greatly improving the signal quality. The results are comparable to those obtained with cumyloxy radicals since the  $n \rightarrow \pi^*$  state of the carbonyl has a similar electronic structure to alkoxy radicals. Representative results from measurements of the reactivity of  $\text{PhSO}_2\text{H}$  with the benzophenone triplet ( $^3\text{BP}$ ) are shown in Fig. 4C. Interestingly, the pseudo first order rate constants from the decay traces yielded a non-linear relationship with sulfinic acid concentration. This suggested that self-association of the sulfinic acid in H-bonded dimers may retard the rate by precluding access to the labile H-atom. Indeed, the data could be easily fit to a non-linear equation accounting for the contribution of the competing equilibrium that defines the amount of sulfinic acid available for reaction (Fig. 4D). Doing so for the data in Fig. 4C yielded  $k_{\text{H}} = (1.7 \pm 0.3) \times 10^9$  M<sup>-1</sup> s<sup>-1</sup> and  $K_{\text{d}} = (6.4 \pm 2.2) \times 10^{-3}$  M. Corresponding data obtained with cumyloxy radicals under otherwise identical conditions was *ca.* 2-fold slower, yielding  $k_{\text{H}} = (6.9 \pm 1.3) \times 10^8$  M<sup>-1</sup> s<sup>-1</sup>.







**Fig. 4** (A) Excitation of benzophenone to its triplet state and photolytic generation of cumyloxy radicals from dicumylperoxide. (B) The HAT rate constants for PhSO<sub>2</sub>H (red) and TrptSOH (black) with cumyloxy radicals plotted as a function of medium  $\beta_2^H$ . (C) Dependence of the pseudo-first order rate on the concentration of PhSO<sub>2</sub>H upon excitation of BP in benzene at 25 °C and non-linear fit; inset: example decay of <sup>3</sup>BP. (D) Contribution of sulfenic acid dimerization to the observed HAT rate constant. (E) The HAT rate constants for PhSO<sub>2</sub>H (black), TrptSO<sub>2</sub>H (red) and TrptSOH (blue), with <sup>3</sup>BP, plotted as a function of medium  $\beta_2^H$  (F and G) calculated (CBS-QB3) transition state structures and associated free energy barrier, estimated rate constant and reaction free energy for HAT between a model alkoxy radical (*t*-BuO<sup>•</sup>) and PhSO<sub>2</sub>H and *t*-BuSOH.

Given the retardation of the H-atom transfer from PhSO<sub>2</sub>H to <sup>3</sup>BP and cumyloxy by self-association, and the foregoing results with alkyl radicals, it was expected that the H-atom transfer kinetics would again be highly solvent dependent. Therefore, our investigations were expanded to include more (CH<sub>3</sub>CN, EtOAc) and less (PhCl) H-bond accepting solvents, the results of which are summarized in Fig. 4E. Plotting these data according to the Ingold–Abraham equation yields  $\alpha_2^H = 0.65$ , in good agreement with the value of 0.61 derived from the alkyl radical kinetic data. The rate constant for the reaction of TrptSO<sub>2</sub>H with <sup>3</sup>BP was also determined for comparison. In CH<sub>3</sub>CN, where self-association does not contribute,  $k_H = (1.8 \pm 0.1) \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ , a factor of  $\sim 3$  greater than that measured for PhSO<sub>2</sub>H ( $k_H = 6.5 \pm 0.2) \times 10^7 \text{ M}^{-1} \text{ s}^{-1}$ , consistent with its slightly lower O–H BDE, and implying that the triptycene moiety imparts little to no steric effect on its fast HAT activity.

For comparison, the kinetics of HAT from TrptSOH to <sup>3</sup>BP and cumyloxy were also determined. In benzene,  $k_H = (2.3 \pm 0.1) \times 10^9$  and  $(1.4 \pm 0.2) \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$  for <sup>3</sup>BP and cumyloxy, respectively. Consistent with the foregoing observations with the sulfinic acids, and trends in HAT reactions, in general, the rate constant for HAT to <sup>3</sup>BP is slightly greater than for cumyloxy. These kinetics were also explored in different solvents, revealing a significantly smaller effect than for the sulfinic acids (see Fig. 4B and E as well as the ESI<sup>†</sup>) – a surprising result given

the similarity in H-bond acidity of the sulfenic and sulfinic acids (*vide supra*). In fact, when the rate constants for reactions with <sup>3</sup>BP and cumyloxy are plotted against the solvent H-bond basicity, the Ingold–Abraham equation predicts  $\alpha_2^H = 0.23$  and 0.26, respectively – less than half the value measured directly by IR spectroscopy ( $\alpha_2^H = 0.54$ )<sup>31</sup> or determined above from the kinetics of reactions with alkyl radicals ( $\alpha_2^H = 0.57$ ). We surmise that a competing electron transfer reaction is enabled by the more polar (H-bond accepting) solvents.<sup>43</sup>

The CBS-QB3-calculated TS structures for HAT from PhSO<sub>2</sub>H and *t*-BuSOH to a model alkoxy radical (*t*-butyloxy) are shown in Fig. 4F and G. Consistent with experiment, the calculated free energy barriers are very small;  $\Delta G^\ddagger = 6.6$  and 4.7 kcal mol<sup>-1</sup> for PhSO<sub>2</sub>H and *t*-BuSOH, respectively, corresponding to  $k = 2.4 \times 10^9$  and  $5.3 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$ , respectively. Obviously, the exergonicity of these reactions is so great ( $\Delta G^0 = -29.2$  and  $-38.3 \text{ kcal mol}^{-1}$ , respectively) that the kinetics have converged to be essentially diffusion-controlled.

### Reactions with peroxy radicals

The reactivity of sulfinic acids to peroxy radicals was explored *via* inhibited co-oxidation of PBD-BODIPY (Fig. 5A).<sup>44</sup> PBD-BODIPY reports on the reaction progress of the autoxidation and its consumption is conveniently monitored by conventional spectrophotometry. Since its reactivity toward peroxy radicals






Fig. 5 (A and B) PBD-BODIPY serves as the signal carrier in hydrocarbon autoxidations, enabling determination of rate constants ( $k_{\text{inh}}$ ) and reaction stoichiometries ( $n$ ) for reactions of inhibitors with chain-carrying peroxy radicals. (C) Co-oxidations of 1-hexadecene (2.8 M) and PBD-BODIPY (10  $\mu\text{M}$ ) initiated by AIBN (6 mM) at 37  $^\circ\text{C}$  (dashed black trace) and inhibited by 10  $\mu\text{M}$  of TrptSOH (red), TrpSO<sub>2</sub>H (blue), and PhSO<sub>2</sub>H (green). (D) The inhibition rate constants for TrptSOH, plotted as a function of medium  $\beta_2^{\text{H}}$ . (E–H) Calculated (CBS-QB3) TS structures for HAT between model sulfinic and sulfenic acid and model peroxy radical (MeOO $\cdot$ ) and associated free energy barriers and rate constants estimated using transition state theory, HOMO visualized. TS structures were confirmed to link H-bonded pre- and post-reaction complexes by intrinsic reaction coordinate calculations (see ESI $^\dagger$ ).

has been independently determined (e.g.  $k_{\text{PBD-BODIPY}} = 3790 \text{ M}^{-1} \text{ s}^{-1}$  in 1-hexadecene/chlorobenzene at 37  $^\circ\text{C}$ , see ESI $^\dagger$ ), the inhibition rate constant ( $k_{\text{inh}}$ ) of an added peroxy radical-trapping antioxidant and the stoichiometry of the inhibition reactions ( $n$ ) can be easily derived from the data as in Fig. 5B.

Despite extensive experimentation under various conditions and co-substrates, such as 1-hexadecene and styrene, we were unable to observe any inhibition of autoxidation by the sulfinic acids, suggesting that they do not efficiently react with the chain-carrying peroxy radicals. Representative data are shown in Fig. 5C, where co-oxidations carried out in the presence of PhSO<sub>2</sub>H and TrptSO<sub>2</sub>H can be compared to TrptSOH, which has previously been shown to undergo very fast reactions with peroxy radicals ( $k = 3.0 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$ ).<sup>30</sup> Since sulfonyl radicals are known to readily add to alkenes (e.g.  $k = 7 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$  for 2-vinylnaphthalene),<sup>45</sup> which would afford an alkyl radical that could carry the autoxidation chain, inhibited autoxidations of substrates lacking a double bond (THF and dioxane) were also carried out, but yielded similar results. As expected, the peroxy radical-trapping kinetics of the sulfenic acid was highly solvent-dependent, yielding  $\alpha_2^{\text{H}} = 0.46$  when the data was plotted

according to the Ingold–Abraham relationship (Fig. 5D), consistent with our previous report.<sup>31</sup>

To provide insight on the lack of reactivity of the sulfinic acids to peroxy radicals, we calculated the transition state structures for the reactions of *t*-BuSO<sub>2</sub>H and *t*-BuSOH with a model peroxy radical (methylperoxy). Two low energy structures were identified for each system (Fig. 5E–H), one wherein the substituents on the oxygen atoms between which the H-atom is transferred are oriented *anti* with respect to one another, and the other wherein they are *syn*. The *syn* TS structures were significantly lower in energy in both cases and, consistent with the experimental observations, the predicted barriers for the reaction of *t*-BuSO<sub>2</sub>H were much higher than those for *t*-BuSOH. These results suggest that the inhibition rate constants for sulfinic acids are roughly seven orders of magnitude lower than for corresponding sulfinic acids, and as such, consistent with our observations that the sulfinic acids are incapable of inhibiting the co-oxidations that were carried out.

Insight to the stark contrast in reactivity of the sulfinic and sulfenic acids is provided upon consideration of the HOMO of



the transition states which are shown in Fig. 5E and F for the preferred *syn* TS and in Fig. 5G and H for the *anti* TS. Although there is clearly overlap between the  $\pi^*$  of the peroxy radical and the lone pair of the sulfur atom that leads to the preference for the *syn* TS in the reactions of both the sulfinic and sulfenic acids, this interaction is smaller in the sulfinic acid ( $\Delta\Delta G^\ddagger = 2.7 \text{ kcal mol}^{-1}$ ) compared to the sulfenic acid ( $\Delta\Delta G^\ddagger = 5.1 \text{ kcal mol}^{-1}$ ), presumably due to the lower sulfur lone pair energy in the sulfinic acid. Indeed, natural bond orbital analyses indicate that the interaction energy of the sulfur lone pair and peroxy radical  $\pi^*$  drops from  $3 \text{ kcal mol}^{-1}$  in the *syn* TS of the sulfenic acid reaction to  $0.4 \text{ kcal mol}^{-1}$  in the *syn* TS of the sulfinic acid reaction.

In addition to the foregoing efforts, we performed transient absorption spectroscopic experiments to directly monitor the reaction of peroxy radicals with sulfinic acids. The cumyl radicals generated from photolysis of dicumylketone (see above) were used to produce cumylperoxy radicals simply by carrying out the photolysis under an atmosphere of  $\text{O}_2$ . Although cumylperoxy radicals do not exhibit a visible absorption – precluding observation of their decay in the presence of sulfinic acids – the product sulfonyl radicals absorb at  $350 \text{ nm}$ . Nevertheless, the data acquired over a wide range of concentrations (validated through comparing with the equivalent alkyl radicals – see ESI† for conditions), suggested no sulfonyl radical formation on the timescale of the experiment.

## Discussion

Thiols, the precursors to both sulfenic and sulfinic acids,<sup>32,46,47</sup> are eminently useful H-atom transfer agents,<sup>48</sup> serving as terminal (stoichiometric) reductants in a number of transformations, including the thiol-ene coupling<sup>49,50</sup> and its aerobic counterpart, the TOCO reaction.<sup>51,52</sup> We recently showed that the oxidation of a thiol to a sulfenic acid<sup>32</sup> renders it a better H-atom donor – on thermodynamic grounds (*i.e.* alkyl sulfenic acid SO-H BDE  $\sim 70 \text{ kcal mol}^{-1}$  *vs.* alkyl thiol S-H BDE  $\sim 87 \text{ kcal mol}^{-1}$ )<sup>31</sup> and kinetic grounds (*e.g.*  $k \sim 10^7 \text{ M}^{-1} \text{ s}^{-1}$  for reactions of alkyl sulfenic acids with peroxy radicals *vs.*  $<10^3 \text{ M}^{-1} \text{ s}^{-1}$  for alkyl thiols).<sup>30</sup> However, the greater stability of the sulfinyl radicals precludes useful chemistry. To the best of our knowledge, there is no equivalent of the thiol-ene coupling or TOCO reaction with a sulfenic acid, although these could be

very useful reactions! In the preceding pages we have expanded our previous studies to show that, despite their significantly weaker bonds, sulfenic acids are only slightly more inherently reactive towards alkyl and alkoxy radicals compared to thiols. Moreover, because sulfenic acids are such good H-bond donors ( $\alpha_2^{\text{H}} \sim 0.55$ ) they react more slowly than thiols ( $\alpha_2^{\text{H}} \sim 0-0.1$ )<sup>53</sup> in good H-bond accepting solvents.

Further oxidation of the sulfenic acid to a sulfinic acid enables production of the more reactive sulfonyl radical that can be used for productive chemistry. Indeed, Lei and co-workers recently demonstrated the sulfinic acid equivalent of a TOCO reaction,<sup>12</sup> another example that sulfinic acids are gaining in popularity. The foregoing results enable us to place the reactivity of sulfinic acids into proper context; the alkyl sulfinic S(O)O-H BDE  $\sim 78 \text{ kcal mol}^{-1}$  infers that the stability of the sulfonyl radical is roughly midway between that of thiyl and sulfinyl radicals. Nevertheless, sulfinic acids are similarly inherently reactive to both alkyl and alkoxy radicals as sulfenic acids and thiols. Importantly, like sulfenic acids, sulfinic acids are excellent H-bond donors ( $\alpha_2^{\text{H}} \sim 0.63$ ), and as such, they are relatively poor H-atom donors compared to thiols in good H-bonding solvents. Thus, sulfinic acids are expected to be useful as H-atom transfer agents only in non-H-bonding solvents;<sup>54</sup> otherwise, thiols are likely to be a much better choice. Indeed, Lei's TOCO-like reaction and Nicewicz's use of sulfinic acids as catalytic H-atom donors in photocatalytic alkene functionalizations are optimal in weak H-bond accepting chlorinated hydrocarbons ( $\beta_2^{\text{H}} \sim 0.15$ ).<sup>55</sup>

To complete the sulfur oxyacid series, we used CBS-QB3 to predict the O-H BDE in a sulfonic acid. The computed result of  $107.4 \text{ kcal mol}^{-1}$  for *t*-BuSO<sub>3</sub>H underscores why these compounds are unreactive to any of the radicals we have investigated here. For ease of reference, a summary of our results is presented in Fig. 6.

In light of the similar inherent reactivity of thiols, sulfenic acids, and sulfenic acids to alkyl and alkoxy radicals, the difference in the peroxy radical reactivity between thiols and sulfenic acids ( $>4$  orders of magnitude) is quite significant. In one sense, this is intuitive given that the reactions are far less exergonic (*e.g.* compare  $\Delta G^\circ \sim -16$  and  $\sim 0 \text{ kcal mol}^{-1}$  for reactions of sulfenic acids and thiols with peroxy radicals *versus*  $\Delta G^\circ = -34$  and  $-18 \text{ kcal mol}^{-1}$  for reactions of sulfenic acids and thiols with primary alkyl or alkoxy radicals).





	thiol	sulfenic acid	sulfinic acid	sulfonic acid	
	$\text{R-S-H}$	$\text{R-S-O-H}$	$\text{R-S(=O)-O-H}$	$\text{R-S(=O)(=O)-O-H}$	
					BDEs in kcal/mol
					rate constants in $\text{M}^{-1}\text{s}^{-1}$
<b>BDE</b>	<b>~87</b>	<b>~70</b>	<b>~78</b>	<b>~107</b>	
$k_{\text{H}} (+ \text{R}^\bullet)$	high $10^6$	high $10^6$	low $10^6$	unreactive	previously reported
$k_{\text{H}} (+ \text{RO}^\bullet)$	high $10^7$	low $10^9$	mid $10^8$	unreactive	this work
$k_{\text{H}} (+ \text{ROO}^\bullet)$	$<10^3$	$\sim 10^7$	$<10^3$	unreactive	

Fig. 6 Comparison of H-atom transfer thermochemistry and kinetics for thiols and each of the thiol-derived sulfur oxyacids.






However, previous work from our group<sup>31,56,57</sup> suggests that there is more to it; specifically, that HAT to peroxy radicals is enabled by secondary orbital interactions, wherein filled  $\pi$  or non-bonding orbitals on the group attached to the atom from which the H-atom is being transferred interact with the proximal end of the peroxy radical  $\pi^*$  MO.<sup>73</sup> Upon oxidation of the electron-rich sulfur atom in the sulfenic acid to a sulfinic acid, this secondary orbital interaction is expected to be significantly reduced. Indeed, our calculations predict that the sulfinic acid is not only less reactive to peroxy radicals than the sulfenic acid, but is even less reactive than thiols, despite significantly more favourable reaction energetics (of *ca.* 10 kcal mol<sup>-1</sup>). Moreover, both PhSO<sub>2</sub>H and TrptSO<sub>2</sub>H were incapable of inhibiting the autoxidation of styrene and 1-hexadecene, substrates whose autoxidation is propagated by peroxy radicals.<sup>58</sup>

At first glance, the poor HAT reactivity of sulfinic acids toward peroxy radicals would seem to be the best evidence to date that secondary orbital interactions contribute to the reactivity of good peroxy radical-trapping agents (*i.e.* antioxidants). However, it must be acknowledged that there is another explanation for the lack of inhibition of the autoxidation: the sulfinic acids themselves autoxidize. This has been noted to be particularly facile in water,<sup>59</sup> where it is likely to be initiated and propagated by single electron transfer reactions, eventually producing a sulfonate:



There is also precedence for sulfinic acid autoxidation in non-polar organic solvents,<sup>60</sup> which presumably propagates *via* H-atom transfer *in lieu* of single electron transfer (In $\cdot$  is an arbitrary initiating radical):



Presented in this way, sulfinic acid autoxidation is analogous to aldehyde autoxidation, which eventually yields carboxylic acids *via* Baeyer–Villiger oxidation of the aldehyde with the autoxidation product peracid.

The increasing popularity of sulfinic acids (and/or their corresponding sulfinate salts) to carry out sulfonylations of multiple bonds or serve as a terminal reducing agent is surprising in light of their purported facile autoxidation. The autoxidation depends on the rate and equilibrium position of

eqn (6) as well as the follow-up H-atom transfer reaction (eqn (7)). The rate constant for the reaction of sulfonyl radicals with O<sub>2</sub> has been reported to be 1 × 10<sup>9</sup> M<sup>-1</sup> s<sup>-1</sup> in water/MeOH.<sup>61</sup> If the reaction were this fast in organic solvent, reactions involving sulfonyl radicals would be impossible under aerobic conditions.<sup>12,13,17</sup> For example, under the reaction conditions developed by the Lei group for oxidative functionalization of alkynes,<sup>13</sup> both the addition of sulfonyl radicals and the addition of oxygen would have effective rates of around 1 × 10<sup>7</sup> s<sup>-1</sup>. Therefore, yields of 80% would be impossible, since the sulfinic acid (present in excess) would simply autoxidize. Surprisingly, there is no data for the reaction of sulfonyl radicals with O<sub>2</sub> in the non-aqueous solvents in which synthetic organic transformations are generally carried out. Moreover, there are no data on the position of the RSO<sub>2</sub> $\cdot$ /O<sub>2</sub> equilibrium and the H-atom abstraction from a sulfinic acid by a sulfonylperoxy radical.

To provide some insight on the rate and equilibrium position of RSO<sub>2</sub> $\cdot$ /O<sub>2</sub> in organic solvents, we carried out additional CBS-QB3 calculations. The addition of oxygen to MeSO<sub>2</sub> $\cdot$  and PhSO<sub>2</sub> $\cdot$  is predicted to be exergonic by 2.9 kcal mol<sup>-1</sup>, with barriers of  $\Delta G^\ddagger = 11.0$  and 10.0 kcal mol<sup>-1</sup>, corresponding to rate constants of  $k = 2.8 \times 10^6$  and  $1.4 \times 10^7$  M<sup>-1</sup> s<sup>-1</sup>, respectively, at 37 °C.<sup>62</sup> These results suggest that water and/or protic solvents accelerate the reaction and increase the driving force – perhaps due to stronger H-bonding interactions in the product sulfonylperoxy than the starting sulfonyl.

Interestingly, if the sulfonylperoxy radical forms, it is predicted to be extremely reactive; the calculated barrier for H-atom abstraction from MeSO<sub>2</sub>H by MeS(O)<sub>2</sub>OO $\cdot$  is predicted to be  $\Delta G^\ddagger = 7.0$  kcal mol<sup>-1</sup>, corresponding to a rate constant of  $1.9 \times 10^9$  M<sup>-1</sup> s<sup>-1</sup>. This would lead to very fast consumption of sulfinic acid.<sup>63</sup> Recall, the predicted barrier for H-atom abstraction from MeSO<sub>2</sub>H by MeOO $\cdot$  is nine orders of magnitude slower!

To corroborate these predictions, we photolyzed *p*-toluene-sulfonyl iodide in benzene and monitored the decay of the *p*-toluenesulfonyl radical at 350 nm in the presence and absence of O<sub>2</sub>. Under an atmosphere of N<sub>2</sub>, the tosyl radicals decayed with  $k = (7.3 \pm 0.5) \times 10^8$  M<sup>-1</sup> s<sup>-1</sup>, consistent with the kinetics of their disproportionation to sulfinyl and sulfonyloxy radicals obtained by EPR.<sup>35</sup> The tosyl radicals decayed at a greater rate when analogous experiments were carried out under an air atmosphere, yielding an apparent rate constant of  $k = (1.0 \pm 0.2) \times 10^9$  M<sup>-1</sup> s<sup>-1</sup>. Assuming this rate constant is the superposition of the disproportionation and O<sub>2</sub> addition reactions, a rate constant for the latter can be estimated to be  $k = (9 \pm 4) \times 10^5$  M<sup>-1</sup> s<sup>-1</sup>, in remarkably good agreement with the value predicted by CBS-QB3, and most importantly, >1000-fold slower than in aqueous solution.<sup>64</sup> With a rate constant in this range, typical propagating reactions of sulfonyl radicals (*e.g.* addition to double and triple bonds,  $k = 10^7$  to  $10^9$  M<sup>-1</sup> s<sup>-1</sup>)<sup>45</sup> can compete effectively with oxygen addition, which under typical synthetic conditions<sup>65</sup> translates to a difference of 3–3.5 orders of magnitude in reaction rates.<sup>66</sup> An indication that optimized synthetic procedures depend strongly on these kinetics is highlighted by the significant drop in yield (>50%)





that is observed upon switching from an air atmosphere to exclusively O<sub>2</sub>.<sup>13</sup>

## Conclusions

Sulfinic acids are very good H-atom donors which lead to highly reactive sulfonyl radicals having a wide range of synthetic applications. Their HAT reactivity is quite similar to that of sulfenic acids, however, because of their increased BDE of  $\sim 78$  (vs. 70) kcal mol<sup>-1</sup> they are typically slightly slower with rate constants of at  $2 \times 10^6$  M<sup>-1</sup> s<sup>-1</sup> with alkyl radicals and  $5 \times 10^8$  M<sup>-1</sup> s<sup>-1</sup> with alkoxy radicals. A surprising property is the slow reaction of sulfinic acids with peroxy radicals, which could not be measured, and was calculated to take place with a rate constant of  $\sim 1$  M<sup>-1</sup> s<sup>-1</sup>, which underscores the role of secondary orbital interactions in facilitating H-atom transfer to peroxy radicals. Although sulfinic acids are known to readily autoxidize, this has not deterred recent synthetic endeavours. We have found that the key propagating reaction of the sulfonyl radical with O<sub>2</sub> is greatly slowed in organic solvents relative to water to enable these reactions to proceed. Similar to sulfenic acids, the more oxidized sulfinic acids also show high H-bond acidity and therefore a very large modulation of their HAT activity in different solvents. This should be taken into account when employing them in catalytic and/or chain reactions.

## Experimental section

### General

Reagents were obtained from commercial suppliers and used as received, unless indicated otherwise. Column chromatography was carried out with 40–63  $\mu$ m, 230–400 mesh silica gel. <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on a Bruker AVANCE spectrometer operating at 400 MHz and 101 MHz, respectively, unless indicated otherwise. High-resolution mass spectra were obtained on a Kratos Concept Tandem mass spectrometer (EI) and Micromass Q-TOF (ESI). TrpSO<sub>2</sub>H, PhSO<sub>2</sub>H and dicumylketone were synthesized as described in the ESI.† TrpSOH,<sup>28</sup> PBD-BODIPY,<sup>31</sup> STY-BODIPY,<sup>44</sup> NMBHA,<sup>67</sup> tosyl iodide,<sup>68</sup> as well as the GC standards for the alkyl radical clock kinetics<sup>69</sup> were synthesized following previously reported procedures. Chlorobenzene was dried over 3 Å molecular sieves before use. Solvents for kinetics experiments were of anhydrous or HPLC quality. UV-visible spectra were measured with a Cary 100 spectrophotometer equipped with a thermostatted 6 × 6 multi-cell holder.

### Radical equilibration EPR (REqEPR)

Electron paramagnetic resonance (EPR) spectra were recorded on a Bruker EMXplus (X-band) spectrometer equipped with an ER 4119HS cavity. The (composite) spectra were fitted and the radical concentration was determined using the quantitative EPR package of the Bruker Xenon software. The full procedure to determine thermodynamic data can be found in the ESI.† To generate the radicals, 5% (v/v) di-*tert*-butyl peroxide was added to the solutions in benzene (with or without 10% v/v *t*-BuOH)

under nitrogen and the EPR cavity irradiated with a Hamamatsu LC5 Hg-Xe lamp (150 W) *via* a 3.5 mm quartz light guide. REqEPR experiments with sulfonyl radicals were performed under continuous irradiation.

### Laser flash photolysis

Nanosecond transient absorption experiments were performed on an LFP-112 spectrometer (Luzchem, Canada) using a EX10 (GAM Laser, USA) XeCl excimer laser (308 nm, *ca.* 10 mJ per pulse, *ca.* 12 ns pulse width). The transient absorption data were recorded in a quartz cuvette (1 cm × 1 cm) equipped with a septum. Sample concentrations were adjusted to yield an absorbance of 0.2–0.3 at 308 nm and the solutions were bubbled with nitrogen for 10 minutes before measurement. The rate constants for H-atom abstraction ( $k_{\text{H}}$ ) were determined under pseudo-first-order conditions and calculated according to  $k_{\text{obs}} = k_0 + k_{\text{H}}[\text{H-donor}]$ .

### Competition kinetics

Alkyl radical clock kinetics was determined using a previously reported procedure by Valgimigli and co-workers.<sup>40</sup> The Barton (PTOC) ester synthesis is described in the ESI† and the standards were synthesized as described previously.<sup>69</sup> Briefly, vials were loaded with the appropriate amount of sulfinic acid (150–400 mM) and capped with a septum. THF (950  $\mu$ L) was added followed by 1 eq. of methanesulfonic acid and the vials were wrapped in aluminum foil before adding the PTOC ester (20 mM). The vials were purged with nitrogen and the aluminum foil was removed before the vials were exposed to a sodium lamp (400 W) at a distance of 40 cm for 1 h at room temperature (*ca.* 25 °C). Aliquots of 50  $\mu$ L were taken from each vial and added to a GC vial containing hexylbenzene standard solution (50  $\mu$ L, 40 mM). The solution was diluted with 900  $\mu$ L of acetonitrile for a total volume of 1 mL per vial. The samples (4  $\mu$ L splitless injections) were analyzed by GC-MS on an Agilent HP-5 ms column (30 m × 0.25 mm × 0.25  $\mu$ m) with a constant He flow of 1.1 mL min<sup>-1</sup> using the following temperature profile: (inlet temperature was set to 150 °C): 90 °C, hold 1 min, 2 °C min<sup>-1</sup> to 120 °C, hold 1 min, 5 °C min<sup>-1</sup> to 165 °C, hold 1 min, 20 °C min<sup>-1</sup> to 300 °C, hold 5 min. The method yielded retention times of 10.9, 22.9, 23.4, and 25.4 min for hexylbenzene, 2-*tert*-butylnaphthalene, 2-isobutylnaphthalene, 2-(2-methylallyl)naphthalene and 2-(2-methylprop-1-en-1-yl)naphthalene, respectively.

### Inhibited autoxidations

Inhibited autoxidations were performed as previously described.<sup>56</sup> The procedure for 1-hexadecene autoxidation in chlorobenzene is transcribed below. 1-Hexadecene (2.0 mL) was added to a 3.5 mL cuvette along with PhCl (0.44 mL). The cuvette was then placed in the thermostated sample holder of a UV-vis spectrophotometer and equilibrated to 37 °C for approximately 10 min. The PBD-BODIPY probe (12.5  $\mu$ L, 2.0 mM solution in 1,2,4-trichlorobenzene) was added followed by AIBN (50  $\mu$ L, 300 mM in chlorobenzene) and the solution was thoroughly mixed. The absorbance at 588 nm was monitored for



10 min after which 10  $\mu\text{L}$  of a solution of the test antioxidant was added. The solution was then mixed and the absorbance readings resumed. The resulting data were processed as previously reported.<sup>44,56</sup> PMC, which has an established stoichiometry of 2,<sup>70</sup> was used as a standard to derive the rate of initiation ( $R_i = 1.3 \times 10^{-9} \text{ M s}^{-1}$ ) and propagation rate constant for the dye ( $k_{\text{PBD-BODIPY}} = 3790 \text{ M}^{-1} \text{ s}^{-1}$ ).

### Calculations

Electronic structure calculations were carried out using the CBS-QB3 complete basis set method<sup>71</sup> as implemented in the Gaussian 16 suite of programs.<sup>72</sup> Stationary points were identified as minima or maxima by calculation of second derivatives. Maxima were verified to be transition states for the relevant H-atom transfer reactions by carrying out intrinsic reaction coordinate calculations in both the forward and reverse directions. Rate constants were calculated *via* transition state theory at 25 °C, except for reactions with peroxy radicals (37 °C, for comparison with experiment).

### Conflicts of interest

There are no conflicts to declare.

### Acknowledgements

This work was supported by grants from the Natural Sciences and Engineering Research Council of Canada and the Canada Foundation for Innovation and through generous access to the computational resources of the Centre for Advanced Computing (<https://cac.queensu.ca>). JPC acknowledges the support of the Ontario Graduate Scholarships program. We are grateful to Dr Marco Mazzonna and Prof. Osvaldo Lanzalunga (University of Rome, Sapienza) for their early efforts to understand the kinetics of the reactions of sulfenic acids with alkoxy radicals.

### Notes and references

- 1 C. Chatgililoglu, M. P. Bertrand and C. Ferreri, in *S-Centered Radicals*, ed. Z. B. Alfassi, Wiley, 1999, pp. 311–354.
- 2 R. S. Glass, *Top. Curr. Chem.*, 2018, **376**, 1–42.
- 3 *Oxidation and Antioxidants in Organic Chemistry and Biology*, ed. E. T. Denisov and I. B. Afanas'ev, CRC Press, 2005.
- 4 Y. Jiang and T.-P. Loh, *Chem. Sci.*, 2014, **5**, 4939–4943.
- 5 A. Garcia-Dominguez, S. Mueller and C. Nevado, *Angew. Chem., Int. Ed.*, 2017, **56**, 9949–9952.
- 6 C. S. Gloor, F. Denes and P. Renaud, *Angew. Chem., Int. Ed.*, 2017, **56**, 13329–13332.
- 7 T. Taniguchi, A. Idota and H. Ishibashi, *Org. Biomol. Chem.*, 2011, **9**, 3151–3153.
- 8 J. M. Fang and M. Y. Chen, *Tetrahedron Lett.*, 1987, **28**, 2853–2856.
- 9 N. Mantrand and P. Renaud, *Tetrahedron*, 2008, **64**, 11860–11864.
- 10 E. Block, M. Aslam, V. Eswarakrishnan, K. Gebreyes, J. Hutchinson, R. Iyer, J. A. Laffitte and A. Wall, *J. Am. Chem. Soc.*, 1986, **108**, 4568–4580.
- 11 They also figure prominently in polymer chemistry,<sup>11a-c</sup> and air pollution.<sup>11d</sup> (a) V. Percec, B. Barboiu and H. J. Kim, *J. Am. Chem. Soc.*, 1998, **120**, 305–316; (b) C. Gorsche, M. Griesser, G. Gescheidt, N. Moszner and R. Liska, *Macromolecules*, 2014, **47**, 7327–7336; (c) G. Gryn'ova, T. Guliasvili, K. Matyjaszewski and M. L. Coote, *Aust. J. Chem.*, 2013, **66**, 308–313; (d) A. Mardyukov and P. R. Schreiner, *Acc. Chem. Res.*, 2018, **51**, 475–483.
- 12 Q. Lu, J. Zhang, F. Wei, Y. Qi, H. Wang, Z. Liu and A. Lei, *Angew. Chem., Int. Ed.*, 2013, **52**, 7156–7159.
- 13 Q. Lu, J. Zhang, G. Zhao, Y. Qi, H. Wang and A. Lei, *J. Am. Chem. Soc.*, 2013, **135**, 11481–11484.
- 14 Z. Huang, Q. Lu, Y. Liu, D. Liu, J. Zhang and A. Lei, *Org. Lett.*, 2016, **18**, 3940–3943.
- 15 A. J. Perkowski and D. A. Nicewicz, *J. Am. Chem. Soc.*, 2013, **135**, 10334–10337.
- 16 Z. Yuan, H.-Y. Wang, X. Mu, P. Chen, Y.-L. Guo and G. Liu, *J. Am. Chem. Soc.*, 2015, **137**, 2468–2471.
- 17 Y. Ning, Q. Ji, P. Liao, E. A. Anderson and X. Bi, *Angew. Chem., Int. Ed.*, 2017, **56**, 13805–13808.
- 18 Y. Xi, B. Dong, E. J. McClain, Q. Wang, T. L. Gregg, N. G. Akhmedov, J. L. Petersen and X. Shi, *Angew. Chem., Int. Ed.*, 2014, **53**, 4657–4661.
- 19 F. Chen, Q. Meng, S.-Q. Han and B. Han, *Org. Lett.*, 2016, **18**, 3330–3333.
- 20 T. C. Johnson, B. L. Elbert, A. J. M. Farley, T. W. Gorman, C. Genicot, B. Lallemand, P. Pasau, J. Flasz, J. L. Castro, M. MacCoss, D. J. Dixon, R. S. Paton, C. J. Schofield, M. D. Smith and M. C. Willis, *Chem. Sci.*, 2018, **9**, 629–633.
- 21 A. U. Meyer, K. Strakova, T. Slanina and B. Koenig, *Chem.–Eur. J.*, 2016, **22**, 8694–8699.
- 22 Gilbert and co-workers have estimated  $k \sim 10^6 \text{ M}^{-1} \text{ s}^{-1}$  for the reaction of methanesulfinic acid with methyl radicals, both of which are formed in the reaction of DMSO with hydroxyl radicals.<sup>36</sup>
- 23 J. G. Calvert, C. C. Badcock, H. W. Sidebottom, G. W. Reinhardt and E. K. Damon, *J. Am. Chem. Soc.*, 1971, **93**, 3115–3121.
- 24 A. G. Davies, B. P. Roberts and B. R. Sanderson, *J. Chem. Soc., Perkin Trans. 2*, 1973, 626–630.
- 25 S. W. Benson, *Chem. Rev.*, 1978, **78**, 23–35.
- 26 F. De Vleeschouwer, V. Van Speybroeck, M. Waroquier, P. Geerlings and F. De Proft, *J. Org. Chem.*, 2008, **73**, 9109–9120.
- 27 A. Gunturu, R. Asatryan and J. W. Bozzelli, *J. Phys. Org. Chem.*, 2011, **24**, 366–377.
- 28 A. J. McGrath, G. E. Garrett, L. Valgimigli and D. A. Pratt, *J. Am. Chem. Soc.*, 2010, **132**, 16759–16761.
- 29 V. Vaidya, K. U. Ungold and D. A. Pratt, *Angew. Chem., Int. Ed.*, 2009, **48**, 157–160.
- 30 R. Amorati, P. T. Lynett, L. Valgimigli and D. A. Pratt, *Chem.–Eur. J.*, 2012, **18**, 6370–6379.
- 31 Z. Zielinski, N. Presseau, R. Amorati, L. Valgimigli and D. A. Pratt, *J. Am. Chem. Soc.*, 2014, **136**, 1570–1578.



- 32 J.-P. R. Chauvin and D. A. Pratt, *Angew. Chem., Int. Ed.*, 2017, **56**, 6255–6259.
- 33 M. Lucarini, P. Pedrielli, G. F. Pedulli, S. Cabiddu and C. Fattuoni, *J. Org. Chem.*, 1996, **61**, 9259–9263.
- 34 C. Chatgililoglu, B. C. Gilbert and R. O. C. Norman, *J. Chem. Soc., Perkin Trans. 2*, 1979, 770–775.
- 35 J. E. Bennett, G. Brunton, B. C. Gilbert and P. E. Whittall, *J. Chem. Soc., Perkin Trans. 2*, 1988, 1359–1364.
- 36 B. C. Gilbert, R. O. C. Norman and R. C. Sealy, *J. Chem. Soc., Perkin Trans. 2*, 1975, 303–308.
- 37 The previously calculated value was  $75.7 \text{ kcal mol}^{-1}$  for *t*-butylsulfenic acid<sup>27</sup> which shows a significant difference to  $\text{TrptSO}_2\text{H}$ .
- 38 Previous CBS-QB3 calculations on *t*-butylsulfenic acid yielded  $68.6 \text{ kcal mol}^{-1}$ , indicating that the three aryl substituents bonded to the carbon which bears the sulfenic acid in  $\text{TrptSOH}$  clearly impart a slight electron-withdrawing effect, destabilizing the electron-poor sulfinyl radical and driving up the O–H BDE.
- 39 R. Amorati, M. Lucarini, V. Mugnaini, G. F. Pedulli, F. Minisci, F. Recupero, F. Fontana, P. Astolfi and L. Greci, *J. Org. Chem.*, 2003, **68**, 1747–1754.
- 40 R. Leardini, M. Lucarini, G. F. Pedulli and L. Valgimigli, *J. Org. Chem.*, 1999, **64**, 3726–3730.
- 41 D. W. Snelgrove, J. Luszyk, J. T. Banks, P. Mulder and K. U. Ingold, *J. Am. Chem. Soc.*, 2001, **123**, 469–477.
- 42 A rate constant for the reaction of hydroxyl radicals with methanesulfenic acid has been reported ( $6.0 \times 10^{-9} \text{ M}^{-1} \text{ s}^{-1}$ ).<sup>36</sup>
- 43 ET reactions are more facile in more polar solvents, and will therefore increase when moving to higher  $\beta_2^{\text{H}}$  values in the solvent diagram. This will lower the slope of the overall rate constant  $k$  and therefore lead to a decrease in  $\alpha_2^{\text{H}}$ . This effect correlates with the reduction potential of the involved radical and is therefore strongest with benzophenone ( $E^\circ = 1.27 \text{ V}$ )<sup>43a</sup> and alkoxy radicals ( $E^\circ = 0.9 \text{ V}$ ),<sup>43b</sup> but not present with alkyl radicals ( $E^\circ = -1.73 \text{ V}$ ).<sup>43c</sup> (a) K. McNeill and S. Canonica, *Environ. Sci.: Processes Impacts*, 2016, **18**, 1381–1399; (b) G. Merenyi, J. Lind and L. J. Engman, *J. Chem. Soc., Perkin Trans. 2*, 1994, 2551–2553; (c) D. D. M. Wayner, D. J. McPhee and D. Griller, *J. Am. Chem. Soc.*, 1988, **110**, 132–137.
- 44 E. A. Haidasz, A. T. M. Van Kessel and D. A. Pratt, *J. Org. Chem.*, 2016, **81**, 737–744.
- 45 C. Chatgililoglu, O. Mozziconacci, M. Tamba, K. Bobrowski, G. Kciuk, M. P. Bertrand, S. Gastaldi and V. I. Timokhin, *J. Phys. Chem. A*, 2012, **116**, 7623–7628.
- 46 M. Hugo, L. Turell, B. Manta, H. Botti, G. Monteiro, L. E. S. Netto, B. Alvarez, R. Radi and M. Trujillo, *Biochemistry*, 2009, **48**, 9416–9426.
- 47 K. Goto, M. Holler and R. Okazaki, *J. Am. Chem. Soc.*, 1997, **119**, 1460–1461.
- 48 F. Denes, M. Pichowicz, G. Povie and P. Renaud, *Chem. Rev.*, 2014, **114**, 2587–2693.
- 49 M. J. Kade, D. J. Burke and C. J. Hawker, *J. Polym. Sci., Part A: Polym. Chem.*, 2010, **48**, 743–750.
- 50 C. E. Hoyle, A. B. Lowe and C. N. Bowman, *Chem. Soc. Rev.*, 2010, **39**, 1355–1387.
- 51 M. S. Kharasch, W. Nudenberg and G. J. Mantell, *J. Org. Chem.*, 1951, **16**, 524–532.
- 52 H. H. Szmant, A. J. Mata, A. J. Namis and A. M. Panthananickal, *Tetrahedron*, 1976, **32**, 2665–2680.
- 53 M. H. Abraham, R. J. Abraham, J. Byrne and L. Griffiths, *J. Org. Chem.*, 2006, **71**, 3389–3394.
- 54 At elevated concentrations in solvents of low H-bond accepting ability, the reactivity of sulfenic acids to highly reactive species (e.g. alkoxy radicals) may, however, be diminished due to self-association (dimerization).
- 55 M. H. Abraham, P. L. Grellier, D. V. Prior, J. J. Morris and P. J. Taylor, *J. Chem. Soc., Perkin Trans. 2*, 1990, 521–529.
- 56 J.-P. R. Chauvin, M. Griesser and D. A. Pratt, *J. Am. Chem. Soc.*, 2017, **139**, 6484–6493.
- 57 D. Hu and D. A. Pratt, *Chem. Commun.*, 2010, **46**, 3711–3713.
- 58 During our investigations we found one further example of a significant difference in HAT reactivity between sulfenic and sulfenic acids: the phenylthiyl radical – for further details see the ESI.†
- 59 K. Sehested and J. Holcman, *Radiat. Phys. Chem.*, 1996, **47**, 357–360.
- 60 L. Horner and O. H. Basedow, *Justus Liebigs Ann. Chem.*, 1958, **612**, 108–131.
- 61 M. Tamba, K. Dajka, C. Ferreri, K.-D. Asmus and C. Chatgililoglu, *J. Am. Chem. Soc.*, 2007, **129**, 8716–8723.
- 62 The corresponding reaction of  $\text{MeSO}^\bullet$  is  $12.2 \text{ kcal mol}^{-1}$  endergonic.
- 63 To test this assumption, we attempted to monitor the sulfenic acid during the autoxidation. To do so, we employed a 9-triptycenesulfenic acid featuring a fluorine atom at the other bridgehead position such that the reaction could be followed directly by  $^{19}\text{F}$  NMR. We recently showed that the sulfenic and sulfonic acids of this fluorinated triptycene have distinct chemical shifts.<sup>32</sup> Although the concentration of the sulfenic acid was increased up to 10 mM in order to facilitate detection by NMR, the experiment revealed rapid consumption of the sulfenic acid on the time-scale of the experiment, which would prevent them to inhibit the autoxidation.
- 64 This indicates that the formation of sulfonyl radicals should be visible in LFP experiments with cumyloxy radicals if they would react at rates similar to sulfenic acid ( $10^7 \text{ M}^{-1} \text{ s}^{-1}$ ) or other comparable antioxidants (such as  $\alpha$ -tocopherol, which has a comparable BDE and reacts at  $10^6 \text{ M}^{-1} \text{ s}^{-1}$ ). Therefore, the LFP results strongly suggest that the reaction of alkylperoxy radicals with sulfenic acids are as slow as predicted by calculations ( $<10^3 \text{ M}^{-1} \text{ s}^{-1}$ ).
- 65 [Substrate (alkene or alkyne)] = 50 mM and [oxygen] = 2–7 mM.
- 66 Of course, the additions must also compete with the disproportionation of sulfonyl radicals, but this is quite slow in comparison due to the very low concentrations of these transient intermediates compared to the alkene/alkyne substrate.



- 67 G. X. Ortiz, B. N. Hemric and Q. Wang, *Org. Lett.*, 2017, **19**, 1314–1317.
- 68 G. L. Edwards, C. A. Muldoon and D. J. Sinclair, *Tetrahedron*, 1996, **52**, 7779–7788.
- 69 J. J. Hanthorn, R. Amorati, L. Valgimigli and D. A. Pratt, *J. Org. Chem.*, 2012, **77**, 6895–6907.
- 70 G. W. Burton, T. Doba, E. Gabe, L. Hughes, F. L. Lee, L. Prasad and K. U. Ingold, *J. Am. Chem. Soc.*, 1985, **107**, 7053–7065.
- 71 J. A. Montgomery Jr, M. J. Frisch, J. W. Ochterski and G. A. Petersson, *J. Chem. Phys.*, 1999, **110**, 2822–2827.
- 72 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. V. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery Jr, J. E. Peralta, F. Ogliaro, M. J. Bearpark, J. J. Heyd, E. N. Brothers, K. N. Kudin, V. N. Staroverov, T. A. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. P. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman and D. J. Fox, *Gaussian 16*, Revision A.03; Wallingford, CT, 2016.
- 73 It is very difficult to definitively tease out whether these types of reactions should be described as a HAT or proton-coupled electron transfer (PCET). However, simply considering the TS geometries in Fig. 5, the authors are inclined to describe them as HAT-like. Since both the sulfinyl/sulfonyl and peroxy radicals are  $\pi$  radicals, a HAT TS would feature the H-atom moving along a coordinate that is orthogonal to the planes made up by the carbon, sulfur and oxygen of the sulfinyl/sulfonyl radical and the carbon, oxygen and oxygen of the peroxy radical. In contrast, in a purely PCET process, the proton and electron are, by definition, transferred between different pairs of (nominally orthogonal) orbitals; the proton between the lone pairs on the terminal oxygen atoms of the sulfinyl/sulfonyl and peroxy radicals and the unpaired electron is delocalized across both  $\pi^*$  orbitals. The predicted structures are more consistent with the former description than the latter. This is perhaps easiest to see in the SOMOs, where there is a node on the H-atom (see ESI<sup>†</sup>).
- 74 N. Nakamura, *J. Am. Chem. Soc.*, 1983, **105**, 7172–7173.

