


Natural Product Reports

rsc.li/npr



Themed issue: Engineering of Cell Factories for the Production of Natural Products
Guest Editor: Tilmann Weber

ISSN 0265-0568



ROYAL SOCIETY
OF CHEMISTRY

Celebrating
IYPT 2019




REVIEW ARTICLE

Harald Gross, Marnix H. Medema *et al.*
Computer-aided re-engineering of nonribosomal peptide
and polyketide biosynthetic assembly lines



Cite this: *Nat. Prod. Rep.*, 2019, 36, 1249

Computer-aided re-engineering of nonribosomal peptide and polyketide biosynthetic assembly lines

Mohammad Alanjary, ^{†a} Carolina Cano-Prieto, ^{†b} Harald Gross ^{*b} and Marnix H. Medema ^{*a}

Covering: 2014 to 2019

Nonribosomal peptide synthetases (NRPSs) and polyketide synthases (PKSs) have been the subject of engineering efforts for multiple decades. Their modular assembly line architecture potentially allows unlocking vast chemical space for biosynthesis. However, attempts thus far are often met with mixed success, due to limited molecular compatibility of the parts used for engineering. Now, new engineering strategies, increases in genomic data, and improved computational tools provide more opportunities for major progress. In this review we highlight some of the challenges and progressive strategies for the re-design of NRPSs & type I PKSs and survey useful computational tools and approaches to attain the ultimate goal of semi-automated and design-based engineering of novel peptide and polyketide products.

Received 14th March 2019

DOI: 10.1039/c9np00021f

rsc.li/npr

1. NRPSs and PKSs: enzymatic factories for compound production
2. Strategies for re-engineering PKS/NRPS systems
 - 2.1 Precursor directed biosynthesis
 - 2.2 Domain exchanges
 - 2.3 Specificity code mutations
 - 2.4 Starter units and tailoring modifications
 - 2.5 Multi-module exchange
3. Computational tools for re-engineering biosynthetic assembly lines
 - 3.1 Search and collection of parts
 - 3.2 Optimization and refactoring
 - 3.3 Toward automated cluster design
4. Conclusion and future outlook
5. Conflicts of interest
6. References

1. NRPSs and PKSs: enzymatic factories for compound production

Using biology to produce high-value compounds has been a boon to humanity; cell factories are an integral production method to supply antibiotics and other useful natural products to the world, with a market value estimated at several hundreds

of billions of USD per year.¹ For example, complex structures such as vancomycin require over 40 steps for total synthesis,² while production in *Amycolatopsis orientalis* can be made in continuous batch cultures with relative ease.³ Along with optimization efforts,^{4,5} cell factories can be a more economic strategy for mass production or for the generation of new leads, especially for the variety of natural products with intricate structures.

Non-ribosomal peptide synthetases (NRPSs) and polyketide synthases (PKSs) can be described as the factory ‘machinery’ for many natural products. Engineering these multi-modular systems has been a ‘holy grail’ for synthetic biology due to their modular nature and the endless combinatorial design possibilities available through module deletions, insertions, duplications or exchanges. Leveraging these systems can thus accelerate discovery of high-value compounds, especially to help refill the waning antimicrobial pipeline.⁶ However, ever since the earliest attempts in the 1990s, modifying PKSs and NRPSs on demand has turned out to be much more challenging than first expected: due to the complex molecular interactions among modules and domains, their engineering requires deep understanding of protein–protein interactions and domain specificities. For this reason, many engineering successes have led to low yields impractical for industrial production. Now, new breakthroughs are finally providing solutions to leverage these systems, opening up countless new opportunities that can be exploited more effectively with the help of computer-aided design tools. Here, we review state-of-the-art strategies for engineering of modular PKSs and NRPSs, as well as the computational tools that can be used to support and accelerate this process.

^aBioinformatics Group, Wageningen University, Wageningen, The Netherlands. E-mail: marnix.medema@wur.nl

^bDepartment of Pharmaceutical Biology, Pharmaceutical Institute, Eberhard Karls Universität Tübingen, Tübingen, Germany. E-mail: harald.gross@uni-tuebingen.de

[†] Authors contributed equally.



Modular PKSs as well as NRPSs constitute complex enzymatic assembly lines comprised of multiple enzymatic components that are responsible for initiation, elongation and termination of polyketide or peptide chains to produce the core scaffolds of natural products. In the NRPS context, each module catalyzes the addition and modification of a specific amino acid and successively extends the peptide in an assembly line manner. Each module consists of at least three domains that define a minimal module: the adenylation (A) domain, the thiolation (T) domain and the condensation (C) domain. The A domain selects, activates and transfers a specific amino acid to the T domain. Subsequently, the C domain catalyzes the peptide bond formation of the amino acid tethered to the T-domain of the same module and one of the preceding module. The integrated amino acids can be further modified concerning their absolute configuration at the C α atom by epimerization

domains (E) and the degree of C- or N-methylation by methyltransferases (MT). Furthermore, heterocyclization can be performed by cyclization domains (Cy), while redox-active domains (Ox, Red) determine their redox state. A loading module typically comprises an A and a T domain, an elongation module a C–A–T set, while a termination module contains a C–A–T and thioesterase domain (TE). The latter is responsible for the release, or optionally, macrocyclization of the peptide product. Thus, a vast variety of compounds can be generated based on different composition and arrangements of these elements. With only a few exceptions,^{7–9} the organization and order of modules corresponds to the amino acid sequence of the peptide product (co-linearity rule). Thanks to the pioneering work of the Lipmann^{10,11} and the Laland¹² groups in the 1960s and later on by Stachelhaus and coworkers, who discovered the specificity conferring code of the A domains,¹³ the logic of NRPSs is



Dr Mohammad Alanjary obtained his PhD in bioinformatics at the University of Tübingen, Germany, with a focus on antibiotic resistance. Previously he aided in the launch of the first commercial semi-conductor gene-sequencing platform and developed several procedures for performance optimization while working at IonTorrent. He is currently working at Wageningen University to develop

computational methods leveraging comparative genomics for natural product discovery and engineering.



Harald Gross is a pharmacist by training and obtained his PhD degree in 2004 from the University of Bonn, Germany, followed by 2 years of postdoctoral research at the Oregon State University in Corvallis/OR, USA and at the Scripps Institution of Oceanography in La Jolla/CA, USA. Back at the University of Bonn, he established in 2006 his own independent research group where he was mainly working on

the genomics and secondary metabolism of Pseudomonas bacteria. Since 2012 he is Full Professor at the University of Tübingen. His research interests are in the field of genome-driven discovery of microbial secondary metabolites, biosynthesis research and genetic engineering.



Carolina Cano Prieto was born in 1984. She obtained her bachelors degree in Biology at University of Granada (Spain) followed by a MSc in Microbiology under the Spanish Educational Minister fellowship. She completed her PhD studies in 2015 under the supervision of Prof. José Antonio Salas and Dr Carlos Olano at University of Oviedo where her work was focused in identification of the

polyketide biosynthesis produced by streptomycetes. Since 2016 she is a post-doctoral researcher in the lab of Prof. Dr Harald Gross at the University of Tübingen. There she currently works on the characterization and modification of NRPS biosynthetic gene clusters in Pseudomonas and Bacillus strains and with the development of biosynthetic gene clusters using synthetic biology tools.



Marnix Medema is an Assistant Professor of Bioinformatics at Wageningen University, The Netherlands. In 2013, he completed his PhD in the groups of Eriko Takano and Rainer Breitling, at the University of Groningen. During his PhD he spent time in the lab of Michael Fischbach at the University of California, San Francisco, as a visiting research fellow. Following a postdoctoral

fellowship in the group of Frank Oliver Glöckner at the Max Planck Institute for Marine Microbiology in Bremen, Germany, he joined Wageningen University in 2015. His group develops and applies computational tools to understand bacterial, fungal and plant natural product biosynthesis from a genomic perspective.



reasonably understood and set the stage for engineering of the NRPS assembly lines.

In a similar fashion to NRPSs, modular PKSs synthesize polyketides through the stepwise elongation of the starter unit of 2-, 3- or 4-carbon units molecule such as acetyl-CoA, propionyl-CoA, butyryl-CoA, and their activated derivatives, malonyl-, methylmalonyl-, and ethylmalonyl-CoA extender units.¹⁴ Three main types of PKSs have been described: type I PKSs, type II PKSs and type III PKSs.¹⁵ In this review, we will focus on modular type I PKSs. The minimal module of these is also formed by three core domains: the acetyltransferase (AT) domain which selects for the extender unit and transfers it to the acyl carrier protein (ACP) domain. Finally, a ketosynthase (KS) domain catalyzes the condensation reaction between two modules – analogous to the C-domain in NRPSs. Elongation modules comprise all three core domains, while the loading module lacks a KS domain and the terminal module contains a TE domain, which is responsible for the release of the linear polyketide chain or of the release with macrocyclization. The sole use of KS–AT–ACP-modules leads to β -keto chains. The additional integration of a ketoreductase (KR) domain converts the keto-functionality into a β -OH group, which can be eliminated by a dehydratase (DH) domain to give an α - β unsaturated alkene, which in turn can be reduced to a single bond with the help of an enoyl-reductase (ER) domain. Thus, the final PKS assembly line can be defined as KS–AT–(DH–ER–KR)–ACP. For more detailed information on NRPS and PKSs, we refer the reader to several excellent reviews.^{16–21}

2. Strategies for re-engineering PKS/NRPS systems

Various strategies have been employed to leverage NRPS/PKS systems, ranging from combinatorial design to direct refactoring of processing steps. The following is a brief overview of

these tactics with some of their successes and challenges (Fig. 1).

2.1 Precursor directed biosynthesis

Precursor directed biosynthesis (PDB) and mutasynthesis were among the first re-engineering attempts applied on PKSs and NRPSs. PDB is based on feeding of alternative monomers to change the final product by leveraging domain promiscuity. This strategy is based on the frequent observation of major and minor products.^{22,23} In an NRPS context, the reason is that the A domains possess a natural relaxed substrate flexibility, *e.g.* a Glx-specific domain recognizes, and also to a minor extent activates, Asx. Likewise the Leu/Val/Ile-specific domains recognize their respective closely related member(s) of the branched amino acid family. Occasionally, A-domains display an intriguing promiscuity: *e.g.*, the putatively Pro-specific A domain of the pyreudione NRPS accepts not only proline derivatives, but also ring-extended residues.²⁴ This has proven to be an effective strategy to diversify NRPS products, for example with the addition of a fluorinated non-proteinogenic amino acid to the lipopeptide iturin.²⁵ However, empirically the yields of the obtained new compounds are commonly low because the synthetic amino acids compete with the natural amino acids for cellular uptake as well as A-domain recognition. To overcome this problem, the mutasynthesis concept can be integrated,^{26,27} where the microorganism is deficient in the synthesis of the natural precursor and relies on the fed synthetic precursor to complete the secondary metabolite.²⁸ Successful examples include *e.g.* the alteration of the regular dihydroxyphenylglycine and β -OH-Tyr units in the glycopeptide antibiotic balhimycin with methoxylated and fluoro-derivatives, respectively.^{29,30}

In the same way, PDB has been applied to PKSs. One of the first such experiments conducted with the 6-deoxyerythronolide B biosynthesis pathway led to the new analogue 15-fluoroethyl-6-deoxyerythronolide B.³¹ Similarly, mutasynthesis approaches have also been applied to PKSs. An example of this is the work on the antitumoral polyketide geldanamycin by Eichner and coworkers,³² in which variants of the natural 3-amino-5-hydroxybenzoic acid starter unit were incorporated into the polyketide product.

2.2 Domain exchanges

Considering the modular structure of the NRPS, it is most appealing to simply exchange a module with another. Such exchanges have been demonstrated at the level of a single A domain,³³ combined C–A or A–T domains³⁴ and finally of complete C–A–T modules.³⁵ However, with a few exceptions, the success of these experiments remained modest due to reduced product yields. These swaps worked best if domains are exchanged with those from the same biosynthetic gene cluster (BGC) or from closely related BGCs that encode the production of molecules within the same compound family. The most extensive and successful example of this approach represents the work from Baltz and coworkers, who engineered the NRPS assembly line of the clinically applied lipopeptide antibiotic daptomycin in 2006. They replaced single or multiple C–A–T




Fig. 1 Various strategies for re-engineering PKS/NRPS systems including: (a) precursor-directed biosynthesis to leverage domain promiscuity. (b) Domain editing to re-program monomer specificities. (c) Domain exchanges to replace partial or whole modules. (d) Multi-module exchanges to produce chimeric clusters. (e) Insertion or deletion of domains. (f) Post processing enzyme addition or deletion.



modules with modules from the same BGC or from the closely related NRPSs for the calcium-dependent antibiotics (CDA) and A54145.³⁶ The authors demonstrated that 8 of the 13 AA positions in daptomycin can be modified by module exchange, which led to a combinatorial library of 40 daptomycin derivatives.³⁷ In a recent review, Baltz exposed the failure and success in daptomycin NRPS engineering and suggested that C–A linkers are not flexible while T–C linkers and A–T are.³⁸ In summary, these impressive results showed that while domain swapping is possible, there are also considerable limitations due to the lack of understanding of inter-modular communications and downstream specificity filters.

A recent breakthrough in the understanding of the NRPS logic was achieved by contributions of the Bode group. During investigations of the flexibility of inter-domain linkers, they observed that C–A linkers are indeed more flexible than previously suggested. Two separate structural parts form these linkers, and 22 N-terminal amino acids appear to mediate the interactions between C and A domains. Based on these findings, they defined a new concept of an exchange unit (XU). An XU consists of sets of A–T–C (or A–T–C/E) domains instead of canonical C–A–T modules. The consequent application of these rules resulted in the successful *de novo* design of the NRPS of xenotetrapeptides.³⁹ However, the yields decreased drastically, maintaining the common problem encountered in previous NRPS engineering attempts. In a follow-up work, Bode and coworkers refined and expanded their model and recognized the importance of the involved C-domains. In general, C domains catalyze the condensation of the downstream T-domain-bound amino acid (donor substrate) with the activated upstream T-domain-bound amino acid or peptide (acceptor substrate). For the fusion, donor and acceptor substrates have to be coordinated in their respective C_{acceptor} - and C_{donor} -subdomains. The authors established that particularly the acceptor site is very selective for the nature of the side chain and chirality. However, the A domains have been considered to act as the primary specificity determinants in NRPSs. Bode and coworkers picked up the hypothesis that both subdomains display specificity for the corresponding amino acids and act as gatekeepers.⁴⁰ Thus, an exchange that respects involved C_{acceptor} - and C_{donor} -subdomains, as well as native protein–protein interactions adjacent to an A–T-bidomain, yielded a redefined exchange unit (referred as XUC, see Fig. 2a). During their experiment series, the Bode group located the exact dissection position in the linker region of the C-domain, so that it can be complemented with the respected counterpart for combinatorial biosynthesis and stays fully functional. The application of these rules led to the production of the target compounds without a loss in yields, a breakthrough for NRPS engineering.⁴¹

Similar to the exchange of domains and modules in NRPSs, many engineering attempts for PKSs have been based on swapping or insertion of AT domains. For the erythromycin assembly line, new derivatives have been obtained by such domain insertions.⁴³ DEBS-PKS engineering has been performed in the native producer (*Saccharopolyspora erythraea*), but also in heterologous hosts (*Streptomyces coelicolor* CH1999




Fig. 2 General domain and module organization of (a) non-ribosomal peptide synthetase (NRPS) and (b) polyketide synthase (PKS).⁴² Classic module definitions are highlighted in pink and redefined modules, such as XUC, are shown in red.

and *S. lividans* K4-114). For example, Oliynyk and co-workers exchanged the AT of the module 1 of erythromycin PKS by the AT of module 2 of the rapamycin PKS, resulting in an extension unit exchange from methylmalonyl-CoA to malonyl-CoA and obtaining a novel triketide lactone instead of a lactone without methylation in the C-4.⁴⁴ Similar approaches have recently also been used for the engineering of polyketides related to biofuel production.⁴⁵ However these single domain exchanges have been known for some time to introduce complications in downstream compatibility⁴⁶ and result in lower product yields. Yuzawa and coworkers (2016) related this problem with the sequence of the linkers among domains, specifically, between AT and KR domains and KS and AT domains. The AT–KR linker region is denominated by the authors as post AT-linker (PAL) region and can be further subdivided in PAL1 and PAL2, while the KS–AT linker region was designated as KAL. In-depth analysis of the linker sequences led to the identification of significant consensus sequences between the linker regions of DEBS-PKS and epothilone (EPO) PKS. The authors conducted diverse swapping experiments using different variants of swapped regions and finally resolved that the best region for the swapping is the region formed by KAL–AT–PAL1, while PAL2 was kept native. The production yield showed a minimal decrease compared to wild type. Subsequently, the group confirmed these rules using the lipomycin PKS assembly line and showed successful exchanges using modules from 5 different PKSs. With yields on par or greater than wild-type production, this illustrates that this approach is both robust and effective.⁴⁷

Like for NRPSs, recent advances in PKS engineering have led to the redefinition of the domain organisation of PKS modules. PKS modules were previously organised into KS–AT–(DH–ER–KR)–ACP modules, but recent phylogenetic analysis by Abe *et al.* has highlighted that processing enzymes co-migrate during the assembly line with the KS domain downstream of the ACP.^{42,48,49} Thus, new PKS modules would be termed AT + (DH + ER + KR) +



ACP + KS. These updated boundary definitions have already shown to have a beneficial effect in the rational design of a PKS to produce homoaureothin.⁵⁰ These findings are a milestone in engineering NRPS/PKS systems and open the possibility for realizing total *de novo* design of clusters.

2.3 Specificity code mutations

Another approach to maintain inter-domain compatibility is to simply alter the specificity-coding regions of a domain. Structural biology investigations with the initial A domain of the gramicidin NRPS with L-Phe and AMP revealed the eight key amino acid residues in the active site of the A-domain¹³ which allows the substrate specificity prediction straight from the DNA level. This 8 AA-containing code, also referred as “Stachelhaus-code” was later on refined.⁵¹ While it needs to be adapted when applied to NRPS BGCs originating from certain bacterial genera or from fungi,⁵² it still forms the basis for today’s bioinformatics-driven prediction of the products of NRPS gene clusters. This specificity code can also be modified to alter the chemical outcome of an assembly line. A successful example for A-domain code mutations is provided by the Micklefield group, who reprogrammed the 10th calcium-dependent antibiotic (CDA)-synthetase module to recognize Gln and Me-Gln instead of Glu.⁵³ This has also recently been shown to be effective in altering non-promiscuous domains to accept unnatural extender units.⁵⁴ Furthermore, besides rationally guided specificity code mutagenesis, it is also conceivable to perform directed evolution experiments, *i.e.* mutant libraries are generated and screened for enhanced properties such as an increased bioactivity of the final metabolite or an increased or different A-domain-selectivity. For details about the progress in this field, the reader is referred to other reviews.⁵⁵

Regarding PKS systems, it is also established to change the specificity of the AT-domains concerning natural substrates by mutagenesis. However, recently, some studies are taking a step forward trying to incorporate non-natural extender units. Based on the mutations which were made in module 6 of DEBS-PKS⁵⁴ the Williams group repeated the same mutations in the pikromycin synthase cluster.⁵⁶ Initially, they did not obtain satisfactory results with the mutation of Try-Arg in the corresponding AT domains but with the site-specific mutation of Try to Val at position 755, they achieved the desired substrate shift; they were also able to produce derivatives with the non-natural units propargyl-, ethyl- and allylmalonyl-CoA in robust yields.⁵⁴ The incorporation of above mentioned non-natural units enables a further derivatization through semisynthetic chemistry, especially by click chemistry. In this way, the polyketide of interest can be conjugated with other molecules such as a further pharmacophore-containing moiety or dyes and would expand in this way the possibilities for drug discovery and diverse chemical-biological applications with the polyketidic compound.

2.4 Starter units and tailoring modifications

Other factors to take into account are the engineering of the starter domains or tailoring domains. The swapping of loading

modules (LM) in PKSs has proven successful. For example, Leadlay and coworkers exchanged the 6-deoxyerythronolide LM for the avermectin LM, and Long *et al.* exchanged the DEBS LM for the oleandomycin LM.^{57,58} NRPS initiation modules have also been modified with increasingly positive results. These starting modules are attractive areas for redesign because they omit upstream restraints that confound similar extender module redesigns.⁵⁹ Challenges still remain however with downstream compatibility. For example, several failed attempts at reprogramming tyrocidine initiation modules were shown to result from incompatibilities between PCP and E-domain interfaces.⁶⁰

Further successful domain engineering was achieved regarding internal tailoring domains, for example KR domain replacements with a tri-domain to form a new PKS.⁶¹ Various studies have shown positive swapping between two different KR domains in isolation, however the engineering of DH domains depends on the KR domains that precede them because DH domains are sensitive to the stereochemical configuration of the substrate.⁶² Tailoring domains can also be embedded as subdomains within functional NRPS domains, as seen with methylation interrupted A-domains. These proved to be a boon to NRPS design by allowing site-specific and selective methylation of monomer side chains (O- or S-) or at N termini.⁶³ Additionally, exogenous reactions can also be incorporated to diversify compounds; examples include halogenation, glycosylation, acylation and, sulfation tailoring enzymes.²⁸

2.5 Multi-module exchange

To govern the protein–protein interactions between modules that are part of separate polypeptides, not only the interactions between ACP–KS domain pairs are important, but also sets of small specialized N- and C-terminal regions called docking domains (DDs) for PKSs and communication (COM) domains for NRPSS. These interactions have been shown to be a dominant cause for lowered product turnover rates, overshadowing the impact of substrate recognition in chimeric clusters.⁶⁴ By manipulation of these regions, examples of combinatorial clusters have been generated^{65,66} or re-designed to modulate products of single-module NRPS systems.⁶⁷ DDs ensure the correct PKS assembly into a functional enzyme. Studies on virginiamycin biosynthesis show that the DDs are essential for the correct assembly of the enzymatic complex and finally for the communication between the KS and ACP domains. For successful engineering of PKSs, the identification of the interaction regions and the elucidation of the mechanism involved in the communication between the domains or modules is required.⁶⁸

For PKS engineering, the studies have been based on two research lines: modification of AT domains and docking domains (DDs). Principally, the engineering of ATs has concentrated on the swapping of solely AT domains or the complete module. Furthermore, direct mutagenesis to change the precursors’ specificity of the targeted AT has been conducted. All of these concepts were applied extensively to the erythromycin PKS (syn. 6-deoxyerythronolide B synthase =



DEBS). One example is the exchange of module 1 and 2 (load methylmalonyl-CoA) of DEBS by specific malonyl-CoA modules of *Streptomyces hygroscopicus* ATCC 29253 and the rapamycin module 14 PKS.^{68,69} New derivatives were obtained, however, again with strongly decreased yield. It is noteworthy to mention that the drop in production was highly dependent on the swapping position of the PKS assembly line. It appears that the linker region is of considerable importance to mediate communication between heterologous modules. A recent study showed that the yields of a chimeric DEBS system can be improved by introducing non-native docking domains between the native ones; this apparently added flexibility between processing and condensing domains⁷⁰, which led to improved yields to near-native amounts.

3. Computational tools for re-engineering biosynthetic assembly lines

3.1 Search and collection of parts

Extensive genetic and biosynthetic parts catalogs have been growing in recent years, such as the iGEM Registry of Standard Biological Parts.⁷¹ In addition, databases of BGCs are helping to provide valued comparative context for design efforts. Some of the most successful computational approaches to NRPS/PKS redesign have relied on comparative analysis of related BGCs, which can help decipher optimal fusion/recombination points, identify new biosynthetic parts, and improve our understanding of module specificities.^{39,41,72} Furthermore, large-scale comparative analyses have shed light on natural evolutionary patterns that can guide future efforts and approaches.⁷³ For instance, conglomerate BGCs show significant acquisition and refactoring of cluster fragments that encode functionally independent elements. These sub-clusters, responsible for similar moieties in the final product (including, *e.g.*, the biosynthesis of non-proteinogenic amino acids or the synthesis and transfer of modified sugar units), are shared across a variety of distinct BGCs and can potentially be leveraged for combinatorial redesign of new chimeric clusters.⁷³ Similar lessons in BGC evolution have been shown on laboratory time scales, as seen with sequencing comparisons of PKSs that underwent deletion and recombination events.⁷⁴ These experiments showed analogs that produced near-native product yields, further underscoring the benefit of re-defined PKS modules as building blocks for engineering. Comparative approaches have also aided traditional combinatorial design; a classic example is the generation of 154 bimodular PKSs based on raw material from eight related PKS assembly lines.⁷⁵ To take advantage of this perspective, cataloging and rapid search methods are required. Fortunately, progress in recent years has made this process easier and more automated through the use of computational tools, which are increasingly necessary as genomic data continues to grow.

With maturing methods for BGC detection⁷⁶ and increasing amounts of genomic data, there is a growing need to effectively navigate available BGC sequences. Several databases and tools

facilitate the cataloging and comparison of known and putative BGCs. The recently updated antiSMASH database contains predicted gene clusters for ~25 000 representative complete and high-quality draft bacterial genomes, whereas the IMG Atlas of Biosynthetic gene Clusters contains a higher number of BGCs by also including lower-quality genomes as well as metagenomes.^{77,78} Both databases provide various search options to quickly identify biosynthetic parts of interest; for example, one can quickly query the antiSMASH database to download all adenylation domains specific for hydroxyphenylglycine in BGCs with a high similarity to the vancomycin BGC. The MiBIG repository provides data on >1800 BGCs of known function, with annotations and evidence codes that document functions of gene clusters, enzymes and domains; these are all downloadable in a computer-readable JSON format, which makes it possible to quickly screen for enzymes or domains of interest.^{77–79} The NORINE database is yet another resource for cluster mining, which contains extensive catalogs of NRPS/PKS monomers.⁸⁰ This resource has served as an aid for retro-biosynthetic structure matching, which has recently become an automated process using the rBAN server.⁸¹ While much of the putative BGC data (over 95%)⁸² have yet to be associated with a known product, they can be leveraged through gene and structure similarity methods to aid with collection of similar clusters. Besides sequence identity-based methods such as clusterblast and MultiGeneBlast,^{83,84} tools that allow large-scale comparison of BGCs based on architectural similarity are recently coming to light. BiG-SCAPE⁸⁵ is one such tool, which incorporates Pfam domain similarity as demonstrated previously.⁸⁶ Multi-locus phylogenetic analysis of gene cluster families (GCFs) using CORASON⁸⁵ can be used to provide even higher-resolution analysis of the relationships between groups of similar BGCs. Biocompass⁸⁷ is another tool to perform BGC networking, with a focus on identifying syntenic blocks of sub-clusters. These approaches allow for a rapid collection of related BGCs, or GCFs with similar subunits, even for understudied or novel BGCs (Fig. 3).

In addition to genomic comparisons, structure prediction and search methods can help with collecting similar known compounds or screen for desired products. One interesting result of structure comparison is a virtual library and generator of polyketides, PKS enumerator, which is based on several biologically active macrolides.⁸⁸ While current structure prediction methods still require advancements in specialized cases (*e.g.* iterative PKS domains, external tailoring, macrocyclization), they are capable of accurately predicting core chemical scaffolds quite well for NRPS/PKS systems. Detection tools such as antiSMASH and PRISM include recent advancements in predicting structures from BGCs, such as enhanced NRPS specificity detection using SANDPUMA and improved exploration of product permutations, respectively.^{72,89,90} In addition to defining module and domain specificities, these predictions can be used to help corroborate putative GCFs or find a starting cluster for engineering a desired compound. Intuitive web servers such as ChemSpider,⁹¹ SIMCOMP,⁹² and the NP Atlas⁹³ provide user-friendly search interfaces for global and sub-structure searches; results are also enhanced with






Fig. 3 (a) Example chemical network using NP Atlas (left), BiG-SCAPE network (right), and structures using the lipopeptide iturin as a query. (b) Modules of iturin-like clusters illustrate parts mining for different substrate specificities.

similarity scoring, moiety highlighting, and exploration in a chemical network. For any compounds of interest, the MIBiG repository can then be queried to identify whether a BGC has been characterized for it. Based on these connections, a larger set of similar clusters can be compared to ultimately help identify useful parts for constructing new clusters.

Another useful product of comparative analysis is the recent NRPS-linker database.⁹⁴ This resource houses a catalog of inter-module linkers (IMLs) taken from NRPS clusters found in all publicly available genomes and a parser to extract IMLs from query sequences. Protein–protein linkers have been shown to play an important role in multi-domain dynamics and flexibility,⁹⁵ and they potentially encode for important module compatibility elements. The database analyses showed IMLs have significant structured elements and were consistently associated to A-domain specificity pairs, suggesting they are crucial to module compatibility.⁹⁴ Thus, conservation of IMLs may aid in module swapping. Furthermore, the authors noticed consistent phylogenetic conservation of IMLs, which implies host optimization of engineered NRPSs might benefit by using IMLs taken from related phyla. This is an interesting unexploited resource, as it expands parts mining to also include a library of “mortar” that can help ‘glue together’ desired combinations of BioBricks.

3.2 Optimization and refactoring

Concerted evolution (the homogenization of domain sequences within assembly lines through recombination) and family-

specific adaptations provide other important evolutionary lessons for BGC refactoring.⁷³ This underscores a need to custom-tailor re-engineered components, which includes codon optimization for host expression or protein level changes *via* directed evolution to maximize compatibility. Prioritizing the exchange of modules or domains for ones with closer phylogenetic proximity and respecting integral interfaces can potentially increase the success of redesigning native clusters. While phylogenetic proximity is not an absolute requirement, as interkingdom hybrid BGCs have been observed,⁹⁶ it is likely to increase the chances of success. Improvements *via* codon optimization, reviewed elsewhere,⁹⁷ illustrate this point. Many web services can perform this optimization as well as account for mRNA secondary structure and GC content;⁹⁷ these functions are also included in DNA supplier workflows, such as IDT’s codon optimization tool (<https://www.idtdna.com/codonopt>). One recent study on the effect of codon context revealed that the influence of neighboring codons can limit the rate of translation,⁹⁸ and may be an additional measure to consider. CCtool (<http://algo.tenju.edu/ccool/>) is a recent server that aims to address this consideration, albeit with limited host selection.⁹⁹

Optimizing amino acid sequence *via* directed evolution experiments is another level of refinement that can have major benefits, as protein–protein interactions or substrate specificities can be enhanced, by *e.g.* increasing their specificity. One example shows a drastic 500 fold improvement of enterobactin production after only 3 rounds of mutations.¹⁰⁰ As the number of generated variants and testing of engineered BGCs can be



Table 1 List of tools to aid with parts collection, directed evolution, homology modeling, and automated pathway design

	Description	URL	Year (latest)	Ref.
Databases & comparative tools				
antismashDB	Antismash-detected BGCs from public genomes	http://antismash-db.secondarymetabolites.org	2018	77
MiBIG	Experimentally verified BGCs	http://mibig.secondarymetabolites.org	2015	79
IMG-ABC	Clusterfinder-detected BGCs from all deposited genomes	http://img.jgi.doe.gov/abc	2015	78
NRPS-linker	Parser and database of Inter Module Linkers (IMLs) for NRPSs	http://nrps-linker.unc.edu	2019	94
NPATlas	Collection of known natural products with structural similarity searches	http://www.npatlas.org	2018	93
SIMCOMP	Structural similarity search tool	http://www.genome.jp/tools/simcomp	2010	92
BiG-SCAPE	Calculates similarity between BGCs and groups into gene cluster families	http://bigscape-corason.secondarymetabolites.org	2018	85
CORASON	Groups gene clusters using multi-locus phylogeny methods	http://bigscape-corason.secondarymetabolites.org	2018	85
Biocompass	Calculates similarity between BGCs with a focus on shared syntenic sub-clusters	http://np-omix.github.io/BioCompass	2017	87
Norine	Database and analysis tools for NRPS clusters and monomers	http://bioinfo.lifl.fr/norine	2016	80
rBAN	Retro-biosynthetic analysis of nonribosomal peptides	http://bioinfo.cristal.univ-lille.fr/rban	2019	81
PKS Enumerator	Virtual database and generator of macrolide polyketides	http://www.fourches-laboratory.com/single-post/2018/09/04/PKS-Enumerator	2018	88
Directed evolution tools				
SCHEMA	Minimization of disruptions to local amino acids	http://github.com/mattasmith/SCHEMA-RASPP	2002	106
HotSpot	Energy minimization scoring from homology modeling results	http://loschmidt.chemi.muni.cz/hotspotwizard	2018	107
Wizard 3.0	Substrate binding approximation method	http://www.cs.duke.edu/donaldlab/osprey.php	2013/2009	101 and 102
CADEE	Reactivity approximation using empirical valence-bond theory	http://github.com/kamerlinlab/cadee	2017	108
Homology search/modeling tools				
IntFOLD	3D modeling with additional built in features: quality assessment, ligand binding, and disorder prediction	http://www.reading.ac.uk/bioinf/IntFOLD	2015	109
RaptorX	Uses closely related homologs and structural data to create context specific models. Ranked 1st in CASP12 for contact point prediction	http://raptorx.uchicago.edu	2015	110
HHpred	Remote homology search with alignment	http://toolkit.tuebingen.mpg.de/#/tools/hhpred	2005	111
MODELLER	Search, model, and evaluation of models in one pipeline	http://salilab.org/modeller	2003	112
PHYRE2	Homology search and additional applications such as loop refinement and variant analysis	http://www.sbg.bio.ic.ac.uk/phyre2	2015	113
ROBETTA	Server suite with <i>de novo</i> prediction and template based 3d model generation	http://rosetta.bakerlab.org	2011	114
SWISS-MODEL	Server suite with variety of model building applications including quaternary structure prediction	http://swissmodel.expasy.org	2018	115
I-TASSER	Part of suite of software that offers function prediction and contact mapping with consistent high rankings in CASP competitions	http://zhanglab.ccmb.med.umich.edu/I-TASSER	2018	116
Automated parts/pathway mining tools				
ClusterCAD	Automated PKS parts discovery engine for <i>de novo</i> design of polyketides	http://clustercad.jbei.org	2018	117
RetroPATH	Automated pathway design using known and permissive enzymes	http://www.jfaulon.com/category/retropath	2014	118
genoCAD	Designer for DNA/vectors	http://www.genocad.com	2009	119

quite time consuming, it is important to use the most educated mutations as possible. Using computational methods to reduce this search space is a valuable solution, as illustrated with the redesign of an NRPS A domain using the K* algorithm, implemented in the OSPREY application,¹⁰¹ which evaluates protein

variants that improve or maintain protein flexibility and ligand binding *via* energy minimization methods.¹⁰² Many tools have been developed, and covered thoroughly in previous reviews,^{103–105} that aim to predict the impact of variants using different criteria: structure consistency, protein stability,






Fig. 4 Example 3D structure generated from the C-domain sequence of module 2 of Gramicidin synthetase using MODELLER with PDB:2JGP as the template. Identified regions: C-donor (orange), alpha-5 helix (blue), linker (red), and C-acceptor (grey).

evolutionary consistency, and coverage of mutant library space (Table 1).

Unfortunately some of these, such as quantum mechanics/molecular mechanics (QM/MM) methods, can be impractical for high-throughput screening due to the lengthy compute times required. A recent addition, CADEE, aims to bridge this gap between fast heuristic methods and higher accuracy QM/MM methods using an empirical valence-bond (EVB) approach.¹⁰⁸ Although these methods still struggle to accurately predict multi-site influence on structure, they can help to avoid severely deleterious mutations and save valuable experimental time.

Rational design constraints, such as focusing on certain hotspot regions, can further complement these methods. One way to highlight these areas is to interrogate the protein structure. Utilization of the growing number of available structures in the Protein Data Bank¹²⁰ for comparative structural analysis can help indicate areas that might be tolerant to change. For example, the T domain central to NRPSs is now known to maintain a relatively consistent conformation throughout its different catalytic states.¹²¹ This was shown through comparative analysis of 18 structures at various catalytic states and further supports that tight interaction of T and surrounding domains and linkers are less amenable to change. Structural context not only informs subdomain lobes and flexible regions, but can also identify binding pocket residues, domain interface locations, and contact points. For example, homology modeling helped to confirm a linker fusion point that enabled the generation of several chimeric variants of antimycin.¹²² These structural homology approaches also led to the identification of a successful recombination site in the GameXPeptide-producing NRPS GxpS of *Photorhabdus luminescens*. This

involved altering the alpha-5 helix into a consistent consensus sequence of all XUs involved, which resulted in the production of several peptide variants.⁴¹ Fig. 4 illustrates how these regions can be identified using homology modeling with tools like MODELLER.¹¹²

With more structures coming to light, for example the first methyltransferase-interrupted A-domain,¹²³ more accurate template-based sequence-to-structure modeling can be leveraged for design efforts. For example, using this analysis, a mutation of a key residue in TioS from the thiocoraline biosynthetic pathway was shown to abolish methylation activity while retaining yields nearly identical to the wild-type.⁶³

Homology modeling can be performed with a variety of web servers and standalone tools (Table 1). The bi-annual Critical Assessment of techniques for protein Structure Prediction (CASP) competitions has helped to mature these methods, showing a significant increase in model accuracy, particularly for template-based methods;¹²⁴ additionally, refinement and template-free modeling have also seen improvements. While much work is still required to further predict protein-protein docking, and ultimately the impact of mutations, these models have helped to identify fruitful areas for directed evolution.¹²⁵

3.3 Toward automated cluster design

Collection of optimal parts for individual modules, domains or enzymes is an important step for successful re-factoring experiments. Currently, homology tools such as BLAST¹²⁶ and HMMer¹²⁷ are largely employed to extract these elements, but steps toward automated identification of useful elements based on a (desired) chemical structure are being employed. ClusterCAD¹¹⁷ is one such tool that focuses on cataloging and highlighting of PKS elements (collected from MIBiG), with the ultimate goal of *de novo* design of clusters. In addition to identifying the best starting BGC most similar to a desired structure, ClusterCAD aims to identify a parsimonious number of domain swaps, deletions and additions based on sequence homology to known clusters. This was shown to automatically select similar components for a previously validated re-engineering of an adipic acid producing cluster.¹²⁸

A more generalized tool, RetroPATH,¹²⁹ aims to identify reactions beyond PKS and NRPS systems. This can be useful for PKS/NRPS engineering by identifying tailoring enzymes or improving the precursor supply of a pathway. The webserver genoCAD¹³⁰ is another application that aims to leverage known context-free parts for designing DNA and vectors; this also has been shown fruitful for the design of plant expression systems.¹³¹ Despite requiring manual curation and optimization of these automated pipelines, these applications provide a foundation for automated design and ultimately can be matured to account for other constraints such as compatibility of compound intermediates.

4. Conclusion and future outlook

Traditional approaches for cell-factory design have largely focused on optimization of native production or expression in



model hosts. Beyond these goals, synthetic biology aims to provide the plans and parts for *de novo* pathway construction, “BioBricks”, that can produce any desired compound. Some examples are the construction of synthetic pathways from plant polyketides: olivetolic acid in *Escherichia coli*¹³² and the biosynthesis of opioids, cannabinoids, and biofuels in yeast.^{133–135} The field is gradually progressing toward the ability to biosynthesize valued compounds “from scratch” in a design-based fashion.¹³⁶ Considering the complexity of genomic regulation, multiple catalytic pathways, and environmental interactions of a cell, it is clear this goal will not be met overnight. Detailed reviews on synthetic biology have illustrated this long, and worthwhile, road for the design, build, and test hurdles toward this end.^{137–139} Although examples of successful computer-aided redesign efforts are discouragingly few, it is promising to see that many barriers for employing these methods are reducing. As accessible web interfaces and automated methods are coming to light, we can expect a greater number of experiments leveraging these approaches and improved generations of these tools from these results. While these methods still have room to mature, and account for more complex design restraints, they have already demonstrated great utility in aiding the rational design of biological machinery.

Here, we highlight another longstanding goal within the general scope of cell factory design: manipulation of modular NRPS and PKS enzyme factories. Although many attempts to design these systems have led to mere proof-of-concept levels of production, a new understanding of inter-module connections, new computational tools, and increasing comparative data analysis provide exciting new opportunities. With the redefinition of module boundaries, it has been shown that combinatorial design can lead to novel products with near-native yields; fully leveraging catalogs of such redefined modules is currently still an underexplored space, but will be made easier with new computational tools. Several applications can help with this step of parts mining, including gene cluster and structural similarity networking. While much of this process still remains manual and can benefit greatly from optimization and custom-tailoring efforts, tools to automate collection and assembly of components are already underway. These applications, such as ClusterCAD for PKSs, currently do not take into account potential incompatibilities *via* protein–protein interactions or substrate specificities but provide the groundwork for searching and structuring a *de novo* BGC in an automated pipeline. Future addition of NRPSs as well as the updated module definitions (exchange units) to ClusterCAD or similar tools, as well as integration with refactoring tools for inserting regulatory elements and optimizing codon usage, will enable even more full-fledged computer-aided design of PKS and NRPS assembly lines, especially if this can be supported by *e.g.* drag-and-drop functionality to assemble new polypeptides *in silico*. As parts collections continue to expand, we are confident that these tools can eventually help solve remaining challenges and thus facilitate the dreams of combinatorial design of these wonderful modular systems to allow accessing a vast chemical space through biosynthesis.

5. Conflicts of interest

MHM is a member of the Scientific Advisory Board of Hexagon Bio.

6. References

- 1 A. M. Davy, H. F. Kildegaard and M. R. Andersen, *Cell Syst.*, 2017, **4**, 262–275.
- 2 D. A. Evans, M. R. Wood, B. W. Trotter, T. I. Richardson, J. C. Barrow and J. L. Katz, *Angew. Chem., Int. Ed.*, 1998, **37**, 2700–2704.
- 3 J. J. McIntyre, A. T. Bull and A. W. Bunch, *Biotechnol. Bioeng.*, 1996, **49**, 412–420.
- 4 K. S. Lee, B. M. Lee, J. H. Ryu, D. H. Kim, Y. H. Kim and S.-K. Lim, *Lett. Appl. Microbiol.*, 2016, **63**, 222–228.
- 5 J. Thykaer, J. Nielsen, W. Wohlleben, T. Weber, M. Gutknecht, A. E. Lantz and E. Stegmann, *Metab. Eng.*, 2010, **12**, 455–461.
- 6 E. Kalkreuter and G. J. Williams, *Curr. Opin. Microbiol.*, 2018, **45**, 140–148.
- 7 S. Lautru, R. J. Deeth, L. M. Bailey and G. L. Challis, *Nat. Chem. Biol.*, 2005, **1**, 265–269.
- 8 D. Reimer, K. N. Cowles, A. Proschak, F. I. Nollmann, A. J. Dowling, M. Kaiser, R. French-Constant, H. Goodrich-Blair and H. B. Bode, *ChemBioChem*, 2013, **14**, 1991–1997.
- 9 A. C. Ross, Y. Xu, L. Lu, R. D. Kersten, Z. Shao, A. M. Al-Suwailem, P. C. Dorrestein, P.-Y. Qian and B. S. Moore, *J. Am. Chem. Soc.*, 2013, **135**, 1155–1162.
- 10 F. Lipmann, *Science*, 1971, **173**, 875–884.
- 11 F. Lipmann, W. Gevers, H. Kleinkauf and R. J. Roskoski, *Adv. Enzymol. Relat. Areas Mol. Biol.*, 1971, **35**, 1–34.
- 12 T. L. Berg, L. O. Froholm and S. G. Laland, *Biochem. J.*, 1965, **96**, 43–52.
- 13 T. Stachelhaus, H. D. Mootz and M. A. Marahiel, *Chem. Biol.*, 1999, **6**, 493–505.
- 14 J. Lau, H. Fu, D. E. Cane and C. Khosla, *Biochemistry*, 1999, **38**, 1643–1651.
- 15 B. Shen, *Curr. Opin. Chem. Biol.*, 2003, **7**, 285–295.
- 16 A. Koglin and C. T. Walsh, *Nat. Prod. Rep.*, 2009, **26**, 987–1000.
- 17 G. H. Hur, C. R. Vickery and M. D. Burkart, *Nat. Prod. Rep.*, 2012, **29**, 1074–1098.
- 18 R. D. Sussmuth and A. Mainz, *Angew. Chem., Int. Ed. Engl.*, 2017, **56**, 3770–3821.
- 19 M. Klaus and M. Grninger, *Nat. Prod. Rep.*, 2018, **35**, 1070–1081.
- 20 T. Robbins, Y.-C. Liu, D. E. Cane and C. Khosla, *Curr. Opin. Struct. Biol.*, 2016, **41**, 10–18.
- 21 K. J. Weissman, *Nat. Prod. Rep.*, 2016, **33**, 203–230.
- 22 J. M. Crawford, C. Portmann, R. Kontnik, C. T. Walsh and J. Clardy, *Org. Lett.*, 2011, **13**, 5144–5147.
- 23 Y. Xie, Q. Cai, H. Ren, L. Wang, H. Xu, B. Hong, L. Wu and R. Chen, *J. Nat. Prod.*, 2014, **77**, 1744–1748.
- 24 M. Klapper, D. Braga, G. Lackner, R. Herbst and P. Stallforth, *Cell Chem. Biol.*, 2018, **25**, 659–665.



- 25 S. Moran, D. K. Rai, B. R. Clark and C. D. Murphy, *Org. Biomol. Chem.*, 2009, **7**, 644–646.
- 26 A. Kirschning and F. Hahn, *Angew. Chem., Int. Ed. Engl.*, 2012, **51**, 4012–4022.
- 27 R. J. M. Goss, S. Shankar and A. A. Fayad, *Nat. Prod. Rep.*, 2012, **29**, 870–889.
- 28 M. Winn, J. K. Fyans, Y. Zhuo and J. Micklefield, *Nat. Prod. Rep.*, 2016, **33**, 317–347.
- 29 S. Weist, B. Bister, O. Puk, D. Bischoff, S. Pelzer, G. J. Nicholson, W. Wohlleben, G. Jung and R. D. Sussmuth, *Angew. Chem., Int. Ed. Engl.*, 2002, **41**, 3383–3385.
- 30 S. Weist, C. Kittel, D. Bischoff, B. Bister, V. Pfeifer, G. J. Nicholson, W. Wohlleben and R. D. Sussmuth, *J. Am. Chem. Soc.*, 2004, **126**, 5942–5943.
- 31 S. L. Ward, R. P. Desai, Z. Hu, H. Gramajo and L. Katz, *J. Ind. Microbiol. Biotechnol.*, 2007, **34**, 9–15.
- 32 S. Eichner, H. G. Floss, F. Sasse and A. Kirschning, *ChemBioChem*, 2009, **10**, 1801–1805.
- 33 M. Crüsemann, C. Kohlhaas and J. Piel, *Chem. Sci.*, 2013, **4**, 1041–1045.
- 34 M. J. Calcott, J. G. Owen, I. L. Lamont and D. F. Ackerley, *Appl. Environ. Microbiol.*, 2014, **80**, 5723–5731.
- 35 S. Zobel, S. Boecker, D. Kulke, D. Heimbach, V. Meyer and R. D. Sussmuth, *ChemBioChem*, 2016, **17**, 283–287.
- 36 K. T. Nguyen, D. Kau, J.-Q. Gu, P. Brian, S. K. Wrigley, R. H. Baltz and V. Miao, *Mol. Microbiol.*, 2006, **61**, 1294–1307.
- 37 R. H. Baltz, *ACS Synth. Biol.*, 2014, **3**, 748–758.
- 38 R. H. Baltz, *J. Ind. Microbiol. Biotechnol.*, 2018, **45**, 1003–1006.
- 39 K. A. J. Bozhüyük, F. Fleischhacker, A. Linck, F. Wesche, A. Tietze, C.-P. Niesert and H. B. Bode, *Nat. Chem.*, 2017, **10**, 275.
- 40 K. Bloudoff and T. M. Schmeing, *Biochim. Biophys. Acta, Proteins Proteomics*, 2017, **1865**, 1587–1604.
- 41 K. A. J. Bozhüyük, A. Linck, A. Tietze, J. Kranz, F. Wesche, S. Nowak, F. Fleischhacker, Y. N. Shi, P. Grün and H. B. Bode, *Nat. Chem.*, 2019, 1755–4349, DOI: 10.1038/s41557-019-0276-z.
- 42 A. T. Keatinge-Clay, *Angew. Chem., Int. Ed.*, 2017, **56**, 4658–4660.
- 43 C. J. Rowe, I. U. Böhm, I. P. Thomas, B. Wilkinson, B. A. M. Rudd, G. Foster, A. P. Blackaby, P. J. Sidebottom, Y. Roddis, A. D. Buss, J. Staunton and P. F. Leadlay, *Chem. Biol.*, 2001, **8**, 475–485.
- 44 M. Oliynyk, M. J. Brown, J. Cortes, J. Staunton and P. F. Leadlay, *Chem. Biol.*, 1996, **3**, 833–839.
- 45 W. Cai and W. Zhang, *Curr. Opin. Biotechnol.*, 2018, **50**, 32–38.
- 46 M. Hans, A. Hornung, A. Dziarnowski, D. E. Cane and C. Khosla, *J. Am. Chem. Soc.*, 2003, **125**, 5366–5374.
- 47 S. Yuzawa, K. Deng, G. Wang, E. E. K. Baidoo, T. R. Northen, P. D. Adams, L. Katz and J. D. Keasling, *ACS Synth. Biol.*, 2016, 7–10.
- 48 D. A. Vander Wood and A. T. Keatinge-Clay, *Proteins: Struct., Funct., Bioinf.*, 2018, **86**, 664–675.
- 49 L. Zhang, T. Hashimoto, B. Qin, J. Hashimoto, I. Kozono, T. Kawahara, M. Okada, T. Awakawa, T. Ito, Y. Asakawa, M. Ueki, S. Takahashi, H. Osada, T. Wakimoto, H. Ikeda, K. Shin-Ya and I. Abe, *Angew. Chem., Int. Ed. Engl.*, 2017, **56**, 1740–1745.
- 50 Y. Sugimoto, K. Ishida, N. Traitcheva, B. Busch, H.-M. Dahse and C. Hertweck, *Chem. Biol.*, 2015, **22**, 229–240.
- 51 M. Rottig, M. H. Medema, K. Blin, T. Weber, C. Rausch and O. Kohlbacher, *Nucleic Acids Res.*, 2011, **39**, W362–W367.
- 52 D. Kalb, G. Lackner and D. Hoffmeister, *Appl. Environ. Microbiol.*, 2014, **80**, 6175–6183.
- 53 M. Styles, J. Thirlway, M. Al Nakeeb, C. P. Smith, R. Lewis, J. Micklefield, L. Nunns and A.-W. Struck, *Angew. Chem., Int. Ed.*, 2012, **51**, 7181–7184.
- 54 E. Kalkreuter, J. M. CroweTipton, A. N. Lowell, D. H. Sherman and G. J. Williams, *J. Am. Chem. Soc.*, 2019, **141**, 1961–1969.
- 55 H. Kries and A. Stanišić, *ChemBioChem*, 2019, **20**, 1347, DOI: 10.1002/cbic.201800750.
- 56 B. Vögeli, K. Geyer, P. D. Gerlinger, S. Benkstein, N. S. Cortina and T. J. Erb, *Cell Chem. Biol.*, 2018, **25**, 833–839.e4.
- 57 A. F. Marsden, B. Wilkinson, J. Cortes, N. J. Dunster, J. Staunton and P. F. Leadlay, *Science*, 1998, **279**, 199–202.
- 58 P. F. Long, C. J. Wilkinson, C. P. Bisang, J. Cortes, N. Dunster, M. Oliynyk, E. McCormick, H. McArthur, C. Mendez, J. A. Salas, J. Staunton and P. F. Leadlay, *Mol. Microbiol.*, 2002, **43**, 1215–1225.
- 59 A. S. Brown, M. J. Calcott, J. G. Owen and D. F. Ackerley, *Nat. Prod. Rep.*, 2018, **35**, 1210–1228.
- 60 U. Linne, S. Doekel and M. A. Marahiel, *Biochemistry*, 2001, **40**, 15824–15834.
- 61 S. Yuzawa, J. D. Keasling and L. Katz, *J. Antibiot.*, 2016, **69**, 494–499.
- 62 J. F. Barajas, J. M. Blake-Hedges, C. B. Bailey, S. Curran and J. D. Keasling, *Synth. Syst. Biotechnol.*, 2017, **2**, 147–166.
- 63 T. A. Lundy, S. Mori and S. Garneau-Tsodikova, *ACS Synth. Biol.*, 2018, **7**, 399–404.
- 64 M. Klaus, M. P. Ostrowski, J. Austerjost, T. Robbins, B. Lowry, D. E. Cane and C. Khosla, *J. Biol. Chem.*, 2016, **291**, 16404–16415.
- 65 M. Hahn and T. Stachelhaus, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 275–280.
- 66 C. Chiochini, U. Linne and T. Stachelhaus, *Chem. Biol.*, 2006, **13**, 899–908.
- 67 C. Hacker, X. Cai, C. Kegler, L. Zhao, A. K. Weickmann, J. P. Wurm, H. B. Bode and J. Wöhnert, *Nat. Commun.*, 2018, **9**, 4366.
- 68 E. Musiol-kroll and W. Wohlleben, *Antibiotics*, 2018, **7**, 62, DOI: 10.3390/antibiotics7030062.
- 69 X. Ruan, A. Pereda, D. L. Stassi, D. Zeidner, R. G. Summers, M. Jackson, A. Shivakumar, S. Kakavas, M. J. Staver, S. Donadio and L. Katz, *J. Bacteriol.*, 1997, **179**, 6416–6425.
- 70 M. Klaus, A. D. D'Souza, A. Nivina, C. Khosla and M. Grninger, *ACS Chem. Biol.*, 2017, **6**, 139–147, DOI: 10.1021/acscchembio.8b01060.



- 71 R. Kelwick, L. Bowater, K. H. Yeoman and R. P. Bowater, *FEMS Microbiol. Lett.*, 2015, **362**, 16, DOI: 10.1093/femsle/fnv129.
- 72 M. G. Chevette, F. Aicheler, O. Kohlbacher, C. R. Currie and M. H. Medema, *Bioinformatics*, 2017, **33**, 3202–3210.
- 73 M. H. Medema, P. Cimerancic, A. Sali, E. Takano and M. A. Fischbach, *PLoS Comput. Biol.*, 2014, **10**, 12, DOI: 10.1371/journal.pcbi.1004016.
- 74 A. Wlodek, S. G. Kendrew, N. J. Coates, A. Hold, J. Pogwizd, S. Rudder, L. S. Sheehan, S. J. Higginbotham, A. E. Stanley-Smith, T. Warneck, M. Nur-E-Alam, M. Radzom, C. J. Martin, L. Overvoorde, M. Samborsky, S. Alt, D. Heine, G. T. Carter, E. I. Graziani, F. E. Koehn, L. McDonald, A. Alanine, R. M. Rodriguez Sarmiento, S. K. Chao, H. Ratni, L. Steward, I. H. Norville, M. Sarkar-Tyson, S. J. Moss, P. F. Leadlay, B. Wilkinson and M. A. Gregory, *Nat. Commun.*, 2017, **8**, 1206.
- 75 H. G. Menzella, R. Reid, J. R. Carney, S. S. Chandran, S. J. Reisinger, K. G. Patel, D. A. Hopwood and D. V. Santi, *Nat. Biotechnol.*, 2005, **23**, 1171.
- 76 T. Weber and H. U. Kim, *Synth. Syst. Biotechnol.*, 2016, **1**, 69–79.
- 77 K. Blin, V. Pascal Andreu, E. L. C. de los Santos, F. Del Carratore, S. Y. Lee, M. H. Medema and T. Weber, *Nucleic Acids Res.*, 2018, **47**, 625–630.
- 78 M. Hadjithomas, I.-M. A. Chen, K. Chu, A. Ratner, K. Palaniappan, E. Szeto, J. Huang, T. B. K. Reddy, P. Cimerancic, M. A. Fischbach, N. N. Ivanova, V. M. Markowitz, N. C. Kyrpides and A. Pati, *mBio*, 2015, **6**, e00932.
- 79 M. H. Medema, R. Kottmann, P. Yilmaz, M. Cummings, J. B. Biggins, K. Blin, I. de Bruijn, Y. H. Chooi, J. Claesen, R. C. Coates, P. Cruz-Morales, S. Duddela, S. Dusterhus, D. J. Edwards, D. P. Fewer, N. Garg, C. Geiger, J. P. Gomez-Escribano, A. Greule, M. Hadjithomas, A. S. Haines, E. J. N. Helfrich, M. L. Hillwig, K. Ishida, A. C. Jones, C. S. Jones, K. Jungmann, C. Kegler, H. U. Kim, P. Kotter, D. Krug, J. Masschelein, A. V. Melnik, S. M. Mantovani, E. A. Monroe, M. Moore, N. Moss, H.-W. Nuttmann, G. Pan, A. Pati, D. Petras, F. J. Reen, F. Rosconi, Z. Rui, Z. Tian, N. J. Tobias, Y. Tsunematsu, P. Wiemann, E. Wyckoff, X. Yan, G. Yim, F. Yu, Y. Xie, B. Aigle, A. K. Apel, C. J. Balibar, E. P. Balskus, F. Barona-Gomez, A. Bechthold, H. B. Bode, R. Borris, S. F. Brady, A. A. Brakhage, P. Caffrey, Y.-Q. Cheng, J. Clardy, R. J. Cox, R. De Mot, S. Donadio, M. S. Donia, W. A. van der Donk, P. C. Dorrestein, S. Doyle, A. J. M. Driessen, M. Ehling-Schulz, K.-D. Entian, M. A. Fischbach, L. Gerwick, W. H. Gerwick, H. Gross, B. Gust, C. Hertweck, M. Hofte, S. E. Jensen, J. Ju, L. Katz, L. Kaysser, J. L. Klassen, N. P. Keller, J. Kormanec, O. P. Kuipers, T. Kuzuyama, N. C. Kyrpides, H.-J. Kwon, S. Lautru, R. Lavigne, C. Y. Lee, B. Linqun, X. Liu, W. Liu, A. Luzhetskyy, T. Mahmud, Y. Mast, C. Mendez, M. Metsa-Ketela, J. Micklefield, D. A. Mitchell, B. S. Moore, L. M. Moreira, R. Muller, B. A. Neilan, M. Nett, J. Nielsen, F. O'Gara, H. Oikawa, A. Osbourn, M. S. Osburne, B. Ostash, S. M. Payne, J.-L. Pernodet, M. Petricek, J. Piel, O. Ploux, J. M. Raaijmakers, J. A. Salas, E. K. Schmitt, B. Scott, R. F. Seipke, B. Shen, D. H. Sherman, K. Sivonen, M. J. Smanski, M. Sosio, E. Stegmann, R. D. Sussmuth, K. Tahlan, C. M. Thomas, Y. Tang, A. W. Truman, M. Viaud, J. D. Walton, C. T. Walsh, T. Weber, G. P. van Wezel, B. Wilkinson, J. M. Willey, W. Wohlleben, G. D. Wright, N. Ziemert, C. Zhang, S. B. Zotchev, R. Breitling, E. Takano and F. O. Glockner, *Nat. Chem. Biol.*, 2015, **11**, 625–631.
- 80 M. Pupin, Q. Esmael, A. Flissi, Y. Dufresne, P. Jacques and V. Leclere, *Synth. Syst. Biotechnol.*, 2016, **1**, 89–94.
- 81 E. Ricart, V. Leclère, A. Flissi, M. Mueller, M. Pupin and F. Lisacek, *J. Cheminf.*, 2019, **11**, 13.
- 82 J. J. J. van der Hooft, J. Wandy, M. P. Barrett, K. E. V. Burgess and S. Rogers, *Proc. Natl. Acad. Sci. U. S. A.*, 2016, **113**, 13738–13743.
- 83 M. H. Medema, E. Takano and R. Breitling, *Mol. Biol. Evol.*, 2013, **30**, 1218–1223.
- 84 T. Weber, K. Blin, S. Duddela, D. Krug, H. U. Kim, R. Bruccoleri, S. Y. Lee, M. A. Fischbach, R. Muller, W. Wohlleben, R. Breitling, E. Takano and M. H. Medema, *Nucleic Acids Res.*, 2015, **43**, W237–W243.
- 85 J. Navarro-Muñoz, N. Selem-Mojica, M. Muldowney, S. Kautsar, J. Tryon, E. Parkinson, E. De Los Santos, M. Yeong, P. Cruz-Morales, S. Abubucker, A. Roeters, W. Lokhorst, A. Fernandez-Guerra, L. Teresa Dias Cappelini, R. Thomson, W. Metcalf, N. Kelleher, F. Barona-Gomez and M. H. Medema, *bioRxiv*, 2018, 445270, DOI: 10.1101/445270.
- 86 P. Cimerancic, M. H. Medema, J. Claesen, K. Kurita, L. C. Wieland Brown, K. Mavrommatis, A. Pati, P. a. Godfrey, M. Koehrsen, J. Clardy, B. W. Birren, E. Takano, A. Sali, R. G. Linington and M. a. Fischbach, *Cell*, 2014, **158**, 412–421.
- 87 T. Leao, G. Castêllo, A. Korobeynikov, E. A. Monroe, S. Podell, E. Glukhov, E. E. Allen, W. H. Gerwick and L. Gerwick, *Proc. Natl. Acad. Sci. U. S. A.*, 2017, **114**, 3198–3203.
- 88 P. P. K. Zin, G. Williams and D. Fourches, *J. Cheminf.*, 2018, **10**, 53.
- 89 K. Blin, T. Wolf, M. G. Chevette, X. Lu, C. J. Schwalen, S. A. Kautsar, H. G. Suarez Duran, E. L. C. de Los Santos, H. U. Kim, M. Nave, J. S. Dickschat, D. A. Mitchell, E. Shelest, R. Breitling, E. Takano, S. Y. Lee, T. Weber and M. H. Medema, *Nucleic Acids Res.*, 2017, **45**, W36–W41.
- 90 C. W. Johnston, M. A. Skinnider, N. J. Merwin and N. A. Magarvey, *Nucleic Acids Res.*, 2017, **45**, W49–W54.
- 91 H. E. Pence and A. Williams, *J. Chem. Educ.*, 2010, **87**, 1123–1124.
- 92 M. Hattori, N. Tanaka, M. Kanehisa and S. Goto, *Nucleic Acids Res.*, 2010, **38**, W652–W656.
- 93 R. G. Linington, *NP Atlas*, <https://www.npatlas.org>.
- 94 S. Farag, R. M. Bleich, E. A. Shank, O. Isayev, A. A. Bowers and A. Tropsha, *Bioinformatics*, 2019, btz127, DOI: 10.1093/bioinformatics/btz127.



- 95 W. Wriggers, S. Chakravarty and P. A. Jennings, *Biopolymers*, 2005, **80**, 736–746.
- 96 D. P. Lawrence, S. Kroken, B. M. Pryor and A. E. Arnold, *PLoS One*, 2011, **6**, e28231.
- 97 C. Elena, P. Ravasi, M. E. Castelli, S. Peirú and H. G. Menzella, *Front. Microbiol.*, 2014, **5**, 21.
- 98 F. F. V. Chevance, S. Le Guyon and K. T. Hughes, *PLoS Genet.*, 2014, **10**, e1004392.
- 99 D. Papamichail, H. Liu, V. Machado, N. Gould, J. R. Coleman and G. Papamichail, *IEEE/ACM Trans. Comput. Biol. Bioinf.*, 2018, **15**, 452–459.
- 100 Z. Zhou, J. R. Lai and C. T. Walsh, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 11621–11626.
- 101 P. Gainza, K. E. Roberts, I. Georgiev, R. H. Lilien, D. A. Keedy, C.-Y. Chen, F. Reza, A. C. Anderson, D. C. Richardson, J. S. Richardson and B. R. Donald, *Methods Enzymol.*, 2013, **523**, 87–107.
- 102 C.-Y. Chen, I. Georgiev, A. C. Anderson and B. R. Donald, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**, 3764–3769.
- 103 R. Verma, U. Schwaneberg and D. Roccato, *Comput. Struct. Biotechnol. J.*, 2012, **2**, e201209008.
- 104 M. S. Packer and D. R. Liu, *Nat. Rev. Genet.*, 2015, **16**, 379.
- 105 R. E. Cobb, R. Chao and H. Zhao, *AIChE J.*, 2013, **59**, 1432–1440.
- 106 C. A. Voigt, C. Martinez, Z.-G. Wang, S. L. Mayo and F. H. Arnold, *Nat. Struct. Biol.*, 2002, **9**, 553.
- 107 L. Sumbalova, J. Stourac, T. Martinek, D. Bednar and J. Damborsky, *Nucleic Acids Res.*, 2018, **46**, W356–W362.
- 108 B. A. Amrein, F. Steffen-Munsberg, I. Szeler, M. Purg, Y. Kulkarni and S. C. L. Kamerlin, *IUCr*, 2017, **4**, 50–64.
- 109 L. J. McGuffin, J. D. Atkins, B. R. Salehe, A. N. Shuid and D. B. Roche, *Nucleic Acids Res.*, 2015, **43**, W169–W173.
- 110 M. Kallberg, H. Wang, S. Wang, J. Peng, Z. Wang, H. Lu and J. Xu, *Nat. Protoc.*, 2012, **7**, 1511–1522.
- 111 J. Söding, A. Biegert and A. N. Lupas, *Nucleic Acids Res.*, 2005, **33**, W244–W248.
- 112 A. Fiser and A. B. T.-M. in E. Šali, in *Macromolecular Crystallography, Part D*, Academic Press, 2003, vol. 374, pp. 461–491.
- 113 L. A. Kelley, S. Mezulis, C. M. Yates, M. N. Wass and M. J. E. Sternberg, *Nat. Protoc.*, 2015, **10**, 845–858.
- 114 D. E. Kim, D. Chivian and D. Baker, *Nucleic Acids Res.*, 2004, **32**, W526–W531.
- 115 A. Waterhouse, C. Rempfer, F. T. Heer, G. Studer, G. Tauriello, L. Bordoli, M. Berton, R. Gumienny, R. Lepore, S. Bienert, T. A. P. de Beer and T. Schwede, *Nucleic Acids Res.*, 2018, **46**, W296–W303.
- 116 C. Zhang, S. M. Mortuza, B. He, Y. Wang and Y. Zhang, *Proteins*, 2018, **86**(Suppl 1), 136–151.
- 117 C. H. Eng, T. W. H. Backman, C. B. Bailey, C. Magnan, H. Garcia Martin, L. Katz, P. Baldi and J. D. Keasling, *Nucleic Acids Res.*, 2018, **46**, D509–D515.
- 118 P. Carbonell, P. Parutto, C. Baudier, C. Junot and J.-L. Faulon, *ACS Synth. Biol.*, 2014, **3**, 565–577.
- 119 M. J. Czar, Y. Cai and J. Peccoud, *Nucleic Acids Res.*, 2009, **37**, W40–W47.
- 120 S. K. Burley, H. M. Berman, G. J. Kleywegt, J. L. Markley, H. Nakamura and S. Velankar, *Methods Mol. Biol.*, 2017, **1607**, 627–641.
- 121 T. Izoré and M. J. Cryle, *Nat. Prod. Rep.*, 2018, **35**, 1120–1139.
- 122 T. Awakawa, T. Fujioka, L. Zhang, S. Hoshino, Z. Hu, J. Hashimoto, I. Kozono, H. Ikeda, K. Shin-Ya, W. Liu and I. Abe, *Nat. Commun.*, 2018, **9**, 3534.
- 123 S. Mori, A. H. Pang, T. A. Lundy, A. Garzan, O. V. Tsodikov and S. Garneau-Tsodikova, *Nat. Chem. Biol.*, 2018, **14**, 428–430.
- 124 J. Moulton, K. Fidelis, A. Kryshtafovich, T. Schwede and A. Tramontano, *Proteins*, 2018, **86**(Suppl 1), 7–15.
- 125 F. Goedegebuur, L. Dankmeyer, P. Gualfetti, S. Karkehabadi, H. Hansson, S. Jana, V. Huynh, B. R. Kelemen, P. Kruithof, E. A. Larenas, P. J. M. Teunissen, J. Ståhlberg, C. M. Payne, C. Mitchinson and M. Sandgren, *J. Biol. Chem.*, 2017, **292**, 17418–17430.
- 126 S. F. Altschul, W. Gish, W. Miller, E. W. Myers and D. J. Lipman, *J. Mol. Biol.*, 1990, **215**, 403–410.
- 127 R. D. Finn, J. Clements and S. R. Eddy, *Nucleic Acids Res.*, 2011, **39**, W29–W37.
- 128 A. Hagen, S. Poust, T. de Rond, J. L. Fortman, L. Katz, C. J. Petzold and J. D. Keasling, *ACS Synth. Biol.*, 2016, **5**, 21–27.
- 129 B. Delépine, T. Duigou, P. Carbonell and J. Faulon, *Metab. Eng.*, 2018, **45**, 158–170.
- 130 B. Hartnett, C. Gustafsson, J. Peccoud and Y. Cai, *Bioinformatics*, 2007, **23**, 2760–2767.
- 131 A. Coll, M. L. Wilson, K. Gruden and J. Peccoud, *Methods Mol. Biol.*, 2016, **1482**, 219–232.
- 132 Z. Tan, J. M. Clomburg and R. Gonzalez, *ACS Synth. Biol.*, 2018, **7**, 1886–1896.
- 133 S. Galanie, K. Thodey, I. J. Trenchard, M. Filsinger Interrante and C. D. Smolke, *Science*, 2015, **349**, 1095–1100.
- 134 X. Luo, M. A. Reiter, L. d'Espaux, J. Wong, C. M. Denby, A. Lechner, Y. Zhang, A. T. Grzybowski, S. Harth, W. Lin, H. Lee, C. Yu, J. Shin, K. Deng, V. T. Benites, G. Wang, E. E. K. Baidoo, Y. Chen, I. Dev, C. J. Petzold and J. D. Keasling, *Nature*, 2019, **567**, 123–126.
- 135 P. P. Peralta-Yahya, F. Zhang, S. B. del Cardayre and J. D. Keasling, *Nature*, 2012, **488**, 320.
- 136 D. E. Cameron, C. J. Bashor and J. J. Collins, *Nat. Rev. Microbiol.*, 2014, **12**, 381.
- 137 J. C. Blain and J. W. Szostak, *Annu. Rev. Biochem.*, 2014, **83**, 615–640.
- 138 P. Carbonell, A. Currin, A. J. Jervis, N. J. W. Rattray, N. Swainston, C. Yan, E. Takano and R. Breitling, *Nat. Prod. Rep.*, 2016, **33**, 925–932.
- 139 E. Kim, B. S. Moore and Y. J. Yoon, *Nat. Chem. Biol.*, 2015, **11**, 649.

