



Cite this: *RSC Adv.*, 2019, 9, 10781

The interaction of dietary flavonoids with xanthine oxidase *in vitro*: molecular property-binding affinity relationship aspects

Mengmeng Yuan,^{†ac} Yi Liu,^{†a} Aiping Xiao,^a Juan Leng,^a Liping Liao,^a Lei Ma^{*b} and Liangliang Liu^{id *a}

The molecular property–affinity relationships of dietary flavonoids binding to xanthine oxidase were investigated *in vitro* by comparing the binding constants obtained from a fluorescence-quenching method. The inhibitions of dietary flavonoids on xanthine oxidase were also investigated and analyzed, revealing that the binding process was influenced by the structural differences of the flavonoids under investigation. For example, methylation and hydroxylation at the 7- and 5-positions weakened the binding affinities, while hydroxylation at the 3- and 3'-positions mostly improved binding affinities. Glycosylation and hydrogenation of the C₂=C₃ double bond also increased affinities for xanthine oxidase. In addition, galloylated catechins showed higher binding affinities than non-galloylated catechins. Trends in the binding affinities and inhibition of flavonoids during structure modifications were summarized. Affinities for xanthine oxidase and inhibition on xanthine oxidase changed in the opposite direction during the methylation and hydroxylation of flavonoids in the A ring, and the glycosylation and hydrogenation of C₂=C₃. However, affinities and inhibition for xanthine oxidase changed in the same direction during the methylation and hydroxylation of flavonoids in the B ring.

Received 3rd December 2018
 Accepted 25th March 2019

DOI: 10.1039/c8ra09926j

rsc.li/rsc-advances

1. Introduction

Dietary flavonoids exist widely in many plant foods, such as fruits, vegetables, nuts, and tea.^{1,2} Over 10 000 flavonoids have been separated and identified from plants, including anthocyanidins, flavones, flavonols, flavanones, and isoflavones.³ Investigation of flavonoids from dietary sources has attracted considerable interest because of their nutritional and medical effects in humans. Most of their bioactivity is related to their structure;^{4,5} for example, the presence of a 2,3 double bond in conjugation with a 4-oxo group and a catechol unit (1,2-dihydroxybenzene) is usually required for antioxidant activity in flavonoids. Methylation, hydroxylation, and glycosylation of flavonoids also affect their absorption, metabolism, and bioactivities *in vivo*.^{6,7} Furthermore, structural differences in flavonoids strongly affect the binding process with plasma proteins, which are linked to the chemistry of the C ring, the number and distribution of hydroxyl groups, and their substitutions on the A and B rings.⁸ According to reports, the

methylation of the free hydroxyl groups in the flavonoids dramatically increases their intestinal absorption and metabolic resistance and transport of flavonoids.⁹

Fluorescence-quenching spectroscopy is an important method to determine the binding parameters between ligands and bio-macromolecules.^{10,11} It refers to the decrease in the quantum yield of fluorescence from a fluorophore induced by a variety of molecular interactions with a quencher molecule under excitation at 280 nm.¹² In recent years, studies on fluorescence quenching have been widely reported in the literature. Frazier *et al.* investigated the interaction of four different flavonoids with bovine serum albumin (BSA) by quenching the intrinsic tryptophan fluorescence in order to study the chemical associations.¹³ Xiao *et al.* investigated the interaction of puerarin and BSA by means of fluorescence spectroscopy.¹⁴ Shao *et al.* investigated the structure–affinity relationship in the interactions of different oleanane-type triterpenoids with BSA by spectroscopic analysis. In addition to the research focusing on the binding process, the structure–affinity relationships of bioactive compounds for proteins such as common rat plasma proteins, common human plasma proteins, γ -globulin, total plasma proteins, BSA and HSA were also investigated.^{15–19}

Xanthine oxidase could catalyze the oxidation of hypoxanthine to xanthine and further catalyze the oxidation of xanthine to uric acid, which plays a crucial role in gout.²⁰ Excess accumulation of uric acid in serum will lead to hyperuricemia, which is a risk factor for cardiovascular disease, gout and other

^aInstitute of Bast Fiber Crops, Chinese Academy of Agricultural Sciences, Changsha 410205, China. E-mail: liuliangliang@caas.cn

^bZhengzhou Research Base, State Key Laboratory of Cotton Biology, Zhengzhou University, China. E-mail: malei@caas.cn

^cCollege of Chemistry and Chemical Engineering, Central South University, Changsha 410083, China

[†] These authors contributed equally to this work.



metabolic disorders.²¹ Thus, the investigation of xanthine oxidase inhibitors could be an effective strategy in the prevention of these metabolic disorders.²² Fluorescence-quenching spectroscopy was also used in research of xanthine oxidase. By analyzing fluorescence parameters, considerable information relating to structural changes in enzymes could be obtained.²³ Rashidi *et al.* first studied the interaction of quercetin and bovine milk xanthine oxidase using a fluorescence-quenching method.²⁴ Lin *et al.* reported that five dietary flavonoids could interact with xanthine oxidase at more than one binding site through fluorescence-quenching data.²⁰ However, there were no reports on structure–affinity relationships in the interactions of different flavonoids with xanthine oxidase.

In our previous research, xanthine oxidase inhibitors from natural plants were screened and identified.^{25,26} The binding properties of tectoridin, daidzin, ononin, biochanin A, gardenin B, and eupatorin for xanthine oxidase were also studied. Based on these results, we became interested in detailed research on the binding between activity compounds and xanthine oxidase, and a series of flavonoids with similar structures were selected






for research. In the present study, xanthine oxidase was used as a modeling protein in order to investigate the interaction between small molecules and protein, and 35 dietary flavonoids (shown in Table 1) were selected to investigate the interaction with xanthine oxidase.

2. Experimental

2.1. Materials

Xanthine oxidase powder was purchased from Yuanye Biotechnology Co. (Shanghai, China). Flavone (99.0%), 7-OH flavone (98.0%), chrysin (98.0%), biochanin A (98.0%), naringin (98.0%), and myricetin (98.0%) were bought from Alfa Aesar (Thermo Fisher Scientific, MA, USA). Formononetin (99.0%) and genistein (99.0%) were acquired from Acros Organics (Fisher Scientific, PA, USA). Baicalin (98.0%), baicalein (98.0%), wogonin (98.0%), apigenin (98.0%), luteolin (98.0%), hispidulin (98.0%), tangeretin (98.0%), nobiletin (98.0%), galangin (98.0%), kaempferide (98.0%), kaempferol (98.0%), kaempferitrin (98.0%), quercetin (98.0%), fisetin (98.0%), rutin (98.0%),

Table 1 Chemical structures of the 35 investigated dietary flavonoids

Subclass	Name	Substitutions			Structure
		OH	OCH ₃	Others	
Flavones	Flavone				
	7-OH flavone	7			
	Chrysin	5,7			
	Baicalin	5,6			
	Baicalein	5,6,7			
	Wogonin	5,7	8		
	Apigenin	5,7,4'			
	Luteolin	5,7,3',4'			
	Hispidulin	5,7,4'	6		
	Tangeretin		5,6,7,8,4'		
	Nobiletin		5,6,7,8,4',5'		
Flavonols	Galangin	3,5,7			
	Kaempferide	3,5,7	4'		
	Kaempferol	3,5,7,4'			
	Kaempferitrin	5,4'		3,7-Dirhamnoside	
	Quercetin	3,5,7,3',4'			
	Myricetin	3,5,7,3',4',5'			
	Fisetin	3,7,3',4'			
	Rutin	5,7,3',4'		3- α -L-Rham-1,6-D-Glc	
	Formononetin	7	4'		
	Genistein	5,7,4'			
Isoflavones	Daidzein	7,4'			
	Biochanin A	5,7	4'		
	Tectorigenin	5,7,4'	6		
	Puerarin	7,4'		8-C-Glucoside	
	Dihydromyricetin	3,5,7,3',4',5'			
Flavanone	Naringenin	5,7,4'			
	Naringin	5,4'		7-Neohesperidose	
Flavanonol	C (2,3-trans)	3,5,7,4',5'			
	EC (2,3-cis)	3,5,7,4',5'			
	ECG (2,3-cis)	5,7,4',5'		3-Gallate	
	EGC (2,3-cis)	3,5,7,3',4',5'			
	EGCG (2,3-cis)	5,7,3',4',5'		3-Gallate	
	GC (2,3-trans)	3,5,7,3',4',5'			
	GCG (2,3-trans)	5,7,3',4',5'		3-Gallate	



daidzein (98.0%), tectorigenin (98.0%), puerarin (98.0%), dihydromyricetin (98.0%), naringenin (98.0%), catechin (C; 98.0%), (–)-gallocatechin (GC; 98.0%), epicatechin, (EC; 98.0%), (–)-epicatechin gallate (ECG; 98.0%), (–)-epigallocatechin (EGC; 98.0%), (–)-epigallocatechin gallate (EGCG; 98.0%), and (–)-gallocatechin gallate (GCG; 98.0%) were commercially purchased from Yuanye Biotechnology Co. (Shanghai, China). High-performance liquid chromatography (HPLC) grade acetonitrile was purchased from Merck KGaA (Darmstadt, Germany). Ultrapure water (18.2 MΩ cm resistivity) was obtained from an ELGA water purification system (ELGA Berkefeld, Veolia, Germany). All other reagents and solvents were of analytical grade.

2.2. Apparatus

Fluorescence spectra were recorded on a fluorometer (Hitachi F-7000, Tokyo, Japan) using 1.0 cm of quartz cuvette. The thermostatic control of incubation was completed with a thermostat water bath (MS7-H550-Pro, DLAB Scientific Inc., Beijing, China) controlled at 25 °C. The air conditioner was set at 25 °C to minimize interference during detection. Ultraviolet-visible (UV-vis) spectra and inhibition tests were recorded on an UV-vis spectrophotometer using 1.0 cm of quartz cuvette (UV2700, Shimadzu, Kyoto, Japan). Qualitative analysis of dietary flavonoids was completed by HPLC analysis on an Agilent 1260 HPLC system (Agilent Technologies, Santa Clara, CA, USA) equipped with a C₁₈ reverse-phase column (Waters Xbridge™, 250 mm × 4.6 mm i.d., 5 μm, Milford, MA, USA).

2.3. Inhibition test

First, 20 μL of xanthine oxidase solution (3.70 nmol L⁻¹) and a sample with a different volume (10 to 100 μL) were mixed in a quartz cuvette,²⁶ and water was added to make the total volume of mixture up to 1.0 mL. Then, 1.0 mL of xanthine solution (0.45 mmol L⁻¹) was added to start the enzymatic reaction. The cuvette was rapidly moved into a UV-vis spectrophotometer, and the absorbance at 295 nm was monitored. The same amount of water was used as in the blank, under the same conditions. The inhibition of xanthine oxidase could be calculated using the following formula:

$$\text{Inhibition \%} = (1 - \Delta A_s / \Delta A_b) \times 100\% \quad (1)$$

where ΔA_s and ΔA_b are the increase of absorbance for the sample and blank, respectively. Inhibition of the sample was expressed as the concentration of sample needed to inhibit 50% of enzymatic activity (IC₅₀). The detection limit of xanthine oxidase activity was 1.0 U L⁻¹. The detection limit of related inhibitors was in the range 0.05 nmol L⁻¹ to 5.0 mmol L⁻¹. All assays were performed with three replicates.

2.4. Fluorescence spectra analysis

The fluorescence spectra of xanthine oxidase and the quenching effects of samples were performed according to procedures reported in previous literature.^{27,28} Various volumes of samples (0–100 μL, 1.0 mmol L⁻¹ in methanol) were mixed with 2.0 mL of

xanthine oxidase solution (1.0 μmol L⁻¹ in phosphate buffered saline). After incubation at 25 °C for 5 min, the fluorescence emission spectra of samples were recorded from 300 to 450 nm under an excitation wavelength at 280 nm using a fluorometer. All assays were performed with three replicates.

2.5. Data processing

The fluorescence quenching of xanthine oxidase with samples were described by the Stern–Volmer formula shown below:^{18,29}

$$F_0/F = 1 + K_q\tau_0[Q] = 1 + K_{sv}[Q] \quad (2)$$

where F_0 and F are the fluorescence intensity of xanthine oxidase at 335 nm in the absence and presence of sample, $[Q]$ is the concentration of the sample, K_q is the quenching rate constant, τ_0 is the average lifetime (6.2 ns), and K_{sv} is the dynamic quenching constant. The binding constants were calculated according to the double-logarithm formula:

$$\log_{10}[(F_0 - F)/F] = \log_{10} K_a + n \log_{10}[Q] \quad (3)$$

where K_a is the binding constant and n is the number of binding sites per enzyme molecule. According to formulas (2) and (3), the values of “ F_0/F ” and “ $(F_0 - F)/F$ ” could be observed in each “[Q]” of samples. Then, the linear regression equation between the “ F_0/F ” values and “[Q]” values was obtained, and the slope means “ K_{sv} ”. The linear regression equation between the “ $\log_{10}[(F_0 - F)/F]$ ” and “ $\log_{10}[Q]$ ” values could be obtained. The slope means “ n ” and the means “ $\log_{10} K_a$ ”.

2.6. Statistical analysis

The experiments under each condition were performed in triplicate. The experimental errors were less than 5.0%. The results were presented as the means ± standard deviation (SD), and analyzed by one-way analysis of variance (ANOVA) and Duncan's *post hoc* analysis. $p < 0.05$ was considered to be statistically significant.

3. Results and discussion

3.1. Inhibition of flavonoids on xanthine oxidase

The inhibitions of 35 dietary flavonoids on xanthine oxidase were investigated. Inhibition was expressed as IC₅₀ values in Table 2, and all 35 dietary flavonoids exhibited some degree of inhibition on xanthine oxidase except kaempferide. Quercetin showed the highest inhibitory activity at 0.0001 μmol L⁻¹, while puerarin showed the lowest inhibitory activity at 2.1074 μmol L⁻¹. Therefore, these dietary flavonoids showed good inhibition on xanthine oxidase.

3.2. Fluorescence-quenching effects

Fluorescence quenching is a common bioanalytical method widely used in investigating the interaction between biological macromolecules and small molecules. The fluorescence emission spectra of xanthine oxidase with the addition of various dietary flavonoids were monitored, and six of these flavonoids



Table 2 Affinities, inhibitions and property parameters of flavonoids for xanthine oxidase^a

Name	Inhibition IC ₅₀ (μmol L ⁻¹)	Affinity				TPSA (Å ²)	X log P ₃
		K _{sv} (10 ⁶)	K _q (10 ¹³)	log ₁₀ K _a	<i>n</i>		
Flavone	0.3464	0.0552	0.8903	6.3124	1.3695	26.3	3.6
7-OH flavone	0.1505	0.0360	0.5806	4.4265	0.9676	46.5	3.6
Chrysin	0.0138	0.0195	0.3145	2.5792	0.6025	66.8	2.1
Baicalin	0.0267	0.3668	5.9161	8.9699	1.8782	183.0	1.1
Baicalein	0.0258	0.1981	3.1951	7.3561	1.5312	87.0	1.7
Wogonin	0.0174	—	—	—	—	76.0	3.0
Apigenin	0.0007	0.0285	0.4596	2.1428	0.4524	87.0	1.7
Luteolin	0.0003	0.0926	1.4935	6.2828	1.3087	107.0	1.4
Hispidulin	0.0159	0.0989	1.5951	6.1617	1.2571	96.2	1.7
Tangeretin	1.7885	0.1498	2.4161	5.1591	1.0022	72.4	3.0
Nobiletin	0.2647	0.9637	15.5435	14.019	2.8371	81.7	3.0
Galangin	0.0090	0.0276	0.4451	4.0235	0.9038	87.0	2.3
Kaempferide	ND	—	—	—	—	96.2	2.2
Kaempferol	0.0050	0.0225	0.3629	4.5889	1.0535	107.0	1.9
Kaempferitrin	0.2788	0.0929	1.4983	7.0817	1.5204	225.0	-0.1
Quercetin	0.0001	0.0477	0.7693	4.9449	1.0614	127.0	1.5
Myricetin	0.0086	0.0294	0.4741	4.7891	1.0764	148.0	1.2
Fisetin	0.0063	0.0729	1.1758	6.4458	1.3663	107.0	2.0
Rutin	1.0333	0.0532	0.8580	5.9546	1.2904	266.0	-1.3
Formononetin	0.0719	0.0147	0.2370	2.8576	0.6866	55.8	2.8
Genistein	0.2508	0.0537	0.8661	6.3981	1.3911	87.0	2.7
Daidzein	0.1799	0.0230	0.3709	5.4529	1.2554	66.8	2.5
Biochanin A	0.1570	0.0359	0.5790	0.3808	0.7052	76.0	3.0
Tectorigenin	0.1426	0.0644	1.0387	3.0234	0.6283	96.2	2.6
Puerarin	2.1074	0.0290	0.4677	5.2024	1.1801	157.0	0.0
Dihydromyricetin	0.0428	0.0470	0.7580	5.6369	1.2262	148.0	1.1
Naringenin	0.1436	0.0568	0.9161	5.8127	1.2510	87.0	2.4
Naringin	0.3267	0.0846	1.3645	7.6780	1.6838	225.0	-0.5
C (2,3-trans)	0.2591	—	—	—	—	110.0	0.4
EC (2,3-cis)	0.2774	—	—	—	—	110.0	0.4
ECG (2,3-cis)	0.2610	0.0193	0.3112	4.7240	1.099	177.0	1.5
EGC (2,3-cis)	0.2510	—	—	—	—	131.0	0.0
EGCG (2,3-cis)	0.2805	0.0226	0.3645	5.3153	1.2461	197.0	1.2
GC (2,3-trans)	0.2819	—	—	—	—	131.0	0.0
GCG (2,3-trans)	0.0063	0.0178	0.2871	4.3766	1.0301	197.0	1.2

^a TPSA: values were obtained from <https://pubchem.ncbi.nlm.nih.gov/compound/>, and the units are Å².

are shown in Fig. 1. Xanthine oxidase solution showed an apparent fluorescence emission peak at 335 nm, and the fluorescence intensities reduced gradually with increasing addition of dietary flavonoids in XO solution.

In order to evaluate and compare the quenching effects of 35 dietary flavonoids, Stern–Volmer plots for fluorescence quenching were used to find further quenching information. The Stern–Volmer plots of six flavonoids are shown in Fig. 2, and the calculated parameters of 35 dietary flavonoids are demonstrated in Table 2. As shown in Fig. 2 and Table 2, the degrees of fitting of the Stern–Volmer plots were greater than 0.9, reflecting that the calculations of K_{sv} and K_q were acceptable. High linearities of the Stern–Volmer plots indicated that one quenching mode played a key role in the interaction, and K_q was the factor indicating the efficiency of quenching. When K_q is apparently greater than 2.0×10^{10} L mol⁻¹ s⁻¹, it could be considered as static quenching;³⁰ otherwise it would be considered dynamic quenching. In this study, the values of K_q of 35 dietary flavonoids were much

greater than 2.0×10^{10} L mol⁻¹ s⁻¹. The lowest value of K_q was 0.237 L mol⁻¹ s⁻¹ for formononetin. Therefore, based on this, the quenching of 35 dietary flavonoids could be considered as static quenching.

3.3. Binding parameters

For static quenching, the number of binding sites per protein molecule (n) and binding constants (K_a) could be calculated by fitting the double-logarithm curves. The values of n and K_a are also shown in Table 2 and the plots of $\log[(F_0 - F)/F]$ versus $\log[Q]$ of six representative flavonoids are shown in Fig. 3. The degrees of fitting of these calculations were higher than 0.9882, and the linearity of curves indicated that the calculations were acceptable.³¹ The plots of $\log_{10} K_a$ versus n of flavonoids are shown in Fig. 4. The values of $\log_{10} K_a$ were linear to the number of binding sites (n) (fitting curve: $\log_{10} K_a = 0.182n + 0.212$, $R^2 = 0.9498$). This indicates that the equation used in this study was suitable for the interaction between flavonoids and xanthine oxidase. Thirty-five dietary






Fig. 1 Quenching effects of (a) quercetin, (b) myricetin, (c) daidzein, (d) puerarin, (e) dihydromyricetin, and (f) ECG (2,3-*cis*) on xanthine oxidase fluorescence spectra. $\lambda_{\text{ex}} = 280 \text{ nm}$; xanthine oxidase, $1.0 \mu\text{mol L}^{-1}$; the addition of sample was 0 (black), 20 (red), 40 (green), 60 (blue), 80 (light blue), and 100 μL (purple).




Fig. 2 Stern–Volmer plots for xanthine oxidase fluorescence quenching by quercetin, myricetin, daidzein, puerarin, dihydromyricetin, and ECG (2,3-*cis*). The degrees of fitting were 0.9978 for quercetin, 0.9972 for myricetin, 0.9893 for daidzein, 0.9895 for puerarin, 0.9906 for dihydromyricetin, and 0.9994 for ECG.




Fig. 3 Plots of $\log[(F_0 - F)/F]$ versus $\log[Q]$ for quercetin, myricetin, daidzein, puerarin, dihydromyricetin, and ECG (2,3-*cis*). The degrees of fitting were 0.9985 for quercetin, 0.9991 for myricetin, 0.9987 for daidzein, 0.9964 for puerarin, 0.9996 for dihydromyricetin and 0.9988 for ECG.

flavonoids showed a wide range of $\log_{10} K_a$ values (from 0.3808 to 14.019), and further comparison of $\log_{10} K_a$ values would be described.




Fig. 4 Relationship between affinities ($\log_{10} K_a$) and the number of binding sites (n).

3.4. Structure–affinity relationship of flavonoids-xanthine oxidase interactions

3.4.1. Methylation. Table 3 lists the effects of an additional methoxy group (hydrogen to methoxy group and hydroxyl group to methoxy group) in flavonoids on the affinities and inhibition for xanthine oxidase. Generally, the addition of a methoxy group in flavonoids weakened the binding affinities for xanthine oxidase by 268 to 10^6 times. At the extreme, the affinity of genistein for xanthine oxidase was found to be 10^6 times higher than that of biochanin A, which was the methylated form at the 4'-position. This kind of trend was reported in the interaction between flavonoids and bovine γ -globulin.⁹ The methylation increased the hydrophobicity of flavonoids, and the hydrophobic interaction plays an important role in binding to enzymes.³² However, the methylation of flavone, apigenin, and tangeretin increased the affinities for xanthine oxidase. For the inhibition on xanthine oxidase, the addition of a methoxy group in flavonoids mostly increased the inhibitory activities.

3.4.2. Hydroxylation. Table 4 showed the effect of hydroxylation of flavonoids on affinities for xanthine oxidase. It could be seen that the hydroxylation on rings A, B, and C of flavonoids affected the binding affinities for xanthine oxidase. Hydroxylation on the 7-position of flavone and flavanone weakened binding affinities, while hydroxylation on the 5-position of flavonoids mostly (3 out of 4) decreased binding affinities. However, hydroxylation on the 3-position (2 out of 3) and 3'-position (2 out of 2) of flavonoids mostly (3 out of 4) improved binding affinities. In terms of inhibition, hydroxylation on the 7-, 3', and 4'-positions of flavonoids enhanced inhibitory activities on xanthine oxidase, while hydroxylation on the 3-position (2 out of 3) of flavonoids improved inhibitory activities.

3.4.3. Glycosylation. The effect of glycosylation of flavonoids on the affinities for xanthine oxidase was investigated (Table 5). Herein, the involved sugar moieties were rhamnose, glucose-rhamnose, and neohesperidose. The 3,7-glycosylation of kaempferol increased the affinity of kaempferol for xanthine oxidase by 311 times. 3-Glycosylation of quercetin increased the affinity by 10.2 times, 7-glycosylation of naringenin increased the affinity by 73.3 times, while the inhibition of kaempferol, quercetin, and naringenin decreased after glycosylation.



Table 3 Effects of methylation of flavonoids on affinities

Class	Ring	Position	Example	Affinity	Inhibition	
Flavone	A	6H → OCH ₃	Apigenin → hispidulin	↑	↓	
		8H → OCH ₃	Genistein → tectorigenin	↓	↑	
		3'H → OCH ₃	Chrysin → wogonin	↓	↑	
Flavonol	B	4'OH → OCH ₃	Tangeretin → nobiletin	↑	↑	
		4'H → OCH ₃	Kaempferol → kaempferide	↓	↓	
Isoflavone	A	6H → OCH ₃	Galangin → kaempferide	↓	↓	
		B	4'OH → OCH ₃	Daidzein → tectorigenin	↓	↑
			4'OH → OCH ₃	Daidzein → formononetin	↓	↑
			Genistein → biochanin A	↓	↑	

Table 4 Effects of hydroxylation of flavonoids on affinities

Class	Ring	Position	Example	Affinity	Inhibition	
Flavone	A	5H → OH	7-OH flavone → chrysin	↓	↑	
		6H → OH	Chrysin → baicalein	↑	↓	
		7H → OH	Flavone → 7-OH flavone	↓	↑	
	B	3'H → OH	Baicalin → baicalein	↓	↑	
		4'H → OH	Apigenin → luteolin	↑	↑	
		3H → OH	Chrysin → apigenin	↓	↑	
Flavonol	C	3H → OH	Chrysin → galangin	↑	↑	
			Apigenin → kaempferol	↑	↓	
			Luteolin → quercetin	↓	↑	
	A	5H → OH	Fisetin → quercetin	↓	↑	
		B	3'H → OH	Kaempferol → quercetin	↑	↑
			4'H → OH	Galangin → kaempferol	↑	↑
Isoflavone	A	5'H → OH	Quercetin → myricetin	↓	↓	
		5H → OH	Daidzein → genistein	↑	↓	
Flavanone	A		Formononetin → biochanin A	↓	↓	
		7H → OH	Naringin → naringenin	↓	↑	

3.4.4. Hydrogenation of the C₂=C₃ double bond. The C₂=C₃ double bond in conjugation with a 4-oxo group plays a very important role in the affinity for proteins.³³ It was found in Table 5 that the hydrogenation of the C₂=C₃ double bond of flavonoids increased the binding affinities for xanthine oxidase. The affinities of dihydromyricetin and naringenin for xanthine oxidase were 7- and 4676-times higher than those of myricetin and dihydromyricetin apigenin, respectively. This trend is opposite to the affinities for α -amylase and total plasma proteins.^{17,33} The inhibition of dihydromyricetin and naringenin also decreased after hydrogenation of the C₂=C₃ double bond.

Table 5 Effects of glycosylation and hydrogenation of the C₂=C₃ double bond of flavonoids on affinities

	Example	Affinity	Inhibition
Glycosylation	Kaempferol → kaempferitrin	↑	↓
	Quercetin → rutin	↑	↓
	Naringenin → naringin	↑	↓
Hydrogenation of C ₂ =C ₃	Myricetin → dihydromyricetin	↑	↓
	Apigenin → naringenin	↑	↓

3.4.5. Catechins. Catechins are the major active components in green tea leaves, including (–)-C, (–)-GC, EC, (–)-ECG, (–)-EGC, (–)-EGCG, and (–)-GCG. The affinities between catechins and xanthine oxidase were determined, and the binding constants (log₁₀ K_a) of ECG, EGCG, and GCG for xanthine oxidase were found to be 4.7240, 5.3153, and 4.3766, respectively (Table 2). However, C, EC, EGC, and GC were not able to quench the fluorescence of xanthine oxidase, and there was no binding constant value. This indicated the galloylated catechins have higher binding affinities for xanthine oxidase than non-galloylated catechins, and the presence of the galloyl moiety became the key factor. In addition, the affinity of gallocatechin gallate with a 2,3-*trans* structure was lower than that of gallocatechin gallate with a 2,3-*cis* structure. In terms of inhibition, the changes in inhibitory activities of catechins were not apparent.

3.5. Comparison of trends in the binding affinities and inhibition

Table 6 summarizes and compares trends in the binding affinities and inhibition of flavonoids during the structure modifications previously mentioned. In the methylation and hydroxylation of flavonoids in the A ring, the changes in



Table 6 Comparison of trends in the binding affinities and inhibition of flavonoids for xanthine oxidase

Modification	Ring	Samples	Affinity		Inhibition		Trends	
			↑	↓	↑	↓	Accordance	Opposite
Methylation	A	4	1	3	3	1	0	4
	B	5	1	4	1	4	3	2
Hydroxylation	A	8	2	6	4	4	1	7
	B	5	3	2	4	1	4	1
	C	3	2	1	2	1	1	2
Glycosylation		3	3	0	0	3	0	3
Hydrogenation of C ₂ =C ₃		2	2	0	0	2	0	2

affinities and inhibition were almost all opposite (11 out of 12), while in the same structure modifications of flavonoids in the B ring, the changes of affinities and inhibition were almost all in accordance (7 out of 10). Moreover, in the glycosylation and hydrogenation of C₂=C₃ of flavonoids, the changes of affinities and inhibition were all opposite. Based on these findings, it might be concluded that the affinities for xanthine oxidase and the inhibition on xanthine oxidase would change in the opposite direction during methylation and hydroxylation of flavonoids in the A ring, and glycosylation and hydrogenation of C₂=C₃. However, the affinities and inhibition for xanthine oxidase would change in the same direction during methylation and hydroxylation of flavonoids in the B ring.

3.6. Relationship of partition coefficients and the binding affinities

The partition coefficient is a quantitative descriptor of lipophilicity and is important in related pharmacokinetic research. The partition coefficient values ($X \log P_3$) of flavonoids could be obtained through the PubChem website (<https://pubchem.ncbi.nlm.nih.gov/compound/>). As shown in Fig. 5, the $X \log P_3$ values of flavonoids were inversely proportional to the corresponding $\log_{10} K_a$ values. The linear regression curve was $X \log P_3 = -0.073 \log_{10} K_a +$




Fig. 5 Relationship between binding constants ($\log_{10} K_a$) and partition coefficients ($X \log P_3$) of flavonoids.




Fig. 6 Relationship between binding constants ($\log_{10} K_a$) and TPSA of flavonoids.

2.154 ($R = 0.151$). The trend was the same as in the reported literature,³⁴ while the degree of fitting was a little lower.

3.7. Relationship of topological polar surface area (TPSA) and binding affinities

The TPSA was defined as the sum of surfaces of polar atoms in a molecule. This property has been shown to correlate with the human intestinal absorption, Caco-2 monolayers permeability and blood-brain barrier penetration.¹⁹ The relationship between TPSA and the binding affinity for proteins and enzymes were often studied. Herein, the TPSA values of flavonoids were also obtained through the PubChem website (<https://pubchem.ncbi.nlm.nih.gov/compound/>). As a result, TPSA values of flavonoids were proportional to $\log_{10} K_a$ values (Fig. 6). The linear regression curve was $TPSA = 5.090 \log_{10} K_a + 92.621$ ($R = 0.205$), which is consistent with reported trend.³² This might illustrate that flavonoids with high TPSA bind tightly with xanthine oxidase.

4. Conclusions

Based on the investigation of the interactions between 35 dietary flavonoids and xanthine oxidase, much analysis has been performed on the relationship between the structure properties of dietary flavonoids and their affinities for xanthine oxidase.



Some structural influences on the affinities were concluded; methylation of hydroxyl groups weakened the affinities for xanthine oxidase and hydroxylation at the 7- and 5-positions weakened the binding affinities. However, hydroxylation at the 3- and 3'-positions mostly improved binding affinities. Glycosylation hydrogenation of the C₂=C₃ double bond also increased the affinities for xanthine oxidase. In addition, galloylated catechins showed higher binding affinities than non-galloylated catechins. The trends in the binding affinities and inhibition of flavonoids during the structure modifications were summarized; the affinities for xanthine oxidase and inhibition on xanthine oxidase change in the opposite direction during the methylation and hydroxylation of flavonoids in the A ring, and the glycosylation and hydrogenation of C₂=C₃. However, the affinities and inhibition for xanthine oxidase change in the same direction during the methylation and hydroxylation of flavonoids in the B ring. In order to further explore the effects of methylation in terms of the binding and inhibition of proteins, some flavonoids and their methylation derivatives were selected as the studied samples. The inhibition and binding parameters, using further spectroscopic methods, inhibition kinetics and docking analyses, of several enzymes were subject to investigation.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was supported by the risk assessment of agricultural products quality and safety project (GJFP2019010) and the Natural Science Foundation of Hunan Province (2017JJ3348).

References

- H. Cao, L. Chen and J. Xiao, *Mol. Biol. Rep.*, 2010, **38**, 2257–2262.
- W. Jiang and M. Hu, *RSC Adv.*, 2012, **2**, 7948–7963.
- J. Xiao, G. Kai, K. Yamamoto and X. Chen, *Crit. Rev. Food Sci. Nutr.*, 2013, **53**, 818–836.
- F. Yang, Y. Zhao, G. Kai and J. Xiao, *Food Funct.*, 2011, **2**, 137–141.
- H. Rimac, Ž. Debeljak, D. Šakić, T. Weitner, M. Gabričević, V. Vrček, B. Zorc and M. Bojić, *RSC Adv.*, 2016, **6**, 75014–75022.
- J. Xiao, L. Chen, F. Yang, C. Liu and Y. Bai, *J. Hazard. Mater.*, 2010, **182**, 696–703.
- J. Xiao, T. Chen, L. Chen, H. Cao, F. Yang and Y. Bai, *J. Inorg. Biochem.*, 2010, **104**, 1148–1155.
- J. B. Xiao, Y. R. Zhao, H. Wang, Y. M. Yuan, F. Yang, C. Zhang and G. Y. Kai, *Integr. Biol.*, 2011, **3**, 1087–1094.

- J. Xiao, G. Kai, F. Yang, C. Liu, X. Xu and K. Yamamoto, *Mol. Nutr. Food Res.*, 2011, **55**, S86–S92.
- J. Xiao, H. Cao, Y. Wang, K. Yamamoto and X. Wei, *Mol. Nutr. Food Res.*, 2010, **54**, S253–S260.
- B. Tu, Z.-F. Chen, Z.-J. Liu, R.-R. Li, Y. Ouyang and Y.-J. Hu, *RSC Adv.*, 2015, **5**, 73290–73300.
- J. R. Lakowicz and B. R. Masters, *Naturwissenschaften*, 1991, **78**, 456.
- A. Papadopoulou, R. J. Green and R. A. Frazier, *J. Agric. Food Chem.*, 2005, **53**, 158–163.
- J. Xiao, J. Shi, H. Cao, S. Wu, F. Ren and M. Xu, *J. Pharm. Biomed. Anal.*, 2007, **45**, 609–615.
- H. Cao, L. Chen and J. Xiao, *Mol. Biol. Rep.*, 2011, **38**, 2257–2262.
- J. Xiao, H. Cao, Y. Wang, K. Yamamoto and X. Wei, *Mol. Nutr. Food Res.*, 2010, **54**, S253–S260.
- J. Xiao, Y. Zhao, H. Wang, Y. Yuan, F. Yang, C. Zhang and G. Kai, *Integr. Biol.*, 2011, **3**, 1087–1094.
- J. Xiao, Y. Zhao, H. Wang, Y. Yuan, F. Yang, C. Zhang and K. Yamamoto, *J. Agric. Food Chem.*, 2011, **59**, 10747–10754.
- F. Yang, Y. Zhao, G. Kai and J. Xiao, *Food Funct.*, 2011, **2**, 137–141.
- Y. Dong, H. Huang, M. Zhao, D. Sun-Waterhouse, L. Lin and C. Xiao, *J. Funct. Foods*, 2016, **24**, 26–36.
- A. Nagao, M. Seki and H. Kobayashi, *Biosci., Biotechnol., Biochem.*, 1999, **63**, 1787–1790.
- T. Hatano, T. Yasuhara, R. Yoshihara, Y. Ikegami, M. Matsuda, K. Yazaki, I. Agata, S. Nishibe, T. Noro, M. Yoshizaki and T. Okuda, *Planta Med.*, 1991, **57**, 83–84.
- J. Xiao, F. Wang, J. Liu, L. Wang, G. Kai and X. Yu, *Mol. Biosyst.*, 2011, **7**, 2452–2458.
- F. Rasoulzadeh, H. N. Jabary, A. Naseri and M. R. Rashidi, *Spectrochim. Acta, Part A*, 2009, **72**, 190–193.
- L. L. Liu, A. P. Xiao, L. Ma and D. F. Li, *J. Braz. Chem. Soc.*, 2017, **28**, 360–366.
- L. Liu, M. Yuan, S. Huang, J. Li, D. Li and L. Zhao, *Appl. Sci.*, 2018, **8**, 158.
- X. Tang, P. Tang and L. Liu, *Molecules*, 2017, **22**, 1036.
- G. H. Wang, X. Y. Huang, D. Pei, W. D. Duan, K. J. Quan, X. T. Li and D. L. Di, *New J. Chem.*, 2016, **40**, 3885–3891.
- B. Tu, Z.-J. Liu, Z.-F. Chen, Y. Ouyang and Y.-J. Hu, *RSC Adv.*, 2015, **5**, 106171–106181.
- S. Li, L. Tang and H. Bi, *Luminescence*, 2016, **31**, 442–452.
- J. Xiao, E. Capanoglu, A. R. Jassbi and A. Miron, *Crit. Rev. Food Sci. Nutr.*, 2016, **56**, S29–S45.
- J. Xiao, H. Cao, T. Chen, F. Yang, C. Liu and X. Xu, *Biochimie*, 2011, **93**, 134–140.
- J. Xiao, G. Kai, X. Ni, F. Yang and X. Chen, *Mol. Biosyst.*, 2011, **7**, 1883–1890.
- J. Xiao, T. Chen, H. Cao, L. Chen and F. Yang, *Mol. Nutr. Food Res.*, 2010, **55**, 310–317.

