




Cite this: *RSC Adv.*, 2019, 9, 7961

Hyperpatulones A–F, polycyclic polyprenylated acylphloroglucinols from *Hypericum patulum* and their cytotoxic activities†

Zhong-Nan Wu,^a Qian-Wen Niu,^a Yu-Bo Zhang,^{ab} Ding Luo,^a Qing-Guo Li,^c Ying-Ying Li,^a Guang-Kai Kuang,^a Li-Jun He,^a Guo-Cai Wang ^{*ab} and Yao-Lan Li^{*a}

Six new compounds, hyperpatulones A–F (1–6), along with ten additional known related derivatives (7–16), were isolated from *Hypericum patulum* (Guttiferae). Their structures were elucidated by extensive analysis of spectroscopic data (IR, UV, HRESIMS, 1D and 2D NMR), X-ray crystallography, electronic circular dichroism (ECD) spectroscopy and Rh₂(OCOFCF₃)₄-induced ECD. All compounds were tested for their cytotoxic activities on human HepG-2, HeLa, MCF-7, and A549 cell lines via 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. Compound 5 exhibited significant cytotoxicities against HepG-2, HeLa and A549 cell lines with IC₅₀ values of 9.52 ± 0.27, 11.87 ± 0.22 and 12.63 ± 0.12 μM, respectively.

Received 12th January 2019
 Accepted 2nd March 2019

DOI: 10.1039/c9ra00277d

rsc.li/rsc-advances

Introduction

Hypericum patulum (Guttiferae) is well known as “Jinsimej” in China, and is distributed mainly in southwest China, such as Guizhou, Sichuan and Yunnan Provinces.¹ The herbs of *H. patulum* are used as a traditional medicine to clear heat, cool blood, relax tendons and activate collaterals, and to treat gonorrhoea, hepatitis, colds, etc.^{2–6} Modern pharmacological investigations demonstrated that the plants of the genus *Hypericum* possessed anti-depression,^{7–11} anti-tumor,^{12–18} anti-bacterial,^{19,20} anti-viral,^{17,18,21–23} and liver protective activities.²⁴ Previous phytochemical studies on these plants showed that derivatives of polycyclic polyprenylated acylphloroglucinols (PPAPs), which possessed a highly oxygenated bicyclo[3.3.1]nonane-2,4,9-trione or other related core decorated with C₅H₉ or C₁₀H₁₇ (prenyl or geranyl) side chains, were the main bioactive components.^{7,13–19,21,24,25}

In this paper, we report the isolation and structural elucidation of six new PPAPs (1–6) (Fig. 1), together with ten known

ones (7–16). Their structures were elucidated using spectroscopic data, X-ray crystallography, ECD spectroscopy and Rh₂(OCOFCF₃)₄-induced ECD. Moreover, compounds 1–16 were evaluated for their cytotoxic activities on human HepG-2, HeLa, MCF-7, and A549 cell lines using the MTT assay. Among them, compound 5 shows significant cytotoxicities toward HepG-2, HeLa and A549 cell lines (IC₅₀ = 9.52 ± 0.27, 11.87 ± 0.22 and 12.63 ± 0.12 μM).

Results and discussion

The 95% EtOH extract of *Hypericum patulum* was subjected to liquid–liquid fractionation to afford a petroleum ether (PE)-soluble fraction and an ethyl acetate (EtOAc)-soluble fraction. The PE fraction was separated by silica gel column chromatography, Sephadex LH-20 and preparative HPLC to obtain six new compounds (1–6) and ten known ones (7–16).

Compound 1 was isolated from CH₃OH as colorless crystals with [α]_D²⁵ +39.6 (*c* 1.0, MeOH). Its molecular formula was deduced as C₃₈H₅₀O₆ on the basis of ¹³C NMR and HRESIMS (*m/z* 625.3515 [M + Na]⁺, calcd for C₃₈H₅₀NaO₆ 625.3500) data. IR spectroscopy suggested the presence of hydroxyl (3456 cm⁻¹), carbonyl (1716 cm⁻¹) and aromatic double bond (1624, 1450 cm⁻¹) groups. The NMR data of 1 (Table S1 and S2, ESI†) indicated the presence of an enolized 1,3-dicarbonyl ether group (δ_C 193.8, C-9; 116.3, C-8; 172.9, C-7), an unconjugated carbonyl carbon (δ_C 205.0, C-1), a methylene (δ_C 38.8, C-5), a methine (δ_C 43.2, C-4), and three quaternary carbons at δ_C 79.7 (C-2), 60.2 (C-6), and 49.7 (C-3), which suggested that 1 was a polycyclic polyprenylated acylphloroglucinol.^{7,26,27} Besides the above carbons, signals for eight methyls, six methylenes, nine methines and six quaternary carbons were observed. The NMR spectroscopic data of 1 resembled those of 32-*epi*-hyperforatin

^aInstitute of Traditional Chinese Medicine & Natural Products, Guangdong Province Key Laboratory of Pharmacodynamic Constituents of TCM and New Drugs Research, College of Pharmacy, Jinan University, Guangzhou 510632, People's Republic of China. E-mail: twanguocai@jnu.edu.cn; tliyl@jnu.edu.cn

^bIntegrated Chinese and Western Medicine Postdoctoral Research Station, Jinan University, Guangzhou 510632, People's Republic of China

^cSchool of Pharmaceutical Sciences, Guangzhou University of Chinese Medicine, Guangzhou 510006, China

† Electronic supplementary information (ESI) available: ECD spectra of the [Rh₂(OCOFCF₃)₄] complexes of compounds 1–4 with the intrinsic ECD spectrum subtracted, calculated and experimental ECD spectra of 1–6, detailed HRESIMS, UV, IR, 1D, 2D NMR data of compounds 1–6. CCDC 1865373. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c9ra00277d






Fig. 1 Chemical structures of 1–6.

E.²⁷ The main differences were that the absence of the 2-methylpropanoyl group [δ_{H} 2.00 (CH), 1.04 (CH₃), 0.96 (CH₃); δ_{C} 211.5 (C=O), 43.0 (CH), 21.8 (CH₃), 20.8 (CH₃)], and the presence of a benzoyl group [δ_{H} 7.41 (2CH), 7.36 (CH), 7.20 (2CH); δ_{C} 194.2 (C=O), 137.1 (C), 132.3 (CH), 128.3 (2CH), 128.1 (2CH)] in **1** (Fig. 1), which implied that the 2-methylpropanoyl group in 32-*epi*-hyperforatin **E** was replaced by a benzoyl group in **1**. This was confirmed by the ¹H–¹H COSY cross-peaks between H-13/15 (δ_{H} 7.20) and H-12/16 (δ_{H} 7.41)/H-14 (δ_{H} 7.36), as well as the HMBC cross-peaks from H-12/16 to C-10 (δ_{C} 194.2)/C-14 (δ_{C} 132.3) (Fig. 2). The relative stereochemistry of **1** resembled those of 32-*epi*-hyperforatin **E**, basing on the NOESY correlations of Me-17 (δ_{H} 1.17) with H-5b (δ_{H} 1.63)/H-24, H-5b with H-34 and of H-5a (δ_{H} 2.10) with H-35 (δ_{H} 4.61) (Fig. 3). The absolute configuration at C-20 was confirmed by the induced ECD of the *in situ* formed [Rh₂(OCOCF₃)₄] complex.^{28,29} According to the bulkiness rule,^{28–30} the 20*S* configuration of **1** was confirmed by the Cotton effect (positive E band) of the Rh complex (Fig. S1, ESI[†]). Additionally, the absolute configuration of **1** was unequivocally confirmed by X-ray crystallography (Fig. 4, CCDC 1865373) and ECD calculations (Fig. S2, ESI[†]), allowing the assignment of the absolute configuration of **1** as 2*R*, 3*R*, 4*S*, 6*S*,

20*S*, 35*S* (Fig. 1). Based on the above analysis, the structure of **1** was elucidated and named hyperpatulone **A**.

Compound **2** was isolated as a colorless oil with $[\alpha]_{\text{D}}^{25} +41.7$ (*c* 1.0, MeOH). The HRESIMS of compound **2** showed an $[\text{M} + \text{Na}]^+$ ion peak at *m/z* 625.3506 (calcd for C₃₈H₅₀NaO₆, 625.3500), consistent with the molecular formula of C₃₈H₅₀O₆. Compounds **2** and **1** were separated by using chiral HPLC over a CHIRALPAK IC column. And the NMR spectroscopic data of **2** (Table S1 and S2, ESI[†]) was almost identical to those of **1**, which indicated that **2** possessed the same planar structure as that of **1**. However, compound **2** showed a negative E band in the *in situ* [Rh₂(OCOCF₃)₄] complex-induced ECD spectrum (Fig. S1, ESI[†]), which is different from that of **1**, suggesting an 20*R* configuration in compound **2**. Thus, structure **2** was established, and named hyperpatulone **B**.

Compound **3** had the molecular formula C₃₈H₅₂O₇, which was assigned by HRESIMS (*m/z* 643.3640 $[\text{M} + \text{Na}]^+$, calcd for C₃₈H₅₂O₇Na, 643.3605). According to its 1D NMR spectra (Tables S1 and S2, ESI[†]), compound **3** has the same skeleton as that of **1** except for the C-18–C-23 side chain. The differences between them were the absence of a terminal double bond (δ_{C}




Fig. 2 Key ¹H–¹H COSY and HMBC correlations of **1**, **3**, **5** and **6**.




Fig. 3 Key NOESY correlations of **1**, **3**, **5** and **6**.






Fig. 4 X-ray ORTEP drawing of 1.

147.9, 111.2) between C-21 and C-22, but the presence of one additional oxygenated quaternary carbon (δ_C 73.1) and one additional methyl group (δ_C 26.4) in 3, and the chemical shifts of C-18, 19, 20, 23 shifted from δ_C 32.7, 32.0, 76.5, 17.8 in 1 to δ_C 34.2, 28.2, 79.5, 23.8 in 3, which indicated the olefinic carbons (C-21, C-22) in 1 were replaced by a tertiary alcohol hydroxy group and a methyl group in 3. This was confirmed by the HMBC cross-peaks from H-20 (δ_H 3.26)/H-22 (δ_H 1.18)/H-23 (δ_H 1.12) to C-21 (δ_C 73.1) (Fig. 2). NOESY correlations of Me-17 (δ_H 1.18) with H-5b (δ_H 1.63)/H-24, of H-5b with H-34 and of H-5a (δ_H 2.10) with H-35 (δ_H 4.62) indicated that the relative configuration of 3 was identical to that of 1 (Fig. 3). The *in situ* $[\text{Rh}_2(\text{OCOCF}_3)_4]$ complex-induced ECD spectrum of 3 exhibited a positive E band for a 20S configuration (Fig. S3, ESI[†]). Therefore, structure 3 was determined and named hyperpatulone C.

The molecular formula of 4 was established to be $\text{C}_{33}\text{H}_{42}\text{O}_6$ by its HRESIMS m/z 643.3626 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{33}\text{H}_{52}\text{O}_7\text{Na}$, 643.3605). The NMR data (Tables S1 and S2, ESI[†]) of 4 showed lots of similarities to those of 3, suggesting that 4 and 3 possessed the same planar structure. The only difference between 4 and 3 was the orientation of H-20, which was determined by a negative E band for a 20R configuration in the *in situ* $[\text{Rh}_2(\text{OCOCF}_3)_4]$ complex-induced ECD spectrum of 4 (Fig. S3, ESI[†]). Accordingly, compound 4 was elucidated and named hyperpatulone D.

The molecular formula $\text{C}_{33}\text{H}_{42}\text{O}_6$ of compound 5 was assigned by HRESIMS (m/z 557.2893 $[\text{M} + \text{Na}]^+$, calcd for $\text{C}_{33}\text{H}_{42}\text{O}_6\text{Na}$, 557.2874). The 1D NMR data (Tables S1 and S3, ESI[†]) of 5 showed lots of similarities to those of hyperascryone G,¹⁸ with a 6/6/5 tricyclic spiro ring system. The natural occurring polyprenylated spirocyclic acylphloroglucinol derivatives (PSAPs), with a 6/6/5 tricyclic spiro ring system, were a special subgroup of PPAPs. Detailed comparison of the NMR spectra of 5 with those of hyperascryone G indicated the absence of a 3-

methylbutanoyl group [δ_H 3.04 and 2.88 (CH_2), 2.30 (CH), 1.01 (CH_3), 0.98 (CH_3); δ_C 197.6 ($\text{C}=\text{O}$), 45.9 (CH_2), 27.1 (CH), 22.9 (CH_3), 22.6 (CH_3)] in hyperascryone G, but the presence of a benzoyl group [δ_H 7.44 (2CH), 7.43 (CH), 7.37 (2CH); δ_C 191.5 ($\text{C}=\text{O}$), 136.6 (C), 131.3 (CH), 128.1 (2CH), 127.9 (2CH)] in 5. Thus, it could be deduced that the 3-methylbutanoyl group in hyperascryone G was replaced by a benzoyl group in 5. This was confirmed by the $^1\text{H}-^1\text{H}$ COSY cross-peaks between H-30/32 (δ_H 7.37) and H-29/33 (δ_H 7.44)/H-31 (δ_H 7.43), as well as the HMBC cross-peaks from H-29/33 to C-27 (δ_C 191.5)/C-31 (δ_C 131.3) (Fig. 2). The relative configurations of 5 and hyperascryone G were very similar by analysis of the NOESY correlations between H-18 (δ_H 4.55) and H-22a (δ_H 2.66), between H-23 (δ_H 5.13) and H-17a (δ_H 2.15), between H-8 (δ_H 1.83) and Me-15 (δ_H 1.71)/Me-16 (δ_H 1.18)/H-14a (δ_H 4.78) (Fig. 3). The ECD data obtained for 5 showed positive Cotton effects at λ_{max} 201 and 278 nm and a negative Cotton effect at λ_{max} 242 and 311 nm (Fig. S4, ESI[†]) comparable to those of hyperascryone G.¹⁸ Thus, structure 5 was established, and named hyperpatulone E.

Compound 6 was assigned the molecular formula $\text{C}_{25}\text{H}_{36}\text{O}_6$ by HRESIMS (m/z 455.2412 $[\text{M} + \text{Na}]^+$, calcd for $\text{C}_{25}\text{H}_{36}\text{O}_6\text{Na}$, 455.2404). The 1D NMR data (Tables S1 and S3, ESI[†]) of 6 showed lots of similarities to chipericumun D (**14**).³¹ Detailed comparison of the NMR spectra of 6 with those of chipericumun D indicated the absence of a 2-methylbutanoyl group [δ_H 3.16 (CH), 1.78 and 1.44 (CH_2), 1.22 (CH_3), 0.83 (CH_3); δ_C 205.1 ($\text{C}=\text{O}$), 41.9 (CH), 25.3 (CH_2), 19.5 (CH_3), 12.3 (CH_3)] in chipericumun D, but the presence of a propanoyl group [δ_H 2.98 (CH_2), 0.97 (CH_3); δ_C 201.0 ($\text{C}=\text{O}$), 46.2 (CH_2), 23.0 (CH_3)] in 6. Thus, it could be deduced that the 2-methylbutanoyl group in chipericumun D was replaced by a propanoyl group in 6. The structure was supported by the $^1\text{H}-^1\text{H}$ COSY correlations between H-24 (δ_H 2.98) and Me-25 (δ_H 0.97) together with the HMBC correlations between Me-25 and C-23 (δ_C 201.0) (Fig. 2). The relative configuration of 6 was same as that of chipericumun D with the analysis of the NOESY correlations of H-7a (δ_H 1.91)/H-14a (δ_H 1.45), H-12 (δ_H 1.78)/H-14a, H-8 (δ_H 1.67)/Me-15 (δ_H 0.95), H-7b (δ_H 1.76)/Me-16, H-8/Me-16 (δ_H 1.40), Me-15/H-24 (δ_H 2.98) and H-7b/H-18 (δ_H 4.55) (Fig. 3). In addition, compounds 6 and chipericumun D (**14**) gave closely correlated Cotton effects in the ECD spectrum (Fig. S4, ESI[†]). Thus, structure 6 was established, and named hyperpatulone F.

Ten known compounds were identified as uralodin A (**7**),³² uralodin B (**8**),¹³ attenuatumione H (**9**),²⁶ uralone D (**10**),⁷ uralone I (**11**),⁷ tomoeone A (**12**),¹⁵ tomoeone B (**13**),¹⁵ chipericumun D (**14**),³¹ hyperascryone F (**15**),¹⁸ hypercohone G (**16**),³³ by comparison of their spectroscopic and physical data with those of related literature.

The isolates 1–16 were tested for their cytotoxic activities by MTT assay on human HepG-2, HeLa, MCF-7 and A549 cell lines. Cisplatin was used as the positive control. As shown in Table 1, PSAPs compounds (5–6, 12–16) exhibited more potent cytotoxic activities than other PPAPs compounds (1–4, 7–11), with IC_{50} values of 9.52 ± 0.27 to 42.33 ± 1.91 μM . Especially, compound 5 shows significant cytotoxicities toward HepG-2, HeLa and A549 cell lines ($\text{IC}_{50} = 9.52 \pm 0.27$, 11.87 ± 0.22 and 12.63 ± 0.12 μM).



Table 1 Cytotoxic activities of compounds 1–16

Compounds	IC ₅₀ ^a (μM)			
	HepG-2	HeLa	MCF-7	A549
1	>50	>50	>50	>50
2	>50	>50	46.83 ± 1.26	>50
3	>50	>50	>50	>50
4	>50	45.79 ± 1.21	>50	44.35 ± 0.62
5	9.52 ± 0.27	11.87 ± 0.22	20.83 ± 0.52	12.63 ± 0.12
6	26.73 ± 0.23	39.67 ± 0.27	42.33 ± 1.91	36.89 ± 0.81
7	>50	>50	>50	47.82 ± 1.17
8	41.03 ± 0.68	39.27 ± 1.23	35.72 ± 0.93	42.90 ± 1.04
9	>50	>50	>50	>50
10	>50	42.67 ± 0.42	39.31 ± 0.67	41.32 ± 1.32
11	>50	>50	42.97 ± 1.21	>50
12	30.91 ± 0.25	27.46 ± 0.37	35.29 ± 0.82	21.78 ± 0.57
13	35.67 ± 0.49	29.67 ± 0.21	31.44 ± 0.95	32.47 ± 0.31
14	22.83 ± 0.53	25.59 ± 0.32	26.92 ± 0.58	27.41 ± 0.71
15	29.38 ± 0.28	24.39 ± 0.28	27.37 ± 0.53	23.76 ± 0.17
16	19.28 ± 0.37	28.59 ± 0.35	22.91 ± 0.32	17.92 ± 0.23
Cisplatin ^b	5.9 ± 0.45	4.7 ± 0.17	6.7 ± 0.61	5.1 ± 0.21

^a IC₅₀ values of 1–16 were detected by MTT assay after incubation for 48 h; data are expressed as mean ± SD. ^b Positive control.

Experimental

General experimental procedures

Optical rotations were obtained on a JASCO P-1020 polarimeter. UV spectra were recorded using a JASCO V-550 UV/VIS spectrophotometer. CD spectra were measured on a JASCO J-810 spectrometer. 1D and 2D NMR spectra were recorded on Bruker AV-500 NMR spectrometers with TMS as an internal standard. HRESIMS analyses were recorded on an Agilent 6210 ESI/TOF mass spectrometer. Column chromatography (CC) was performed with Silica gel (Qingdao Marine Chemical Plant, Qingdao, P. R. China), ODS (50 μm, YMC, Kyoto, Japan) and Sephadex LH-20 (Pharmacia Biotech, Uppsala, Sweden). Preparative HPLC was conducted on a Cosmosil C₁₈ preparative column (5 μm, 20 × 250 mm) equipped with a G1311C pump and a G1315D photodiode array detector (Agilent Technologies, CA, USA). All chemical reagents were purchased from Tianjin Damao Chemical Company (Tianjin, P. R. China).

Plant material

The whole plant of *Hypericum patulum* was collected in Guizhou Province of China, in August of 2016 and authenticated by Zhenqiu Mai, the senior engineer of Guangdong Province. A voucher specimen (no. 20160817) was deposited in the Institute of Traditional Chinese Medicine & Natural Products, Jinan University, Guangzhou, China.

Extraction and isolation

The dried and powdered herbs of *Hypericum patulum* (12 kg) were extracted under reflux with 95% EtOH (30 L × 3) at room temperature. The combined ethanol extract was concentrated to afford a residue (654 g), which was suspended in water (4 L)

and then extracted with petroleum ether (PE) (4 L × 3) and ethyl acetate (EtOAc) (4 L × 3). The PE extract (217 g) was subjected to silica gel column chromatography, eluting with PE-EtOAc (100 : 0 to 0 : 1, v/v) to yield seven fractions (Fr. A–F). Fr. C (18.7 g) was further applied to a silica gel CC with PE/EtOAc (10 : 1 to 1 : 1, v/v) to afford five subfractions (Fr. C1–C5). Fr. C2 (1.5 g) was purified by Sephadex LH-20 (CHCl₃/MeOH, 2 : 1, v/v) and further separated by preparative HPLC (MeOH/H₂O, 70 : 30, v/v) to yield compounds 1 (21.2 mg), 2 (18.5 mg), 7 (11.7 mg) and 8 (16.9 mg). Fr. C3 (7.5 g) was purified by ODS CC and Sephadex LH-20 to obtain compounds 3 (12.7 mg), 4 (13.9 mg), 9 (19.7 mg), 10 (15.7 mg) and 11 (13.2 mg). Fr. D (21.9 g) was applied to ODS CC using a MeOH/H₂O gradient (40 : 60 to 100 : 0, v/v) to afford five subfractions (Fr. D.1–D.5). Fr. D.3 (3.6 g) was further purified by Sephadex LH-20 CC (CHCl₃/MeOH, 1 : 1, v/v) and preparative HPLC (MeOH/H₂O, 80 : 20, v/v) and to yield compounds 5 (9.5 mg), 12 (21.3 mg) and 13 (25.1 mg). Fr. D.4 (5.8 g) was separated by preparative HPLC (MeOH/H₂O, 80 : 20, v/v) to yield compounds 6 (15.8 mg) and 14 (11.9 mg). Fr. D.5 (4.9 g) was purified by preparative HPLC (MeOH/H₂O, 80 : 20, v/v) to achieve compounds 15 (9.2 mg) and 16 (15.2 mg).

Hyperpatulone A (1). Colorless needle crystals (MeOH); mp 116–117 °C; [α]_D²⁵ +39.6 (c 1.0, MeOH); UV (MeOH) λ_{max} 203, 250 and 277 nm; IR (KBr) ν_{max} 3456, 2981, 2931, 1716, 1693, 1624, 1450, 1369, 1227 cm⁻¹; ¹H and ¹³C NMR spectroscopic data, see Tables S1 and S2 (ESI[†]); HRESIMS *m/z* 625.3515 [M + Na]⁺ (calcd for C₃₈H₅₀NaO₆: 625.3500).

X-ray crystallographic analysis of 1 (Table S4, ESI[†]). C₃₈H₅₀O₆, *M* = 602.78, orthorhombic, space group *P*2₁2₁2₁; *a* = 19.2963(4) Å, *b* = 16.3762(4) Å, *c* = 11.0039(2) Å, α = 90°, β = 90°, γ = 90°, *V* = 3477.23(13) Å³, *T* = 100.00(10) K, *Z* = 4, *D*_{calcd} = 1.151 g m⁻³, *F*(000) = 1304.0. The final *R* values were *R*₁ = 0.0809, *wR*₂ = 0.2257, and the goodness of fit on *F*² was equal to 1.156. Flack parameter = 0.0(2). The crystal data of compound 1 was deposited with the Cambridge Crystallographic Data Centre (CCDC 1865373, <http://www.ccdc.cam.ac.uk/>).

Hyperpatulone B (2). Colorless oil; [α]_D²⁵ +41.7 (c 1.0, MeOH); UV (MeOH) λ_{max} 204, 248 and 274 nm; IR (KBr) ν_{max} 3448, 2970, 2924, 1724, 1693, 1620, 1446, 1369, 1227 cm⁻¹; ¹H and ¹³C NMR spectroscopic data, see Tables S1 and S2 (ESI[†]); HRESIMS *m/z* 625.3506 [M + Na]⁺ (calcd for C₃₈H₅₀NaO₆: 625.3500).

Hyperpatulone C (3). Colorless oil; [α]_D²⁵ +56.6 (c 1.0, MeOH); UV (MeOH) λ_{max} 204, 247 and 274 nm; IR (KBr) ν_{max} 3425, 2977, 2931, 1705, 1600, 1442, 1389, 1273, 1215, 1119 cm⁻¹; ¹H and ¹³C NMR spectroscopic data, see Tables S1 and S2 (ESI[†]); HRESIMS *m/z* 643.3640 [M + Na]⁺ (calcd for C₃₈H₅₂O₇Na: 643.3605).

Hyperpatulone D (4). Colorless oil; [α]_D²⁵ +51.8 (c 1.0, MeOH); UV (MeOH) λ_{max} 203, 248 and 275 nm; IR (KBr) ν_{max} 3413, 2974, 2927, 1709, 1604, 1446, 1381, 1281, 1219, 1122 cm⁻¹; ¹H and ¹³C NMR spectroscopic data, see Tables S1 and S2 (ESI[†]); HRESIMS *m/z* 643.3626 [M + Na]⁺ (calcd for C₃₈H₅₂O₇Na: 643.3605).

Hyperpatulone E (5). Colorless oil; [α]_D²⁵ -37.5 (c 1.0, MeOH); UV (MeOH) λ_{max} 208, 242 and 353 nm; IR (KBr) ν_{max} 3413, 2965, 2927, 2877, 1716, 1612, 1454, 1376, 1269, 1153 cm⁻¹; ¹H and ¹³C NMR spectroscopic data, see Tables S1 and S3 (ESI[†]); HRESIMS *m/z* 557.2893 [M + Na]⁺ (calcd for C₃₃H₄₂O₆Na: 557.2874).



Hyperpatulone F (6). Colorless oil; $[\alpha]_D^{25} +22.8$ (c 1.0, MeOH); UV (MeOH) λ_{\max} 203, 216, 249 and 279 nm; IR (KBr) ν_{\max} 3410, 2970, 2935, 2877, 1720, 1662, 1612, 1454, 1385, 1319, 1273, 1211, 1153, 1107 cm^{-1} ; ^1H and ^{13}C NMR spectroscopic data, see Tables S1 and S3 (ESI[†]); HRESIMS m/z 455.2412 $[\text{M} + \text{Na}]^+$ (calcd for $\text{C}_{25}\text{H}_{36}\text{O}_6\text{Na}$, 455.2404).

Cell culture

Human HepG-2, HeLa, MCF-7, and A549 cells were obtained from the Human Virology Institute of Sun Yat-Sen University. Cells were maintained in RPMI-1640 medium (Gibco, USA) supplemented with 10% fetal bovine serum (Gibco, USA) and 1% penicillin/streptomycin at 37 °C with 5% CO_2 for 24 h.

Cytotoxic assay *in vitro*

Four selected human cancer cell lines at the logarithmic phase were seeded in 96-well plates at 5×10^3 cells per well, respectively. After incubating for 24 h, cells were treated with various concentrations of compounds 1–16 and incubated at 37 °C for 48 h. Then, the medium of each well was removed and 5 mg mL^{-1} MTT (30 μL) was added. After incubating for 4 h, the supernatant of each well was removed and DMSO (200 μL) was added to dissolve the formazan produced in the cells. The absorbance was recorded using an enzyme immunoassay reader (Thermo Labsystems Multiskan MK3) at 570 nm. The IC_{50} was calculated by the Bliss method: inhibitory rate = [(absorbance of the test group – absorbance of the blank control)/(absorbance of the control group – absorbance of the blank control)] \times 100.

Conclusions

In summary, six new PPAPs derivatives, hyperpatulone A–F (1–6), together with ten known analogs, were obtained from the dried herbs of *Hypericum patulum*. Their structures were determined by spectroscopic data, X-ray crystallography, ECD spectrum and $\text{Rh}_2(\text{OCOCF}_3)_4$ -induced ECD. Moreover, compounds 1–16 were evaluated for their cytotoxic activities on human HepG-2, HeLa, MCF-7, and A549 cell lines using the MTT assay. Compound 5 shows significant cytotoxicity toward HepG-2, HeLa and A549 cell lines ($\text{IC}_{50} = 9.52 \pm 0.27$, 11.87 ± 0.22 and 12.63 ± 0.12 μM).

Conflicts of interest

The authors declare no competing financial interest.

Acknowledgements

This work was supported by grants from the National Natural Science Foundation of China (No. 81803376, 81673319, 81673670), the Science and Technology Planning Project of Guangdong Province (No. 2016A030303011, 2016B030301004), and China Postdoctoral Science Foundation (No. 2017M620405).

Notes and references

- Q. L. Wu, S. P. Wang, L. W. Wang, J. S. Yang and P. G. Xiao, *Nat. Prod. Res. Dev.*, 1997, **10**, 15–18.
- H. F. Lv, Q. G. Chu and H. Z. Hu, *Chin. Tradit. Herb. Drugs*, 2002, **33**, 1135–1138.
- Z. Y. Xiao and Q. Mu, *Nat. Prod. Res. Dev.*, 2007, **19**, 344–355.
- Z. Q. Yin, Y. Wang, D. M. Zhang, W. C. Ye and S. X. Zhao, *Chinese Wild Plant Resources*, 2004, **23**, 6–11.
- Y. H. Cui and J. Li, *J. Northeast Agric. Univ.*, 2006, **37**, 105–110.
- L. S. Zhang, G. P. Dong and G. M. Liu, *J. Chin. Med. Mater.*, 2009, **32**, 224–226.
- Z. B. Zhou, Z. R. Li, X. B. Wang, J. G. Luo and L. Y. Kong, *J. Nat. Prod.*, 2016, **79**, 1231–1240.
- E. Ernst, J. I. Rand, J. Barnes and C. Stevinson, *Eur. J. Clin. Pharmacol.*, 1998, **54**, 589–594.
- A. Singer, M. Wonnemann and W. E. Müller, *J. Pharmacol. Exp. Ther.*, 1999, **290**, 1363–1368.
- V. Butterweck, F. Petereit, H. Winterhoff and A. Nahrstedt, *Planta Med.*, 1998, **64**, 291–294.
- S. S. Chatterjee, S. K. Bhattacharya, M. Wonnemann, A. Singer and W. E. Muller, *Life Sci.*, 1998, **63**, 499–510.
- D. Albert, I. Zundorf, T. Dingermann, W. E. Muller, D. Steinhilber and O. Werz, *Biochem. Pharmacol.*, 2002, **64**, 1767–1775.
- X. Q. Chen, Y. Li, X. Cheng, K. Wang, J. He, Z. H. Pan, M. M. Li, L. Y. Peng, G. Xu and Q. S. Zhao, *Chem. Biodiversity*, 2010, **7**, 196–204.
- L. H. Hu and K. Y. Sim, *Tetrahedron*, 2000, **56**, 1379–1386.
- W. Hashida, N. Tanaka, Y. Kashiwada, M. Sekiya, Y. Ikeshiro and Y. Takaishi, *Phytochemistry*, 2008, **69**, 2225–2230.
- J. J. Zhang, X. W. Yang, J. Z. Ma, Y. Ye, X. L. Shen and G. Xu, *Tetrahedron*, 2015, **71**, 8315–8319.
- S. A. T. Fobofou, K. Franke, G. Sanna, A. Porzel, E. Bullita, P. L. Colla and L. A. Wessjohann, *Bioorg. Med. Chem.*, 2015, **23**, 6327–6334.
- H. C. Zhu, C. M. Chen, J. J. Liu, B. Sun, G. Z. Wei, Y. Li, J. W. Zhang, G. M. Yao, Z. W. Luo, Y. B. Xue and Y. H. Zhang, *Phytochemistry*, 2015, **115**, 222–230.
- T. Naonobu, Y. Yuki, T. Yutaka and K. Yoshiki, *Org. Lett.*, 2016, **18**, 5360–5363.
- H. Jayasariya, A. M. Clark and C. J. D. Mc, *J. Nat. Prod.*, 1991, **54**, 1314–1320.
- S. A. T. Fobofou, C. R. Harmon, A. H. N. Lonfouo, K. Franke, S. M. Wright and L. A. Wessjohann, *Phytochemistry*, 2016, **124**, 108–113.
- A. K. George, P. Dan, T. John and C. Susan, *Biochem. Biophys. Res. Commun.*, 1990, **172**, 149–153.
- J. Zhao, Z. P. Zhang, H. S. Chen and X. H. Chen, *Acta Pharmacol. Sin.*, 1998, **33**, 67–72.
- W. Gao, W. Z. Hou, J. Zhao, F. Xu, L. Li, F. Xu, H. Sun, J. G. Xing, Y. Peng, X. L. Wang, T. F. Ji and Z. Y. Gu, *J. Nat. Prod.*, 2016, **79**, 1538–1547.
- X. W. Yang, Y. Q. Ding, J. J. Zhang, X. Liu, L. X. Yang, X. N. Li, D. Ferreira, L. A. Walker and G. Xu, *Org. Lett.*, 2014, **16**, 2434–2437.



- 26 Z. B. Zhou, Y. M. Zhang, J. G. Luo and L. Y. Kong, *Phytochem. Lett.*, 2016, **15**, 215–219.
- 27 Y. Guo, N. Zhang, C. M. Chen, J. F. Huang, X. N. Li, J. J. Liu, H. C. Zhu, Q. Y. Tong, J. W. Zhang, Z. W. Luo, Y. B. Xue and Y. H. Zhang, *J. Nat. Prod.*, 2017, **80**, 1493–1504.
- 28 J. Frelek and W. J. Szczepek, *Tetrahedron*, 1999, **10**, 1507–1520.
- 29 M. Gerards and G. Snatzke, *Tetrahedron*, 1990, **1**, 221–236.
- 30 L. Liu, H. Gao, X. Chen, X. Cai, L. Yang, L. Guo, X. Yao and Y. Che, *Eur. J. Org. Chem.*, 2010, **17**, 3302–3306.
- 31 S. Abe, N. Tanaka and J. Kobayashi, *J. Nat. Prod.*, 2012, **75**, 484–488.
- 32 N. Guo, X. Q. Chen and Q. S. Zhao, *Acta Bot. Yunnanica*, 2008, **30**, 515–518.
- 33 J. J. Zhang, X. W. Yang, J. Z. Ma, X. Liu, L. X. Yang, S. C. Yang and G. Xu, *Nat. Prod. Bioprospect.*, 2014, **4**, 73–79.

