



Cite this: RSC Adv., 2019, 9, 7218

Received 15th January 2019

Accepted 26th February 2019

DOI: 10.1039/c9ra00350a

[rsc.li/rsc-advances](http://rsc.li/rsc-advances)

## An efficient synthesis of new imidazo[1,2-a]pyridine-6-carbohydrazide and pyrido[1,2-a]pyrimidine-7-carbohydrazide derivatives via a five-component cascade reaction†

Hajar Hosseini and Mohammad Bayat \*

A highly efficient and straightforward synthesis of N-fused heterocyclic compounds including *N'*-(1-(4-nitrophenyl)ethylidene)imidazo[1,2-a]pyridine-6-carbohydrazide and *N'*-(1-(4-nitrophenyl)ethylidene)pyrido[1,2-a]pyrimidine-7-carbohydrazide derivatives is successfully achieved via a five-component cascade reaction utilizing cyanoacetohydrazide, 4-nitroacetophenone, 1,1-bis(methylthio)-2-nitroethylene and various diamines in a mixture of water and ethanol. The new efficient domino protocol involving a sequence of *N,N*-acetal formation, Knoevenagel condensation, Michael reaction, imine-enamine tautomerization and *N*-cyclization as key steps. The merit of this catalyst free approach is highlighted by its easily available starting materials, operational simplicity, clean reaction profile, the use of environmentally benign solvents and tolerance of a wide variety of functional groups.

## Introduction

Imidazopyridines have displayed a broad spectrum of pharmacological and biological activities.<sup>1</sup> Among the diverse derivatives of imidazopyridine, the imidazo[1,2-a]pyridine skeleton is probably the most important structure due to its vital role as a key construction in drugs and biologically active compounds with properties such as anti-inflammatory,<sup>2,3</sup> anti-viral,<sup>4–6</sup> antifungal,<sup>7–9</sup> anticancer,<sup>10</sup> anxiolytic,<sup>11</sup> anti-ulcer,<sup>12</sup> and antiprotozoal.<sup>13</sup> They can be found in marketed drugs such as the clinical anti-ulcer compound zolpidem and alpidem,<sup>14</sup> olprinone,<sup>15</sup> zolimidine,<sup>16</sup> necopidem and saripidem<sup>17</sup> (Fig. 1).

Pyridopyrimidine derivatives are also known as antidepressant,<sup>18</sup> gastrointestinal protective,<sup>19</sup> neurotropic and stress-protecting,<sup>20</sup> and anticancer agents.<sup>21</sup> Pyrido[1,2-a]pyrimidine constitutes the core structure of some marketed drugs, including the antiasthmatic agent pemirolast,<sup>22</sup> the tranquilizer pirenperone,<sup>23</sup> and the antiallergic agent ramastine<sup>24</sup> (Fig. 1).

The design of reactions with minimized number of steps is one of the purpose of modern synthesis. One approach to achieve this goal includes the development of multicomponent methods. Multicomponent reactions (MCRs) offer a wide range of probabilities for the fast generation of functionalized molecules in a single step with high atom economy, minimum time and cost and straight forward experimental procedures.<sup>25</sup> These

benefits are highlights for multicomponent cascade reactions, which involve *in situ* production of an intermediate with a reactive site for subsequent transformations.<sup>26</sup>

By now, various synthetic approaches have been reported to synthesize imidazo[1,2-a]pyridines. The common reactions were the cyclocondensations of 2-aminopyridines with  $\alpha$ -halocarbonyl compounds,<sup>27</sup> 1,3-dicarbonyl compounds,<sup>28</sup>




Fig. 1 Drugs containing the imidazo[1,2-a]pyridine and pyrido[1,2-a]pyrimidine cores.

Department of Chemistry, Faculty of Science, Imam Khomeini International University, Qazvin, Iran. E-mail: bayat\_mo@yahoo.com; m.bayat@sci.ikiu.ac.ir; Tel: +98 28 33780040

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c9ra00350a



nitroolefins or alkynes.<sup>29</sup> Condensation of 2-aminopyridines, aldehydes and isonitriles or alkynes in a one-pot process, was also a convenient method for the synthesis of imidazo[1,2-*a*]pyridines.<sup>30</sup> Furthermore, over the past decade, a number of methods have been described to synthesize pyrido[1,2-*a*]pyrimidines by focusing on traditional two-component condensation of 2-aminopyridines with a variety of bifunctional electrophiles.<sup>31–33</sup>


In recent years some other new synthetic approaches have been developed for the synthesis of tetrahydroimidazo[1,2-*a*]pyridines and tetrahydro-1*H*-pyrido[1,2-*a*]pyrimidine using heterocyclic ketene aminals (HKAs).<sup>34–38</sup> Heterocyclic ketene aminals (HKAs) are efficient synthons for the synthesis of heterocyclic compounds. Reactions of cyclic ketene aminals with a variety of bis-electrophilic compounds have so far been applied to construct five- and six-membered fused heterocyclic structures.<sup>39</sup>

In the process of our efforts to synthesize the new heterocyclic compounds using cyanoacetohydrazide, we report herein an efficient one-pot five-component synthesis of novel imidazo[1,2-*a*]pyridine-6-carbohydrazides and 1*H*-pyrido[1,2-*a*]pyrimidine-7-carbohydrazides *via* *in situ* preparation of nitroketene aminals. To the best of our knowledge, there is no report on the synthesis of these structures.

## Results and discussion

We succeeded in synthesizing two categories of functionalized N-fused heterocyclic structures containing tetrahydroimidazo[1,2-*a*]pyridines-6-carbohydrazides and tetrahydro-1*H*-pyrido[1,2-*a*]pyrimidine-7-carbohydrazides by using of cyanoacetohydrazide **1**, 4-nitroacetophenone **2**, aromatic aldehydes **3**, 1,1-bis(methylthio)-2-nitroethene **4** and various diamines **5**.

Table 1 Optimization conditions for the formation of **6c**<sup>a</sup>



Entry	Solvent	Catalyst (mol%)	Time (h)	Temp (°C)	Yield (%)
1	EtOH	—	24	78	60
2	EtOH	Piperidine	24	78	57
3	EtOH	<i>p</i> -TSA	24	78	No reaction
4	EtOH	AcOH	24	78	No reaction
5	H <sub>2</sub> O	—	24	100	No reaction
6	H <sub>2</sub> O/EtOH (1 : 2, v/v)	—	10	78	65
7	<b>H<sub>2</sub>O/EtOH (1 : 3, v/v)</b>	—	5	<b>78</b>	<b>87</b>
8	CH <sub>3</sub> CN	—	24	82	No reaction

<sup>a</sup> Reagents and conditions: cyanoacetohydrazide (1 mmol), 4-nitroacetophenone (1 mmol), 4-chlorobenzaldehyde (1 mmol), 1,1-bis(methylthio)-2-nitroethene (1 mmol), ethylenediamine (1 mmol), solvent (20 mL), catalyst (0.2 mmol). The bold row represents the best results.

## Optimization of the conditions

Initially, cyanoacetohydrazide **1**, 4-nitroacetophenone **2**, 4-chlorobenzaldehyde **3**, 1,1-bis(methylthio)-2-nitroethene **4** and ethylenediamine **5** were used as model substrates to optimize the reaction conditions.

The experimental results showed when ethanol was used as solvent without any catalyst at reflux conditions, the yield of desired product was 60% (Table 1, entry 1). With piperidine as catalyst, the reaction efficiency did not change significantly (entry 2). With *p*-TSA and acetic acid, the product did not form (entry 3, 4). The use of water or acetonitrile instead of ethanol did not result in the desired product (entry 5, 8), but when the mixture of water and ethanol was used (overall 1 : 2, v/v), the efficiency increased slightly (entry 6). By changing the ratio of solvents (1 : 3, v/v) at reflux conditions the five-component product **6c** was obtained in a yield of 87% within 5 hours (entry 7).


With information obtained from optimization conditions table, we used cyanoacetohydrazide **1**, 4-nitroacetophenone **2**, various aromatic aldehydes **3**, 1,1-bis(methylthio)-2-nitroethene **4** and diamines (ethylenediamine, 1,3-diaminopropane, 2,2-dimethyl-1,3-diaminopropane) **5** as starting materials to synthesize the target compounds **6a–q** (Scheme 1).

The reactions were completed after 4–7 h overall to afford corresponding heterocyclic systems in good to high yields (73–90%). The results are summarized in Table 2.

## Effect of substituents

From the observation of reaction times in Table 2, it was found that with aldehydes containing an electron-withdrawing group on *para* position of ring (nitro and halogens), the reaction rate is the highest and with methoxy group, the rate is the lowest.





Scheme 1 Synthetic scheme for the generation of products **6a-q**.

This reaction was performed with other derivatives of diamines (1,4-diaminobutane and 1,2-diaminocyclohexane) under the same conditions, which did not result in the product. Also the reaction with *ortho* derivatives of benzaldehyde (2-chloro and 2-nitro) did not produce the desired product. Other derivatives of acetophenone (4-chloro, 4-bromo and 4-methoxyacetophenone) were also used which resulted no product formation.

In addition, the reaction with aliphatic ketones instead of 4-nitroacetophenone and aliphatic aldehydes instead of aromatic aldehydes did not lead to the desired products.

It should be noted that these reactions involve two-, three- and four-component impurities. In fact, the most important side product was a four-component structure without participation of 4-nitroacetophenone that was previously synthesized using two equivalents of aldehyde.<sup>34</sup> To achieve the pure product, it was necessary to complete the reaction of cyanoacetohydrazide and 4-nitroacetophenone in a mixture of water and ethanol at reflux conditions in sufficient time (3 hours), then ketene aminal solution and aromatic aldehyde were added at the same time.

### Structure determination

The structures of compounds **6a-q** were deduced from their IR, <sup>1</sup>H NMR, <sup>13</sup>C NMR spectroscopic and Mass spectrometric data (see the ESI†).

The formation of suggested products **6a-q** is clearly verified by the <sup>1</sup>H and <sup>13</sup>C NMR spectra of the crude products. As a representative case the key signals of <sup>1</sup>H and <sup>13</sup>C NMR chemical shifts of 5-amino-7-(3-chlorophenyl)-8-nitro-N'-(1-(4-nitrophenyl)ethylidene)-1,2,3,7-tetrahydroimidazo[1,2-*a*]pyridine-6-carbohydrazide **6a** are presented in Fig. 2.

The <sup>1</sup>H NMR spectrum of **6a** showed two NH groups at  $\delta$  9.16 and 9.43 ppm. The protons of 4-nitrobenzene ring were seen at  $\delta$  7.96 and 8.20 ppm as two doublet signals and the protons of 3-chlorobenzene ring were observed at  $\delta$  6.92–7.31 ppm. The NH<sub>2</sub> group appeared at  $\delta$  8.21 ppm. The proton of CH at pyridine ring was seen at  $\delta$  5.49 ppm. Two protons of two methylene groups appeared at  $\delta$  3.81 and 4.05 ppm. The signal at  $\delta$  2.14 ppm was related to methyl group.

The <sup>1</sup>H-decoupled <sup>13</sup>C NMR spectrum of **6a** indicated 20 distinct resonances in accordance to desired structure. The characteristic signals of four aliphatic carbons (CH<sub>3</sub>, CH and two CH<sub>2</sub> groups) were seen at  $\delta$  13.9, 38.2, 43.6 and 44.7 ppm respectively. Two signals at  $\delta$  80.0 and 107.4 ppm were related to

C=C-CO and C-NO<sub>2</sub> respectively. The carbonyl group appeared at  $\delta$  165.78 ppm (Fig. 2).

The mass spectrum of **6a** displayed the molecular-ion peak at *m/z* 497 in agreement with the proposed structure. The IR spectrum of this compound showed absorption bands at 3486, 3400, 3327 cm<sup>-1</sup> due to NH and NH<sub>2</sub> groups, stretching vibration of aliphatic C-H bands at 2909, and strong absorption of carbonyl group at 1658 and C-N band at 1254 cm<sup>-1</sup>. Two absorption bands due to nitro group appeared at 1519 and 1375 cm<sup>-1</sup>.

### Mechanism

A general reasonable mechanism for the formation of imidazo[1,2-*a*]pyridines-6-carbohydrazides and pyrido[1,2-*a*]pyrimidine-7-carbohydrazides is shown in Scheme 2. On the basis of well-established chemistry of 1,1-bis(methylthio)-2-nitroethene, initially, addition of diamine **5** to 1,1-bis(methylthio)-2-nitroethene **4** leads to the formation of ketene aminal **9**. On the other hand condensation of cyanoacetohydrazide **1** with 4-nitroacetophenone **2** leads to hydrazone **7**. Further, with adding aldehyde **3**, the Knoevenagel condensation affords intermediate **8**. Then, Michael addition of ketene aminal **9** to adduct **8** leads to the intermediate **10**, which undergoes successive imine-enamine tautomerization followed by an intramolecular cyclization via nucleophilic addition of -NH to nitrite group. Finally, imine-enamine tautomerization gives the titled products **6** (Scheme 2).

## Experimental

### Materials

All commercially available reagents and other solvents were purchased from Aldrich and Merck chemical Co. and used without further purification. The NMR spectra were recorded with a Bruker DRX-300 AVANCE instrument (300 MHz for <sup>1</sup>H and 75.4 MHz for <sup>13</sup>C) with DMSO-d<sub>6</sub> as solvent. Chemical shifts are given in ppm ( $\delta$ ), and coupling constant ( $J$ ) are reported in Hertz (Hz). Melting points were measured with an electro-thermal 9100 apparatus. Mass spectra were recorded with an Agilent 5975C VL MSD with Triple-Axis detector operating at an ionization potential of 70 eV. IR spectra were measured with Bruker Tensor 27 spectrometer. Elemental analyses for C, H and N were performed using a PerkinElmer 2004 series [II] CHN elemental analyzer.

### General procedure of the synthesis of tetrahydroimidazo[1,2-*a*]pyridine-6-carbohydrazide and tetrahydro-1*H*-pyrido[1,2-*a*]pyrimidine-7-carbohydrazide derivatives

A mixture of diamine (1 mmol), 1,1-bis(methylthio)-2-nitroethylene (0.165 g, 1 mmol) and 10 mL EtOH in a 50 mL flask was refluxed for 6 hours. In another 50 mL flask the stoichiometric mixture of cyanoacetohydrazide (1 mmol, 0.099 g) and 4-nitroacetophenone (1 mmol, 0.165 g) in H<sub>2</sub>O/EtOH (5 : 5 mL) was stirred at reflux conditions for 3 hours. After this time, TLC shows the consumption of the starting components. Then, aromatic aldehyde (1 mmol) and the first solution (HKA), was



Table 2 Compounds 6a–q<sup>a</sup>

Entry	Aromatic aldehyde	Diamine	Product	Time (h)	Yield (%)	Mp (°C)
1				6	90	231–233
2				5	86	235–237
3				5	87	239–241
4				4	89	243–245
5				5	90	239–240
6				7	78	267–269



Table 2 (Contd.)

Entry	Aromatic aldehyde	Diamine	Product	Time (h)	Yield (%)	Mp (°C)
7				5	80	271–273
8				6	75	284–286
9				5	86	264–266
10				4	88	268–270
11				5	85	265–267
12				4	90	298–300



Table 2 (Contd.)

Entry	Aromatic aldehyde	Diamine	Product	Time (h)	Yield (%)	Mp (°C)
13				7	73	269–271
14				7	75	315–317
15				5	85	317–319
16				5	84	322–324
17				6	75	329–331

<sup>a</sup> The reaction was performed using cyanoacetohydrazide (1 mmol), 4-nitroacetophenone (1 mmol), aromatic aldehyde (1 mmol), 1,1-bis(methylthio)-2-nitroethene (1 mmol), diamine (1 mmol), EtOH (15 mL), H<sub>2</sub>O (5 mL), reflux.

added to this mixture simultaneously. The progress of the reaction was monitored by TLC using ethyl acetate/n-hexane (1 : 1). After completion of the reaction, the precipitated product was collected by filtration and washed with warm ethanol to give the pure products **6a–q** in 73–90% yield.

**5-Amino-7-(3-chlorophenyl)-8-nitro-N'-(1-(4-nitrophenyl)ethylidene)-1,2,3,7-tetrahydroimidazo[1,2-a]pyridine-6-carbohydrazide (6a).** Dark yellow solid; yield: 0.447 g (90%); mp: 231–233 °C; IR (KBr) ( $\nu_{\text{max}}/\text{cm}^{-1}$ ): 3486, 3400, 3327, 2909, 1658, 1519, 1458, 1375, 1254, 785; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>):

Fig. 2  $^1\text{H}$  and  $^{13}\text{C}$  NMR chemical shifts of 6a.

Scheme 2 Proposed mechanism for the formation of products 6.

$\delta$  2.14 (3H, s,  $\text{CH}_3$ ), 3.78–3.84 (2H, m,  $\text{CH}_2$ ), 3.99–4.11 (2H, m,  $\text{CH}_2$ ), 5.49 (1H, s,  $\text{CH}$ ), 6.92–7.31 (4H, m, ArH), 7.96 (2H, d,  $J$  = 8.7 Hz, ArH), 8.20 (2H, d,  $J$  = 8.7 Hz, ArH), 8.21 (2H, br s,  $\text{NH}_2$ ), 9.16 (1H, s, NH), 9.43 (1H, s, NH);  $^{13}\text{C}\{^1\text{H}\}$  NMR (75.4 MHz, DMSO- $d_6$ ):  $\delta$  13.9 ( $\text{CH}_3$ ), 38.2 (CH), 43.6 ( $\text{CH}_2\text{-NH}$ ), 44.7 ( $\text{CH}_2\text{-N}$ ), 80.0 ( $\text{C}=\text{C-CO}$ ), 107.4 ( $\text{C}=\text{C-NO}_2$ ), 123.9, 126.8, 127.1, 127.3, 128.3, 130.3, 132.8, 145.0, 147.3 (Ar), 147.5 (Me-C=N), 148.0 ( $\text{C}=\text{C-NH}$ ), 151.3 ( $\text{C}_{\text{Ar}}\text{-NO}_2$ ), 151.4 ( $\text{C}=\text{C-NH}_2$ ), 165.7 ( $\text{C}=\text{O}$ ); MS (EI, 70 eV):  $m/z$  (%) = 497 (0.008) [ $\text{M}^+$ ], 454 (2), 326 (59), 311 (100), 292 (29), 265 (37), 246 (51), 204 (26), 179 (63), 117 (83), 76 (47), 57 (7); anal. calcd for  $\text{C}_{22}\text{H}_{20}\text{ClN}_7\text{O}_5$ : C, 53.07; H, 4.05; N, 19.69. Found: C, 53.37; H, 3.84; N, 19.65.

**5-Amino-7-(3-fluorophenyl)-8-nitro-N'-(1-(4-nitrophenyl)ethylidene)-1,2,3,7-tetrahydroimidazo[1,2-a]pyridine-6-carbohydrazide (6b).** Orange solid; yield: 0.413 g (86%); mp: 235–237 °C; IR (KBr) ( $\nu_{\text{max}}$ /cm $^{-1}$ ): 3477, 3332, 3401, 1660, 1519, 1462, 1377, 1331, 1254;  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ):  $\delta$  2.15 (3H, s,  $\text{CH}_3$ ), 3.37–3.84 (2H, m,  $\text{CH}_2$ ), 4.02–4.12 (2H, m,  $\text{CH}_2$ ), 5.47 (1H, s, CH), 7.17–7.39 (4H, m, ArH), 7.96 (2H, d,  $J$  = 8.7 Hz, ArH), 8.20 (2H, d,  $J$  = 8.7 Hz, ArH), 8.24 (2H, br s,  $\text{NH}_2$ ), 9.17 (1H, s, NH), 9.45 (1H, s, NH);  $^{13}\text{C}\{^1\text{H}\}$  NMR (75.4 MHz, DMSO- $d_6$ ):  $\delta$  13.9 ( $\text{CH}_3$ ), 38.1 (CH), 43.6 ( $\text{CH}_2\text{-NH}$ ), 44.7 ( $\text{CH}_2\text{-N}$ ), 80.1

( $\text{C}=\text{C-CO}$ ), 107.5 ( $\text{C}=\text{C-NO}_2$ ), 113.7, 115.2, 123.9, 124.4, 127.3, 130.2, 145.0, 147.5 (Ar), 148.5 (Me-C=N), 148.5 ( $\text{C}=\text{C-NH}$ ), 151.4 ( $\text{C}_{\text{Ar}}\text{-NO}_2$ ), 151.5 ( $\text{C}=\text{C-NH}_2$ ), 160.7 ( $\text{C}_{\text{Ar}}\text{-F}$ ), 165.8 ( $\text{C}=\text{O}$ ); MS (EI, 70 eV):  $m/z$  (%) = 481 (0.008) [ $\text{M}^+$ ], 438 (6), 392 (2), 326 (57), 311 (100), 276 (62), 230 (87), 179 (54), 117 (71), 76 (39), 50 (5); anal. calcd for  $\text{C}_{22}\text{H}_{20}\text{FN}_7\text{O}_5$ : C, 54.88; H, 4.19; N, 20.37. Found: C, 54.56; H, 4.04; N, 20.53.

### 5-Amino-7-(4-chlorophenyl)-8-nitro-N'-(1-(4-nitrophenyl)ethylidene)-1,2,3,7-tetrahydroimidazo[1,2-a]pyridine-6-carbohydrazide (6c).

Yellow solid; yield: 0.432 g (87%); mp: 239–241 °C; IR (KBr) ( $\nu_{\text{max}}$ /cm $^{-1}$ ): 3456, 3401, 3339, 2919, 1675, 1522, 1456, 1373, 1255, 761;  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ):  $\delta$  2.15 (3H, s,  $\text{CH}_3$ ), 3.77–3.86 (2H, m,  $\text{CH}_2$ ), 4.02–4.10 (2H, m,  $\text{CH}_2$ ), 5.45 (1H, s, CH), 7.28 (2H, d,  $J$  = 8.4 Hz, ArH), 7.36 (2H, d,  $J$  = 8.4 Hz, ArH), 7.96 (2H, d,  $J$  = 8.7 Hz, ArH), 8.20 (2H, d,  $J$  = 8.7 Hz, ArH), 8.21 (2H, s,  $\text{NH}_2$ ), 9.16 (1H, s, NH), 9.43 (1H, s, NH);  $^{13}\text{C}\{^1\text{H}\}$  NMR (75.4 MHz, DMSO- $d_6$ ):  $\delta$  13.9 ( $\text{CH}_3$ ), 37.8 (CH), 43.6 ( $\text{CH}_2\text{-NH}$ ), 44.7 ( $\text{CH}_2\text{-N}$ ), 80.2 ( $\text{C}=\text{C-CO}$ ), 107.7 ( $\text{C}=\text{C-NO}_2$ ), 123.9, 127.3, 128.5, 129.8, 130.2, 131.3, 145.0 (Ar), 147.5 (Me-C=N), 151.3 ( $\text{C}=\text{C-NH}$ ), 151.4 ( $\text{C}_{\text{Ar}}\text{-NO}_2$ ), 151.8 ( $\text{C}=\text{C-NH}_2$ ), 165.7 ( $\text{C}=\text{O}$ ); anal. calcd for  $\text{C}_{22}\text{H}_{20}\text{ClN}_7\text{O}_5$ : C, 53.07; H, 4.05; N, 19.69. Found: C, 53.28; H, 4.16; N, 19.51.

### 5-Amino-7-(4-nitrophenyl)-8-nitro-N'-(1-(4-nitrophenyl)ethylidene)-1,2,3,7-tetrahydroimidazo[1,2-a]pyridine-6-carbohydrazide (6d).

Light Yellow solid; yield: 0.452 g (89%); mp: 243–245 °C;  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ):  $\delta$  2.18 (3H, s,  $\text{CH}_3$ ), 3.79–3.85 (2H, m,  $\text{CH}_2$ ), 4.04–4.07 (2H, m,  $\text{CH}_2$ ), 5.67 (1H, s, CH), 7.59 (2H, d,  $J$  = 9 Hz, ArH), 7.95 (2H, d,  $J$  = 9 Hz, ArH), 8.10 (2H, d,  $J$  = 8.7 Hz, ArH), 8.20 (2H, d,  $J$  = 8.7 Hz, ArH), 8.27 (2H, s,  $\text{NH}_2$ ), 9.34 (1H, s, NH), 9.50 (1H, s, NH);  $^{13}\text{C}\{^1\text{H}\}$  NMR (75.4 MHz, DMSO- $d_6$ ):  $\delta$  14.1 ( $\text{CH}_3$ ), 38.3 (CH), 43.7 ( $\text{CH}_2\text{-NH}$ ), 44.7 ( $\text{CH}_2\text{-N}$ ), 79.7 ( $\text{C}=\text{C-CO}$ ), 107.1 ( $\text{C}=\text{C-NO}_2$ ), 123.6, 123.9, 124.4, 127.3, 129.6, 145.0, 146.4 (Ar), 147.5 (Me-C=N), 147.6 ( $\text{C}=\text{C-NH}$ ), 151.4 ( $\text{C}_{\text{Ar}}\text{-NO}_2$ ), 151.5 ( $\text{C}_{\text{Ar}}\text{-NO}_2$ ), 153.4 ( $\text{C}=\text{C-NH}_2$ ), 166.0 ( $\text{C}=\text{O}$ ); MS (EI, 70 eV):  $m/z$  (%) = 508 (0.01) [ $\text{M}^+$ ], 326 (40), 311 (72), 257 (32), 204 (14), 179 (100), 117 (59), 76 (7), 57 (12); anal. calcd for  $\text{C}_{22}\text{H}_{20}\text{N}_8\text{O}_7$ : C, 51.97; H, 3.96; N, 22.04. Found: C, 52.13; H, 3.89; N, 22.14.

### 5-Amino-7-(4-bromophenyl)-8-nitro-N'-(1-(4-nitrophenyl)ethylidene)-1,2,3,7-tetrahydroimidazo[1,2-a]pyridine-6-carbohydrazide (6e).

Yellow solid; yield: 0.486 g (90%); mp: 239–240 °C; IR (KBr) ( $\nu_{\text{max}}$ /cm $^{-1}$ ): 3454, 3338, 2919, 1658, 1522, 1462, 1377, 1254, 580;  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ):  $\delta$  2.14 (3H, s,  $\text{CH}_3$ ), 3.79–3.85 (2H, m,  $\text{CH}_2$ ), 4.00–4.07 (2H, m,  $\text{CH}_2$ ), 5.42 (1H, s, CH), 7.30 (2H, m, ArH), 7.39 (2H, m, ArH), 7.94 (2H, d,  $J$  = 7.5 Hz, ArH), 8.19 (2H, d,  $J$  = 7.5 Hz, ArH), 8.20 (2H, s,  $\text{NH}_2$ ), 9.16 (1H, s, NH), 9.43 (1H, s, NH);  $^{13}\text{C}\{^1\text{H}\}$  NMR (75.4 MHz, DMSO- $d_6$ ):  $\delta$  13.9 ( $\text{CH}_3$ ), 37.9 (CH), 43.6 ( $\text{CH}_2\text{-NH}$ ), 44.7 ( $\text{CH}_2\text{-N}$ ), 80.1 ( $\text{C}=\text{C-CO}$ ), 107.6 ( $\text{C}=\text{C-NO}_2$ ), 119.8, 123.8, 127.3, 130.2, 130.6, 131.1, 144.9 (Ar), 147.1 (Me-C=N), 147.4 ( $\text{C}=\text{C-NH}$ ), 151.4 ( $\text{C}_{\text{Ar}}\text{-NO}_2$ ), 151.4 ( $\text{C}=\text{C-NH}_2$ ), 165.7 ( $\text{C}=\text{O}$ ); anal. calcd for  $\text{C}_{22}\text{H}_{20}\text{BrN}_7\text{O}_5$ : C, 48.72; H, 3.72; N, 18.08. Found: C, 48.64; H, 3.78; N, 18.19.

### 5-Amino-7-(4-methoxyphenyl)-8-nitro-N'-(1-(4-nitrophenyl)ethylidene)-1,2,3,7-tetrahydroimidazo[1,2-a]pyridine-6-carbohydrazide (6f).

Dark yellow solid; yield: 0.342 g (78%); mp: 267–269 °C; IR (KBr) ( $\nu_{\text{max}}$ /cm $^{-1}$ ): 3401, 3343, 2919, 2850, 1658,

1514, 1451, 1342, 1257, 1167;  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ):  $\delta$  2.11 (3H, s, CH<sub>3</sub>), 3.69 (3H, s, OCH<sub>3</sub>), 3.74–3.83 (2H, m, CH<sub>2</sub>), 4.02–4.05 (2H, m, CH<sub>2</sub>), 5.28 (1H, s, CH), 6.79 (2H, d,  $J$  = 8.4 Hz, ArH), 7.27 (2H, d,  $J$  = 8.4 Hz, ArH), 7.94 (2H, d,  $J$  = 8.7 Hz, ArH), 8.19 (2H, d,  $J$  = 8.7 Hz, ArH), 8.20 (2H, br s, NH<sub>2</sub>), 8.97 (1H, s, NH), 9.38 (1H, s, NH);  $^{13}\text{C}\{\text{H}\}$  NMR (75.4 MHz, DMSO- $d_6$ ):  $\delta$  13.7 (CH<sub>3</sub>), 37.7 (CH), 43.5 (CH<sub>2</sub>-NH), 44.7 (CH<sub>2</sub>-N), 55.4 (OCH<sub>3</sub>), 80.6 (C=C-CO), 108.2 (C=C-NO<sub>2</sub>), 113.8, 123.9, 127.3, 129.4, 137.4, 145.0 (Ar), 146.4 (Me-C=N), 147.4 (C=C-NH), 151.2 (C<sub>Ar</sub>-NO<sub>2</sub>), 151.4 (C=C-NH<sub>2</sub>), 158.3 (C<sub>Ar</sub>-OCH<sub>3</sub>), 165.5 (C=O); anal. calcd for C<sub>23</sub>H<sub>23</sub>N<sub>7</sub>O<sub>6</sub>: C, 55.98; H, 4.70; N, 19.87. Found: C, 55.93; H, 4.77; N, 19.75.

**5-Amino-7-(4-fluorophenyl)-8-nitro-N'-(1-(4-nitrophenyl)ethylidene)-2,3,7-tetrahydroimidazo[1,2-a]pyridine-6-carbohydrazide (6g).** Light orange solid; yield: 0.384 g (80%); mp: 271–273 °C;  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ):  $\delta$  2.12 (3H, s, CH<sub>3</sub>), 3.80–3.83 (2H, m, CH<sub>2</sub>), 4.03–4.05 (2H, m, CH<sub>2</sub>), 5.42 (1H, s, CH), 6.96–7.23 (4H, m, ArH), 7.95 (2H, d,  $J$  = 9 Hz, ArH), 8.19 (2H, d,  $J$  = 9 Hz, ArH), 8.24 (2H, br s, NH<sub>2</sub>), 9.10 (1H, s, NH), 9.42 (1H, s, NH);  $^{13}\text{C}\{\text{H}\}$  NMR (75.4 MHz, DMSO- $d_6$ ):  $\delta$  13.8 (CH<sub>3</sub>), 37.6 (CH), 43.5 (CH<sub>2</sub>-NH), 44.7 (CH<sub>2</sub>-N), 80.4 (C=C-CO), 107.9 (C=C-NO<sub>2</sub>), 114.8, 116.0, 123.9, 127.3, 127.9, 128.8, 130.1 (Ar), 141.7 (Me-C=N), 142.4 (C=C-NH), 147.4 (C<sub>Ar</sub>-NO<sub>2</sub>), 151.4 (C=C-NH<sub>2</sub>), 165.6 (C=O); anal. calcd for C<sub>22</sub>H<sub>20</sub>FN<sub>7</sub>O<sub>5</sub>: C, 54.88; H, 4.19; N, 20.37. Found: C, 54.55; H, 3.94; N, 20.15.

**5-Amino-8-nitro-N'-(1-(4-nitrophenyl)ethylidene)-7-phenyl-1,2,3,7-tetrahydroimidazo[1,2-a]pyridine-6-carbohydrazide (6h).** Yellow solid; yield: 0.347 g (75%); mp: 284–286 °C;  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ):  $\delta$  2.12 (3H, s, CH<sub>3</sub>), 3.81–3.84 (2H, m, CH<sub>2</sub>), 4.05–4.08 (2H, m, CH<sub>2</sub>), 5.36 (1H, s, CH), 7.12–7.36 (5H, m, ArH), 7.94 (2H, d,  $J$  = 7.8 Hz, ArH), 8.19 (2H, d,  $J$  = 7.8 Hz, ArH), 8.19 (2H, br s, NH<sub>2</sub>), 9.05 (1H, s, NH), 9.40 (1H, s, NH);  $^{13}\text{C}\{\text{H}\}$  NMR (75.4 MHz, DMSO- $d_6$ ):  $\delta$  13.8 (CH<sub>3</sub>), 38.4 (CH), 43.6 (CH<sub>2</sub>-NH), 44.7 (CH<sub>2</sub>-N), 80.6 (C=C-CO), 107.4 (C=C-NO<sub>2</sub>), 123.9, 126.9, 127.3, 128.4, 145.5 (Ar), 146.6 (Me-C=N), 147.5 (C=C-NH), 151.3 (C<sub>Ar</sub>-NO<sub>2</sub>), 151.5 (C=C-NH<sub>2</sub>), 165.7 (C=O); anal. calcd for C<sub>22</sub>H<sub>21</sub>N<sub>7</sub>O<sub>5</sub>: C, 57.02; H, 4.57; N, 21.16. Found: C, 57.37; H, 4.25; N, 21.45.

**6-Amino-8-(4-bromophenyl)-9-nitro-N'-(1-(4-nitrophenyl)ethylidene)-2,3,4,8-tetrahydro-1H-pyrido[1,2-a]pyrimidine-7-carbohydrazide (6i).** Light yellow solid; yield: 0.477 g (86%); mp: 264–266 °C; IR (KBr) ( $\nu_{\text{max}}/\text{cm}^{-1}$ ): 3481, 3403, 3968, 2925, 1645, 1516, 1456, 1340, 1257, 590;  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ):  $\delta$  1.80–1.92 (2H, m, CH<sub>2</sub>), 2.48 (3H, s, CH<sub>3</sub>), 3.58–3.75 (2H, m, CH<sub>2</sub>NH), 3.76–3.92 (2H, m, CH<sub>2</sub>N), 5.53 (1H, s, CH), 7.26 (2H, d,  $J$  = 8.4 Hz, ArH), 7.41 (2H, d,  $J$  = 8.4 Hz, ArH), 7.97 (2H, d,  $J$  = 8.7 Hz, ArH), 8.20 (2H, d,  $J$  = 8.7 Hz, ArH), 8.25 (2H, s, NH<sub>2</sub>), 9.36 (1H, s, NH), 11.49 (1H, s, NH);  $^{13}\text{C}\{\text{H}\}$  NMR (75.4 MHz, DMSO- $d_6$ ):  $\delta$  14.0 (CH<sub>3</sub>), 20.2 (CH<sub>2</sub>), 36.6 (CH), 38.7 (CH<sub>2</sub>-NH), 43.1 (CH<sub>2</sub>-N), 81.4 (C=C-CO), 109.8 (C=C-NO<sub>2</sub>), 119.9, 123.9, 127.4, 130.4, 131.4, 144.1, 144.9 (Ar), 147.6 (Me-C=N), 147.9 (C=C-NH), 150.3 (C<sub>Ar</sub>-NO<sub>2</sub>), 152.6 (C=C-NH<sub>2</sub>), 166.1 (C=O); MS (EI, 70 eV):  $m/z$  (%) = 343 (58), 311 (53), 265 (20), 243 (8), 204 (13), 183 (40), 156 (27), 133 (10), 117 (100), 97 (25), 76 (8), 57 (55); anal. calcd for C<sub>23</sub>H<sub>22</sub>BrN<sub>7</sub>O<sub>5</sub>: C, 49.65; H, 3.99; N, 17.62. Found: C, 49.87; H, 3.50; N, 17.26.

**6-Amino-9-nitro-8-(4-nitrophenyl)-N'-(1-(4-nitrophenyl)ethylidene)-2,3,4,8-tetrahydro-1H-pyrido[1,2-a]pyrimidine-7-carbohydrazide (6j).** Yellowish orange solid; yield: 0.459 g (88%); mp: 268–271 °C; IR (KBr) ( $\nu_{\text{max}}/\text{cm}^{-1}$ ): 3470, 3389, 2929, 1648, 1521, 1458, 1344, 1259;  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ):  $\delta$  1.85–1.94 (2H, m, CH<sub>2</sub>), 2.48 (3H, s, CH<sub>3</sub>), 3.58–3.75 (2H, m, CH<sub>2</sub>NH), 3.76–3.95 (2H, m, CH<sub>2</sub>N), 5.75 (1H, s, CH), 7.55 (2H, d,  $J$  = 8.4 Hz, ArH), 7.97 (2H, d,  $J$  = 8.4 Hz, ArH), 8.09 (2H, d,  $J$  = 8.7 Hz, ArH), 8.19 (2H, d,  $J$  = 8.7 Hz, ArH), 8.31 (2H, s, NH<sub>2</sub>), 9.55 (1H, s, NH), 11.47 (1H, s, NH);  $^{13}\text{C}\{\text{H}\}$  NMR (75.4 MHz, DMSO- $d_6$ ):  $\delta$  14.2 (CH<sub>3</sub>), 20.1 (CH<sub>2</sub>), 37.15 (CH), 38.8 (CH<sub>2</sub>-NH), 43.2 (CH<sub>2</sub>-N), 81.0 (C=C-CO), 109.3 (C=C-NO<sub>2</sub>), 123.8, 127.4, 128.4, 129.1, 144.9, 146.4 (Ar), 147.6 (Me-C=N), 148.4 (C=C-NH), 150.4 (C<sub>Ar</sub>-NO<sub>2</sub>), 152.5 (C<sub>Ar</sub>-NO<sub>2</sub>), 152.8 (C=C-NH<sub>2</sub>), 166.3 (C=O); MS (EI, 70 eV):  $m/z$  (%) = 353 (6), 311 (35), 271 (67), 225 (19), 179 (100), 164 (67), 132 (49), 118 (77), 91 (35), 76 (13), 56 (31); anal. calcd for C<sub>23</sub>H<sub>22</sub>N<sub>8</sub>O<sub>7</sub>: C, 52.87; H, 4.24; N, 21.45. Found: C, 52.57; H, 4.62; N, 21.18.

**6-Amino-8-(4-chlorophenyl)-9-nitro-N'-(1-(4-nitrophenyl)ethylidene)-2,3,4,8-tetrahydro-1H-pyrido[1,2-a]pyrimidine-7-carbohydrazide (6k).** Light brown solid; yield: 0.434 g (85%); mp: 265–267 °C;  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ):  $\delta$  1.80–1.90 (2H, m, CH<sub>2</sub>), 2.17 (3H, s, CH<sub>3</sub>), 3.38–3.92 (4H, m, 2CH<sub>2</sub>N), 5.54 (1H, s, CH), 7.27 (2H, d,  $J$  = 8.7 Hz, ArH), 7.32 (2H, d,  $J$  = 8.7 Hz, ArH), 7.97 (2H, d,  $J$  = 8.7 Hz, ArH), 8.20 (2H, d,  $J$  = 8.7 Hz, ArH), 8.25 (2H, s, NH<sub>2</sub>), 9.37 (1H, s, NH), 11.49 (1H, s, NH);  $^{13}\text{C}\{\text{H}\}$  NMR (75.4 MHz, DMSO- $d_6$ ):  $\delta$  14.0 (CH<sub>3</sub>), 20.2 (CH<sub>2</sub>), 36.5 (CH), 38.7 (CH<sub>2</sub>-NH), 43.1 (CH<sub>2</sub>-N), 81.5 (C=C-CO), 109.8 (C=C-NO<sub>2</sub>), 123.9, 127.4, 128.5, 129.8, 131.4, 143.7, 144.9 (Ar), 147.6 (Me-C=N), 147.9 (C=C-NH), 150.3 (C<sub>Ar</sub>-NO<sub>2</sub>), 152.6 (C=C-NH<sub>2</sub>), 166.1 (C=O); anal. calcd for C<sub>23</sub>H<sub>22</sub>ClN<sub>7</sub>O<sub>5</sub>: C, 53.96; H, 4.33; N, 19.37.

**6-Amino-3,3-dimethyl-9-nitro-8-(4-nitrophenyl)-N'-(1-(4-nitrophenyl)ethylidene)-2,3,4,8-tetrahydro-1H-pyrido[1,2-a]pyrimidine-7-carbohydrazide (6l).** Creamy yellow solid; yield: 0.495 g (90%); mp: 297–300 °C; IR (KBr) ( $\nu_{\text{max}}/\text{cm}^{-1}$ ): 3320, 3199, 2959, 1642, 1518, 1350, 1261;  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ):  $\delta$  0.95 (3H, s, CH<sub>3</sub>), 1.09 (3H, s, CH<sub>3</sub>), 2.19 (3H, s, CH<sub>3</sub>), 3.23–3.32 (2H, m, CH<sub>2</sub>NH), 3.52–3.56 (2H, m, CH<sub>2</sub>N), 5.80 (1H, s, CH), 7.52 (2H, d,  $J$  = 7.5 Hz, ArH), 7.96 (2H, d,  $J$  = 7.5 Hz, ArH), 8.11 (2H, d,  $J$  = 7.8 Hz, ArH), 8.19 (2H, d,  $J$  = 7.8 Hz, ArH), 8.33 (2H, s, NH<sub>2</sub>), 9.54 (1H, s, NH), 11.38 (1H, s, NH);  $^{13}\text{C}\{\text{H}\}$  NMR (75.4 MHz, DMSO- $d_6$ ):  $\delta$  14.2 (CH<sub>3</sub>), 22.7 (CH<sub>3</sub>), 24.5 (CH<sub>3</sub>), 27.7 (CMe<sub>2</sub>), 37.0 (CH), 49.4 (CH<sub>2</sub>-NH), 53.4 (CH<sub>2</sub>-N), 81.0 (C=C-CO), 109.2 (C=C-NO<sub>2</sub>), 123.9, 127.5, 128.9, 144.9, 146.5 (Ar), 147.6 (Me-C=N), 148.5 (C=C-NH), 149.6 (C<sub>Ar</sub>-NO<sub>2</sub>), 152.6 (C<sub>Ar</sub>-NO<sub>2</sub>), 152.9 (C=C-NH<sub>2</sub>), 166.3 (C=O); MS (EI, 70 eV):  $m/z$  (%) = 409 (6), 345 (13), 327 (38), 299 (84), 253 (20), 179 (100), 149 (52), 118 (71), 96 (14), 77 (78), 55 (42); anal. calcd for C<sub>25</sub>H<sub>26</sub>N<sub>8</sub>O<sub>7</sub>: C, 54.54; H, 4.76; N, 20.35. Found: C, 54.68; H, 4.59; N, 20.24.

**6-Amino-8-(3-methoxyphenyl)-3,3-dimethyl-9-nitro-N'-(1-(4-nitrophenyl)ethylidene)-2,3,4,8-tetrahydro-1H-pyrido[1,2-a]pyrimidine-7-carbohydrazide (6m).** Dark yellow solid; yield: 0.390 g (73%); mp: 269–271 °C;  $^1\text{H}$  NMR (300 MHz, DMSO- $d_6$ ):  $\delta$  0.96 (3H, s, CH<sub>3</sub>), 1.09 (3H, s, CH<sub>3</sub>), 2.14 (3H, s, CH<sub>3</sub>), 3.23 (2H, s, CH<sub>2</sub>NH), 3.56 (2H, s, CH<sub>2</sub>N), 3.65 (3H, s, OCH<sub>3</sub>), 5.49



(1H, s, CH), 6.71–6.89 (3H, m, ArH), 7.15 (1H, t,  $J$  = 7.5 Hz, ArH), 7.97 (2H, d,  $J$  = 9 Hz, ArH), 8.20 (2H, d,  $J$  = 9 Hz, ArH), 8.25 (2H, s, NH<sub>2</sub>), 9.23 (1H, s, NH), 11.44 (1H, s, NH); <sup>13</sup>C{<sup>1</sup>H} NMR (75.4 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  13.8 (CH<sub>3</sub>), 22.6 (CH<sub>3</sub>), 24.6 (CH<sub>3</sub>), 27.6 (CMe<sub>2</sub>), 36.2 (CH), 49.4 (CH<sub>2</sub>-NH), 53.3 (CH<sub>2</sub>-N), 55.4 (OCH<sub>3</sub>), 82.0 (C=C-CO), 110.2 (C=C-NO<sub>2</sub>), 114.0, 115.3, 123.9, 127.4, 128.0, 133.2, 136.5, 144.9 (Ar), 147.3 (Me-C=N), 147.5 (C=C-NH), 149.4 (C<sub>Ar</sub>-NO<sub>2</sub>), 152.6 (C=C-NH<sub>2</sub>), 158.4 (C<sub>Ar</sub>-OCH<sub>3</sub>), 165.8 (C=O); anal. calcd for C<sub>25</sub>H<sub>29</sub>N<sub>7</sub>O<sub>6</sub>: C, 58.31; H, 5.46; N, 18.31. Found: C, 58.50; H, 5.34; N, 18.27.

**6-Amino-8-(4-methoxyphenyl)-3,3-dimethyl-9-nitro-N'-(1-(4-nitrophenyl)ethylidene)-2,3,4,8-tetrahydro-1H-pyrido[1,2-*a*]pyrimidine-7-carbohydrazide (6n).** Yellow solid; yield: 0.401 g (75%); mp: 315–317 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  0.96 (3H, s, CH<sub>3</sub>), 1.09 (3H, s, CH<sub>3</sub>), 2.13 (3H, s, CH<sub>3</sub>), 3.26–3.33 (2H, m, CH<sub>2</sub>NH), 3.55–3.65 (2H, s, CH<sub>2</sub>N), 3.75 (3H, s, OCH<sub>3</sub>), 5.41 (1H, s, CH), 6.79 (2H, d,  $J$  = 8.7 Hz, ArH), 7.23 (2H, d,  $J$  = 8.7 Hz, ArH), 7.96 (2H, d,  $J$  = 9 Hz, ArH), 8.20 (2H, d,  $J$  = 9 Hz, ArH), 8.24 (2H, s, NH<sub>2</sub>), 9.62 (1H, s, NH), 11.45 (1H, s, NH); <sup>13</sup>C{<sup>1</sup>H} NMR (75.4 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  13.8 (CH<sub>3</sub>), 22.6 (CH<sub>3</sub>), 24.1 (CH<sub>3</sub>), 27.7 (CMe<sub>2</sub>), 36.3 (CH), 49.2 (CH<sub>2</sub>-NH), 53.3 (CH<sub>2</sub>-N), 55.4 (OCH<sub>3</sub>), 82.0 (C=C-CO), 110.2 (C=C-NO<sub>2</sub>), 114.0, 115.3, 123.9, 127.4, 128.9, 133.2, 136.5, 144.9 (Ar), 147.3 (Me-C=N), 147.5 (C=C-NH), 149.4 (C<sub>Ar</sub>-NO<sub>2</sub>), 152.6 (C=C-NH<sub>2</sub>), 158.4 (C<sub>Ar</sub>-OCH<sub>3</sub>), 165.8 (C=O); anal. calcd for C<sub>25</sub>H<sub>29</sub>N<sub>7</sub>O<sub>6</sub>: C, 58.31; H, 5.46; N, 18.31. Found: C, 58.68; H, 5.15; N, 18.23.

**6-Amino-8-(4-chlorophenyl)-3,3-dimethyl-9-nitro-N'-(1-(4-nitrophenyl)ethylidene)-2,3,4,8-tetrahydro-1H-pyrido[1,2-*a*]pyrimidine-7-carbohydrazide (6o).** Brownish yellow solid; yield: 0.458 g (85%); mp: 317–319 °C; IR (KBr) ( $\nu_{\text{max}}$ /cm<sup>-1</sup>): 3324, 3204, 2925, 2861, 1643, 1524, 1480, 1353, 1263, 755; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  0.93 (3H, s, CH<sub>3</sub>), 1.08 (3H, s, CH<sub>3</sub>), 2.16 (3H, s, CH<sub>3</sub>), 3.23 (2H, s, CH<sub>2</sub>NH), 3.54 (2H, m, CH<sub>2</sub>N), 5.58 (1H, s, CH), 7.29 (4H, s, ArH), 7.96 (2H, d,  $J$  = 8.7 Hz, ArH), 8.20 (2H, d,  $J$  = 8.7 Hz, ArH), 8.27 (2H, s, NH<sub>2</sub>), 9.36 (1H, s, NH), 11.40 (1H, s, NH); <sup>13</sup>C{<sup>1</sup>H} NMR (75.4 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  14.0 (CH<sub>3</sub>), 22.6 (CH<sub>3</sub>), 24.5 (CH<sub>3</sub>), 27.7 (CMe<sub>2</sub>), 37.0 (CH), 50.4 (CH<sub>2</sub>-NH), 51.5 (CH<sub>2</sub>-N), 81.5 (C=C-CO), 109.7 (C=C-NO<sub>2</sub>), 123.9, 124.0, 127.5, 128.0, 128.5, 128.8, 129.1, 129.6 (Ar), 143.7 (Me-C=N), 144.6 (C=C-NH), 148.0 (C<sub>Ar</sub>-NO<sub>2</sub>), 152.9 (C=C-NH<sub>2</sub>), 166.0 (C=O); MS (EI, 70 eV):  $m/z$  (%) = 539 (0.007) [M]<sup>+</sup>, 368 (4), 311 (30), 288 (100), 223 (13), 179 (57), 149 (27), 117 (37), 102 (20), 77 (42), 55 (27); anal. calcd for C<sub>25</sub>H<sub>26</sub>ClN<sub>7</sub>O<sub>5</sub>: C, 55.61; H, 4.85; N, 18.16. Found: C, 55.90; H, 4.68; N, 18.04.

**6-Amino-8-(4-bromophenyl)-3,3-dimethyl-9-nitro-N'-(1-(4-nitrophenyl)ethylidene)-2,3,4,8-tetrahydro-1H-pyrido[1,2-*a*]pyrimidine-7-carbohydrazide (6p).** Yellow solid; yield: 0.491 g (84%); mp: 322–324 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  1.03 (3H, s, CH<sub>3</sub>), 1.08 (3H, s, CH<sub>3</sub>), 2.16 (3H, s, CH<sub>3</sub>), 3.22–3.35 (2H, m, CH<sub>2</sub>NH), 3.42–3.55 (2H, m, CH<sub>2</sub>N), 5.56 (1H, s, CH), 7.24 (2H, d,  $J$  = 8.4 Hz, ArH), 7.43 (2H, d,  $J$  = 8.4 Hz, ArH), 7.96 (2H, d,  $J$  = 9 Hz, ArH), 8.20 (2H, d,  $J$  = 9 Hz, ArH), 8.29 (2H, s, NH<sub>2</sub>), 9.38 (1H, s, NH), 11.41 (1H, s, NH); <sup>13</sup>C{<sup>1</sup>H} NMR (75.4 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  14.0 (CH<sub>3</sub>), 22.6 (CH<sub>3</sub>), 24.5 (CH<sub>3</sub>), 27.6 (CMe<sub>2</sub>), 36.4 (CH), 49.4 (CH<sub>2</sub>-NH), 53.4 (CH<sub>2</sub>-N), 81.4 (C=C-CO), 109.7 (C=C-NO<sub>2</sub>), 120.0, 123.9, 127.5, 128.0, 130.0, 131.4, 132.0 (Ar), 144.1 (Me-C=N), 147.6 (C=C-NH), 149.5 (C<sub>Ar</sub>-NO<sub>2</sub>), 152.8 (C=C-

NH<sub>2</sub>), 166.1 (C=O); anal. calcd for C<sub>25</sub>H<sub>26</sub>BrN<sub>7</sub>O<sub>5</sub>: C, 51.38; H, 4.48; N, 16.78. Found: C, 51.18; H, 4.66; N, 16.47.

**6-Amino-3,3-dimethyl-9-nitro-N'-(1-(4-nitrophenyl)ethylidene)-8-phenyl-2,3,4,8-tetrahydro-1H-pyrido[1,2-*a*]pyrimidine-7-carbohydrazide (6q).** Light yellow solid; yield: 0.378 g (75%); mp: 329–331 °C; <sup>1</sup>H NMR (300 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  0.95 (3H, s, CH<sub>3</sub>), 1.09 (3H, s, CH<sub>3</sub>), 2.14 (3H, s, CH<sub>3</sub>), 3.19–3.26 (2H, m, CH<sub>2</sub>NH), 3.56 (2H, s, CH<sub>2</sub>N), 5.50 (1H, s, CH), 7.11–7.32 (5H, m, ArH), 7.96 (2H, d,  $J$  = 8.7 Hz, ArH), 8.20 (2H, d,  $J$  = 8.7 Hz, ArH), 8.26 (2H, s, NH<sub>2</sub>), 9.26 (1H, s, NH), 11.44 (1H, s, NH); <sup>13</sup>C{<sup>1</sup>H} NMR (75.4 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  13.9 (CH<sub>3</sub>), 22.6 (CH<sub>3</sub>), 24.6 (CH<sub>3</sub>), 27.7 (CMe<sub>2</sub>), 37.0 (CH), 49.4 (CH<sub>2</sub>-NH), 53.3 (CH<sub>2</sub>-N), 81.9 (C=C-CO), 110.0 (C=C-NO<sub>2</sub>), 120.0, 123.9, 127.5, 128.0, 130.0, 131.4, 132.0 (Ar), 144.1 (Me-C=N), 147.5 (C=C-NH), 150.5 (C<sub>Ar</sub>-NO<sub>2</sub>), 152.8 (C=C-NH<sub>2</sub>), 166.0 (C=O); MS (EI, 70 eV):  $m/z$  (%) = 505 (0.03) [M]<sup>+</sup>, 447 (0.09), 334 (14), 300 (25), 283 (19), 254 (100), 179 (20), 156 (89), 128 (41), 103 (26), 77 (42), 55 (11); anal. calcd for C<sub>25</sub>H<sub>27</sub>N<sub>7</sub>O<sub>5</sub>: C, 59.40; H, 5.38; N, 19.40. Found: C, 59.56; H, 5.25; N, 19.34.

## Conclusion

A green and convenient approach to easy construction of novel and highly substituted imidazo[1,2-*a*]pyridines and pyrido[1,2-*a*]pyrimidines has been developed by annulation of heterocyclic ketene aminals (HKAs) and a three-component product of cyanoacetohydrazide, 4-nitroacetophenone and aromatic aldehydes. The reactions are performed in a mixture of EtOH and H<sub>2</sub>O at reflux conditions in the absence of any catalyst. The present reaction shows considerable properties such as high regioselectivity, cascade one-pot methodology, short reaction times, easy purification of products, high yields, avoiding the use of catalyst and high atom economy.

## Conflicts of interest

The authors declare no competing financial interest.

## Acknowledgements

Financial support of this research from Imam Khomeini International University, Iran is gratefully acknowledged.

## Notes and references

- 1 A. Bagdi and S. Santra, *Chem. Commun.*, 2015, **51**, 1555.
- 2 A. Chiacchio, M. Rimoli and L. Avvione, *Arch. Pharm. Pharm. Med. Chem.*, 1998, **331**, 273.
- 3 S. Kotovskaya and Z. Baskakova, *Pharm. Chem. J.*, 2005, **39**, 574.
- 4 L. Almirante, L. Polo, A. Mugnaini, E. Provinciali, P. Rugarli, A. Biancotti, A. Gamba and W. Murmann, *J. Med. Chem.*, 1965, **8**, 305.
- 5 G. Puerstinger and J. Paeshuyse, *Bioorg. Med. Chem. Lett.*, 2007, **17**, 390.
- 6 M. Lhassani and O. Chavignon, *Eur. J. Med. Chem.*, 1999, **34**, 271.



- 7 J. Kaminsky, C. Puchalski and D. Solomon, *J. Med. Chem.*, 1989, **32**, 1686.
- 8 J. Kaminsky and A. Doweyko, *J. Med. Chem.*, 1997, **40**, 427.
- 9 Y. Rival, G. Grassy and A. Taudou, *Eur. J. Med. Chem.*, 1991, **26**, 13.
- 10 H. Huang and X. Ji, *Org. Lett.*, 2013, **15**, 24.
- 11 J. J. Kaminski and A. M. Doweyko, *J. Med. Chem.*, 1997, **40**, 427.
- 12 M. Bollini, J. J. Casal, D. E. Alvarez, L. Boiani, M. Gonzalez, H. Cerecetto and A. M. Bruno, *Bioorg. Med. Chem.*, 2009, **17**, 1437.
- 13 R. J. Boerner and H. Moller, *J. Psychopharmacother*, 1997, **4**, 145.
- 14 S. Kataev and B. Syropyatov, *Pharm. Chem. J.*, 2004, **38**, 181.
- 15 S. Z. Langer, S. Arbillia, J. Benavides and B. Scatton, *Adv. Biochem. Psychopharmacol.*, 1990, **46**, 61.
- 16 K. Mizushige, T. Ueda, K. Yukiiri and H. Suzuki, *Cardiovasc. Drug Rev.*, 2002, **20**, 163.
- 17 L. Almirante, L. Polo, A. Mugnaini, E. Provinciali, P. Rugarli, A. Biancotti, A. Gamba and W. Murmann, *J. Med. Chem.*, 1965, **817**, 305.
- 18 L. E. J. Kennis, F. P. Bischoff, C. J. Mertens, C. J. Love, F. A. F. Van den Keybus, M. Braeken and J. E. Leysen, *Bioorg. Med. Chem. Lett.*, 2000, **10**, 71.
- 19 J. Knoll, K. Gyires and I. Hermecz, *Drugs Exp. Clin. Res.*, 1987, **13**, 253.
- 20 S. Rapolu, M. Alla, R. J. Ganji, V. Saddabapu, C. Kishor, V. R. Bommena and A. Addlagatta, *Med. Chem. Commun.*, 2013, **4**, 817.
- 21 Y. Yanagihara, H. Kasai, T. Kawashima and T. Shida, *Jpn. J. Pharmacol.*, 1988, **48**, 91.
- 22 R. L. Smith, R. J. Barrett and E. J. Sanders-Bush, *J. Pharmacol. Exp. Ther.*, 1995, **275**, 1050.
- 23 F. Awouters, J. Vermeire, F. Smeyers, P. Vermote, R. Van Beek and C. J. E. Niemegeers, *Drug Dev. Res.*, 1986, **8**, 95.
- 24 M. M. Kozlovskaya, A. N. Inozemtsev, S. V. Nikitin, A. G. Gochmuradov, R. A. Yakushev and R. Chabak-Gorbach, *Bull. Exp. Biol. Med.*, 1995, **119**, 291.
- 25 J. D. Sunderhaus and S. F. Martin, *Chem.-Eur. J.*, 2008, **15**, 1300.
- 26 (a) K. C. Nicolaou, D. J. Edmonds and P. G. Bulger, *Angew. Chem., Int. Ed.*, 2006, **45**, 7134; (b) F. Shi, W. Tan, R. Y. Zhu, G. J. Xing and S. J. Tu, *Adv. Synth. Catal.*, 2013, 355, 1605; (c) K. M. Wang, S. J. Yan and J. Lin, *Eur. J. Org. Chem.*, 2014, **6**, 1129–1145; (d) W. Dai, X. L. Jiang, Q. Wu, F. Shi and S. J. Tu, *J. Org. Chem.*, 2015, **80**, 5737; (e) X. X. Sun, C. Li, Y. Y. He, Z. Q. Zhu, G. J. Mei and F. Shi, *Adv. Synth. Catal.*, 2017, **359**, 2660; (f) L. Wang, L. X. Shi, L. Liu, Z. X. Li, T. Xu, W. J. Hao, G. Li, S. J. Tu and B. Jiang, *J. Org. Chem.*, 2017, **82**, 3605.
- 27 S. Ulloora, A. V. Adhikari and R. Shabaraya, *Chin. Chem. Lett.*, 2013, **24**, 853.
- 28 X. Wang, L. Ma and W. Yu, *Synthesis*, 2011, **15**, 2445.
- 29 R. L. Yan, H. Han and C. Ma, *J. Org. Chem.*, 2012, **77**, 2024.
- 30 S. K. Guchhait and A. L. Chandgude, *J. Org. Chem.*, 2012, **77**, 4438.
- 31 L. Peng, X. Gao, L. Duan, X. Ren, D. Wu and K. Ding, *J. Med. Chem.*, 2011, **54**, 7729.
- 32 S. A. Shiba, A. K. Ei-Ziaty, N. K. El-Aasar and H. A. Al-Saman, *J. Chem. Res.*, 2008, **9**, 500.
- 33 R. L. Sharma, D. Kour, J. Singh, S. Kumar, P. Gupta, S. Gupta, B. Kour and A. Sachar, *J. Heterocycl. Chem.*, 2008, **45**, 1775.
- 34 M. Bayat and F. S. Hosseini, *Tetrahedron Lett.*, 2017, **58**, 1616.
- 35 H. Hosseini and M. Bayat, *RSC Adv.*, 2018, **8**, 41218.
- 36 C. Y. Yu, P. H. Yang, M. X. Zhao and Z. T. Huang, *Synlett*, 2006, 1835.
- 37 W. Y. Xu, Y. M. Jia, J. K. Yang and Z. T. Huang, *Synlett*, 2010, 1682.
- 38 H. Wang, Y. Wang, D. Liang, L. Liu, J. Zhang and Q. Zhu, *Angew. Chem., Int. Ed.*, 2011, **50**, 5678.
- 39 (a) K. M. Wang, S. J. Yan and J. Lin, *Eur. J. Org. Chem.*, 2014, 1129; (b) P. H. Yang, *Res. Chem. Intermed.*, 2016, **42**, 5617; (c) L. Zhang, J. Dong, X. Xu and Q. Liu, *Chem. Rev.*, 2016, **116**, 287; (d) F. Sun, F. Zhu, Z. Shao and Z. Li, *Synlett*, 2015, **26**, 2306.

