# **RSC Advances**

## PAPER

Check for updates

Cite this: RSC Adv., 2019, 9, 16035

Cytotoxic metabolites from the endophytic fungus *Chaetomium globosum* 7951<sup>†</sup>

Fang Wang,<sup>‡ab</sup> Wuli Zhao,<sup>‡b</sup> Conghui Zhang,<sup>b</sup> Shanshan Chang,<sup>b</sup> Rongguang Shao,<sup>b</sup> Jianguo Xing,<sup>c</sup> Minghua Chen,<sup>b</sup> \*<sup>bc</sup> Yixuan Zhang<sup>b</sup> \*<sup>a</sup> and Shuyi Si\*<sup>b</sup>

The following compounds were isolated from acetate extracts of *Chaetomium globosum* 7951 solid cultures: demethylchaetocochin C (1) and chaetoperazine A (3), two new epipolythiodioxopiperazine (ETP) alkaloids, a novel pyridine benzamide, 4-formyl-*N*-(3'-hydroxypyridin-2'-yl) benzamide (6), and three known ETP derivatives (2, 4, and 5). The structures of these compounds were determined using extensive spectroscopic data analysis. Compounds 1–3, and 6, inhibited the growth of MCF-7, MDA-MB-231, H460 and HCT-8 cells with an IC<sub>50</sub> of 4.5 to 65.0  $\mu$ M.

Received 9th April 2019 Accepted 12th May 2019 DOI: 10.1039/c9ra02647a

rsc.li/rsc-advances

### Introduction

Novel bioactive secondary metabolites have been identified in endophytic fungi.<sup>1-3</sup> The *Chaetomium* genus, belonging to the Chaetomiaceae family, contains more than 100 species derived from terrestrial and marine habitats.<sup>4</sup> *Chaetomium globosum* is a species of the *Chaetomium* genus. The isolation of the cytotoxic chaetoglobosins A and B was reported by Sekita and coworkers in 1973.<sup>5</sup> Since then more than 200 metabolites including chaetoglobosins, diketopiperazines, tetramic acids, bis(3-indolyl)-benzoquinones, azaphilones, pyranones, xanthones, anthraquinones, orsellides, steroids, and terpenoids were identified in *C. globosum* cultures. Some of these metabolites exhibit cytotoxic, antibacterial, antimalarial, and antiviral activities.<sup>6</sup>

Epipolythiodioxopiperazine (ETP) alkaloids, with either polysulphide bridges or thiomethyl groups, represent an important family of bioactive secondary metabolites, which are toxic to cancer cell lines.<sup>7</sup> About 20 ETPs have been identified in the *Chaetomium* genus.<sup>6</sup> During our search for novel and bioactive compounds from microorganisms,<sup>8-10</sup> we identified the endophytic fungus, *Chaetomium globosum* 7951, which has cytotoxic activity towards human breast cancer cell lines. *Chaetomium globosum* 7951 comes from the root of *Panax notoginseng*, a traditional Chinese medicine. Chemical investigations of the solid fermentation of the *Chaetomium globosum* 7951 strain led to the identification of 2 new ETP alkaloids, 3 known analogs, and a new pyridine benzamide. Herein, the isolation, structural determination, and cytotoxicity of these compounds are described.

### Results and discussion

The molecular formula of compound 1, a white amorphous powder, is C32H34N6O6S4 according to the (+)-HRESIMS data, with 19 degrees of unsaturation. The IR spectrum displayed the hydroxy or amino (3359 cm<sup>-1</sup>), methyl (2921 cm<sup>-1</sup>), and carbonyl (1680 cm<sup>-1</sup>) functionalities. The <sup>1</sup>H NMR spectrum (Table 1) showed two *ortho*-disubstituted benzene rings at  $\delta_{\rm H}$  6.72 (1H, d), 6.77 (1H, t), 7.03 (1H, t), 7.08 (1H, t), 7.17 (1H, t), 7.22 (1H, d), 7.50 (1H, d) and 7.63 (1H, d), and a trisubstituted double bond at  $\delta_{\rm H}$ 7.07 (1H, s) in the lower field. In addition, four methylene groups  $(\delta_{\rm H} 3.61, 3.04, 4.04, 3.21, 4.21, 4.32, 3.73, and 3.43)$ , one methine group ( $\delta_{\rm H}$  6.07), and four isolated methyl groups ( $\delta_{\rm H}$  2.12, 2.28, 2.77, and 3.11) were observed in the higher field. According to the <sup>13</sup>C NMR and DEPT spectra analyses, in addition to the structural features above, there were also four carbonyls at  $\delta_{\rm C}$  161.1, 164.3, 165.3, and 165.6, and five quaternary carbons at  $\delta_{\rm C}$  65.0, 72.9, 73.2, 73.7, and 77.2. The spectral data, combined with the molecular formula, suggested that compound 1 is an analog of epipolythiodioxopiperazine. Extensive analysis of the NMR data indicates similarities in chemical shifts to chaetocochin C,11 including the absence of one methyl group in 1. HMBC correlations of NH-5' with C-1', C-3', C-4', C-5', and C-6', and of N-CH<sub>3</sub>-2' with C-1', and C-3' indicated that N-CH<sub>3</sub>-5' in chaetocochin C was replaced by NH in 1. The ROESY correlations of H-5 with H-9', and of 3'-S-Me with 6'-S-Me, combined with the CD effects at  $\lambda_{max}$  nm  $(\Delta \varepsilon)$  240 (+7.2), 274 (-0.89) and 305 (+3.6),<sup>11</sup> and based on the similar biogenetic perspective of chaetocochin C, which revealed the absolute configuration of 1 was shown in Fig. 1.





View Article Online

View Journal | View Issue

<sup>&</sup>lt;sup>a</sup>School of Life Science and Biopharmaceutics, Shenyang Pharmaceutical University, Shenyang 110016, China. E-mail: zhangyxzsh@163.com

<sup>&</sup>lt;sup>b</sup>Institute of Medicinal Biotechnology, Chinese Academy of Medical Sciences, Peking Union Medical College, Beijing 100050, China. E-mail: mingsunlight@sina.com; sisyimb@hotmail.com

<sup>&</sup>lt;sup>c</sup>Key Laboratory for Uighur Medicine, Institute of Materia Medica of Xinjiang Uygur Autonomous Region, Urumqi 830004, China

<sup>†</sup> Electronic supplementary information (ESI) available: 1D, 2D NMR, HRESIMS and IR spectra for compounds **1**, **3**, and **6**. See DOI: 10.1039/c9ra02647a

<sup>‡</sup> These authors have contributed equally to this work.

No.	1			3	3	
	$\delta_{ m H}$	$\delta_{\rm C}$ , type	No.	$\delta_{ m H}$	$\delta_{\rm C}$ , type	
1		165.3, C	1		162.7, C	
2-N-Me	3.11, s	27.7, CH <sub>3</sub>	2-N-OMe	3.65, s	61.0, CH	
3	,	77.2, C	3	4.36, t (2.4)	63.5, CH	
4		161.1, C	4		165.1, C	
5	6.07, d (1.8)	80.0, CH	NH-5	8.58, s	,	
6a		149.4, C	6		66.5, C	
7	6.72, d (7.8)	110.1, CH	6-S-Me	2.13, s	12.5, C	
8	7.17, t (7.8)	130.5, CH	7	3.58, d (14.4)	33.7, CH	
9	6.77. t (7.8)	118.8. CH		3.20. d (14.4)	, , ,	
10	7.50. d (7.8)	125.9. CH	8		107.2. C	
10a		126.9. C	9	7.19. d (2.4)	125.0. CH	
10b		73.2. C	NH-10	10.90. 8	,	
11	4.04. d (15.6)	42.1. CH <sub>2</sub>	10a	,	135.6. C	
	3.21. d (15.6)	,2	11	7.28. d (7.8)	111.1. CH	
12	0.21, a (10.0)	73.7. C	12	7.02, t (7.8)	120.7. CH	
13	4.32. dd (12.6. 4.8)	58.7. CH <sub>2</sub>	13	6.93. t (7.8)	118.3. CH	
10	4.21, dd (12.6, 6)	0007, 0112	14	7.58. d (7.8)	118.9. CF	
OH-13	$5.93 \pm (6)$		149		127.9 C	
1'	3.30, 2 (0)	165.6 C	15	$3.58 \text{ ov}^b$	58.5 CH	
2'-N-Me	2.77 8	28.4 CH	0H-15	493 t (54)	50.5, 011	
3'	2.77, 5	72 9 C	011 10	1.50; ( (0.1)		
3'-S-Me	2.12. 8	12.3, CH				
4'	2.12, 0	164 3 C				
- NH-5/	9.06 \$	104.0, 0				
6'	5.00, 5	65 0 C				
6'-S-Me	2.28 s	13.9 CH-				
7'	2.20, 3	33 9 CH				
/	3.01, d(15.4)	$55.5, 011_2$				
8′	3.04, u (13.4)	107 4 C				
0 0′	7.07 s	107.4, C				
9 10/2	7.07, 5	120.8, CH				
10 a 11 <sup>7</sup>	7.22 d (7.8)	133.3, C				
11	7.22, u(7.8)	110.0, CH				
12/ 12/	7.00, 1(7.0)	121.4, UH 110.1 CH				
13 1 <i>4</i> /	7.03, t (7.0)	119.1, UI				
14 14/a	7.03, u (7.8)	120.0, CH				
14 d	2.72 dd (10.9 c)	$130.3, \mathbf{U}$				
10	3./3, uu (10.8, 6)	$62.8, CH_2$				
01145	3.43, dd (10.8, 4.8)					
OH-15'	4.90, t (6)					

<sup>*a*</sup> NMR data ( $\delta$ ) were measured at 600 MHz for <sup>1</sup>H and at 150 MHz for <sup>13</sup>C in DMSO-*d*<sub>6</sub>. The assignments were based on <sup>1</sup>H–<sup>1</sup>H COSY, HSQC, and HMBC experiments. <sup>*b*</sup> *J*-value was not determined due to overlapped signals.

The molecular formula of compound **3** is  $C_{16}H_{19}N_3O_4S$  according to the HRESIMS data. The IR spectrum displayed absorptions bands at 3393, 3194, 2921, and 1675 cm<sup>-1</sup>, suggesting the presence of amino or hydroxyl, methyl, and carbonyl groups. The <sup>1</sup>H-NMR spectrum (Table 1) showed 3-substituted indole moiety signals at  $\delta_H$  6.94 (1H, t), 7.02 (1H, t), 7.19 (1H, d), 7.29 (1H, d) and 7.58 (1H, d), two methylene groups at  $\delta_H$  3.58 (1H, d), 3.20 (1H, d), and 3.56 (2H, m), a methine group at  $\delta_H$  4.36 (1H, t), one isolated methyl at  $\delta_H$  2.13 (3H, s), and a methoxyl group at 3.65 (3H, s). In addition, two carbonyls at  $\delta_C$  162.7 and 165.1, and a quaternary carbon at  $\delta_C$  66.7 were identified *via* the <sup>13</sup>C-NMR spectrum. HMBC correlations of H-7 with C-7, C-8, and C-10a; NH-5 ( $\delta_H$  8.58) with C-1, C-6, and C-7; and S-Me-6 with C-6, revealed an  $\alpha$ -S-methyl-substituted tryptophan residue. In addition, <sup>1</sup>H-<sup>1</sup>H COSY relationships between

H-3/H<sub>2</sub>-15/OH-15, in combination with the HMBC relationships between H-3 and H-15 with C-16, indicate a serine residue. Meanwhile, the association of NH-5 with C-3 and H-3 with C-1 in the HMBC spectrum suggests that the serine and tryptophan residues form a diketopiperazine ring. Finally, the methoxyl group is located at N-3, as indicated by the molecular formula and the chemical shift at  $\delta_{\rm C}$  61.0. Thus, compound **3** was proposed as shown in Fig. 2. DP4+ analysis of the <sup>1</sup>H and <sup>13</sup>C NMR data indicates  $3S^*, 6S^*$ -3 appeared agreement with the experimental NMR data with 100% probability (Tables S3– S7†).<sup>12,13</sup> Based on the common biosynthetic origin, the absolute configuration at C-3 and C-6 is probably to be the same as cyclo-L-Trp-L-Ser.<sup>14</sup> In addition, the calculated optical rotation (OR) value<sup>8</sup> (+56.1) of (3*S*, 6*S*)-3 (Table S8†) is similar to the experiment OR value (+80.0), which supports the above speculation.



Fig. 1 The structures of compounds 1–6.



The molecular formula of compound 6, a white amorphous powder, is  $C_{13}H_{10}N_2O_3$ , in accordance with the HRESIMS at m/z243.0769  $[M + H]^+$  (calculated for C<sub>13</sub>H<sub>11</sub>N<sub>2</sub>O<sub>3</sub>, 243.0770). Amino or hydroxyl (3388, 3189 cm<sup>-1</sup>), conjugated carbonyl  $(1690 \text{ cm}^{-1})$ , and aromatic ring  $(1621, 1556, \text{ and } 1453 \text{ cm}^{-1})$ groups were observed in the IR spectrum. The <sup>1</sup>H-NMR spectrum (Table 2) suggests a *para*-substituted phenyl at  $\delta_{\rm H}$  8.03 (2H, d) and 8.17 (2H, d). Three aromatic proton signals at  $\delta_{\rm H}$  7.21 (1H, dd), 7.33 (1H, dd), and 7.95 (1H, d) and three exchanged protons at  $\delta_{\rm H}$  10.64 (1H, s), 10.10 (1H, s), and 9.88 (1H, s) are also observed in the <sup>1</sup>H-NMR spectrum. HMBC relationships of H-7 with C-3 and C-5, H-2 and H-6 with C-10, and NH-8 with C-8 suggest a 4-formylbenzamide unit in 6. The <sup>1</sup>H-<sup>1</sup>H COSY correlations display an isolated spin system as H-4'/H-5'/H-6'. Meanwhile, the HMBC relationships of H-6' with C-2' and OH-3' with C-2', C-3', and C-4', combined with the molecular composition and chemical shifts, revealed a 2-substituted pyridin-3-ol

 Table 2
 NMR spectroscopic data of  $6^a$ 

	6	6		
No.	$\delta_{ m H}$	$\delta_{\rm C}$ , type		
1		138.8, C		
2	8.17, d (8.4)	128.7, CH		
3	8.03, d (8.4)	129.4, CH		
4	, , ,	138.1, C		
5	8.03, d (8.4)	129.4, CH		
6	8.17, d (8.4)	128.7, CH		
7	10.10, s	193.0, CH		
8		165.2, C		
NH-8	10.64, s	,		
2'		147.7, C		
3'		139.9, C		
OH-3'	9.88, s	,		
4'	7.33, dd (8.4, 1.2)	124.7, CH		
5'	7.21, dd (8.4, 4.2)	123.2, CH		
6′	7.95, d (5.4)	138.6, CH		

<sup>*a*</sup> NMR data ( $\delta$ ) were measured at 600 MHz for <sup>1</sup>H and at 150 MHz for <sup>13</sup>C in DMSO-*d*<sub>6</sub>. The assignments were based on <sup>1</sup>H–<sup>1</sup>H COSY, HSQC, and HMBC experiments.

moiety. Finally, the correlation of NH-8 with C-2' in the HMBC spectrum demonstrates that the above two units are linked *via* NH-8 to C-2'. Thus, compound **6** is 4-formyl-*N*-(3'-hydroxypyr-idin-2'-yl) benzamide.

In addition to compounds **1**, **3**, and **6**, the known dethiotetra(methylthio)chetomin (2),<sup>15</sup> chetoseminudin B (4),<sup>16</sup> and chetoseminudin C (5),<sup>16</sup> were also isolated from the *Chaetomium globosum* 7951. The cytotoxic effects of these compounds against human cancer cell lines were evaluated. Compounds **1–** 3 and **6** inhibited the growth of MCF-7, MDA-MB-231, H460, and HCT-8 cells (IC<sub>50</sub> from 4.5 to 65.0  $\mu$ M). Compounds **4** and 5 were inactive (IC<sub>50</sub> > 100  $\mu$ M) (Table 3, Fig. 3).

### Experimental

### General experimental procedures

See the ESI.<sup>†</sup>

### Microorganism and fermentation

The fungus *Chaetomium globosum* 7951 was isolated from the fresh healthy roots of *Panax notoginseng* gathered in Wenshan,

Table 3	Cytotoxicity	against human	cancer	cell lines of 1–6
---------	--------------	---------------	--------	-------------------

	$IC_{50} (\mu M)$					
Compd.	MCF-7	MDA-MB-231	H460	HCT-8		
1	$20.1\pm2.5$	$50.3 \pm 3.6$	$7.0\pm0.8$	$30.3 \pm 3.9$		
2	$60.5\pm7.0$	$61.2\pm5.6$	$9.4\pm0.7$	$4.5\pm0.5$		
3	$30.3\pm2.8$	$50.4\pm5.0$	$65.0\pm6.0$	$41.9\pm5.0$		
4	>100	>100	>100	>100		
5	>100	>100	>100	>100		
6	$18.0 \pm 1.5$	$25.2\pm2.8$	>100	>100		
Cisplatin	$36.0\pm3.0$	$28.0\pm3.0$	$9.0\pm0.6$	$3.5\pm0.2$		
Doxorubicin	$0.5\pm0.02$	$0.3\pm0.03$	$6.2\pm0.3$	$0.3\pm0.02$		



Fig. 3 H460, HCT8, MCF-7 and MDA-MB-231 cells were treated with the indicated concentrations of **1–3** and **6**, and cell survival was detected by the CCK8 assay. A dose-dependent curve was depicted.

Yunnan Province, China, in 2015. The strain, which was assigned the accession no. SUB5310227, was identified using nuclear 18S rDNA sequences (GenBank: MK625020) and deposited in the Microbiology Laboratory at Shenyang Pharmaceutical University. *C. globosum* 7951, an endophytic fungus, was grown on PDA at 26 °C for 6 days, and then flushed (sterilized water) into 250 ml Erlenmeyer flasks containing 50 g rice and autoclaved (121 °C, 30 min). The fermentation was then incubated at 26 °C for 40 days.

#### **Extraction and isolation**

The cultures (10 kg) were extracted three times with methanol and then filtered. The filtrate was concentrated and three extractions were performed with equal volumes of EtOAc. The EtOAc layer was evaporated with reduced pressure resulting in a crude broth extract (12.2 g). The extract was separated into 15 fractions (A–O) using silica gel column chromatography with a  $CH_2Cl_2/MeOH$  gradient elution. Fraction H was purified using Sephadex LH-20 gel column chromatography with  $CH_2Cl_2/$ MeOH (1 : 1), resulting in subfractions H1–H8. Subfraction H5 was subjected to semi-preparative HPLC with 50% acetonitrile elution into 0.1% trifluoroacetic acid to isolate compound **1**. Fraction J was purified by ODS  $C_{18}$  with a gradient of methanol in water (10–100%) to give six subfractions (J1–J6). Subfraction J3 was subjected to preparative TLC using  $CH_2Cl_2/MeOH$  (20 : 1) and then subjected to semi-preparative HPLC with 30% acetonitrile/H<sub>2</sub>O (0.1% CF<sub>3</sub>COOH) as the mobile phase to generate compound **6**. Fraction K was separated into fractions using silica gel CC with CH<sub>2</sub>Cl<sub>2</sub>/MeOH (50 : 1), resulting in fractions K1–K3. Fraction K1 was purified with semi-preparative HPLC with 19% acetonitrile into an aqueous 0.1% trifluoroacetic acid solution, resulting in compound **3**.

**Demethylchaetocochin C** (1). White amorphous powder;  $[\alpha]_D^{20}$  61.0 (*c* 0.3, MeOH); UV (MeOH)  $\lambda_{max}$  219, 286 nm; IR  $\nu_{max}$  3359, 3193, 2921, 2851, 1680, 1468, 1425, 1207, 1140, 1061, 1027, 722 cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 600 MHz) data and <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>, 150 MHz) data, see Table 1. (+)-HR-ESIMS *m/z* 727.1499 [M + H]<sup>+</sup> (calcd for C<sub>32</sub>H<sub>35</sub>N<sub>6</sub>O<sub>6</sub>S<sub>4</sub>, 727.1501).

**Chaetoperazine A (3).** White amorphous powder;  $[\alpha]_{D}^{20} 80.0 (c 0.3, MeOH);$  UV (MeOH)  $\lambda_{max}$  199, 273 nm; IR  $\nu_{max}$  3393, 3194, 2922, 2850, 1675, 1424, 1205, 1141, 801, 749, 723 cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO- $d_6$ , 600 MHz) data and <sup>13</sup>C NMR (DMSO- $d_6$ , 150 MHz) data, see Table 1. (+)-HR-ESIMS m/z 372.0994 [M + Na]<sup>+</sup> (calcd for C<sub>16</sub>H<sub>19</sub>N<sub>3</sub>O<sub>4</sub>NaS, 372.0994).

**4-Formyl-N-(3'-hydroxypyridin-2'-yl) benzamide (6).** White amorphous powder; UV (MeOH)  $\lambda_{max}$  254, 325 nm; IR  $\nu_{max}$  3388, 3189, 2921, 2850, 1690, 1621, 1556, 1453, 1387, 1320, 1210, 1142, 1052, 1029, 1011, 838, 724 cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 600 MHz) data and <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>, 150 MHz) data, see Table 2. (+)-HR-ESIMS *m*/*z* 243.0769 [M + H]<sup>+</sup> (calcd for C<sub>13</sub>H<sub>11</sub>N<sub>2</sub>O<sub>3</sub>, 243.0770).

#### ORs and NMR calculation of 3

See the ESI.<sup>†</sup>

#### Cytotoxicity assay

A CCK colorimetric assay was used to measure the cytotoxicity of compounds **1–6** was in human breast adenocarcinoma cell (MCF-7 and MDA-MB-231), human large cell lung cancer cell (H460) and human cecal adenocarcinoma cell (HCT-8). All of the cell lines were obtained from ATCC. Cells ( $5 \times 10^3$  cells per mL) were added to 96-well culture dishes and grown for 24 h (5% CO<sub>2</sub>, 37 °C) followed by the addition of fresh medium (100 µL) and the test compound. After an additional 48 h, the media was removed and fresh media with 10% CCK solution was added. The cells were incubated for 1 h (37 °C) and then the optical density at 450 nm was determined. Each assay was replicated six times. IC<sub>50</sub> values for each cell line were determined.

### Conclusions

In conclusion, two new ETPs alkaloids (**1**, and **3**), a new pyridine benzamide (**6**), and three known ETPs compounds were identified in the endophytic fungus *Chaetomium globosum* 7951. These new compounds moderately inhibit the human breast cancer cells (MCF-7 and MDA-MB-231) and human ileocecal adenocarcinoma (HCT-8) growth. The new compound **1** significantly exhibits cytotoxic against the human lung cancer cell (H460).

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

Financial support from the National Natural Science Foundation of China (NNSFC, 81302675), CAMS Innovation Fund for Medical Sciences (CIFMS, 2016-I2M-2-002, and 2018-I2M-3-005), the Drug Innovation Major Project of China (Grant No. 2018ZX09735001-002 and 2018ZX09711001-007), PUMC Youth Fund and the Fundamental Research Funds for the Central Universities (3332015133), Natural Science Foundation of Beijing Municipality (7144224), and Tianshan Cedar Project in Xinjiang Uygur Autonomous Region (2017XS10).

### Notes and references

- 1 S. E. Helaly, B. Thongbai and M. Stadler, *Nat. Prod. Rep.*, 2018, **35**, 992–1014.
- 2 L. Chen, Q. Y. Zhang, M. Jia, Q. L. Ming, W. Yue, K. Rahman, L. P. Qin and T. Han, *Crit. Rev. Microbiol.*, 2016, **42**, 454–473.
- 3 H. Nisa, A. N. Kamili, I. A. Nawchoo, S. Shafi, N. Shameem and S. A. Bandh, *Microb. Pathog.*, 2015, **82**, 50–59.
- 4 N. Fatima, S. A. Muhammad, I. Khan, M. A. Qazi, I. Shahzadi, A. Mumtaz, M. A. Hashmi, A. K. Khan and T. Ismail, *Acta Physiol. Plant.*, 2016, **38**, 136.
- 5 S. Sekita, K. Yoshihira, S. Natori and H. Kuwano, *Tetrahedron Lett.*, 1973, **14**, 2109–2112.

- 6 Q. Zhang, H. Q. Li, S. C. Zong, J. M. Gao and A. L. Zhang, *Mini Rev. Med. Chem.*, 2012, **12**, 127–148.
- 7 C. S. Jiang and Y. W. Guo, *Mini Rev. Med. Chem.*, 2011, **11**, 728-745.
- 8 M. Chen, R. Wang, W. Zhao, L. Yu, C. Zhang, S. Chang, Y. Li, T. Zhang, J. Xing, M. Gan, F. Feng and S. Si, *Org. Lett.*, 2019, 21, 1530–1533.
- 9 M.-H. Chen, S.-S. Chang, B. Dong, L.-Y. Yu, Y.-X. Wu, R.-Z. Wang, W. Jiang, Z.-P. Gao and S.-Y. Si, *RSC Adv.*, 2018, **8**, 5138–5144.
- 10 M. Chen, Y. Wu, Y. He, Y. Xu, Y. Li, D. Li, T. Feng, L. Yu, B. Hong, W. Jiang and S. Si, *Bioorg. Med. Chem. Lett.*, 2015, 25, 562–565.
- 11 G. Y. Li, B. G. Li, T. Yang, J. F. Yan, G. Y. Liu and G. L. Zhang, *J. Nat. Prod.*, 2006, **69**, 1374–1376.
- 12 N. Grimblat, M. M. Zanardi and A. M. Sarotti, *J. Org. Chem.*, 2015, **80**, 12526–12534.
- 13 Y. N. Wang, G. Y. Xia, L. Y. Wang, G. B. Ge, H. W. Zhang,
  J. F. Zhang, Y. Z. Wu and S. Lin, *Org. Lett.*, 2018, 20, 7341– 7344.
- 14 L. Li, D. Li, Y. Luan, Q. Gu and T. Zhu, *J. Nat. Prod.*, 2012, 75, 920–927.
- 15 I. Marmouzi, M. El Abbes Faouzi, N. Saidi, Y. Cherrah, N. Rehberg, S. S. Ebada, W. Ebrahim, R. Kalscheuer and P. Proksch, *Chem. Nat. Compd.*, 2017, 53, 1208–1211.
- 16 H. Fujimoto, M. Sumino, E. Okuyama and M. Ishibashi, J. Nat. Prod., 2004, 67, 98–102.