



Cite this: *RSC Adv.*, 2019, 9, 22343

Phosphonic acid mediated practical dehalogenation and benzylation with benzyl halides†

Jing Xiao,^a Yonghao Ma,^a Xiaofang Wu,^a Jing Gao,^a Zilong Tang^a and Li-Biao Han^{a,b}

Received 25th June 2019
 Accepted 16th July 2019

DOI: 10.1039/c9ra04770k

rsc.li/rsc-advances

For the first time, by using H₃PO₃/I₂ system, various benzyl chlorides, bromides and iodides were dehalogenated successfully. In the presence of H₃PO₃, benzyl halides underwent electrophilic substitution reactions with electron-rich arenes, leading to a broad range of diarylmethanes in good yields. These transformations feature green, cheap reducing reagents and metal-free conditions. A possible mechanism was proposed.

Dehalogenation and reactions of benzylation with benzyl halides are important transformations in synthetic organic chemistry. Herein we report a simple and green method for dehalogenation and benzylation with benzyl halides under metal-free conditions. Thus, under the H₃PO₃/I₂ system, dehalogenation took place readily with benzyl chlorides, bromides and iodides to produce the corresponding benzyl derivatives in high yields. In addition, in the absence of I₂, benzyl halides underwent electrophilic substitution reactions with electron-rich arenes, generating diarylmethanes efficiently.

Over the past few decades, significant efforts have been devoted to the reduction of halides.¹ Among them, a variety of reducing reagents were applied to dehalogenation reactions successfully. Some of the most representative reducing agents include metal hydrides such as tri-*n*-butyltin hydrides,² organosilanes,³ sodium borohydrides⁴ and lithium aluminium hydride.⁵ Besides, catalytic hydrogenation is also an important approach to the reduction of halides. For example, dehalogenation catalyzed by transition-metals such as nickel,⁶ palladium,^{3b,7} ruthenium,⁸ rhodium,⁹ indium,^{2a,10} iron¹¹ and cobalt¹² *etc.* have gained increased attention. Although great progress has been achieved in dehalogenation reactions, nontoxic, green, simple and cheap reducing systems still need to be explored (Scheme 1).

Recently, our group was interested in the use of H₃PO₃ in synthetic methodologies due to its low-toxicity, ready availability


and cheap cost.¹³ It is estimated that a million tons of phosphonic acid was generated as the side-product from the process of electroless nickel plating using H₃PO₂ and other chemical processes. Although most of the H₃PO₃ was laid idle as waste, we are still fascinated by the great potential ability in reduction reactions since H₃PO₃ has a reactive P(O)H bond that might serve as a green H source. Thus, for the first time, the H₃PO₃/I₂ combination was used for the reduction of benzyl halides, generating a variety of benzyl derivatives in good yields. This reducing system has the advantages of being green, cheap and metal-free.

Our investigation began with the reduction of benzyl bromide in the presence of H₃PO₃ and I₂. As shown in Table 1, reduction of benzyl bromide took place in benzene at 100 °C for 22 h, generating the desired product toluene in 48% GC yield (entry 1). Next, the temperature of the reaction was examined (entry 2 and 3). The suitable temperature was 120 °C and gave the product in 76% yield. Further screening of the amount of I₂ or H₃PO₃ failed to improve the reaction efficiency (entry 4–6). Among the solvents examined, polar solvents such as chlorobenzene, CH₃CN and dioxane gave poor results (entry 7–9). To our delight, a weak polar solvent DCE (1,2-dichloroethane) was a relatively good solvent with a yield of 69% (entry 10). When prolonging the reaction time to 36 h, DCE gave a better result than benzene with a yield of 92% (entry 11 and 12). The yield of the product could not be increased by elevating the temperature to 130 °C in DCE (entry 13).

^aKey Laboratory of Theoretical Organic Chemistry and Functional Molecule of Ministry of Education, School of Chemistry and Chemical Engineering, Hunan University of Science and Technology, Xiangtan 411201, China


^bNational Institute of Advanced Industrial Science and Technology (AIST), Tsukuba, Ibaraki 305-8565, Japan. E-mail: libiao-han@aist.go.jp; Fax: +81-29-861-6344; Tel: +81-298-61-4855

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c9ra04770k



Scheme 1 Phosphonic acid mediated dehalogenation and benzylation with benzyl halides.




Table 1 Optimization of the reaction conditions^a


Entry	H ₃ PO ₃ (equiv.)	I ₂ (equiv.)	Solvent	Temp/°C	Yield (%)
1	2.0	10%	Benzene	100	48%
2	2.0	10%	Benzene	120	76%
3	2.0	10%	Benzene	130	75%
4	2.0	15%	Benzene	120	74%
5	2.0	5%	Benzene	120	70%
6	3.0	10%	Benzene	120	64%
7	2.0	10%	PhCl	120	21%
8	2.0	10%	CH ₃ CN	120	7%
9	2.0	10%	Dioxane	120	26%
10	2.0	10%	DCE	120	69%
11	2.0	10%	DCE	120	92% ^b
12	2.0	10%	Benzene	120	61% ^b
13	2.0	10%	DCE	130	82%

^a Conditions: a mixture of **1a** (0.3 mmol), H₃PO₃, and I₂ in solvent (0.6 mL) was stirred at indicated temperature for the 22 h. Yield based on GC yield using dodecane as an internal standard. ^b 36 h.

With the optimized reaction conditions in hand, the scope of the reaction with respect to benzyl halides was examined (Table 2). Benzyl bromide was reduced to the corresponding product in 84% GC yield, albeit 38% isolated yield was obtained due to its low boiling point (**2a**). Bromides bearing methyl or phenyl groups at the *para*-position on benzene were reduced smoothly to afford the product in high yields (**2b** and **2c**). Interestingly, substrates containing halogens such as F, Cl and Br at the *para*-position on benzene stayed intact during the dehalogenation

Table 2 Reduction of the benzyl bromide derivatives with H₃PO₃/I₂^a



2a : 38% (84%)	2b : 41% (92%)	2c : 98%	2d : 32% (89%)
2e : 49% (98%)	2f : 68% (97%)	2g : 26% (72%)	2h : 61%
2i : 52%	2j : 39% (95%)	2k : 97%	2l : 78%

^a Conditions: a mixture of **1** (0.6 mmol), H₃PO₃, and I₂ in DCE was stirred at 120 °C for 36 h. Isolated yield based on three parallel reactions. Yield in parenthesis are GC yield.

(**2d–2f**). Other electron-deficient bromides having CF₃, C(O)OMe and NO₂ groups were also tolerated well in this reductive system (**2g–2i**). Secondary and tertiary benzyl bromides such as (1-bromoethyl)benzene and (bromomethanetriyl)tribenzene also underwent this reduction successfully generating the product in 95% and 97% yield, respectively (**2j** and **2k**). When 1-(bromomethyl)naphthalene was subjected into the reaction, 78% yield of the product was obtained (**2l**). Unfortunately, alkyl and aryl bromides did not go through this dehalogenation reaction.

We next extended the scope of benzyl halides under similar conditions. As shown in Table 3, benzyl chloride also proceeded smoothly (**2a**), and various reduction products were obtained. Chlorides bearing substituents (Me, F, Cl and Br) were suitable for this reduction reaction, generating the corresponding product in good yields (**2b**, **2d–2f**). With regard to electron-withdrawing group under H₃PO₃/I₂ system, we observed that CF₃, CN and NO₂ substituents were tolerated in good yields (**2g**, **2m** and **2i**). Furthermore, secondary chlorides such as (1-chloroethyl)benzene and chlorodiphenylmethane were also worked well in this transformation, providing the product in high yields (**2j** and **2n**). Gratefully, large steric hindrance triphenylmethane chloride was reduced readily under the present condition in 97% yield (**2k**). Finally, we conducted the dehalogenation of benzotrichloride and benzyl iodide to explore possible reduction of polyhalogenated substrates and iodides. Indeed, **2a** was formed in 41% and 88% yield, respectively.

During the above studies of dehalogenation of benzyl halides, we found that interestingly, when the reaction was performed in the absence of I₂, electrophilic substituent

Table 3 Reduction of the benzyl chloride derivatives with H₃PO₃/I₂^a


2a : 25% (80%)	2b : 34% (71%)	2d : 22% (65%)	2e : 37% (73%)
2f : 65% (84%) ^b	2g : 20% (54%) ^c	2m : 53% ^d	2i : 42%
2j : 41% (98%)	2n : R ¹ = Ph; R ² = H, 99%; R ¹ = Ph; R ² = Cl, 77%	2k : R ¹ = R ² = Ph; 97%	2a : 41% ^e 25% (88%) ^f

^a Conditions: a mixture of **3** (0.6 mmol), H₃PO₃, and I₂ in benzene was stirred at 100 °C for 36 h. Isolated yield based on three parallel reactions. Yield in parenthesis are GC yield. ^b 120 °C. ^c 4.0 equiv. H₃PO₃. ^d 130 °C. ^e Benzotrichloride and 5.0 equiv. H₃PO₃ was used. ^f Benzyl iodide was used.



reaction took place to affording diarylmethane in excellent yields. Diarylmethane are widely used in organic synthesis, pharmaceutical development and material science.¹⁴ Among the methods reported, Friedel–Crafts (FC) benzylation of arenes¹⁵ and transition-metal catalysed cross coupling reactions¹⁶ are the important approaches to preparing these compounds. However, those methods are usually need equivalent expensive Lewis acids, metal and not readily available materials. Although various strategies for the metal-free access to diarylmethanes have been explored,¹⁷ a more simple, practical, and greener method for their preparation are highly desirable.

As shown in Table 4, various arenes were subjected into this electrophilic substitution reaction. Thus, in the presence of H₃PO₃, benzyl chlorides and bromides underwent this transformation successfully with benzene or mesitylene to afford the corresponding diarylmethanes in good to excellent yields (**2n** and **4a**). Gratifying, electron-rich arenes such as toluene, phenol and anisole were also compatible in this reaction conditions with isomer products generated (**4b–4d**). In addition, when chlorobenzene was used as the substrate, good yield of the corresponding product was observed (**4e**). Subsequently, a series of benzyl halides with benzene were investigated in the presence of H₃PO₃. Benzyl halides bearing methyl, phenyl, fluoride, chloride and bromide led to good to high yield of the products (**4b**, **4f–4h**).

Table 4 Benzylation of arenes with benzyl halides^a

2n: X=Cl; 84%; X=Br; 87% ^b	4a: X=Cl; 95%; X=Br; 96% ^b	4b: X=Cl; 95%; o:m:p = 46:8:46; X=Br; 78% ^b ; o:m:p = 43:7:50
4c: X=Cl 99%; o:p = 64:36 X=Br 98% ^b ; o:p = 55:45	4d: X=Cl; 94%; o:p = 49:51 X=Br; 92% ^b ; o:p = 43:57	4e: X=Cl; 72% ^c o:m:p = 47:2:51
4b: X=Cl; 84%; X=Br; 72%; ^{b,c}	4f: X=Br; 59% ^{b,c}	4g: X=Cl; 99%; X=Br; 74% ^{b,c}
4e: X=Cl; 92% ^{b,c} X=Br; 56% ^{b,c}	4h: X=Cl; 81% ^{b,c} X=Br; 53% ^{b,c}	4i: X=Br; 35% ^d


^a Conditions: a mixture of **1** or **3** (0.6 mmol) and H₃PO₃ in arenes was stirred at 130 °C for 24 h. Isolated yield based on three parallel reactions. Isomer ratio was determined by GC. ^b 36 h. ^c 150 °C. ^d 160 °C.

It is worth noting that benzyl bromide with electron-drawing group such as methyl ester was also applicable in this reaction, albeit in moderate yield of the product (**4i**). Unfortunately, the reaction gave trace product with benzyl halides having strong electron-withdrawing groups such as nitro or trifluoromethyl group.

To highlight the synthetic utility of these transformation, some gram-scale reactions were conducted. As shown in Scheme 2, 12 mmol (chloromethylene)dibenzene **3n** was reduced smoothly in the presence of 2.0 equiv. H₃PO₃ and 10 mol% I₂ in benzene at 130 °C for 36 h, generating 1.956 g product in 97% isolated yield (eqn (1)). Besides, we were pleased to find that the gram scale synthesis of diphenylmethane by reaction of benzyl chloride with benzene was also achieved (eqn (2)). The above results revealed that these reactions were practical and have the potential to scale up.


Finally, some control experiments were conducted to evaluate the different halides and electron effect of these reactions. 4-Methylbenzyl chloride and 4-methylbenzyl bromide were reduced to the product with 20% and 31% substrate conversion, respectively (eqn (3)). Benzyl bromides bearing electron-donating group methyl, H and electron-withdrawing group CF₃ were also subjected into the reaction in one pot, and the corresponding product were obtained in 17%, 9% and trace yield with a large amount of substrate recovered, respectively (eqn (4)). Similarly, the benzylation of benzene by the above three bromides affording the corresponding diarylmethanes in a downward trend in yields (eqn (5)). The above results revealed that benzyl bromides were more reactive than benzyl chloride in dehalogenation reactions. Besides, in the same type of halides, electron-donating group having more positive influence than electron-deficient whether in reduction or benzylation reactions, which may suggest that these reactions were involving a carbon cation mechanism (Scheme 3).

Based on the above results and previous reports,¹³ a possible mechanism was proposed in Scheme 4. HI was generated *in situ* by reaction of H₃PO₃ and I₂, halogen exchange between benzyl halides and HI afforded intermediate **A**, further reduction of **A** by HI to give the corresponding product (eqn (a)). For benzylation of arenes by halides, we deduced that carbon cation intermediate **B** was generated *via* C–X bond cleavage in the presence of H₃PO₃, **B** underwent electrophilic substitution reaction with arenes to afford the corresponding diarylmethanes (eqn (b)).




Scheme 2 Gram-scale reactions.





Scheme 3 Control experiments.



Scheme 4 Possible mechanism.

Conclusions

In summary, we report a practical method for dehalogenation of benzyl halides and preparation of diarylmethanes by using a H_3PO_3 system. Various benzyl halides could be readily reduced by the $\text{H}_3\text{PO}_3/\text{I}_2$ combination for the first time. In the absence of I_2 , electrophilic substitution reactions between benzyl halides and arenes occurred smoothly, furnishing diarylmethanes in good yields. This method provides a simple, cheap and green approach for the reduction of benzyl halides and synthesis of diarylmethanes.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements


JX is thankful for a postdoc fellowship from AIST and financial support from the National Natural Science Foundation of China (21703061) and the Natural Science Foundation of Hunan province (2017JJ3081).

Notes and references

- 1 (a) F. Alonso, I. P. Beletskaya and M. Yus, *Chem. Rev.*, 2002, **102**, 4009–4092; (b) G. Chelucci, S. Baldino, G. A. Pinna

and G. Pinna, *Curr. Org. Chem.*, 2012, **16**, 2921–2945; (c) C. E. Castro, *Reviews of Environmental Contamination and Toxicology*, Springer, New York, 1998, 155, pp. 1–67; (d) G. L. Larson, *Org. React.*, 2008, **71**, 5–119; (e) T. Hennebel, J. Benner, P. Clauwaert, L. Vanhaecke, P. Aelterman, R. Callebaut, N. Boon and W. Verstraete, *Biotechnol. Lett.*, 2011, **33**, 89–95; (f) Á. F. Cañete, C. O. Salas and F. C. Zacconi, *Molecules*, 2013, **18**, 398–407.

- 2 (a) K. Inoue, A. Sawada, I. Shibata and A. Baba, *Tetrahedron Lett.*, 2001, **42**, 4661–4663; (b) A. Rahm, R. Amardeil and M. Degueil-Castaing, *J. Organomet. Chem.*, 1989, **371**, 4–8; (c) M. Pereyre, J. P. Quintard and A. Rahm, *Reduction of Organic Halides in Tin in Organic Synthesis*, Butterworth, London, 1986.
- 3 (a) C. Chatgililoglu, *Acc. Chem. Res.*, 1992, **25**, 188–194; (b) R. Boukherroub, C. Chatgililoglu and G. Manuel, *Organometallics*, 1996, **15**, 1508–1510; (c) J. Yang and M. Brookhart, *Adv. Synth. Catal.*, 2009, **351**, 175–187.
- 4 S. Krishnamurthy and H. C. Brown, *J. Org. Chem.*, 1980, **45**, 849–856.
- 5 S. Krishnamurthy and H. C. Brown, *J. Org. Chem.*, 1982, **47**, 276–280.
- 6 (a) A. F. Barrero, E. J. Alvarez-Manzaneda, R. Chahboun, R. Meneses and J. L. Romera, *Synlett*, 2001, **4**, 485–488; (b) S. Zinovyev, A. Perosa, S. Yufit and P. Tundo, *J. Catal.*, 2002, **211**, 347–354; (c) J. M. Khurana, S. Kumar and B. Nand, *Can. J. Chem.*, 2008, **86**, 1052–1054.
- 7 (a) T. Hara, K. Mori, M. Oshiba, T. Mizugaki, K. Ebitania and K. Kaneda, *Green Chem.*, 2004, **6**, 507–509; (b) G. L. Larson and J. L. Fry, *Org. React.*, 2008, **71**, 1–737; (c) T. Hara, T. Kaneta, K. Mori, T. Mitsudome, T. Mizugaki, K. Ebitania and K. Kaneda, *Green Chem.*, 2007, **9**, 1246–1251.
- 8 (a) A. Jana, J. Mondal, P. Borah, S. Mondal, A. Bhaumik and Y. Zhao, *Chem. Commun.*, 2015, **51**, 10746–10749; (b) M. C. Haibach, B. M. Stoltz and R. H. Grubbs, *Angew. Chem., Int. Ed.*, 2017, **56**, 15123–15126.
- 9 K. Fujita, M. Owaki and R. Yamaguchi, *Chem. Commun.*, 2002, 2964–2965.
- 10 Á. F. Cañete, C. O. Salas and F. C. Zacconi, *Molecules*, 2013, **18**, 398–407.
- 11 H. Guo, K.-i. Kanno and T. Takahashi, *Chem. Lett.*, 2004, **33**, 1356–1357.
- 12 (a) B. Sahoo, A. E. Surkus, M. M. Pohl, J. Radnik, M. Schneider, S. Bachmann, M. Scalone, K. Junge and M. Beller, *Angew. Chem., Int. Ed.*, 2017, **56**, 11242–11247; (b) F. Ungváry and L. Markó, *J. Organomet. Chem.*, 1980, **193**, 379–382.
- 13 J. Xiao and L.-B. Han, *Tetrahedron*, 2019, **75**, 3510–3515.
- 14 (a) J. S. Kim and D. T. Quang, *Chem. Rev.*, 2007, **107**, 3780–3799; (b) Y.-Q. Long, X.-H. Jiang, R. Dayam, T. Sanchez, R. Shoemaker, S. Sei and N. Neamati, *J. Med. Chem.*, 2004, **47**, 2561–2573; (c) W. N. Washburn, *J. Med. Chem.*, 2009, **52**, 1785–1794; (d) T. Brotin and J.-P. Dutasta, *Chem. Rev.*, 2009, **109**, 88–130.
- 15 (a) C. Friedel and J. M. Crafts, *Compt. Rend.*, 1877, **84**, 1450; (b) G. A. Olah, S. J. Kuhn and S. H. Flood, *J. Am. Chem. Soc.*,



- 1962, **84**, 1688–1695; (c) M. Rueping and B. J. Nachtsheim, *Beilstein J. Org. Chem.*, 2010, **6**, 6, DOI: 10.3762/bjoc.6.6.
- 16 (a) H. Jiang, S.-C. Sha, S. A. Jeong, B. C. Manor and P. J. Walsh, *Org. Lett.*, 2019, **21**, 1735–1739; (b) J. Xiao, T. Chen and L.-B. Han, *Org. Lett.*, 2015, **17**, 812–815; (c) J. Xiao, a. J. Yang, T. Chen and L.-B. Han, *Adv. Synth. Catal.*, 2016, **358**, 816–819; (d) B. Liégault, J. L. Renaud and C. Bruneau, *Chem. Soc. Rev.*, 2008, **37**, 290–299.
- 17 (a) R.-J. Tang, T. Milcent and B. Crousse, *J. Org. Chem.*, 2018, **83**, 14001–14009; (b) G. Schafer and J. W. Bode, *Angew. Chem., Int. Ed.*, 2011, **50**, 10913–10916; (c) J. Zhu, M. Perez and D. W. Stephan, *Angew. Chem., Int. Ed.*, 2016, **55**, 8448–8451; (d) P. A. Champagne, Y. Benhassine, J. Desroches and J. F. Paquin, *Angew. Chem., Int. Ed.*, 2014, **53**, 13835–13839.

