


 Cite this: *RSC Adv.*, 2019, 9, 41639

# Sophorolipid exhibits antifungal activity by ROS mediated endoplasmic reticulum stress and mitochondrial dysfunction pathways in *Candida albicans*<sup>†</sup>

 Farazul Haque,<sup>‡§<sup>a</sup></sup> Nitish Kumar Verma,<sup>‡<sup>a</sup></sup> Mohammad Alfatah,<sup>¶<sup>b</sup></sup> Swati Bijlani<sup>||<sup>b</sup></sup> and Mani Shankar Bhattacharyya<sup>||<sup>‡\*<sup>a</sup></sup></sup>

In the present study, we investigated the mechanism of cell death in *C. albicans* due to treatment with sophorolipid (SL). SL is an extracellular glycolipid biosurfactant produced by various species of non-pathogenic yeasts and is known to inhibit the growth and biofilm formation of *C. albicans*. This study revealed that treatment of *C. albicans* cells with SL increases the ROS production and expression of oxidative stress-related genes significantly (*SOD1*, *CAT1*). Increased ROS level within the cells causes ER stress and release of Ca<sup>2+</sup> in the cytoplasm and alteration of the mitochondrial membrane potential (MMP). Quantitative real time-polymerase chain reaction (qRT-PCR) data showed that SL also upregulates the Endoplasmic Reticulum (ER) stress marker *HAC1*. Flow cytometric analysis (AnnexinV/PI) indicated that the cell death may have occurred due to necrosis which was further confirmed by LDH release assay and transmission electron microscopy (TEM). Further experiments with the null mutant  $\Delta$  *hog1* strain of *C. albicans* SC5314 indicated the activation of the osmotic stress response pathway (HOG-MAPK) and *SAP9*. This study gave an insight into the mechanism of cell death initiation by glycolipids and indicated that further modification of these molecules can lead to the development of new therapeutic agent against *C. albicans*.

 Received 19th September 2019  
 Accepted 1st December 2019

DOI: 10.1039/c9ra07599b

[rsc.li/rsc-advances](http://rsc.li/rsc-advances)

## Introduction

Members of the genus *Candida* are known to be some of the most common opportunistic human pathogens which cause infection to the superficial mucosa and are major causative agents for life threatening systemic and blood level infections.<sup>1,2</sup> *Candida albicans* (*C. albicans*) is the fourth most common causative agent of candidiasis and is associated with very high mortality rates (about 50%).<sup>3,4</sup> Depending on the environmental conditions *C. albicans* can grow either as unicellular budding yeast or in the form of filamentous pseudohypha and hypha.<sup>5,6</sup>

Both yeast and hyphal forms of pathogen invade the bloodstream and cause invasive candidiasis.<sup>7</sup> Currently, three classes of antifungal drugs (azoles, echinocandins, and polyenes) are being used for the treatment of candidiasis.<sup>3</sup> However, severe toxicity, adverse side effects, high cost and development of drug resistance have emerged to be the pitfalls of these therapies.<sup>3</sup> Thus, there is an urgent need to develop new antifungal agents for effective therapy of candidiasis. Research is in progress to synthesize effective antifungal agents by derivatizing and improving the properties of existing antifungal drugs, by screening of novel compounds or by developing new chemical entities against potential fungal targets.<sup>8–10</sup> Presently, several new molecules are in the various stages of their development as potential antifungal drug candidate.<sup>11</sup>

Sophorolipid (SL), a glycolipid biosurfactant (Bs) is produced extracellularly by several non-pathogenic yeasts.<sup>12,13</sup> It is one of the most promising and attractive Bs found in nature and has been reported to have antimicrobial,<sup>14</sup> antiviral, anti-spermi-cidal,<sup>15</sup> anticancer<sup>16,17</sup> and immunomodulatory activities.<sup>18</sup> In a previous study, we have demonstrated the inhibitory effect of SL against biofilm formation of *C. albicans* by downregulating the hyphae associated genes.<sup>19</sup> SL interacts synergistically with antifungal drugs (amphotericin B and fluconazole) to kills planktonic cells, inhibit biofilm formation and eradicate

<sup>a</sup>Biochemical Engineering Research & Process Development Centre (BERPDC), CSIR-Institute of Microbial Technology (IMTECH), Sector-39A, Chandigarh 160036, India. E-mail: manishb@imtech.res.in; Fax: +91-172-2695215; Tel: +91-172-6665313

<sup>b</sup>Yeast Molecular Biology Laboratory, CSIR-Institute of Microbial Technology (IMTECH), Sector-39A, Chandigarh 160036, India

<sup>†</sup> Electronic supplementary information (ESI) available. See DOI: 10.1039/c9ra07599b

<sup>‡</sup> Contributed equally to this work.

<sup>§</sup> Current address: Department of Immunology and Genomic Medicine, Graduate School of Medicine, Kyoto University, Japan.

<sup>¶</sup> Current address: Chemical Genomics Lab, Bioinformatics Institute, Agency for Science, Technology and Research (A\*STAR), Singapore 138673.

<sup>||</sup> Current address: School of Molecular Cell Biology and Biotechnology, Tel Aviv University, Ramat Aviv, Israel.



already formed biofilm.<sup>19</sup> However, the mechanism of cell death remained unexplored. Existing literature reports suggest that SL mediates bacterial cell death by damaging/impairing the plasma membrane leading to pore formation and causing cell lysis.<sup>20,21</sup> SL also induces apoptosis in H7402 human liver cancer cells by blocking the cell cycle at G1 phase, activating caspase-3, and increasing  $\text{Ca}^{2+}$  level in the cytoplasm.<sup>22</sup> Since SL induces different cascades of cellular activities leading to cell death, here; we have investigated the mechanism of action of the molecule causing cell death in *C. albicans*. Our investigations suggest that treatment of *C. albicans* with SL enhances ROS production, upregulates ROS specific genes (*CAT1*, *SOD1*) and activates pathways leading to oxidative stress. Moreover, as evidenced by the higher transcript level of *HAC1* (a marker of ER stress) SL causes ER stress triggering efflux of  $\text{Ca}^{2+}$  from ER into cytoplasm and the activation of high osmolarity glycerol (HOG) mitogen-activated protein kinase (MAPK) pathway which is known to be involved in the cell's responses to oxidative<sup>23</sup> and osmotic stresses.<sup>24</sup> All these stress conditions finally lead to necrosis and cell death.

## Material and methods

### Strain, media and growth condition

The wild type strain of *C. albicans* SC5314<sup>25</sup> and null  $\Delta$  *hog1* mutant was used in this study. Strains were preserved in glycerol (as frozen stocks) at  $-80\text{ }^{\circ}\text{C}$  and revived on YPD agar medium (1% Bacto yeast extract, 2% Bacto peptone, 2% glucose and 2% Bacto agar). Culture was grown in YPD broth at  $30\text{ }^{\circ}\text{C}$  with shaking (200 rpm). Production and extraction of SL were carried out as reported earlier (production, isolation, purification and characterization has been briefly described in ESI).<sup>†19,26</sup> The stock solution of SL was prepared in dimethyl sulfoxide (DMSO, Sigma), and stored at  $-20\text{ }^{\circ}\text{C}$  until use. In each assay, SL was dissolved in DMSO in such that the concentration of DMSO does not exceed to 5%.

### Quantification of reactive oxygen species (ROS), mitochondrial membrane potential (MMP) and intracellular calcium ion ( $\text{Ca}^{2+}$ ) level

Cells from primary overnight culture were harvested, washed and inoculated in 2 mL fresh YPD medium with an  $\text{OD}_{600}$  of 0.5. Different concentrations of SL (100 and  $200\text{ }\mu\text{g mL}^{-1}$ ) along with control that contains carrier (DMSO) were added to the cells and incubated for 5 h at  $30\text{ }^{\circ}\text{C}$  with shaking condition (200 rpm). After incubation, cells were centrifuged and pellet was washed twice with  $1\times$  phosphate buffer saline (PBS, pH 7.4) and resuspended in  $1\times$  PBS, pH 7.4. ROS-specific dye  $\text{H}_2$ -DCFDA (Molecular probe) was added to the cells at a final concentration of  $2\text{ }\mu\text{M}$  and further incubated for 30 min at  $30\text{ }^{\circ}\text{C}$ . Cells were then washed twice with PBS and analysed by BD Accuri™ C6 flow cytometer. MMP was determined using mitochondria-specific voltage-dependent dye (DiOC<sub>6</sub>) (Molecular probe). Cells were stained with 5 mM DiOC<sub>6</sub> for 30 min at  $30\text{ }^{\circ}\text{C}$ , washed twice with PBS and analysed by flow cytometer. The  $\text{Ca}^{2+}$  level was determined using Fluo-3-acetoxymethyl ester (Fluo-3-AM)

dye (Molecular probe). BAPTA which is an intracellular  $\text{Ca}^{2+}$  chelator was added to the cells at a final concentration of 10 mM, 1 h prior to SL treatment. Cells were incubated with Fluo-3-AM (5 mM) for 15 min at  $30\text{ }^{\circ}\text{C}$ , washed twice with PBS and analysed by flow cytometer. In all the above flow cytometric assays, fluorescence was measured in FL1 channel and for each sample,  $10^4$  cells were analysed. Three independent experiments were performed with two replicates each time.

### qRT-PCR analysis

Effect of SL on the expression of *CAT1*, *SOD1*, *HOG1*, *HAC1* and *SAP9* was evaluated by two-step quantitative real time polymerase chain reaction (qRT-PCR). Total RNA of the cells was extracted from SL treated ( $100\text{ }\mu\text{g mL}^{-1}$  and  $200\text{ }\mu\text{g mL}^{-1}$ ) and untreated samples in RPMI-1640 medium using hot phenol/chloroform extraction method.<sup>27</sup> cDNA was synthesized from DNase I treated total RNA using iScript™ cDNA Synthesis Kit (BIO-RAD) as per manufacturer's instructions. Primers for target genes (*CAT1*, *SOD1*, *HOG1*, *HAC1* and *SAP9*) and housekeeping internal control gene (*ACT1*) were designed using Gene Runner software (Table 1) and synthesized from Sigma. cDNA template (100 ng), gene specific primers (200 nM) and iQ™ SYBR® Green Supermix (BIO-RAD) were added to the reaction mixture in accordance with the manufacturer's instructions. qRT-PCR was performed in Mastercycler® eprealplex Real-time PCR system. The following parameters were used for qRT-PCR: an initial denaturation at  $95\text{ }^{\circ}\text{C}$  (3 min), followed by 40 cycles of denaturation ( $95\text{ }^{\circ}\text{C}/1\text{ min}$ ), annealing ( $58\text{ }^{\circ}\text{C}/30\text{ s}$ ), and extension ( $72\text{ }^{\circ}\text{C}/20\text{ s}$ ), and melting-curve analysis starting from initial temperature  $50\text{ }^{\circ}\text{C}$  to  $95\text{ }^{\circ}\text{C}$ , with gradual increase in  $0.5\text{ }^{\circ}\text{C}/15\text{ second}$ . The specificity of the primers was confirmed by melting curve analysis. The generated CT values of target genes were normalized to the CT value of housekeeping *ACT1* gene. Relative expression fold changes were evaluated by  $\Delta\Delta\text{C}_T$  method using  $2^{-\Delta\Delta\text{C}_T}$  formula.<sup>28</sup>

### Annexin V and propidium iodide (PI) staining

Annexin V-FITC 488 and PI staining was performed according to assay kit protocol (Life Technology, India). Briefly, cells were treated with different concentrations of SL as mentioned above. Cells were washed with cold PBS and resuspended in  $500\text{ }\mu\text{L}$  annexin binding buffer together with  $5\text{ }\mu\text{L}$  each of Annexin V and PI ( $10\text{ mg mL}^{-1}$ ). The cells were incubated for 30 min at room temperature. Stained cells were directly acquired and analysed by flow cytometry. Three independent experiments were done with two replicates each time.

### Lactate dehydrogenase release assay

The reduction of  $\text{NAD}^+$  to NADH in the cell medium was determined by lactate dehydrogenase (LDH) activity kit (Pierce, Thermo fisher scientific). Briefly,  $\sim 2 \times 10^6$  cells in  $100\text{ }\mu\text{L}$  of YPD medium were dispensed in 96 well plate. Different concentrations of SL were added to each well and incubated the plate for 0, 12 and 24 h at  $30\text{ }^{\circ}\text{C}$  and further processed as described by manufacturer protocol. Followed by incubation for defined time-points, absorbance was measured using



Table 1 List of the primers used in the qRT-PCR study

Primer	Sequence (5'-3')	$T_m$ (°C)	Amplified product size (bp)
CaHOG1-R	GGTCAGAAAAGGTTTCATGAC	53.0	90
CaHOG1-F	ATGGGAGCATTTGGTTTG	55.0	
CaSOD1-R	AATATGGAACCTCTCAAGGC	51.0	140
CaSOD1-F	TTAAAGCTGTCGCTGTTGTC	54.0	
CaSAP9-R	AGTCCATGACCGAAAGATC	54.0	252
CaSAP9-F	GAGTCCAAAGATGATTTATCCC	52.0	
CaCAT1-R	AAAAACACCATAAGCACCG	54.0	156
CaCAT1-F	CCAATTCCAGAACCATTG	51.0	
CaHAC1-R	ATTAGTTGGACCGGAAGATG	55.0	203
CaHAC1-F	TACAACCAACACATCAACCAG	54.0	
CaACT1-R	GGTTTGAAGCTGCTGGTATTGACC	60.8	135
CaACT1-F	ACGTTTCAGCAATACCTGGGAACATG	60.0	

spectrophotometer (Biotech) at 490 nm. Three independent experiments were performed with three replicates. Values were expressed as mean with their corresponding standard deviations (SD) and statistical significance was analyzed by Student's *t*-test (two-tailed, unequal variance). A *p*-value of <0.05 was considered statistically significant.

#### PI-staining and permeabilization assay

The Propidium Iodide (PI) staining assay was carried out as described by Alfatah *et al.*<sup>29</sup> with slight modification. Briefly, *C. albicans* SC5314 cells of 0.5 OD were treated with different concentration of SL along with DMSO as vehicle control in 2 mL of YPD medium at 30 °C and 200 rpm for 5 h. Further, the treated cells were washed twice and resuspended in 1× PBS. Furthermore, in the dark condition, cells were incubated in PI stain (5 µg mL<sup>-1</sup>) at 30 °C for 30 min with shaking. The uptake of PI stain in the cells was visualized by Confocal Laser Scanning Microscopy (CLSM) at 60× magnification.

#### Transmission electron microscopy (TEM)

The integrity of the cell's plasma membrane was visualised by TEM study. Briefly, cells were treated with SL as mentioned above. Samples were prepared as described by Sangetha *et al.*, with slight modifications.<sup>30,31</sup> After incubation with SL, cells were centrifuged and fixed with 2% glutaraldehyde at room temperature. After fixation, cells were washed with sodium phosphate buffer and post fixed with 1% OsO<sub>4</sub> for 2 h at 4 °C. The cells were then dehydrated in series of ethanol concentrations. Negative staining was performed to visualise the lipid membrane.

#### Time-kill curve assay

For time kill curve essay, *C. albicans* SC5314 and  $\Delta$  *hog1* mutant cells were inoculated in YPD medium at starting OD<sub>600</sub> of 0.3 (about  $6 \times 10^5$  cells per mL) and allowed to grow overnight. SL alone (200 µg mL<sup>-1</sup>) and/or in combination with different concentrations of sorbitol (0.25 M and 0.5 M) (Sigma Aldrich) were added to cell suspension. Dimethyl sulfoxide (DMSO)

(Sigma Aldrich) was used as vehicle control and the cell suspension was cultured at 30 °C, 200 rpm. Cells without SL were used as a negative control. Starting from 0 h time-point, OD of cell suspension was measured every 2 h intervals, using spectrophotometer (Hitachi, U2900). Six independent experiments were performed and plot using graphpad prism software.

#### Statistical significance


Statistical significance between treated and control groups was analyzed by Student's *t*-test (two-tailed, unequal variance). A *p*-value of <0.05 was considered statistically significant.

## Results

### Measurement of ROS production and overexpression of *SOD1* and *CAT1* in *C. albicans* treated with SL

Literature reports suggest that SL induces cell death by activating various stress response pathway and ultimately by cell lysis.<sup>20-22</sup> Therefore, we hypothesized that SL mediated cell death of *Candida albicans* occurs due to the activation of multiple stress response pathways. Cellular stress due to external stimuli is manifested by ROS production within the cell. Therefore, we determined the ROS production in *C. albicans* treated with SL, by measuring the fluorescence of ROS-specific dye H<sub>2</sub>-DCFDA. SL treatment enhanced the ROS generation in a concentration-dependent manner (Fig. 1A). In the presence of 100 µg mL<sup>-1</sup> SL and 200 µg mL<sup>-1</sup> SL, ROS generation was increased by 3.9-fold (45% of total cell population) and 6.5-fold (74% of cell population), respectively, as compared to control (11.5% of cell population) (Fig. 1A). Furthermore, higher level of superoxide dismutase (SOD) and catalase (CAT) production is known to be the general cellular response to mitigate the effect of high level of cellular ROS. Therefore, we studied the cellular expression level of *CAT1* and *SOD1* by quantitative RT-PCR. Treatment of SL was found to increase the *CAT1* expression by 2 fold and 3 fold (Fig. 1B) and *SOD1* transcript levels by 4 fold and 6 fold (Fig. 1C). (when treated with 100 and 200 µg mL<sup>-1</sup> of SL separately, in each case).





**Fig. 1** Sophorolipid (SL) induced reactive oxygen species (ROS) production and overexpression of stress-related genes in *C. albicans* SC5314. (A) The strain was grown in YPD medium for 5 h in the absence or presence of indicated concentrations of SL and stained with ROS specific dye, H<sub>2</sub>-DCFDA. Samples were analyzed by flow cytometry in FL1 channel. (B and C) Expression of stress-related genes *CAT1* and *SOD1* were assessed by using qRT-PCR. Results represent the average of three independent experiments  $\pm$ SD. \* $p < 0.05$  when compared with the SL untreated controls.

### Alteration of mitochondrial membrane potential (MMP) in the presence of SL

Mitochondria play pivotal role in cellular homeostasis, energy metabolism, ATP synthesis and protein synthesis and import. Maintenance of MMP is essential for normal cellular function.<sup>32–34</sup> Here, we studied the effect of SL on alteration of MMP of *C. albicans* by measuring the fluorescence of a voltage-dependent dye DiOC<sub>6</sub>, using flow cytometry. It was found that 50% and 71% of cell population exhibited mitochondrial membrane depolarization at 100  $\mu\text{g mL}^{-1}$  SL and 200  $\mu\text{g mL}^{-1}$  SL concentration, respectively, thereby, indicating a dose-dependent effect (Fig. 2). These results suggest that SL affect the mitochondrial physiological integrity leading to the loss of its membrane potential which could be the reason of enhanced ROS production.

### SL causes endoplasmic reticulum (ER) stress and elevates intracellular Ca<sup>2+</sup> level in *C. albicans*

Endoplasmic reticulum (ER) works as the Ca<sup>2+</sup> store home within the cellular environment. Upon ER stress, Ca<sup>2+</sup> is released to the cytoplasm and subsequently enters to the mitochondria.<sup>34,35</sup> Although Ca<sup>2+</sup> mediated signalling helps the cells to cope up with the ER stress, it impairs the mitochondrial function leading to elevated ROS production.<sup>35,36</sup> Since ER and mitochondria interplay through Ca<sup>2+</sup> signalling during stress condition,<sup>34–36</sup> we tested the effect of SL on intracellular Ca<sup>2+</sup> level. The intracellular level of Ca<sup>2+</sup> was measured by Fluo-3-AM.

The labelled calcium indicator Fluo-3-AM exhibits an increase in fluorescence upon binding to Ca<sup>2+</sup>. We found that cells treated with 100  $\mu\text{g mL}^{-1}$  SL and 200  $\mu\text{g mL}^{-1}$  SL showed increased Ca<sup>2+</sup> level by 4.9-fold (38.4% of cell population) and 6.2-fold (49.3% of cell population), respectively, as compared to control (7.9% of cell population) (Fig. 3, panel A). However, Fluo-3-AM fluoresces was quenched when the cells were treated with BAPTA (Ca<sup>2+</sup> chelator; Fig. 3A). These results strongly suggested that SL causes ER stress which commits the release of stored Ca<sup>2+</sup> into the cytosol.

ROS disturbs protein folding and induces endoplasmic reticulum (ER) stress activating unfolded protein response (UPR) pathway. UPR elements are related to various ER-resident molecular chaperones and protein folding catalyst genes, classically known as Ire1-*HAC1* signalling elements. Thus, the bZIP transcription factor *HAC1* may be considered as a marker of ER stress. Here, we checked the expression of *HAC1* in SL treated *C. albicans* cells by q-RT-PCR. The expression of *HAC1* was found to be up-regulated by 5 fold and 7.6 fold when treated with 100 and 200  $\mu\text{g mL}^{-1}$  of SL (Fig. 3B). Activation of *HAC1* also signals the activation of many other downstream processes including Ca<sup>2+</sup> leakage from the ER lumen.

### SL mediates cell death through necrosis in *C. albicans*

Since SL enhanced the ROS production; we examined the downstream effectors of ROS for apoptosis and necrosis by Annexin/PI assay. Annexin binds the surface exposed






Fig. 2 Sophorolipid (SL) induces the depolarization of mitochondrial membrane potential (MMP) of *C. albicans*. Cells were grown in YPD medium for 5 h in the presence of indicated concentrations of SL and stained with mitochondria-specific voltage-dependent dye (DiOC<sub>6</sub>). Samples were analyzed by flow cytometry in FL1 channel.

phosphatidylserine (an apoptosis marker) and gives positive signals, whereas, PI only enters dead or damaged cells (necrosed) and is excluded by the viable cells. When cells were treated with 100  $\mu\text{g mL}^{-1}$  SL,  $\sim 22\%$  of cell population was PI-positive (upper left column, Fig. 4B), however, only a small number of cells were found to be annexin positive (upper right

column, Fig. 4B) against the control (Fig. 4A). The PI-positive cell population was further increased to 27% when cells were treated with 200  $\mu\text{g mL}^{-1}$  SL, whereas, almost negligible number of annexin positive cells were observed at this concentration (Fig. 4C). These results suggest that SL causes necrosis in *C. albicans* which attributes to cells death.




Fig. 3 Sophorolipid (SL) treatment induces ER stress and increases the intracellular calcium ion ( $\text{Ca}^{2+}$ ) level in *C. albicans*. (A) Indicated concentrations of SL was added to the cells directly or pre-incubated with 10 mM BAPTA, a  $\text{Ca}^{2+}$  chelator and grown in YPD. (B) Expression of *HAC1*, a marker of ER stress was determined by qRT-PCR. The result represents the average of three independent experiments  $\pm$ SD.  $*p < 0.05$  when compared with the SL untreated controls.



### SL causes membrane permeabilization in *C. albicans*

Permeabilization of the plasma membrane is a key signature of cell necrosis. This event can be quantified by measuring the release of intracellular enzyme lactate dehydrogenase (LDH) into the extracellular milieu. To further ensure the role of SL in necrosis, we performed LDH release assay. The released LDH can be quantified by a coupled enzymatic reaction. First, LDH catalyzes the conversion of lactate to pyruvate by reduction of  $\text{NAD}^+$  to NADH. Second, diaphorase uses NADH to reduce a tetrazolium salt (INT) into red formazan product. Therefore, the level of formazan formation is directly proportional to the amount of released LDH into the medium. When cells were incubated with  $50 \mu\text{g mL}^{-1}$  SL for 12 and 24 h, the percentage LDH activity was increased by 4.8-fold and 6.5-fold, respectively (Fig. 5), as compared to control. The percentage LDH activity was further increased by 9.7-fold and 13.7-fold as compared to control when cells were incubated with  $100 \mu\text{g mL}^{-1}$  SL for 12 and 24 h.

Permeabilization of the cell membrane due to SL treatment can be observed by staining the treated cells with PI followed by visualization under CLSM (Fig. 6A). Due to SL treatment cell membrane integrity is lost and PI enters into the cells through the permeabilized cell membrane. The results showed that cells treated  $100 \mu\text{g mL}^{-1}$  and  $200 \mu\text{g mL}^{-1}$  of SL had 62% and 72% permeabilized cell respectively, compared to the 9% PI positive cells in untreated (control) condition (Fig. 6B).

We also carried out TEM analysis to visualize the effect of SL on *C. albicans* plasma membrane. We found that SL disturbed the integrity of the plasma membrane (Fig. 6C) causing pore formation confirming its role in necrosis.

### Activation of HOG1 mitogen-activated protein kinase (MAPK) pathway in response to SL in *C. albicans*

The high glycerol osmolarity (HOG; a branch of Mitogen Activated Protein Kinase) pathway plays a key role in the adaptation to high osmolarity condition. HOG pathway is also known to play an important role in other stress pathways including




Fig. 5 Sophorolipid (SL) increased the release of lactate dehydrogenase (LDH) in *C. albicans*. Cells were grown in the presence of indicated concentrations of SL for 0, 12 and 24 h. Percentage of LDH activity is directly proportional to the percentage of substrate converted.

oxidative stress.<sup>37–39</sup> Therefore, we checked the effect of SL on osmotic stress and the related HOG1 pathway in *C. albicans* and null  $\Delta hog1$  mutant cells. We found that in the presence of SL ( $200 \mu\text{g mL}^{-1}$ ), the growth of *C. albicans* SC5314 was retarded compared to untreated cells (Fig. 7A). Moreover, the growth was rescued when the cells were complemented with 0.25 M and 0.5 M sorbitol in addition to SL. In addition, null  $\Delta hog1$  mutant of *C. albicans* was unable to rescue the growth in presence of both molar concentration of sorbitol suggesting the role of HOG1 in SL mediated killing pathway (Fig. 7B). When the higher concentration of SL ( $500 \mu\text{g mL}^{-1}$ ) was used along with 0.5 M sorbitol on *C. albicans* cells and spotted on YPD plates, it showed increased sensitivity compared to SL ( $500 \mu\text{g mL}^{-1}$ ) and sorbitol (0.5 M) alone (data not shown).

Thereafter, we were curious to know the effect of SL on the expression of HOG1 gene in *C. albicans* and estimated HOG1 expression by qRT-PCR. A significant increase of HOG1




Fig. 4 Sophorolipid (SL) induced necrosis in *C. albicans*. Cells were grown in YPD medium for 5 h in the presence of indicated concentrations of SL and stained with Annexin V-FITC and propidium iodide (PI) and analyzed by flow cytometry.






Fig. 6 Alteration of membrane integrity of *C. albicans* due to the treatment of SL. (A) Confocal laser scanning microscopy of *C. albicans* cells treated with different concentration of SL and stained with PI. (B) The population of PI-positive cells were visualized under CLSM after the treatment with various concentration SL followed by staining with PI. 10 different fields were observed and the PI-positive cells were counted against the normal cells (to obtain the percentage of PI-positive cellular population). (C) Effect of sophorolipid (SL) on *C. albicans* plasma membrane was visualized by Transmission Electron Microscopy (TEM).

expression indicated the involvement of *HOG1* gene to mitigate the SL mediated stress response (Fig. 7C). Furthermore, we also checked the expression level of *SAP9*. Sap9 is a member of Sap family proteases and bound to the fungal cell surface by a glycosylphosphatidylinositol (GPI) anchor motif and is required for cell wall integrity.<sup>40</sup> Impairment of *SAP9* gene alters the adhesion properties and is considered important for *C. albicans* virulence. Therefore, we proceeded to check the effect of SL on *SAP9* expression. Interestingly, we found that *SAP9* expression was increased 1.8 and 3.1 fold after exposure to SL with 100 µg mL<sup>-1</sup> and 200 µg mL<sup>-1</sup> respectively (Fig. 7D).

## Discussion

The inhibitory effect of SL on growth, biofilm formation and biofilm eradication of *C. albicans* has been reported in a previous study.<sup>19</sup> In the present study, we have shown that the treatment of *C. albicans* cells with SL increases the ROS production and up-regulates the expression of *SOD1* and *CAT1*, indicating higher level of oxidative stress and activation of stress responsive machineries. Generally, ROS is produced during normal physiological condition and is involved in various biological processes including regulation of proliferation, activation of gene expression and other cellular responses.<sup>41–43</sup> However, extensive studies suggest that many external stimuli enhance ROS generation within the cell and

excessive ROS level can induce cell death by necrosis or apoptosis.<sup>44–46</sup> Since mitochondria are considered as a major site of ROS generation, excess ROS causes damage to the mitochondrial components perturbing with its membrane integrity. In order to elucidate the change in the mitochondrial membrane, we checked the mitochondrial membrane potential (MMP) by using flow cytometry. Our result indicates that ROS generation after SL treatment is associated with mitochondrial dysfunction. To counteract with excessive ROS production, cells possess various ROS defence systems which include redox buffers and specific antioxidant enzymes such as glutathione peroxidase, superoxide dismutase and catalase. Overexpression of these enzymes is a common mechanism to mitigate oxidative stress. In the present case, overexpression of *SOD1* and *CAT1* indicated higher level of oxidative stress and activation of stress responsive pathways. In that juncture, it has to be mentioned that *SOD1* and *CAT1* mRNA has longer half-life and generally the overexpression of these genes corresponds with the cellular protein level.<sup>47</sup> However in the present study cellular protein level of *CAT1* and *SOD1* was not measured.

The integrity and function of ER are crucial for the production of membrane-bound and secreted proteins and thus, its function is under tight control. Alteration in redox-homeostatic conditions lead to the failure of this control mechanism and the cell undergoes ER stress. Since ER stress is well characterized by overexpression of *HAC1*, a transcription factor involved in






Fig. 7 Growth curve of *C. albicans* SC5314 and null mutant  $\Delta hog1$  strains and Quantitative RT-PCR analysis of *HOG1* and *Sap9* genes. (A and B) *C. albicans* SC5314 and null mutant  $\Delta hog1$  strains were grown in YPD at 30 °C in presence of sophorolipid (200  $\mu\text{g mL}^{-1}$ ) alone and/or combination with different molar ratios of sorbitol. OD<sub>600</sub> was measured after every 2 h in each case. Untreated cells were used as negative control. (C and D) The expression of *HOG1* and *SAP9* was determined by a qRT-PCR assay. The expression level of each gene is displayed after normalization with internal control housekeeping gene *ACT1*. Results represent the average of three independent experiments  $\pm$ SD. \* $p < 0.05$  when compared with the SL untreated controls.

unfolded protein response (UPR) and is considered as a marker for ER stress.<sup>48</sup> In our study, we have demonstrated that *HAC1* expression is increased in SL treated cells. Carvacrol, an antifungal compound is known to exert its antifungal activity by disrupting calcium homeostasis and increasing *HAC1* expression.<sup>49</sup> The results of this study strongly support our findings. ER stress affects  $\text{Ca}^{2+}$  homeostasis by triggering the efflux of  $\text{Ca}^{2+}$  from the ER lumen to the cytoplasm. As a consequence of that, enhance cytosolic  $\text{Ca}^{2+}$  further magnifies ROS generation by entering into the mitochondria. In our study, we also observed a surge in  $\text{Ca}^{2+}$  level in the cytoplasm after the treatment with SL. Furthermore, the elevated intracellular ROS induces oxidative damage to DNA, proteins, lipids and other cellular components such as plasma membrane. Induction of these processes is associated with cellular death either by apoptosis or necrosis. Apoptosis is a well-accepted subroutine of programmed cell death (PCD), whereas, earlier reports described necrosis as an 'accidental' form of cell death based on its presumed unregulated nature.<sup>33,35,50</sup> However, over the past decade extensive evidences have emerged suggesting the role of specific genes in controlling the onset and progression of necrosis.<sup>33,50,51</sup> As demonstrated earlier, an increased level of ROS within the cellular environment leads to lethal consequences such as oxidative stress, apoptosis and necrosis.<sup>32,52,53</sup> In our study, Annexin/PI dependent flow cytometry analysis demonstrates that SL effectively induces necrosis in *C. albicans*. Moreover, one of the hallmarks of necrosis is loss of cellular membrane integrity and leaching out the intracellular enzymes

including LDH. Since the leakage of LDH is considered to be an indicator of necrosis, we performed LDH assay in time and concentration dependent manner of SL; as a consequence, a significant increase in LDH release indicated the damage in cellular membrane integrity confirming the role of SL in necrosis. A similar result was observed by Basak *et al.*, by the action of acidic diacetate SL on the *Candida albicans* and *Salmonella enterica*.<sup>54</sup> In order to further confirm loss of the membrane integrity, Propidium Iodide (PI) staining assay was carried out by treating the cells with different concentrations of SL and then staining them with PI. The PI enters the cell only when it loses its membrane integrity. From the Fig. 6A it is evident that PI was taken up by 62% and 76% of SL (100 and 200  $\mu\text{g mL}^{-1}$ , respectively) treated cells indicating high-level membrane permeabilization and loss of membrane integrity. Furthermore, loss of membrane integrity was clearly visualised by TEM study.

To survive, *C. albicans* like other fungi are able to respond to changes in extracellular osmolarity and adjust their intracellular conditions to prevent itself from osmotic stress.<sup>55</sup> Such adaptation to high osmolarity involves activation of the HOG pathway which is a branch of MAPK (Mitogen Activated Protein Kinase) signal transduction system.<sup>56</sup> The basic physiological role of the HOG pathway is to orchestrate the adaptation of yeast cells against the surrounding medium.<sup>57</sup> In our study, we observed over-expression of *HOG1* in response to SL treatment indicating activation of HOG pathway. Moreover, we checked the effect of SL on osmotic stress and the related HOG1 pathway





in *C. albicans* and null  $\Delta$  *hog1* mutant cells. This study clearly suggests involvement of HOG1 pathway in response to SL mediated stress. As we observed high sensitivity in null  $\Delta$  *hog1* mutant cells as compared to *C. albicans* SC5314 cells. This sensitivity was restored after addition of sorbitol in null  $\Delta$  *hog1* mutant cells. The reason for rescue of growth by sorbitol could be due to the production of glycerol because sorbitol is used as a precursor molecule in the glycerol synthesis leading to cell survival under osmotic stress conditions.<sup>58</sup> Vylkova *et al.*, reported similar results against histatin 5 which shows increased sensitivity to 1 M sorbitol when grown under osmotic-stress conditions.<sup>59</sup> Furthermore, Sap9 is a cell-surface associated protease which helps in maintaining cell wall integrity in *C. albicans*. We hypothesised that Sap9 secretion may represent a defence mechanism of the cell in response to exposure to SL. Similarly, Copping *et al.*, found increased expression of Sap9 in *C. albicans* after fluconazole treatment. Moreover their group established that several antifungal agents can influence the expression of at least one *C. albicans* virulence factor *in vitro*.<sup>60</sup>

## Conclusion

In the present study, we have demonstrated that SL induces ROS formation in *C. albicans* and influence a cascade of stress response related activities in the cell. This process ultimately leads to the killing of the cells by membrane perforation and necrosis. Increased level of ROS formation, induction of ER stress and killing of *C. albicans* cells by necrosis or apoptosis is a common phenomenon that has been found to be activated due to the influence of many drugs.<sup>33</sup> However, the present study for the first time demonstrated the mechanism of cell death due to the treatment of glycolipid. Modification of these molecules can lead to the development of new therapies against fungal infection.

## Abbreviations

ROS	Reactive oxygen species
MMP	Mitochondrial membrane potential
HOG	High osmolarity glycerol
MAPK	Mitogen activated protein kinase

## Author contributions

F. H., M. A., S. B. and N. V. planned, executed the experiments and analysed the data. F. H. and N. V. M. S. B. discussed the data and co-wrote the manuscript.

## Funding

Funds for this project were obtained from CSIR project "OLP0082". F. H. would like to thanks DST-INSPIRE, M. A., S. B. and N. V. would like to thank CSIR, India and DBT, India for their fellowship.

## Conflicts of interest

Authors do not have any competing interest.

## Acknowledgements

Authors are thankful to DST and CSIR for funding and CSIR-IMTECH for providing technical facilities. Authors are extremely thankful to Dr Sneha Lata, JNU, New Delhi and Dr Jesus Pla, Madrid, Spain for providing the mutant strain. Authors are also thankful to Mr Randeep for TEM studies and Mrs Anjali and Mr Ella Bhagyaraj for FACS experiments. We are also thankful to Dr K. Ganesan for allowing us to carry out initial experiments in his lab and also for providing experimental support.

## References

- 1 M. A. Pfaller and D. J. Diekema, *Clin. Microbiol. Rev.*, 2007, **20**, 133–163.
- 2 J. C. O. Sardi, L. Scorzoni, T. Bernardi, A. M. Fusco-Almeida and M. J. S. Mendes Giannini, *J. Med. Microbiol.*, 2013, **62**, 10–24.
- 3 G. D. Brown, D. W. Denning, N. A. R. Gow, S. M. Levitz, M. G. Netea and T. C. White, *Sci. Transl. Med.*, 2012, **4**, 165rv13.
- 4 D. R. Andes, N. Safdar, J. W. Baddley, G. Playford, A. C. Reboli, J. H. Rex, J. D. Sobel, P. G. Pappas and B. J. Kullberg, *Clin. Infect. Dis.*, 2012, **54**, 1110–1122.
- 5 S. Biswas, P. Van Dijck and A. Datta, *Microbiol. Mol. Biol. Rev.*, 2007, **71**, 348–376.
- 6 P. E. Sudbery, *Nat. Rev. Microbiol.*, 2011, **9**, 737–748.
- 7 Z. A. Kanafani and J. R. Perfect, *Clin. Infect. Dis.*, 2008, **46**, 120–128.
- 8 T. Shekhar-Guturja, G. M. K. B. Gunaherath, E. M. K. Wijeratne, J. P. Lambert, A. F. Averette, S. C. Lee, T. Kim, Y. S. Bahn, F. Tripodi, R. Ammar, K. Döhl, K. Niewola-Staszewska, L. Schmitt, R. J. Loewith, F. Proth, D. Sanglard, D. Andes, C. Nislow, P. Coccetti, A. C. Gingras, J. Heitman, A. Gunatilaka and L. ECowen, *Nat. Chem. Biol.*, 2016, **12**, 867–875.
- 9 S. A. Davis, B. M. Vincent, M. M. Endo, L. Whitesell, K. Marchillo, D. R. Andes, S. Lindquist and M. D. Burke, *Nat. Chem. Biol.*, 2015, **11**, 481–487.
- 10 K. C. Gray, D. S. Palacios, I. Dailey, M. M. Endo, B. E. Uno, B. C. Wilcock and M. D. Burke, *Proc. Natl. Acad. Sci.*, 2012, **109**, 2234–2239.
- 11 M. D. Johnson, C. Macdougall, L. Ostrosky-zeichner, J. R. Perfect and J. H. Rex, *Antimicrob. Agents Chemother.*, 2004, **48**, 693–715.
- 12 A. P. Tulloch, J. F. T. Spencer and M. H. Deinema, *Can. J. Chem.*, 1968, **46**, 345–348.
- 13 C. P. Kurtzman, N. P. J. Price, K. J. Ray and T. M. Kuo, *FEMS Microbiol. Lett.*, 2010, **311**, 140–146.
- 14 M. A. Díaz De Rienzo, I. M. Banat, B. Dolman, J. Winterburn and P. J. Martin, *New Biotechnol.*, 2015, **32**, 720–726.



- 15 V. Shah, G. F. Doncel, T. Seyoum, K. M. Eaton, I. Zalenskaya, R. Hagver, A. Azim and R. Gross, *Antimicrob. Agents Chemother.*, 2005, **49**, 4093–4100.
- 16 S. L. Fu, S. R. Wallner, W. B. Bowne, M. D. Hagler, M. E. Zenilman, R. Gross and M. H. Bluth, *J. Surg. Res.*, 2008, **148**, 77–82.
- 17 L. Shao, X. Song, X. Ma, H. Li and Y. Qu, *J. Surg. Res.*, 2012, **173**, 286–291.
- 18 M. H. Bluth, E. Kandil, C. M. Mueller, V. Shah, Y. Y. Lin, H. Zhang, L. Dresner, L. Lempert, M. Nowakowski, R. Gross, R. Schulze and M. E. Zenilman, *Crit. Care Med.*, 2006, **34**, E188.
- 19 F. Haque, M. Alfatah, K. Ganesan and M. S. Bhattacharyya, *Sci. Rep.*, 2016, **6**, 23575.
- 20 K. Kim, D. Yoo, Y. Kim, B. Lee, D. Shin and E. K. Kim, *J. Microbiol. Biotechnol.*, 2002, **12**, 235–241.
- 21 K. Joshi-Navare and A. Prabhune, *BioMed Res. Int.*, 2013, 512495.
- 22 J. Chen, X. Song, H. Zhang, Y. B. Qu and J. Y. Miao, *Appl. Microbiol. Biotechnol.*, 2006, **72**, 52–59.
- 23 R. Alonso-Monge, F. Navarro-Garcia, E. Roman, A. I. Negredo, B. Eisman, C. Nombela and J. Pla, *Eukaryotic Cell*, 2003, **2**, 351–361.
- 24 J. L. Brewster, T. de Valoir, N. D. Dwyer, E. Winter and M. C. Gustin, *Science*, 1993, **259**, 1760–1763.
- 25 S. Sharma, M. Alfatah, V. K. Bari, Y. Rawal, S. Paul and K. Ganesan, *Antimicrob. Agents Chemother.*, 2014, **58**, 2409–2414.
- 26 F. Haque, M. Sajid, S. S. Cameotra and M. S. Battacharyya, *Biofouling*, 2017, **33**, 768–779.
- 27 M. Amin-ul Mannan, S. Sharma and K. Ganesan, *Anal. Biochem.*, 2009, **389**, 77–79.
- 28 K. J. Livak and T. D. Schmittgen, *Methods*, 2001, **25**, 402–408.
- 29 M. Alfatah, J. H. Wong, C. E. Nge, K. W. Kong, K. N. Low, C. Y. Leong, S. Crasta, M. Munusamy, A. M. L. Chang, S. Hoon, S. B. Ng, Y. Kanagasundaram and P. Arumugam, *Sci. Rep.*, 2019, **9**, 710.
- 30 S. Sangetha, Z. Zuraini, S. Suryani and S. Sasidharan, *Micron*, 2009, **40**, 439–443.
- 31 H. H. Lara, D. G. Romero-Urbina, C. Pierce, J. L. Lopez-Ribot, M. J. Arellano-Jiménez and M. Jose-Yacamán, *J. Nanobiotechnol.*, 2015, **13**, 91.
- 32 G. G. Perrone, S. X. Tan and I. W. Dawes, *Biochim. Biophys. Acta, Mol. Cell Res.*, 2008, **1783**, 1354–1368.
- 33 T. Eisenberg, D. Carmona-Gutierrez, S. Büttner, N. Tavernarakis and F. Madeo, *Apoptosis*, 2010, **15**, 257–268.
- 34 B. Hao, S. Cheng, C. J. Clancy and M. H. Nguyen, *Antimicrob. Agents Chemother.*, 2013, **57**, 326–332.
- 35 D. Carmona-Gutierrez, T. Eisenberg, S. Büttner, C. Meisinger, G. Kroemer and F. Madeo, *Cell Death Differ.*, 2010, **17**, 763–773.
- 36 J. D. Malhotra and R. J. Kaufman, *Cold Spring Harbor Perspect. Biol.*, 2011, **3**, a004424.
- 37 M. Hayashi and T. Maeda, *J. Biochem.*, 2006, **139**, 797–803.
- 38 M. Thorsen, Y. Di, C. Tängemo, M. Morillas, D. Ahmadpour, C. Van Der Does, A. Wagner, E. Johansson, J. Boman, F. Posas, R. Wysocki and M. J. Tamás, *Mol. Biol. Cell*, 2006, **17**, 4400–4410.
- 39 E. Bilsland, C. Molin, S. Swaminathan, A. Ramne and P. Sunnerhagen, *Mol. Microbiol.*, 2004, **53**, 1743–1756.
- 40 D. J. Krysan, E. L. Ting, C. Abeijon, L. Kroos and R. S. Fuller, *Eukaryotic Cell*, 2005, **4**, 1364–1374.
- 41 Y. Morel and R. Barouki, *Biochem. J.*, 1999, **342**(Pt 3), 481–496.
- 42 W. Dröge, *Physiol. Rev.*, 2002, **82**, 47–95.
- 43 H. Sauer, M. Wartenberg and J. Hescheler, *Cell. Physiol. Biochem.*, 2001, **11**, 173–186.
- 44 W. Lee and D. G. Lee, *Free Radical Res.*, 2018, **52**, 39–50.
- 45 M. Zhang, W. Chang, H. Shi, Y. Li, S. Zheng, W. Li and H. Lou, *FEMS Yeast Res.*, 2018, 1–10.
- 46 H. Choi and D. G. Lee, *Biochimie*, 2015, **115**, 108–115.
- 47 L. Sun, K. Liao, C. Hang and D. Wang, *PLoS One*, 2017, **12**(2), e0172228.
- 48 T. H. Lee, Y. H. Bae, M. D. Kim and J. H. Seo, *Process Biochem.*, 2012, **47**, 2300–2305.
- 49 J. Chaillot, F. Tebbji, A. Remmal, C. Boone, G. W. Brown, M. Bellaoui and A. Sellam, *Antimicrob. Agents Chemother.*, 2015, **59**, 4584–4592.
- 50 D. E. Christofferson and J. Yuan, *Curr. Opin. Cell Biol.*, 2010, **22**, 263–268.
- 51 N. Vanlangenakker, T. Vanden Berghe, D. V. Krysko, N. Festjens and P. Vandenabeele, *Curr. Mol. Med.*, 2008, **8**, 207–220.
- 52 A. J. Phillips, I. Sudbery and M. Ramsdale, *Proc. Natl. Acad. Sci. U. S. A.*, 2003, **100**, 14327–14332.
- 53 D. J. Jamieson, *Yeast*, 1998, **14**, 1511–1527.
- 54 G. Basak, D. Das and N. Das, *J. Microbiol. Biotechnol.*, 2014, **24**, 87–96.
- 55 A. Blomberg and L. Adler, *Adv. Microb. Physiol.*, 1992, **33**, 145–212.
- 56 S. Hohmann, *FEBS Lett.*, 2009, **583**, 4025–4029.
- 57 S. Hohmann, *Microbiol. Mol. Biol. Rev.*, 2002, **66**, 300–372.
- 58 B. Shen, S. Hohmann, R. G. Jensen and A. H. Bohnert, *Plant Physiol.*, 1999, **121**, 45–52.
- 59 S. Vylkova, W. S. Jang, W. Li, N. Nayyar and M. Edgerton, *Eukaryotic Cell*, 2007, **6**, 1876–1888.
- 60 V. M. S. Copping, C. J. Barelle, B. Hube, N. A. R. Gow, A. J. P. Brown and F. C. Odds, *J. Antimicrob. Chemother.*, 2005, **55**, 645–654.

