Chemical Science

EDGE ARTICLE

Check for updates

Cite this: Chem. Sci., 2019, 10, 508

o All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 25th June 2018 Accepted 15th October 2018

DOI: 10.1039/c8sc02788a

rsc.li/chemical-science

Stereoselective organocatalyzed glycosylations – thiouracil, thioureas and monothiophthalimide act as Brønsted acid catalysts at low loadings[†]

G. A. Bradshaw, ‡ A. C. Colgan, 🕑 ‡ N. P. Allen, I. Pongener, ២ M. B. Boland, Y. Ortin and E. M. McGarrigle D*

Thiouracil catalyzes stereoselective glycosylations with galactals in loadings as low as 0.1 mol%. It is proposed that in these glycosylations thiouracil, monothiophthalimide, and the previously reported catalyst, Schreiner's thiourea, do not operate via a double H-bonding mechanism but rather by Brønsted acid/base catalysis. In addition to the synthesis of 2-deoxyglycosides and glycoconjugates, we report the first organocatalytic synthesis of 1,1'-linked trehalose-type sugars.

Introduction

Thioureas have found many applications as organocatalysts and it has become generally accepted that they activate their substrates through double-hydrogen bonding.1 Prominent examples include hydrogen bonding to anions and carbonyl groups, thus creating more reactive dienophiles, electrophiles or nucleophiles (Fig. 1). Herein we present evidence that thioureas can activate molecules in at least one other way and describe how this opens the way for new organocatalyst designs, with specific applications in glycosylations detailed.²

In 2012 Galan, McGarrigle and co-workers reported³ that use of Schreiner's thiourea^{1f,4} 1 as a catalyst gave high α-selectivity in glycosylations with galactals to form 2-deoxydisaccharides (Scheme 1a).5 Based on experimental evidence, and calculations by Schreiner⁶ on the related THP-protection of alcohols (Scheme 1b), an 'oxyanion hole'-type activation was previously proposed whereby the catalyst activated the alcohol by forming two hydrogen bonds to the oxygen, making the alcohol proton more acidic and thus catalysing C-H bond formation to C-2 of the galactal (Scheme 1c). A re-examination of this mechanistic proposal was prompted by recent work on the THP-protection of alcohols by Varga, Pápai and co-workers suggesting a Brønsted acid mode of action (see below),² challenging the prevailing notion of thiourea catalysts acting as double H-bond donors. We now report the development of a new glycosylation catalyst that is >600-fold cheaper than 1 and is active at loadings as low as 0.1 mol% in the synthesis of disaccharides, glycoconjugates,

‡ G. A. B. and A. C. C. contributed equally.

and trehalose-type sugars. Our results support a Brønsted acid/ base mechanism for our new catalysts and for Schreiner's thiourea in these glycosylations.

Results and discussion

In contrast to the generally accepted mode of action for thiourea catalysts as H-bond donors, calculations by Varga, Pápai and coworkers favoured a Brønsted acid mode of action in the THP-



Fig. 1 Selected examples of proposed interactions in thiourea catalysis via double H-bonding: (a) enolate binding in Pápai's model for enolate addition to nitroalkenes using Takemoto's catalyst,^{1d,e} (b) proposed complex between Schreiner's catalyst and an N-acyloxazolidinone in Diels-Alder reaction;^{1f} (c) ketone binding in Jacobsen's ketone cyanosilylation;¹⁹ (d) cyanide binding in Jacobsen's enantioselective Strecker reaction.^{1/}



View Article Online

View Journal | View Issue

Centre for Synthesis & Chemical Biology, UCD School of Chemistry, University College Dublin, Belfield, Dublin 4, Ireland. E-mail: eoghan.mcgarrigle@ucd.ie

[†] Electronic supplementary information (ESI) available: Detailed experimental procedures, characterisation and copies of NMR spectra. See DOI: 10.1039/c8sc02788a



Scheme 1 (a) Galan and McGarrigle's thiourea-catalyzed glycosylation reaction;³ (b) Schreiner's THP-protection of alcohols;⁶ (c) original proposal with catalyst acting as a double hydrogen-bond donor to alcohol;³ (d) analogue of Pápai² proposal with catalyst acting as a Brønsted acid and accepting a H-bond from the alcohol.

protection of alcohols.² An analogous transition state for our glycosylation reaction is represented in Scheme 1d. In contrast to Scheme 1c, where the N–H bonds need to be in a parallel orientation so as to enable the archetypal double-H-bonding, the alternative proposed transition state (Scheme 1d) doesn't require this parallel orientation. If such a mode of action were operative in our glycosylations, alternative catalysts designs would be feasible.

To test this new hypothesis, we first examined whether 2-thiouracil 2 was capable of catalysing the reaction between selected galactals 37 and diacetone galactose 4 (Table 1). Indeed, using just 1 mol% of thiouracil we found that high yields of disaccharides could be obtained with excellent α-selectivity.7 As 2-thiouracil is >600 times cheaper than Schreiner's catalyst, this is a significant improvement on our previously reported system.8 In fact not all of the thiouracil dissolves at this loading but it wasn't convenient to weigh out smaller amounts of catalyst, as on a ca. 0.5 mmol scale only 0.5 mg catalyst was used; we estimated the amount of dissolved catalyst to be <0.25 mol% under these conditions (see ESI[†]). In fact on larger scale the catalyst loading could be lowered further (vide infra). Galactals 3a-c worked well, as they had with thiourea 1 as catalyst. Galactal 3d was sluggish but gave a good yield when the reaction was carried out in 2-methyl-THF which allowed a higher reaction temperature.9 Per-acetylated 3e was relatively unreactive, as had been observed with 1. Per-benzylated glucal

 Table 1
 Thiouracil-catalyzed glycosylations, exploring glycal scope^a



^{*a*} 1.2 equiv. of the glycal was used. ^{*b*} Yield of Isolated product. ^{*c*} Determined by ¹H NMR. ^{*d*} Solvent: 2-methyl-THF (reflux = 83 °C). ^{*e*} Isolated with 10–20% inseparable impurities due to Ferrier rearrangement (see ESI for details). ^{*f*} Isolated with *ca*. 10% hydration impurity (*vide infra*). ^{*g*} Ratio prior to chromatography in parentheses.

3f had been problematic previously, so the yield of up to 46% represented progress.³ We didn't pursue this further due to the formation of Ferrier-rearrangement products which were inseparable from the desired glycoside.¹⁰ Rhamnal **3g** also gave an inseparable mixture of Ferrier products and desired disaccharide (Entry 10). McGarrigle, Galan and co-workers previously reported a protecting group strategy for accessing α-linked disaccharides from glucal and rhamnal.^{10α} Fucal **3h** gave a yield of 52% of α-**5h** (entry 11) although in this case a small amount of β-**5h** was also produced. We propose that the greater reactivity of galactals might be due to electrostatic stabilisation of the oxacarbenium ion by the 4-substituent.¹¹ These results are consistent with a common mechanism being operative in thiouracil and thiourea-catalyzed glycosylations.

Table 2 shows the scope of the reaction with respect to the acceptor.¹² The reaction tolerates primary and secondary alcohols at any position on the ring (entries 1–4). In Schreiner's original report⁶ on thiourea-catalyzed THP-protection, it was noted that an amine-containing solid support was not compatible with the reaction. Later studies establishing the pK_a of **1** as 8.5 (DMSO) suggests that this might be due to deprotonation of the catalyst by amines.^{13*a*,*b*} Catalyst **2** has a $pK_a = 7.75$ (H₂O).^{13*c*} Thus we predicted that appropriate protection group choice might enable us to use amino-containing substrates. Indeed, Boc-protected amino acids **6e–g** were glycosylated in excellent yields (entries 5–7). Glycoconjugates **7e**



^{*a*} 1.2 equiv. of the galactal was used and 0.83 M acceptor in CH₂Cl₂ except where stated. ^{*b*} Yield of Isolated product. ^{*c*} Determined by ¹H NMR. ^{*d*} 0.35 M solution w.r.t. acceptor instead of 0.83 M. ^{*e*} 2 equiv. of galactal w.r.t. acceptor was used. ^{*f*} With 17% α,α -9a as an impurity. ^{*g*} Solvent: 2methyl-THF (reflux = 83 °C).

and **7f** thus give access to the 2-deoxy analogues of mucin-type motifs. Entry 7 shows that the reaction also tolerates phenols (although rare, glycosylated tyrosines have been found in nature).¹⁴ Finally, cholesterol also proved a viable substrate (entry 8), opening the way to glycoconjugate synthesis beyond amino acids.

1,1'-Linked sugars and glycosyl sulfonamides

It was noted that the presence of water in the reaction of 3a and 4 was detrimental to yields, and led to the formation of hydrated side-products. This was turned to advantage by replacing the alcohol with water, enabling the synthesis of 1,1'-linked trehalose-like dimer 9a (Scheme 2). Sugars of this type are present in the cell wall of mycobacteria including tuberculosis (TB).¹⁵ The dimer **9a** was obtained as a mixture of $\alpha, \alpha : \alpha, \beta$ 4.6 : 1. No β , β -dimer was detected. No reaction took place in the absence of catalyst, thus this dimerization is also catalyzed by thiouracil. The stereochemical outcome can be rationalized as follows: following addition of water to galactal 3a the resulting hemiacetal 8 rapidly epimerizes, but addition of 8 to a second galactal occurs with excellent *a*-selectivity. We suspect, but cannot prove, that the initial addition of water to the galactal is α -selective. The proposed pathway is supported by the stereoselective addition of hemiacetal 10 to galactal 3a giving a mixture of $\alpha, \alpha : \alpha, \beta \ 1 : 1$. The use of hemiacetals such as **10** thus demonstrates that heterodimeric trehalose-type structures can also be accessed through this methodology. The epimers were separable by chromatography and the disaccharides were deprotected by hydrogenation over Pd/C to give **9b** and **11b**. Thus the method gives easy access to symmetrical and unsymmetrical **1**,1'-linked sugars. Compound **9b** is an inhibitor of mycolyltransesterase enzymes Ag85A, Ag85B and Ag85C which serve as essential mediators of cell envelope function and biogenesis in *Mycobacterium tuberculosis*,^{15b} thus this method may enable access to compounds useful for studying/fighting TB. Further efforts in this regard are ongoing in our laboratory.

Glycosyl sulfonamides have been shown to inhibit growth of tumour cell lines. Reaction with TsNH₂ gave the glycosylated product **12** in 89% yield as a 3.6 : 1 mixture of β : α anomers (Scheme 3), showing that *N*-glycosylation is also possible.¹⁶ The stereochemical outcome may arise from epimerisation of α -**12** to the β -product, indeed the ratio we observe is similar to that reported by Toste.^{16*a*,17}

Exploration of mechanism

With regard to the mechanism, we propose that the catalyst is operating *via* an acid/base-type mechanism, rather than through a double-hydrogen bonding mechanism, as we



Scheme 2 Synthesis of trehalose-type 1,1'-linked dimers and proposed intermediate 8.



originally proposed. The reactions to form 2-deoxyglycosides, 1,1'-linked sugars and glycosylated sulfonamides all require catalyst - no reaction is observed in the absence of catalyst. When a mixture of α/β -5a was resubjected to the reaction conditions the α/β ratio remained constant (Scheme 4). Crossover tests also proved negative (see ESI[†] for details). Thus, the high α -selectivity observed is obtained kinetically, and is not the result of post-reaction anomerisation, except in the case of 1,1linked disaccharides where prolonged reaction times increased the amount of α, α -dimer 9a. The use of D₂O in place of H₂O in the formation of α, α -9a showed stereoselective (>95:5) synaddition of deuterium and oxygen and gives easy access to C2-deuterated $\alpha_1\alpha_2$ -9a. Glycosylation of CD₃OD resulted in a *cis* : *trans* ratio of 95 : 5 for both catalysts 1 and 2 (Scheme 4). This is more *cis*-selective than related reactions of dihydropyrans studied by Varga and Pápai,2a and Zimmerman and Nagorny,18 but goes against an asynchronous concerted mechanism.6 It might be proposed that 2 could be present in solution



Scheme 4 Mechanistic studies on glycosylation: α/β -5a doesn't epimerize under reaction conditions. 13 is a viable catalyst (84% yield of 5a). Reaction of D₂O with 3a gives addition of C–D and C–O bonds to same face of 3a. Schreiner's catalyst and thiouracil give the same outcomes in glycosylation of CD₃OD.

as the tautomer **2B**, and thus fulfil the role of a double Hbonding catalyst. To counter this, monothiophthalimide **13** was synthesized and shown to be a viable stereoselective catalyst at 1 mol% under our standard glycosylation reaction conditions giving 84% of α -**5a** (Scheme 4). At the very least, this demonstrates the existence of a catalytic pathway that does not require a double H-bonding catalyst, and given the calculations of Varga and Pápai, we contend that it is most likely that thiourea **1**, thiouracil **2**, and monothiophthalimide **13** all operate *via* a common mechanism (we are not aware of any other reports of monothioimides as organocatalysts).

Monothiophthalimide 13 has greater solubility in CH₂Cl₂ than thiouracil, thus enabling NMR studies. The chemical shift of the NH proton of 13 changed by 0.4 ppm when alcohol 4 (3.7 equiv.) was added to 13, consistent with complex formation (see ESI[†] for details). In the context of our current proposal we can rationalize these results as follows (Scheme 5): we propose that the catalyst first accepts a H-bond from the alcohol substrate making it a stronger acid. We propose a H-bond to sulfur as uracil does not catalyse the reaction, but several structures are conceivable.19 It is this catalyst-alcohol complex that interacts with the galactal and undergoes proton transfer. Secondly, the anion formed is able to deprotonate the alcohol in the second step of the reaction as C–O bond formation occurs. We note that we had previously discounted³ a 'simple' acid-catalyzed mechanism for this reaction as neither Meldrum's acid ($pK_a = 7.3$ (DMSO)) nor Et₃N·HCl ($pK_a = 9.0$ (DMSO)) catalyzed the reaction.20 It is clear that the chloride anion that would result from $Et_3N \cdot HCl$ would not be sufficiently basic for this role. We suspect that the same is true of the conjugate base of Meldrum's



Scheme 5 Proposed glycosylation mechanism.

acid if complexation with alcohol made it a strong enough acid for the first step. The α -stereoselectivity and *syn*-addition of H/D and O-R are consistent with the catalyst : ROH complex favouring approach from the bottom face and the C-O bond forming before the ion pair has time to rearrange.

Catalyst loading and gram-scale reactions

We conducted larger scale experiments in order to explore the catalyst loadings and demonstrate practicality (Scheme 6). Thiouracil 2 was active at loadings down to 0.01 mol% but for reproducible reaction times we chose 0.1 mol% and carried out a 1 g scale reaction.²¹ The concentration was found to be important on bigger scale (1.7 M w.r.t. acceptor; on smaller scale the initial concentration was 0.8 M but some solvent loss occurred during the experiment so the final concentration was higher – see ESI†). Under these optimized conditions just 0.25 mg of catalyst was required to obtain an isolated yield of 1 g (78%) of 5a.



Scheme 6 Lower catalyst loadings and gram-scale synthesis of 5a.

Conclusions

In summary, we have shown that thiouracil can replace Schreiner's thiourea in glycosylation reactions. Thus 2-thiouracil is a simple, cheap and effective catalyst for α -selective glycosylation with galactals of monosaccharides, steroids, and amino acids at loadings as low as 0.1 mol%. The method also enables the synthesis of symmetrical and unsymmetrical 1,1'-linked disaccharides including examples relevant to *M. tuberculosis*. Further work in this area is underway in our laboratories.

Furthermore, in line with Varga, Pápai and co-worker's publication on thiourea-catalyzed THP-protection, we are supporting a mechanism whereby the thiourea or thiouracil engages *via* general acid/base catalysis rather than acting as a double-hydrogen bond donor. When considering how a thiourea catalyst is activating substrates it is necessary to consider possibilities beyond double H-bonding. These findings should lead to new opportunities to catalyse reactions and to new organocatalyst designs beyond thioureas. We will report in due course on our own efforts in this regard.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

E. M. M. and G. A. B. thank Science Foundation Ireland & the Marie-Curie Action COFUND (11/SIRG/B2154). A. C. C. thanks the Irish Research Council (GOIPG/2013/301), N. P. A. thanks UCD for a Research Demonstratorship. E. M. M. and I. P. thank Science Foundation Ireland (15/CDA/3625). We thank members of EU COST action CM0905 (organocatalysis) and, in particular, Imre Pápai for helpful discussions. We are grateful to Prof Declan Gilheany (UCD) and Dr Kirill Nitikin (UCD) for helpful discussions also.

Notes and references

- 1 (a) G. Jakab and P. R. Schreiner in Comprehensive Enantioselective Organocatalysis, ed. P. Dalko, Wiley-VCH, Weinheim, 2013, ch. 12, vol. 2; (b) K. Hof, K. M. Lippert and P. R. Schreiner in Asymmetric Organocatalysis, ed. K. Maruoka), Thieme, Stuttgart, 2012, ch. 2.2.4, vol. 2; (c) A. G. Doyle and E. N. Jacobsen, Chem. Rev., 2007, 107, 5713; (d) A. Hamza, G. Schubert, T. Soós and I. Pápai, J. Am. Chem. Soc., 2006, 128, 13151; (e) T. Okino, Y. Hoashi and Y. Takemoto, J. Am. Chem. Soc., 2003, 125, 12672; (f) P. R. Schreiner and A. Wittkopp, Org. Lett., 2002, 4, 217; (g) S. J. Zuend and E. N. Jacobsen, J. Am. Chem. Soc., 2007, 129, 15872; (h) S. J. Zuend and E. N. Jacobsen, J. Am. Chem. Soc., 2009, 131, 15358.
- 2 (a) Á. Madarász, Z. Dósa, S. Varga, T. Soós, A. Csámpai and I. Pápai, ACS Catal., 2016, 6, 4379; (b) Á. Madarász, Z. Dósa, S. Varga, T. Soós, A. Csámpai, and I. Pápai, oral presentation, EU Cost Action CM0905 Meeting, Palermo,

May 2014. A thiourea catalyst with one NH was recently described: (c) P. Jovanovic, M. Petkovic, M. Simic, B. Ivkovic and V. Savic, Org. Biomol. Chem., 2016, 14, 6712. See also (d) A. Supady, S. Hecht and C. Baldau, Org. Lett., 2017, 19, 4199. For proposed action of thioureas as Lewis bases see: (e) C. K. Tan, F. Chen and Y.-Y. Yeung, Tetrahedron Lett., 2011, 52, 4892; (f) C. B. Tripathi and S. Mukherjee, J. Org. Chem., 2012, 77, 1592; (g) S. E. Denmark and M. T. Burk, Proc. Natl. Acad. Sci. U. S. A., 2010, 107, 20655; (h) Y. Kobayashi, Y. Nakatsuji, S. Li, S. Tsukuki and Y. Takemoto, Angew. Chem., Int. Ed., 2018, 57, 3646.

- 3 E. I. Balmond, D. M. Coe, M. C. Galan and E. M. McGarrigle, *Angew. Chem., Int. Ed.*, 2012, **51**, 9152.
- 4 (a) E. Kuehle, L. Eue and O. Bayer, BE 613427, 1962, CA59:8698; (b) E. I. Balmond, M. C. Galan, and E. M. McGarrigle, N,N'-Bis[3,5-bis(trifluoromethyl)phenyl]-thiourea, in *Encyclopedia of Reagents for Organic Synthesis, online*, ed. L. A. Paquette, D. Crich, P. L. Fuchs, and G. A. Molander, John Wiley & Sons Ltd., 2014, DOI: 10.1002/047084289X.rn01642; (c) Z. Zhang, Z. Bao and H. Xing, Org. Biomol. Chem., 2014, 12, 3151.
- 5 For leading references on organocatalytic glycosylations see: (a) D. J. Cox, M. D. Smith and A. J. Fairbanks, Org. Lett., 2010, 12, 1452; (b) E. I. Balmond, M. C. Galan and E. M. McGarrigle, Synlett, 2013, 24, 2335; (c) Y. Q. Geng, A. Kumar, H. M. Faidallah, H. A. Albar, I. A. Mhkalid and R. R. Schmidt, Angew. Chem., Int. Ed., 2013, 52, 10089; (d) S. Das, D. Pekel, J. M. Neudorfl and A. Berkessel, Angew. Chem., Int. Ed., 2015, 54, 12479; (e) S. Schmalisch and R. Mahrwald, Org. Lett., 2013, 15, 5854; (f) L. Sun, X. Wu, D.-C. Xiong and X.-S. Ye, Angew. Chem., Int. Ed., 2016, 55, 8041; (g) M. Tanaka, J. Nashida, D. Takahashi and K. Toshima, Org. Lett., 2016, 18, 2288; (h) C. Gouliaras, D. Lee, L. Chan and M. S. Taylor, J. Am. Chem. Soc., 2011, 133, 13926; (i) S. Medina, M. Harper, E. I. Balmond, S. Miranda, G. Crisenza, D. M. Coe, E. M. McGarrigle and M. C. Galan, Org. Lett., 2016, 18, 4222; (j) Y. Park, K. C. Harper, N. Kuhl, E. E. Kwan, R. Y. Liu and E. N. Jacobsen, Science, 2017, 355, 162; (k) R. Williams and M. C. Galan, Eur. J. Org. Chem., 2017, 6247; (l) J.-L. Liu, Y.-T. Zhang, H.-F. Liu, L. Zhou and J. Chen, Org. Lett., 2017, 19, 5272; (m) J. Lee, A. Borovika, Y. Khomutnyk and P. Nagorny, Chem. Commun., 2017, 53, 8976; (n) C. Palo-Nieto, A. Sau, R. Williams and M. C. Galan, J. Org. Chem., 2017, 82, 407; (o) T. Kimura, T. Eto, D. Takahashi and K. Toshima, Org. Lett., 2016, 18, 3190; (p) M. Shaw, Y. Kumar, R. Thakur and A. Kumar, Beilstein J. Org. Chem., 2017, 13, 2385; (q) Y. Singh, T. Wang, S. A. Geringer, K. J. Stine and A. V. Demchenko, J. Org. Chem., 2018, 83, 374; (r) J.-H. Tay, A. J. Argüelles, M. D. DeMars, P. M. Zimmerman, D. H. Sherman and P. Nagorny, J. Am. Chem. Soc., 2017, 139, 8570.
- 6 M. Kotke and P. R. Schreiner, Synthesis, 2007, 779.
- 7 (a) E. Durantie, C. Bucher and R. Gilmour, *Chem.-Eur. J.*, 2012, 18, 8208; (b) We note that care must be taken in the synthesis of donor 3a to avoid an impurity that can act as

a catalyst poison: A. C. Colgan, H. Müller-Bunz and E. M. McGarrigle, *J. Org. Chem.*, 2016, **81**, 11394.

- 8 Price/mol on www.sigmaaldrich.com and www.tcichemicals.com, accessed 22 June 2018.
- 9 CH₂Cl₂ was identified as the best solvent from a solvent screen but the reaction also proceeded in toluene, acetonitrile, THF and 2-MeTHF. 2-Methyl-THF is suggested to be more sustainable (bio-based), and less toxic than CH₂Cl₂: (a) D. F. Aycock, Org. Process Res. Dev., 2007, 11, 156; (b) V. Antonucci, J. Coleman, J. B. Ferry, N. Johnson, M. Mathe, J. P. Scott and J. Xu, Org. Process Res. Dev., 2011, 15, 939; (c) C. M. Alder, J. D. Hayler, R. K. Henderson, A. M. Redman, L. Shukla, L. E. Shuster and H. F. Sneddon, Green Chem., 2016, 18, 3879; (d) V. Pace, P. Hoyos, L. Castoldi, P. Domínguez de María and A. R. Alcántara, ChemSusChem, 2012, 5, 1369.
- 10 Other methods for accessing 2-deoxyglycosides see ref. 5d, j and n and: (a) E. I. Balmond, D. M. Coe, R. W. Alder, E. M. McGarrigle and M. C. Galan, Angew. Chem., Int. Ed., 2014, 53, 8190; (b) J. M. Nogueira, M. Bylsma, D. K. Bright and C. S. Bennett, Angew. Chem., Int. Ed., 2016, 55, 10088; (c) D. Zhu, K. N. Baryal, S. Adhikari and J. Zhu, J. Am. Chem. Soc., 2014, 136, 3172; (d) A. Borovika and P. Nagorny, J. Carbohydr. Chem., 2012, 31, 255; (e) H. Wang, J. Tao, X. Cai, W. Chen, Y. Zhao, Y. Xu, W. Yao, J. Zeng and Q. Wan, Chem.-Eur. J., 2014, 17319; (f) V. P. Verma and C.-C. Wang, Chem.-Eur. J., 2013, 19, 846; (g) G. Zhao and T. Wang, Angew. Chem., Int. Ed., 2018, 57, 6120; (h) T. M. Beale, P. J. Moon and M. S. Taylor, Org. Lett., 2014, 16, 3604; (i) C. H. Marzabadi and R. W. Franck, Tetrahedron, 2000, 56, 8385; (j) D. Hou and T. L. Lowary, Carbohydr. Res., 2009, 344, 1911; (k) C. S. Bennett and M. C. Galan, Chem. Rev., 2018, 118, 7931; (l) D.-M. Yang, Y. Chen, R. P. Sweeney, T. L. Lowary and X.-Y. Liang, Org. Lett., 2018, 20, 2287.
- 11 (a) H. H. Jensen and M. Bols, Acc. Chem. Res., 2006, 39, 259;
 (b) A. Garcia, J. R. Sanzone and K. A. Woerpel, Angew. Chem., Int. Ed., 2015, 127, 12255;
 (c) A. Garcia, D. A. L. Otte, W. A. Salamant, J. R. Sanzone and K. A. Woerpel, Angew. Chem., Int. Ed., 2015, 54, 3061.
- 12 (a) A. Roën, J. I. Padrón and J. T. Vázquez, J. Org. Chem., 2003, 68, 4615; (b) M. Matwiejuk and J. Thiem, Chem. Commun., 2011, 47, 8379; (c) P. J. Garegg, T. Iversen and S. Oscarson, Carbohydr. Res., 1976, 50, C12–C14; (d) L. K. Mydock, M. N. Kamat and A. V. Demchenko, Org. Lett., 2011, 13, 2928; (e) T. Ogawa and T. Kaburagi, Carbohydr. Res., 1982, 103, 53; (f) C.-C. Wang, S. S. Kulkarni, J.-C. Lee, S.-Y. Luo and S.-C. Hung, Nat. Protoc., 2008, 3, 97; (g) P. J. Garegg, H. Hultberg and S. Wallin, Carbohydr. Res., 1982, 108, 97; (h) Z. Zhang, B. Yua and R. R. Schmidt, Synthesis, 2006, 8, 1301; (i) K. Bao, A. Fan, Y. Dai, L. Zhang, W. Zhang, M. Cheng and X. Yao, Org. Biomol. Chem., 2009, 7, 5084.
- 13 (a) X. Li, H. Deng, S. Luo and J. P. Cheng, *Eur. J. Org. Chem.*, 2008, 4350; (b) G. Jakab, C. Tancon, Z. Zhang, K. M. Lippert and P. R. Schreiner, *Org. Lett.*, 2012, 14, 1724; (c) H. N. Christensen, *J. Biol. Chem.*, 1945, 160, 425.

- 14 (a) J. Peters, S. Rudolf, H. Oschkinat, R. Mengele, M. Sumper, J. Kellermann, F. Lottspeich and W. Baumeister, Biol. Chem. Hoppe-Seyler, 1992, 373, 171; (b) A. Halim, G. Brinkmalm, U. Ruetschi, A. Westman-Brinkmalm, E. Portelius, H. Zetterberg, K. Blennow, G. Larson and J. Nilsson, Proc. Natl. Acad. Sci. U. S. A., 2011, 108, 11848; (c) C. Steentoft, S. Y Vakhrushev, M. B. Vester-Christensen, K. T.-B. G. Schjoldager, Y. Kong, E. P. Bennett, U. Mandel, H. Wandall, S. B Levery and H. Clausen, Nat. Methods, 2011, 8, 977; (d) P. O. Larsen, E. Pedersen, H. Sorensen and P. Sorup, Phytochemistry, 1973, 12, 2243; (e) G. Yang, Q. Wang, X. Luo, J. Zhang and J. Tang, Glycoconjugate J., 2012, 29, 453.
- 15 (a) K. J. Welsh, R. L. Hunter and J. K. Actor, Tuberculosis, 2013, 93, S3-S9; (b) K. M. Backus, H. l. Boshoff, C. S. Barry, O. Boutureira, M. K. Patel, F. D'Hooge, S. S. Lee, L. E. Via, K. Tahlan, C. E. Barry III and B. G. Davis, Nat. Chem. Biol., 2011, 7, 228; (c) H. N. Foley, J. A. Stewart, H. W. Kavunja, S. R. Rundell and B. M. Swarts, Angew. Chem., Int. Ed., 2016, 55, 2053; (d) M. Kamariza, P. Shieh, C. S. Ealand, J. S. Peters, B. Chu, F. P. Rodriguez-Rivera, M. R. Babu Sait, W. V. Treuren, N. Martinson, R. Kalscheuer, B. D. Kana and C. R. Bertozzi, Sci. Transl. Med., 2018, 10, eaam6310.
- 16 (a) B. D. Sherry, R. N. Loy and F. D. Toste, J. Am. Chem. Soc., 2004, 126, 4510; (b) P. A. Colinas and R. D. Bravo, Org. Lett., 2003, 5, 4509.
- 17 We thank a referee for this suggestion. A sample of 12 (α : β 1:9) did not show significant epimerisation under our reaction conditions but we were unable to obtain an aenriched sample to carry out a similar test on, and to rule out a decomposition pathway for α -12 that would lead to β enriched product.
- 18 Y. Y. Khomutnyk, A. J. Argüelles, G. A. Winschel, Z. Sun, P. Zimmerman and P. Nagorny, J. Am. Chem. Soc., 2016, 138, 444.

19 H-bonds from NH to the alkoxy fragment may also be present. No cross-peak for the NH was found in a HMBC NMR spectrum. We also considered whether a thioglycoside intermediate could arise from the reaction of catalyst with galactal but NMR studies failed to detect such a species even with excess 13 over 3a (see ESI[†] for details).



- 20 (a) I. M. Kolthoff, M. K. Chantoon and S. Bhowmik, J. Am. Chem. Soc., 1968, 90, 23; (b) E. M. Arnett, S. G. Maroldo, S. L. Schilling and J. A. Harrelson, J. Am. Chem. Soc., 1984, 106, 6759.
- 21 For other examples of low loading organocatalysts see: ref. 6 and: (a) F. Giacalone, M. Gruttadauria, P. Agrigento and R. Noto, Chem. Soc. Rev., 2012, 41, 2406; (b) D. Stevanovicć, G. Bertuzzi, A. Mazzanti, M. Fochi and L. Bernardi, Adv. Synth. Catal., 2018, 360, 893; (c) X. Wu, L. Xue, D. Li, S. Jia, J. Ao, J. Deng and H. Yan, Angew. Chem., Int. Ed., 2017, 56, 13722; (d) D. Bécart, V. Diemer, A. Salaün, M. Oiarbide, Y. R. Nelli, B. Kauffmann, L. Fischer, C. Palomo and G. Guichard, J. Am. Chem. Soc., 2017, 139, 12524; (e) M. Biedrzycki, A. Kasztelan and P. Kwiatkowski, ChemCatChem, 2017, 9, 2453; (f) X. Companyó and J. Bures, J. Am. Chem. Soc., 2017, 139, 8432; (g) J.-L. Li, L. Fu, J. Wu, K.-C. Yang, Q.-Z. Li, X.-J. Gou, C. Peng, B. Han and X.-D. Shen, Chem. Commun., 2017, 53, 6875; (h) M. Yanagi, A. Imayoshi, Y. Ueda, T. Furuta and T. Kawabata, Org. Lett., 2017, 19, 3099.