

Cite this: *Chem. Sci.*, 2019, 10, 2821

All publication charges for this article have been paid for by the Royal Society of Chemistry

# Investigations on *non-classical* silylium ions leading to a cyclobutenyl cation†

Arthur Martens,<sup>1</sup> Marvin Kreuzer, Alexander Ripp, Marius Schneider, Daniel Himmel,<sup>1</sup> Harald Scherer and Ingo Krossing<sup>1\*</sup>

Instead of yielding the desired *non-classical* silylium ions, the reactions of different alkenes/alkynes with several  $[\text{Me}_3\text{Si}]^+$  sources mostly led to oligomerization, or – in the presence of  $\text{Me}_3\text{SiH}$  – hydrosilylation of the alkenes/alkynes. Yet, from the reaction of 2-butyne with *ion-like*  $\text{Me}_3\text{Si}-\text{F}-\text{Al}(\text{OR}^{\text{F}})_3$  ( $\text{R}^{\text{F}} = \text{C}(\text{CF}_3)_3$ ) the salt of the silylated tetramethyl cyclobutenyl cation  $[\text{Me}_4\text{C}_4-\text{SiMe}_3]^+[\text{al}-\text{f}-\text{al}]^-$  ( $[\text{al}-\text{f}-\text{al}]^- = [(\text{R}^{\text{F}}\text{O})_3\text{Al}-\text{F}-\text{Al}(\text{OR}^{\text{F}})_3]^-$ ) was obtained in good yield (NMR, scXRD, Raman, and IR). All the experimental and calculated evidence suggest a mechanism in which **1** was formed *via* a *non-classical* silylium ion as an intermediate. The removal of the  $[\text{Me}_3\text{Si}]^+$  moiety from the cation in **1** was investigated as a means to provide free tetramethyl cyclobutadiene (CBD). However, the addition of  $[\text{NMe}_4]\text{F}$ , in order to release  $\text{Me}_3\text{SiF}$  and form CBD, led to the unexpected deprotonation of the cation. The addition of 4-dimethylaminopyridine to remove the  $[\text{Me}_3\text{Si}]^+$  cation as a Lewis acid/base adduct, led to an adduct with the four-membered ring in the direct neighborhood of the  $\text{Me}_3\text{Si}$  group. By the addition of  $\text{Et}_2\text{O}$  to a solution of **1**, the  $[\text{F}-\text{Al}(\text{OR}^{\text{F}})_3]^-$  anion (and  $\text{Et}_2\text{O}-\text{Al}(\text{OR}^{\text{F}})_3$ ) was generated from the  $[\text{al}-\text{f}-\text{al}]^-$  counterion. Subsequently, the  $[\text{F}-\text{Al}(\text{OR}^{\text{F}})_3]^-$  anion abstracted the  $[\text{Me}_3\text{Si}]^+$  moiety from  $[\text{Me}_4\text{C}_4-\text{SiMe}_3]^+$ , probably releasing CBD. However, due to the immediate reaction of CBD with  $[\text{Me}_4\text{C}_4-\text{SiMe}_3]^+$  and subsequent oligomerization, it was not possible to use CBD in follow-up chemistry.

Received 15th October 2018

Accepted 9th January 2019

DOI: 10.1039/c8sc04591g

rsc.li/chemical-science

## Introduction

Carbocations play an important role as intermediates, *e.g.* in  $\text{S}_{\text{N}}1$  reactions or in Wagner–Meerwein rearrangements.<sup>1</sup> The *classical* carbenium ions, *e.g.*  $[\text{C}(\text{CH}_3)_3]^+$ , feature a tricoordinate electron deficient carbon atom with 2e–2c bonds.<sup>2</sup> While their existence has long been accepted, there has been a long dispute about the existence of the (pentacoordinate) *non-classical* carbonium ions.<sup>3</sup> Only in 2013 the final crystal structure evidence for the existence of *non-classical* carbocations was provided by our group.<sup>4</sup> Motivated by this result, we were interested to see if analogous *non-classical* silylium ions also exist in the condensed phase, *i.e.* silylium ions stabilized by a 3c–2e bond (Scheme 1a and b). Although adducts of  $[\text{Me}_3\text{Si}]^+$  with ethylene and acetylene were detected in the gas phase by mass spectrometry, their structures in condensed phases hitherto remain unknown.<sup>5–7</sup> Yet, quantum-chemical calculations suggest a *non-classical* structure for adducts between  $[\text{Me}_3\text{Si}]^+$  and  $\text{C}_2\text{H}_4/\text{C}_2\text{H}_2$ .


However, it appears that the latter are only intermediates and react further by a methyl shift and formation of a vinyl silylium ion (Scheme 1c).<sup>6,7</sup> By contrast, with substituted alkenes, *classical* carbenium ions as in Scheme 1b were calculated to be favored.<sup>6</sup> Adducts of silylium ions with benzene or toluene could also be considered to possess *non-classical* structures, but crystallographic studies and also quantum-chemical calculations suggest a *classical* structure.<sup>8,9</sup>

The all-silicon analogue to the *non-classical* carbonium ions would be a silylium ion, which is coordinated by a disilene  $\text{R}_2\text{Si}=\text{SiR}_2$  or by a (formal) disilyne  $\text{RSi}\equiv\text{SiR}$ . To the best of our knowledge, for the  $[\text{Si}_3\text{R}_5]^+$  cations no experimental data exist and computational analyses are limited to thermodynamics and do not discuss structural properties.<sup>12</sup> Reactions of silylium ions with di- and trisilenes yielded cyclotetrasilylium ions  $[(\text{RSi})_3\text{SiR}_2]^+$  as part of more complicated rearrangement reactions (Scheme 1d;  $\text{R} = {}^t\text{Bu}_2\text{MeSi}$ ,  ${}^t\text{Bu}$ ).<sup>10,11</sup> In any event, the stabilization of disilenes and disilynes against oligomerization requires large substituents R, which hinder their – *classical* or *non-classical* – coordination to a silylium ion.<sup>10,11,13</sup> Although calculations at the MP2/def2-TZVPP level suggest a *non-classical* adduct between  $[\text{Me}_3\text{Si}]^+$  and the sterically unhindered  $\text{Me}_2\text{Si}=\text{SiMe}_2$ , this disilene would not be isolable due to the discussed oligomerization. Therefore, we set out to synthesize *non-classical* silylium ions by reaction of sources of the small  $[\text{Me}_3\text{Si}]^+$  silylium ion with alkenes or alkynes. With small substituents,

Institut für Anorganische und Analytische Chemie, Freiburger Materialforschungszentrum (FME), Universität Freiburg, Albertstr. 21, 79104 Freiburg, Germany. E-mail: krossing@uni-freiburg.de

† Electronic supplementary information (ESI) available: Experimental details, procedures, weights, and 1D- and 2D-NMR spectra of the reactions are displayed. Details of the quantum chemical calculations are given together with crystallographic details. CCDC 1868136–1868138. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c8sc04591g





**Scheme 1** Classical and non-classical structures of (a) the 2-norbornyl cation and (b) analogous silylium ions (although we refer to “classical silylium ions” here, these are rather to be seen as carbenium ions stabilized by the β-Si effect); (c) the calculated methyl shift for the [Me<sub>3</sub>-Si(C<sub>2</sub>H<sub>2</sub>)<sup>+</sup>] cation;<sup>7</sup> (d) the reaction products of silylium ions with di- and trisilenes.<sup>10,11</sup>



their π bond is more accessible than that in sterically hindered room temperature stable disilenes and disilynes. Additionally, alkenes and alkynes would allow for a homogeneous delocalization of the positive charge among the carbon atoms, which would require more reorganization for the disilenes/disilynes due to their trans-bent structure. However, the fluoride ion affinity (FIA), as a measure of Lewis acidity,<sup>14</sup> of silylium ions is significantly higher than that of carbenium ions (FIA = 952 vs. 836 kJ mol<sup>-1</sup> for [Me<sub>3</sub>E]<sup>+</sup>; E = Si, C; calculated like in ref. 15; BP86/def-SV(P)). As a result, the addition of an alkene or alkyne to a silylium ion may also result in the formation of a carbenium ion that is stabilized in the classical structure by the so-called β-Si effect. This implies a hyperconjugative stabilization due to electron density transfer from the occupied σ(Si-C) orbital into the empty p-orbital of the cationic C atom (Scheme 1b).

## Results and discussion

Before turning to the experiments, we investigated the principle feasibility of the planned reactions by assessing the cations sought for with DFT and *ab initio* calculations and including solvation energies for the polar and weakly basic solvent *ortho*-difluorobenzene (*o*-DFB, ε<sub>r</sub> = 13.4).<sup>16</sup>

### Preliminary assessment of the stabilities of non-classical silylium ions








Calculations on the reaction of the free [Me<sub>3</sub>Si]<sup>+</sup> cation with different alkenes and alkynes were performed at the MP2/def2-TZVPP level of theory according to eqn (1) and (2) (Table 1).



For comparison, the structure of the (classical) benzene complex [Me<sub>3</sub>Si(C<sub>6</sub>H<sub>6</sub>)<sup>+</sup>] is also included in Table 1. In order to evaluate whether the calculated molecules are non-classical silylium ions or classical carbenium ions, we analyzed both relevant C-C-Si bond angles. For a classical carbenium ion, one of these angles is expected to be larger than at least 90°, while for non-classical silylium ions both angles should be (almost) equal and smaller than 90°. The complex [Me<sub>3</sub>Si(Ph<sub>2</sub>C=CPh<sub>2</sub>)<sup>+</sup>] was calculated to be a classical carbenium ion stabilized by the β-Si effect with a C-C-Si angle of 93.8°, possibly due to the resonance of the phenyl moieties and also for steric reasons. The C-C-Si angles in the H<sub>2</sub>C=CH<sub>2</sub> (73.4°), Me<sub>2</sub>C=CMe<sub>2</sub> (72.9°), MeC≡CMe (74.3°) and PhC≡CPh (74.3°) complexes combined with the symmetric C-Si distances suggest a non-classical structure for these cations. Regarding its structure, the [Me<sub>3</sub>Si(HC≡CH)]<sup>+</sup> cation is a special case with asymmetric C-C-Si angles of 78.3° and 71.6°, most likely induced by steric repulsion of the H-atom with one methyl group. Yet, we will refer to its structure as being non-classical. It should be noted that the non-classical structure of this cation with symmetric C-C-Si angles is disfavored by only Δ*G*<sub>gas</sub><sup>0</sup> = 0.01 kJ mol<sup>-1</sup> and therefore these two structures would be expected to be indistinguishable. The structures calculated at the simpler BP86-D3(BJ)/def-TZVP level of theory are similar, except for [Me<sub>3</sub>-Si(PhC≡CPh)]<sup>+</sup>. Here, the PhC≡CPh adduct is calculated to be a classical carbenium ion. It was not possible to calculate similar classical structures for the other adducts, as these are not even local minima or transition states on the respective energy hypersurface. This was exemplarily verified by calculating the energy of [Me<sub>3</sub>Si(MeC≡CMe)]<sup>+</sup> dependent on the C-C-Si angle in the range of 60 to 140° (see the ESI† for details). Therefore, classical starting structures also collapse to the non-classical structures. Thus no clear energy difference between a formally classical and a non-classical structure can be given.



**Table 1** Calculated Gibbs reaction energies  $\Delta G^0$  in  $\text{kJ mol}^{-1}$  and structures of the reaction  $[\text{Me}_3\text{Si}]^+ + \text{L} \rightarrow [\text{Me}_3\text{Si}(\text{L})]^+$  calculated at the MP2/def2-TZVPP level of theory with thermal contributions from BP86-D3(BJ)/def-TZVP calculations. The Gibbs solvation energy in *o*-DFB was calculated using the COSMO model ( $\epsilon_r = 13.4$ )<sup>16</sup> at the BP86-D3(BJ)/def-TZVP level. Scheme: H (light gray), C (dark gray), Si (blue)

Ligand L	$\Delta_r H_{\text{gas}}^0 / \Delta_r G_{\text{gas}}^0$ ( $\Delta G_{o\text{-DFB}}^0$ )	C–C–Si <sup>o</sup> $d_{\text{SiC}}/\text{pm}$	Gas phase structure
H <sub>2</sub> C=CH <sub>2</sub>	−109/−60 (−38)	73.4 236.7	
Me <sub>2</sub> C=CMe <sub>2</sub>	−174/−113 (−56)	72.9 235.6	
Ph <sub>2</sub> C=CPh <sub>2</sub>	−143/−67 (+20)	93.8/54.6 221.6/271.4	
HC≡CH	−98/−53 (−28)	78.3/71.6 231.0/238.4	
MeC≡CMe	−157/−103 (−58)	74.3 228.3	
PhC≡CPh	−188/−132 (−52)	74.3 229.5	
C <sub>6</sub> H <sub>6</sub> <sup>a</sup>	−134/−86 (−39)	98.3/51.3 <sup>b</sup> 217.3/275.8 <sup>b</sup>	


<sup>a</sup> Experimental data from ref. 8. <sup>b</sup>  $C_{\text{ipso}}$  and  $C_{\text{ortho}}$  were used for the measurements.

The gas phase reaction enthalpies  $\Delta_r H_{\text{gas}}^0$  and Gibbs energies  $\Delta_r G_{\text{gas}}^0$  with alkynes become more favorable when replacing the H substituents of the parent alkyne HC≡CH with Me, and even more so with Ph (MP2/def2-TZVPP). This can be explained by an increased stabilization of the positive charge due to hyperconjugation and resonance, respectively ( $\Delta_r H_{\text{gas}}^0 = -98$  (H) vs.  $-157$  (Me) and  $-188$   $\text{kJ mol}^{-1}$  (Ph)). For alkenes, the same trend was expected. However,  $\Delta_r H_{\text{gas}}^0$  of the reaction with Ph<sub>2</sub>C=CPh<sub>2</sub> ( $-143$   $\text{kJ mol}^{-1}$ ) was calculated to be less favored than that with Me<sub>2</sub>C=CMe<sub>2</sub> ( $-174$   $\text{kJ mol}^{-1}$ ). We attribute this to steric reasons, as the phenyl groups cannot be coplanar as in the alkyne case. It should be noted that the *non-classical* structure of  $[\text{Me}_3\text{Si}(\text{Ph}_2\text{C}=\text{CPh}_2)]^+$  with equivalent Si–C distances was calculated to be a transition state for the  $[\text{Me}_3\text{Si}]^+$  migration between both carbon atoms. This transition state is higher in energy than the calculated minimum structure by only  $\Delta G_{o\text{-DFB}}^0 = +7$   $\text{kJ mol}^{-1}$  ( $\Delta G_{\text{gas}}^0 = +11$   $\text{kJ mol}^{-1}$ ). Therefore, it might not be possible to differentiate between the *classical* and *non-classical* structure by NMR spectroscopy (coalescence) or single crystal X-ray diffraction (dynamic disorder).

Including COSMO<sup>17</sup> Gibbs solvation energies, all calculated reaction energies become less favored and the reaction of  $[\text{Me}_3\text{Si}]^+$  with the larger PhC≡CPh was also calculated to be less exergonic than that with the smaller MeC≡CMe. This is attributed to the fact that smaller ions are generally better solvated than larger ones. Since upon reaction with  $[\text{Me}_3\text{Si}]^+$  the product cation always increases in size, reactions (1) and (2) become less favored the larger the ligand. Due to this effect, the formation of  $[\text{Me}_3\text{Si}(\text{Ph}_2\text{C}=\text{CPh}_2)]^+$  was calculated to be endergonic in solution by  $\Delta G_{o\text{-DFB}}^0 = +20$   $\text{kJ mol}^{-1}$ .

### Reactions of Me<sub>3</sub>Si–F–Al(OR<sup>F</sup>)<sub>3</sub> with alkenes and alkynes

Since the calculations suggested that the formation of *non-classical* silylium ions should be possible, in part also in solution, we turned towards their synthesis. Ethylene and acetylene are gases and only weakly bound to  $[\text{Me}_3\text{Si}]^+$  (*cf.* Table 1). Therefore, we performed the reactions of Me<sub>3</sub>Si–F–Al(OR<sup>F</sup>)<sub>3</sub> with the substituted compounds R<sub>2</sub>C=CR<sub>2</sub> and RC≡CR (R = Me, Ph). However, the reactions with Me<sub>2</sub>C=CMe<sub>2</sub>, Ph<sub>2</sub>C=CPh<sub>2</sub> and PhC≡CPh in *o*-DFB or CH<sub>2</sub>Cl<sub>2</sub> and at r.t. or  $-40$  °C did not yield any silylium ions, but only led to exergonic oligomerization of the alkenes and alkynes. However, this signals activation of the unsaturated hydrocarbons by Me<sub>3</sub>Si–F–Al(OR<sup>F</sup>)<sub>3</sub>. By contrast, reactions with MeC≡CMe in *o*-DFB or CH<sub>2</sub>Cl<sub>2</sub> at various temperatures ( $-40$  °C to r.t.; eqn (3)), reproducibly gave a 28% yield of  $[\text{Me}_4\text{C}_4\text{-SiMe}_3]^+[\text{al-f-al}]^-$  ( $[\text{al-f-al}]^- = [(\text{R}^{\text{F}}\text{O})_3\text{Al-F-Al}(\text{OR}^{\text{F}})_3]^-$ ; NMR, scXRD, Raman, IR) after crystallization. This compound will be discussed later in more detail.




### Reactions of [Ph<sub>3</sub>C]<sup>+</sup>[al-f-al]<sup>−</sup> with Me<sub>3</sub>SiH and alkenes/alkynes

By exchanging Me<sub>3</sub>Si–F–Al(OR<sup>F</sup>)<sub>3</sub> for more reactive silylium ions, we hoped to form the desired products faster and thereby prevent oligomerization. In order to generate the  $[\text{Me}_3\text{Si}]^+$  cation as an intermediate, we mixed the starting materials (liquid alkenes or alkynes) with  $[\text{Ph}_3\text{C}]^+[\text{al-f-al}]^-$  in the absence of solvent. An excess of Me<sub>3</sub>SiH was condensed onto this mixture to generate silylium ions or their Me<sub>3</sub>SiH adducts<sup>18</sup> *in situ* and allowed to warm to r.t. Although NMR spectra of these reaction mixtures in CD<sub>2</sub>Cl<sub>2</sub> showed complete conversion of the starting materials, the *non-classical* silylium ions could not be detected. Instead, hydrosilylation of PhC≡CPh and Me<sub>2</sub>C=CMe<sub>2</sub> (see the ESI† for details) and formation of neutral **2** for




$\text{MeC}\equiv\text{CMe}$  were observed (eqn (4)). Upon reaction of  $\text{MeC}\equiv\text{CMe}$  with  $[\text{Ph}_3\text{C}][\text{al-f-al}]^-$  and  $\text{Me}_3\text{SiH}$ , presumably **1** was formed, which then reacted with excess  $\text{Me}_3\text{SiH}$  to yield **2** and  $[\text{Me}_3\text{Si}(\text{solv})]^+$  ( $\text{solv} = \text{Me}_3\text{SiH}, \text{CH}_2\text{Cl}_2$ ). Only after complete conversion of the alkenes and alkynes, the anion was attacked and decomposed.<sup>19</sup> It should be noted that the evolving  $[\text{Me}_3\text{Si-H-SiMe}_3]^+[\text{al-f-al}]^-$  is not isolable as a pure material. Yet, the  $[\text{al-f-al}]^-$  anion was shown to be fairly stable against silylium ions in  $\text{CH}_2\text{Cl}_2$ .<sup>19</sup>



### Reactions of bromo-ethyl/vinyl silanes with $\text{Ag}^+[\text{al-f-al}]^-$

In order to prevent oligomerization of the alkenes and alkynes, we reacted bromo-ethyl and bromo-vinyl silanes with  $\text{Ag}^+[\text{al-f-al}]^-$  in  $\text{CH}_2\text{Cl}_2$  at  $-40^\circ\text{C}$ , respectively. The abstraction of  $\text{Br}^-$  from these silanes by  $\text{Ag}^+$  should result in the formation of ethylene/acetylene adducts of  $[\text{Me}_3\text{Si}]^+$  (cf. eqn (5a) for the bromo-ethyl example). Here the reaction with a second equivalent of an alkene/alkyne is not possible and the *non-classical* silylium ions should become isolable. Upon reaction, no solid  $\text{AgBr}$  visibly formed. Nevertheless, NMR spectra of these reactions did not show signals of the starting silanes, but of  $\text{Me}_3\text{SiBr}$  in both cases. This signal is slightly shifted to a lower field by 2 ppm and suggests a coordination of  $\text{Me}_3\text{SiBr}$  to  $\text{Ag}^+$ .<sup>19</sup> It seems likely that the *non-classical* silylium ions  $[\text{Me}_3\text{Si}(\text{C}_2\text{H}_4/\text{C}_2\text{H}_2)]^+$  were generated *in situ* (eqn (5a)). However, apparently these silylium ions then reacted further with the (solvated)  $\text{AgBr}$  under the formation of  $[(\text{Me}_3\text{SiBr})\text{Ag}(\text{solv})_x]^+$  and release of  $\text{C}_2\text{H}_4/\text{C}_2\text{H}_2$  (see eqn (5b) for the ethene example; the evolving  $\text{C}_2\text{H}_4$  could be identified by gas phase IR spectroscopy).<sup>20</sup>

Similar silver-halosilane adducts were observed and structurally characterized upon reaction of  $t\text{-Bu}_3\text{SiBr}$  with  $\text{Ag}^+[\text{al-f-al}]^-$  – although halide abstraction and  $\text{AgBr}$  formation were desired.<sup>19</sup> Additionally, partial decomposition of the anion and formation of  $\text{Me}_3\text{SiF}$  and  $\text{Me}_3\text{SiCl}$  were observed, which is further evidence for the generation of silylium ions.



### Formation of $[\text{Me}_3\text{Si}(\text{MeC}\equiv\text{CMe})]^+$ vs. $[\text{Me}_4\text{C}_4\text{-SiMe}_3]^+$

The reactions of 2-butyne with  $\text{Me}_3\text{Si-F-Al}(\text{OR}^F)_3$  and silylium ions only yielded **1** and **2**, respectively. Both compounds were formed from two 2-butyne units. This led to the question of whether it is at least theoretically possible to isolate the desired silylium ion  $[\text{Me}_3\text{Si}(\text{MeC}\equiv\text{CMe})]^+$  by these routes, or if such *non-classical* ions reside in shallow potential wells that allow for a simple and low-barrier subsequent reaction with  $\text{MeC}\equiv\text{CMe}$  to give **1**. Therefore, we calculated the reaction mechanism for the formation of  $[\text{Me}_4\text{C}_4\text{-SiMe}_3]^+$  (**P-4**) at the BP86-D3(BJ)/def-TZVP level of theory (Fig. 1). In order to reduce computational cost, we started from 2-butyne and  $[\text{Me}_3\text{Si}]^+$ , instead of  $\text{Me}_3\text{Si-F-Al}(\text{OR}^F)_3$ .

Since 2-butyne itself does not dimerize under standard conditions,<sup>21</sup> the silylation of 2-butyne can be assumed to be the first step in the formation of  $[\text{Me}_4\text{C}_4\text{-SiMe}_3]^+$ . The Gibbs activation energy  $\Delta G^\ddagger$  (**TS1**) for this exothermic process was




Fig. 1 Calculated reaction path, geometries and relative energies in  $\text{kJ mol}^{-1}$  for the formation of  $[\text{Me}_4\text{C}_4\text{-SiMe}_3]^+$  (P-4, black) and the 5-membered ring (P-5, red); BP86-D3(BJ)/def-TZVP.





calculated to be only 10 kJ mol<sup>-1</sup>. The silylated 2-butyne (**11**) then reacts with a second equivalent of 2-butyne to exothermically form the intermediate **12**. We were not able to determine the Gibbs activation energy (**TS2**) for this reaction with the method used. This indicates that this barrier is either very small or not existent at all. Based on thermodynamics, this intermediate should rearrange to yield a 5-membered ring (**P-5**). However, the barrier for this rearrangement (**TS3-5**) was calculated to be 45 kJ mol<sup>-1</sup> – which is almost triple the barrier (**TS3-4**) for the formation of the observed [Me<sub>4</sub>C<sub>4</sub>-SiMe<sub>3</sub>]<sup>+</sup> (**P-4**, 16 kJ mol<sup>-1</sup>). Thus, the calculations are in agreement with the experimental observation and suggest that **P-4** (=1) is the kinetic product.

When Me<sub>3</sub>Si-F-Al(OR<sup>F</sup>)<sub>3</sub> is used for the silylation of 2-butyne, the Gibbs activation energy (transition state similar to **TS1**) for this reaction is expected to be higher, due to its lower reactivity compared to [Me<sub>3</sub>Si]<sup>+</sup>. However, this will not change the entire picture. In conclusion it seems impossible to isolate the silylated 2-butyne (**11**) by use of Me<sub>3</sub>Si-F-Al(OR<sup>F</sup>)<sub>3</sub> or even (non-existent) free [Me<sub>3</sub>Si]<sup>+</sup>, as the reaction with the second molecule of 2-butyne appears to be much faster than the silylation.

### Properties of [Me<sub>4</sub>C<sub>4</sub>-SiMe<sub>3</sub>]<sup>+</sup>[*al-f-al*]<sup>-</sup>

From the reaction solution in *o*-DFB at r.t. single crystals of **1** were obtained by the addition of less polar CH<sub>2</sub>Cl<sub>2</sub>. The molecular cation structure of salt **1** is shown in Fig. 2. With Si-C1-C2/C2' angles being 113°–118°, the stabilization of this cation by the β-silicon effect is not evident. Otherwise, these angles would be expected to lie around 90°.<sup>22</sup>

This cation can be seen as a silylated cyclobutadiene and is closely related to the already known homoaromatic [R<sub>4</sub>C<sub>4</sub>-H]<sup>+</sup>, [R<sub>4</sub>C<sub>4</sub>-Cl]<sup>+</sup>, [<sup>t</sup>Bu<sub>4</sub>C<sub>4</sub>-OH]<sup>+</sup> and the neutral [R<sub>4</sub>C<sub>4</sub>-AlX<sub>3</sub>] (R = H,




Fig. 2 Molecular structure of the cation in [Me<sub>4</sub>C<sub>4</sub>-SiMe<sub>3</sub>][*al-f-al*]<sup>-</sup> (**1**) with thermal ellipsoids set at 50% probability level. The [*al-f-al*]<sup>-</sup> anion was omitted for clarity. Scheme: Si (yellow), C (grey), H (white). Selected distances (pm), bond angles (deg), and torsion angles (deg): Si-C1 192.2(2), C1-C2 151.8(2), C1-C2' 152.6(2), C2-C3 138.9(2), C2'-C3 138.5(2), C2-C2' 179.4(2), Si-C1-C2 117.9(1), Si-C1-C2' 113.0(1), C2-C1-C2' 72.2(1), C2-C3-C2' 80.6(1), and C1-C2-C3-C2' 25.5(1).

Me, <sup>t</sup>Bu, Ph; X = Cl, Br).<sup>23–29</sup> Of these, the protonated cations [R<sub>4</sub>C<sub>4</sub>-H]<sup>+</sup> with R = H, Me, Ph were only characterized by NMR at -40 to -70 °C,<sup>27</sup> while with R = <sup>t</sup>Bu they were shown to be isolable with a large variety of anions, like Br<sup>-</sup> or [SbF<sub>6</sub>]<sup>-</sup>.<sup>26</sup> Also the chlorinated and hydroxylated cations [Ph<sub>4</sub>C<sub>4</sub>-Cl]<sup>+</sup> and [<sup>t</sup>Bu<sub>4</sub>C<sub>4</sub>-OH]<sup>+</sup> and the neutral [Me<sub>4</sub>C<sub>4</sub>-AlCl<sub>3</sub>] were shown to be stable at r.t. and could be isolated.<sup>23,24,26,29</sup>

A comparison of selected structural parameters, calculated NPA and PABOON partial charges and π\*(C2-C3-C2')-orbital energies of the compounds is found in Table 2. In order to allow for a better comparison, these calculations were performed on the methyl-substituted derivatives for all compounds. The [(RSi)<sub>3</sub>SiR<sub>2</sub>]<sup>+</sup> cations (R = <sup>t</sup>Bu<sub>2</sub>MeSi, <sup>t</sup>Bu) can also be seen as silicon analogues of these compounds but will not be discussed.<sup>10,11</sup> When comparing the convolution angles (C1-C2-C2'-C3) of **1**, [<sup>t</sup>Bu<sub>4</sub>C<sub>4</sub>-H]<sup>+</sup>, [Ph<sub>4</sub>C<sub>4</sub>-Cl]<sup>+</sup>, [<sup>t</sup>Bu<sub>4</sub>C<sub>4</sub>-OH]<sup>+</sup> and [Me<sub>4</sub>C<sub>4</sub>-AlX<sub>3</sub>], their absolute values are relatively similar and range between 37.3° and 31.5°, except for [Ph<sub>4</sub>C<sub>4</sub>-Cl][Nb<sub>2</sub>OCl<sub>9</sub>] with a dihedral angle of only 4.3°. However, the dihedral angles in **1** and [Me<sub>4</sub>C<sub>4</sub>-AlX<sub>3</sub>] are positive, while in [<sup>t</sup>Bu<sub>4</sub>C<sub>4</sub>-H]<sup>+</sup>, [<sup>t</sup>Bu<sub>4</sub>C<sub>4</sub>-OH]<sup>+</sup> and [Ph<sub>4</sub>C<sub>4</sub>-Cl]<sup>+</sup> they are negative. This is a consequence of the (C2-)Me groups being bent, which leads to repulsion of the sterically most demanding group. The relatively small dihedral angle in [Ph<sub>4</sub>C<sub>4</sub>-Cl]<sup>+</sup> results from resonance of the phenyl moieties with the (C2-C2') orbital, which is an intermediate between a σ and a π orbital. This leads to an increased C2-C2' distance compared to that in the other compounds (203 vs. ~180 pm).<sup>27,30</sup>

When looking at the bonding π(C2-C3-C2') orbitals of the methylated derivatives of the discussed compounds (Table 2), a correlation between their energies and the partial charge of the Me<sub>4</sub>C<sub>4</sub> moiety is evident, with a higher partial charge leading to lower orbital energies. From a frontier orbital point of view, this implies that the cations [Me<sub>4</sub>C<sub>4</sub>-SiMe<sub>3</sub>]<sup>+</sup>, [R<sub>4</sub>C<sub>4</sub>-H]<sup>+</sup>, [R<sub>4</sub>C<sub>4</sub>-Cl]<sup>+</sup>, and [<sup>t</sup>Bu<sub>4</sub>C<sub>4</sub>-OH]<sup>+</sup> are more electron-deficient than the neutral [Me<sub>4</sub>C<sub>4</sub>-AlX<sub>3</sub>]. This is also evident when simply looking at the total charge of these molecules. Interestingly, the energies of the π\*(C2-C3-C2') orbitals seem to be less affected by the partial charge of the Me<sub>4</sub>C<sub>4</sub> moiety, but mostly by the total charge of the molecules. Therefore, the energy of this orbital is nearly the same for [Me<sub>4</sub>C<sub>4</sub>-AlCl<sub>3</sub>] and Me<sub>4</sub>C<sub>4</sub>, while for the cationic species it is lower by ~4 eV.

### Evaluating the homoaromatic character of [R<sub>4</sub>C<sub>4</sub>-E]<sup>(+)</sup>

Since the structural parameters of the discussed compounds are very similar, especially *d*(C2-C2') and *d*(C2-C3), an unambiguous comparison of the C2-C2' interactions is not possible (except for [Ph<sub>4</sub>C<sub>4</sub>-Cl]<sup>+</sup>, which does not show this interaction). This interaction is a measure of homoaromaticity and can be determined from the difference in NMR chemical shifts between C2/C2' and C3 (Table 3).<sup>26,27</sup> In allylic systems, the terminal C atoms (here C2/C2') bear a positive charge, resulting in the deshielding of these. The interaction between C2 and C2' leads to their shielding, combined with the transfer of the positive charge to C3 (deshielding). For homoaromatic systems



**Table 2** Selected experimental (calculated) properties of different  $[R_4C_4-E]^{(+)}$  compounds (E = SiMe<sub>3</sub>, H, Cl, OH, AlCl<sub>3</sub>, and AlBr<sub>3</sub>). Distances are given in pm, angles are given in ° and orbital energies are given in eV. The calculated values always refer to  $[Me_4C_4-E]^{(+)}$  for better comparability; BP86-D3(BJ)/def-TZVP

1	$[{}^tBu_4C_4-H]^+$ (ref. 26)	$[Ph_4C_4-Cl]^+$ (ref. 23)	$[{}^tBu_4C_4-OH]^+$ (ref. 26)	$[Me_4C_4-AlCl_3]$ (ref. 31)	$[Me_4C_4-AlBr_3]$	$Me_4C_4$
<b>Bond distances (pm) and dihedral angles (deg)</b>						
$d(C1-C2)$	151.8/152.6 (152.7)	152.4 (153.4)	152.9/154.4 (153.2)	152.3/153.8 (154.2)	151.0 (151.5)	(151.6) (159.1)
$d(C2-C2')$	179.4 (183.2)	180.6 (187.5)	203.3 (195.5)	183.3 (190.7)	177.4 (183.1)	(184.4) (208.5)
$d(C2-C3)$	138.5/138.9 (140.2)	140.7 (140.2)	138.7/140.4 (140.6)	139.4/139.6 (140.2)	138.7 (140.2)	(140.2) (134.7)
$C1-C2-C2'-C3$	31.7 (29.3)	-37.3 (-27.7)	-4.3 (-22.1)	-36.3 (-26.2)	31.5 (28.5)	(27.3) (0.0)
<b>NPA/PABOON partial charges for the methyl derivatives <math>[Me_4C_4-EY_x]^{(+)}</math></b>						
$\delta(C1)$	(-0.43/-0.14)	(-0.27/-0.05)	(-0.09/0.09)	(0.26/0.15)	(-0.52/-0.14)	(-0.54/-0.13) (0.00/0.01)
$\delta(C2)$	(0.27/0.21)	(0.28/0.19)	(0.30/0.15)	(0.17/0.15)	(0.27/0.27)	(0.28/0.27) (0.00/0.01)
$\delta(C3)$	(0.00/-0.02)	(0.02/0.01)	(-0.01/0.01)	(0.02/0.03)	(-0.04/-0.05)	(-0.05/-0.05) (0.00/0.01)
$\delta(C1 + 2 \times C2 + C3)$	(0.11/0.26)	(0.31/0.34)	(0.50/0.40)	(0.62/0.48)	(-0.02/0.35)	(-0.03/0.36) (-0.01/0.04)
$\delta(Me_4C_4)$	(0.51/0.82)	(0.75/0.90)	(1.00/1.03)	(1.16/1.01)	(0.25/0.67)	(0.25/0.67) (0.00/0.00)
$\delta(E)$	(0.49/0.18)	(0.25/0.10)	(0.00/-0.03)	(-0.16/-0.01)	(-0.25/-0.67)	(-0.25/-0.67) (-)
<b>Energies of the (anti-)bonding <math>\pi(C2-C3-C2')</math>-orbitals of the methyl derivatives <math>[Me_4C_4-EY_x]^{(+)}</math></b>						
$E_{\pi^*(C2-C3-C2')}$	(-5.60) <sup>a</sup>	(-6.04) <sup>a</sup>	(-6.14) <sup>b</sup>	(-6.02) <sup>c</sup>	(-1.92) <sup>d</sup>	(-1.92) <sup>d</sup> (-1.70) <sup>e</sup>
$E_{\pi(C2-C3-C2')}$	(-10.36) <sup>a</sup>	(-11.75) <sup>a</sup>	(-12.12) <sup>b</sup>	(-11.90) <sup>c</sup>	(-7.33) <sup>d</sup>	(-7.04) <sup>d</sup> (-3.74) <sup>e</sup>

<sup>a</sup> HOMO/LUMO+1. <sup>b</sup> HOMO-2/LUMO+1. <sup>c</sup> HOMO-1/LUMO+1. <sup>d</sup> HOMO-6/LUMO+1. <sup>e</sup> HOMO/LUMO.

a negative value for  $\Delta(\delta^{13}C_{(C2)} - \delta^{13}C_{(C3)})$  is expected, while for allylic systems this difference should be positive.<sup>26</sup>

With  $[H_4C_4-H]^+$  and the cyclopentenylum cation  $[H_6C_5-H]^+$  being the prototypes for homoaromatic and allylic cations, respectively, the homoaromatic character of the discussed compounds and influence of the substituents can be evaluated. Exchanging the H substituents for Me moieties in  $[R_4C_4-H]^+$  leads to hyperconjugation of the C-H bond to C2. As a result, the C2-C2' interactions are decreased and C2 is deshielded. Therefore the difference between  $\delta^{13}C_{(C2)}$  and  $\delta^{13}C_{(C3)}$  is more positive (0.0 vs. -54.1 ppm for  $[H_4C_4-H]^+$ ). The methylated compounds  $[Me_4C_4-SiMe_3]^+$  and  $[Me_4C_4-AlCl_3]$  show similar differences in the shielding of C2 and C3 and therefore are considered to have a similar homoaromatic character as  $[Me_4C_4-H]^+$ . Only  $[Me_4C_4-Cl]^+$  shows a comparatively high deshielding of C2 due to low homoaromaticity, which is a consequence of the hyperconjugation of the  $\sigma(C-Cl)$  orbital to the  $(C2-C2')$  orbital.

When looking at the C2-C2' interactions of  $[{}^tBu_4C_4-H]^+$  and  $[{}^tBu_4C_4-OH]^+$ , they would be expected to be even weaker than in  $[Me_4C_4-H]^+$  and  $[Me_4C_4-Cl]^+$ , respectively, due to the stronger electron-donating properties of the <sup>t</sup>Bu moieties. However, the NMR chemical shifts suggest a significantly increased homoaromatic character. We assign this to the bulkiness of the <sup>t</sup>Bu moieties leading to a repulsion of the C3-<sup>t</sup>Bu and C2-<sup>t</sup>Bu groups. As a result, the C1-C2-C2'-C3 dihedral angle is higher than that for the methylated derivatives (Table 2). Phenyl substituents seem to completely prevent the C2-C2' interactions due to resonance and charge delocalization onto the phenyl residue(s). This follows from the NMR chemical shifts of  $[Ph_4C_4-H]^+$  and from the structural parameters of  $[Ph_4C_4-Cl]^+$  (dihedral angle and  $d(C2-C2')$ ).

Taking a closer look at the dihedral angles in the discussed compounds, a correlation between these and  $\Delta(\delta^{13}C_{(C2)} - \delta^{13}C_{(C3)})$  is evident. The higher the dihedral angle, the stronger the C2-C2' interaction in the compound. For  $[H_4C_4-H]^+$  this angle was calculated to be 33.5° (BP86-D3(BJ)/def-TZVP), which is in agreement with this thesis. Interestingly, the distances  $d(C2-C2')$  and  $d(C2-C3)$  are only slightly affected by these C2-C2' interactions (Table 2).

### Investigations towards release of $Me_4C_4$ from 1

While information on the reactivities of most of the discussed compounds is scarce,  $[R_4C_4-AlX_3]$  was shown to be a source of cyclobutadienes  $R_4C_4$  (CBDs) by use of coordinating solvents, like DMSO, due to abstraction of AlCl<sub>3</sub>.<sup>28,32-34</sup> Besides the discussed AlX<sub>3</sub> adducts cyclobutene dicarboxylic acid anhydrides

**Table 3** Experimental <sup>13</sup>C NMR chemical shifts of  $[R_4C_4-E]^{(+)}$  in ppm (R = H, Me, <sup>t</sup>Bu, and Ph; E = SiMe<sub>3</sub>, H, Cl, OH, AlCl<sub>3</sub>, and AlBr<sub>3</sub>)


Cation	$\delta(C1)$	$\delta(C2/C2')$	$\delta(C3)$	$\delta(C2) - \delta(C3)$
$[Me_4C_4-SiMe_3]^+$	66.8	166.0	170.4	-4.4
$[{}^tBu_4C_4-OH]^+$ (ref. 26)	101.0	161.5	184.9	-23.4
$[Me_4C_4-Cl]^+$ (ref. 27)	76.0	191.5	174.4	+17.1
$[Me_4C_4-AlCl_3]$ (ref. 29)	—	162.0	164.3	-2.3
$[Me_4C_4-H]^+$ (ref. 27)	57.8	171.3	171.3	0.0
$[{}^tBu_4C_4-H]^+$ (ref. 26)	78.5	156.6	196.6	-40.0
$[Ph_4C_4-H]^+$ (ref. 27)	52.5	190.0	152.3	+38.6
$[H_4C_4-H]^+$ (ref. 27)	54.0	133.5	187.6	-54.1
$[H_6C_5-H]^{+a}$ (ref. 27)	48.7	234.7	145.7	+89.0

<sup>a</sup> Cyclopentenylum cation as reference for an allylic cation.



and  $[\text{Fe}(\text{CO})_3(\text{C}_4\text{H}_4)]$  can also be used for the generation of CBDs.<sup>35</sup> CBDs are anti-aromatic and, therefore, unsubstituted CBD undergoes dimerization readily at  $T > 35$  K.<sup>36</sup> Thus, small CBDs have to be generated *in situ*. Only when they bear large substituents, dimerization can be prevented.<sup>37</sup>

By abstraction of the  $[\text{Me}_3\text{Si}]^+$  moiety, **1** might be used as a tetramethylcyclobutadiene ( $\text{Me}_4\text{C}_4$ ) donor as well. The



advantage of **1** over  $[\text{Me}_4\text{C}_4\text{-AlX}_3]$  as a CBD donor would be its increased stability, which allows for storage in a glove box for more than a year. NMR spectra of  $[\text{Me}_4\text{C}_4\text{-AlX}_3]$  show partial coalescence of the signals already at 20 °C.<sup>29</sup> In contrast, all signals of **1** are resolved, including the  $^5J_{\text{H-H}}$  coupling of 0.46 Hz. The increased stability very likely results from the higher Lewis acidity of  $[\text{Me}_3\text{Si}]^+$  over  $\text{AlX}_3$  (FIA = 539 vs. 425/438 kJ mol<sup>-1</sup>, X = Cl/Br). These FIAs were calculated in an environment with the polarity of  $\text{CH}_2\text{Cl}_2$  ( $\epsilon_r = 8.9$ ) using COSMO,<sup>17</sup> in order to account for the different charges of the Lewis acids. The effect is also evident by addition of  $\text{AlX}_3$  to  $[\text{Me}_4\text{C}_4\text{-AlX}_3]$  in order to increase the Lewis acidity, *i.e.* the formation of  $[\text{Me}_4\text{C}_4\text{-Al}_2\text{X}_6]$ . It shows an increased thermal stability and reduced coalescence compared to  $[\text{Me}_4\text{C}_4\text{-AlX}_3]$ .<sup>29</sup>

### Reaction with $[\text{NMe}_4]\text{F}$

Due to the high stability of the Si-F bond, we reacted **1** with  $[\text{NMe}_4]\text{F}$  in *o*-DFB/ $\text{CH}_2\text{Cl}_2$  at r.t., in order to generate  $\text{Me}_3\text{SiF}$  and  $\text{Me}_4\text{C}_4$ . However, only negligible amounts of  $\text{Me}_3\text{SiF}$  formed. The main products were  $[\text{NMe}_4]^+[\text{F}_{1+x}\text{-Al}(\text{OR}^{\text{F}})_{3-x}]^-$  ( $x = 0, 1$ ),  $\text{HOR}^{\text{F}}$  and **3** due to deprotonation of **1** (eqn (6a)). The formation of the anions  $[\text{F}_{1+x}\text{-Al}(\text{OR}^{\text{F}})_{3-x}]^-$  occurs due to the high Lewis acidity of  $[\text{al-f-al}]^-$ .<sup>19</sup> The deprotonation of **1** suggested that a non-charged nucleophile, like 4-dimethylaminopyridine (DMAP), could be helpful to prevent deprotonation of the cation.

### Reaction with DMAP

The reaction of **1** with an excess of DMAP in a mixture of *o*-DFB and  $\text{CH}_2\text{Cl}_2$  at r.t. yielded mainly  $\text{DMAP-Al}(\text{OR}^{\text{F}})_3$ ,  $[\text{F-Al}(\text{OR}^{\text{F}})_3]^-$  (reaction with the anion) and the adduct of  $[\text{Me}_4\text{C}_4\text{-SiMe}_3]^+$  with DMAP (**4**, eqn (6b)). We were able to obtain single crystals of both compounds at -40 °C from the reaction solution (see the ESI† for details). The crystal structure and NMR spectra of **4** showed that the cation is obtained as an enantiomeric mixture, with the DMAP and the  $[\text{Me}_3\text{Si}]^+$  moieties being in *syn*-conformation, *i.e.* in (*S,R*) and (*R,S*) configurations, which is attributed to the orientation of the LUMO of **1** (see Fig. S-60†). A similar adduct between  $\text{Me}_4\text{C}_4\text{-AlCl}_3$  and  $\text{PPhCl}_2$  was already proposed as an intermediate in the synthesis of phosphole oxides.<sup>33</sup> Due to the *syn*-

conformation of the DMAP and  $\text{Me}_3\text{Si}$  moieties it seemed possible to remove both of them in the form of  $[\text{DMAP-SiMe}_3]^+$  under the release of  $\text{Me}_4\text{C}_4$  by gentle warming/heating. However, heating of **4** to 100 °C did not result in the elimination of  $[\text{DMAP-SiMe}_3]^+$ , but only led to the formation of **3**,  $[\text{DMAP-H}]^+[\text{f-al}]^-$  and other unidentified products.

### Reaction with diethylether

Unexpectedly, one of the minor side products in the reaction of **1** with DMAP was  $\text{Me}_3\text{SiF}$ , providing evidence for the reaction of  $[\text{Me}_4\text{C}_4\text{-SiMe}_3]^+$  with a fluoride ion source. Since **1** is stable in *o*-DFB over several days, the Lewis basic  $[\text{F-Al}(\text{OR}^{\text{F}})_3]^-$  anion ( $[\text{f-al}]^-$ )<sup>19</sup> has to be considered as the fluoride ion source. This is in agreement with the findings that the NMR spectra of **1** never showed the presence of  $[\text{f-al}]^-$ , independent of the reaction stoichiometry.<sup>38</sup> Therefore, we dissolved **1** in  $\text{Et}_2\text{O}$  at r.t., which is known to induce the dissociation of  $[\text{al-f-al}]^-$  into  $\text{Et}_2\text{O-Al}(\text{OR}^{\text{F}})_3$  and  $[\text{f-al}]^-$ .<sup>19</sup> NMR spectra of this reaction mixture revealed the formation of  $\text{Me}_3\text{SiF}$  and complete decomposition of **1** (eqn (6c)), but no evidence for the expected dimerization product of  $\text{Me}_4\text{C}_4$  could be found.<sup>34</sup> Instead, the NMR spectra suggested the formation of oligomerization products. When this reaction was performed in the presence of 2-butyne, partial formation of  $\text{C}_6\text{Me}_6$  was observed, similar to the reaction of  $[\text{Me}_4\text{C}_4\text{-AlCl}_3]$  with 2-butyne and DMSO.<sup>34</sup> Therefore, we assume this reaction is (weak) evidence for the intermediate release of CBD from **1**. However, analogous reactions of **1** with other electron-deficient or electron-rich dienophiles ( $\text{MeO}_2\text{C-C}\equiv\text{C-CO}_2\text{Me}$ ,  $\text{Me}_3\text{Si-C}\equiv\text{C-SiMe}_3$ ) never gave the desired Diels-Alder products,<sup>29</sup> but only the known oligomerization products and complete retention of the alkynes.

It should be noted that the reaction solution for the synthesis of **1** also contained  $\text{C}_6\text{Me}_6$  and these oligomerization products. The reason for this is that in the first stage of the reaction of  $\text{Me}_3\text{Si-F-Al}(\text{OR}^{\text{F}})_3$  with 2-butyne the  $[\text{f-al}]^-$  anion is formed. Therefore, **1**,  $[\text{f-al}]^-$  and 2-butyne are present in solution at the same time. Thus, **1** readily reacts with  $[\text{f-al}]^-$  upon its initial synthesis, resulting in its decomposition and accounting for the relatively low yield of 28%. Eventually, the formation of  $[\text{al-f-al}]^-$  from  $[\text{f-al}]^-$  and  $\text{Me}_3\text{Si-F-Al}(\text{OR}^{\text{F}})_3$  suppresses this decomposition reaction.

### Rationalization

Why is there such a difference in the reactivity of alkynes with  $\text{Me}_4\text{C}_4$  released from **1** and from  $[\text{Me}_4\text{C}_4\text{-AlX}_3]$ ?  $\text{Me}_4\text{C}_4$  is an electron-rich diene (high HOMO energy) and therefore should react with electron-deficient dienophiles (low LUMO energy). While the  $\pi$ -LUMO energy of -1.92 eV for  $[\text{Me}_4\text{C}_4\text{-AlX}_3]$  is rather



high and similar to that for free  $\text{Me}_4\text{C}_4$ , it is much lower for **1** with an energy of  $-5.60$  eV (Table 1). For this reason  $[\text{Me}_4\text{C}_4\text{-AlX}_3]$  is a poor dienophile and the released  $\text{Me}_4\text{C}_4$  alone undergoes Diels–Alder reactions with the added alkynes. In contrast, the  $\pi$ -LUMO energy of even very electron-deficient alkynes, like  $\text{F}_3\text{C-C}\equiv\text{C-CF}_3$  ( $-2.70$  eV), is significantly higher than that of **1**. As a result, the evolving  $\text{Me}_4\text{C}_4$  reacts with  $[\text{Me}_4\text{C}_4\text{-SiMe}_3]^+$  instead of other dienophiles, leading to oligomerization. The same problem is likely to arise when using the other cationic cyclobutenyl cations as their  $\pi$ -LUMO energies are expected to be in the same region as those of **1**.

## Conclusion

By the reaction of  $\text{Me}_3\text{Si-F-Al(OR}^{\text{F}})_3$  with different alkenes and alkynes the synthesis of *non-classical* silylium ions was investigated. However, most of these reactions only led to oligomerizations in which such silylium ions may be intermediates; they were assessed by calculations (Table 1). Replacing  $\text{Me}_3\text{Si-F-Al(OR}^{\text{F}})_3$  with silylium ions, generated *in situ* from  $[\text{Ph}_3\text{C}]^+[\text{al-f-al}]^-$  and  $\text{Me}_3\text{SiH}$ , also did not yield the desired *non-classical* silylium ions. Instead, hydrosilylation of the alkenes and alkynes was observed. Preliminary tests suggest that these hydrosilylation reactions can be performed with only catalytic amounts of  $[\text{Ph}_3\text{C}]^+[\text{al-f-al}]^-$ .<sup>39</sup> However, this should be investigated in a separate study by specialists. Halide abstraction reactions from bromo-vinyl/alkyl silanes by  $\text{Ag}^+[\text{al-f-al}]^-$  seemed to be successful, but resulted in the formation of bromo silanes and  $\text{C}_2\text{H}_2/\text{C}_2\text{H}_4$  due to the reaction of the silylium ions with solvated  $\text{AgBr}$ . From the reaction of  $\text{Me}_3\text{Si-F-Al(OR}^{\text{F}})_3$  with 2-butyne, a salt of the stable silylated tetramethyl cyclobutenyl cation  $[\text{Me}_4\text{C}_4\text{-SiMe}_3]^+$  was obtained. Attempts to release the CBD  $\text{Me}_4\text{C}_4$  from this compound did not result in the planned Diels–Alder reactions, but only in oligomerization products. We attribute this to the comparatively low  $\pi$ -LUMO energy of  $[\text{Me}_4\text{C}_4\text{-SiMe}_3]^+$ , leading to the reaction of this cation with the evolving  $\text{Me}_4\text{C}_4$ .

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

This work was supported by the Albert-Ludwigs-Universität Freiburg, the ERC project UniChem and the DFG. We would like to acknowledge the MagRes facility of the University of Freiburg and Fadime Bitgül for the measurement of the NMR spectra and Daniel Kratzert for the valuable discussions of the crystallographic data.

## References

- (a) G. A. Olah and G. K. Surya Prakash, ed. *Carbocation Chemistry*, John Wiley & Sons, Hoboken, New Jersey, 2004; (b) H. Meerwein and K. van Emster, *Ber. Dtsch. Chem. Ges. A/B*, 1922, 55, 2500.
- G. A. Olah, *J. Am. Chem. Soc.*, 1972, 94, 808.
- (a) P. v. R. Schleyer, D. Lenoir, P. Mison, G. Liang, G. K. S. Prakash and G. A. Olah, *J. Am. Chem. Soc.*, 1980, 102, 683; (b) H. C. Brown, *Acc. Chem. Res.*, 1986, 19, 34; (c) S. Winstein and D. S. Trifan, *J. Am. Chem. Soc.*, 1949, 71, 2953; (d) H. C. Brown, *The Nonclassical Ion Problem*, Springer US, Boston, MA, 1977.
- F. Scholz, D. Himmel, F. W. Heinemann, P. v. R. Schleyer, K. Meyer and I. Krossing, *Science*, 2013, 341, 62.
- (a) B. Chiavarino, M. E. Crestoni, J. Lemaire, P. Maitre and S. Fornarini, *J. Chem. Phys.*, 2013, 139, 71102; (b) O. MÓ, M. Yáñez, J.-F. Gal, P. C. Maria and J.-C. Guillemin, *J. Phys. Org. Chem.*, 2002, 15, 509; (c) B. Chiavarino, M. E. Crestoni and S. Fornarini, *Chem. Commun.*, 2002, 1418.
- X. Li and J. A. Stone, *J. Am. Chem. Soc.*, 1989, 111, 5586.
- H.-U. Siehl, S. Brixner, C. Coletti, N. Re, B. Chiavarino, M. E. Crestoni, A. de Petris and S. Fornarini, *Int. J. Mass Spectrom.*, 2013, 334, 58.
- M. F. Ibad, P. Langer, A. Schulz and A. Villinger, *J. Am. Chem. Soc.*, 2011, 133, 21016.
- (a) T. Müller, C. Bauch, M. Ostermeier, M. Bolte and N. Auner, *J. Am. Chem. Soc.*, 2003, 125, 2158; (b) S. Duttwyler, Q.-Q. Do, A. Linden, K. K. Baldrige and J. S. Siegel, *Angew. Chem., Int. Ed.*, 2008, 47, 1719; (c) J. B. Lambert, S. Zhang and S. M. Ciro, *Organometallics*, 1994, 13, 2430; (d) J. B. Lambert, S. Zhang, C. L. Stern and J. C. Huffman, *Science*, 1993, 260, 1917.
- S. Inoue, M. Ichinohe, T. Yamaguchi and A. Sekiguchi, *Organometallics*, 2008, 27, 6056.
- A. Sekiguchi, T. Matsuno and M. Ichinohe, *J. Am. Chem. Soc.*, 2000, 122, 11250.
- A. Burcat and E. Goos, *Int. J. Chem. Kinet.*, 2018, 7, 793.
- (a) A. Sekiguchi, *Science*, 2004, 305, 1755; (b) R. West, M. J. Fink and J. Michl, *Science*, 1981, 214, 1343; (c) I. Bejan and D. Scheschke, *Angew. Chem., Int. Ed.*, 2007, 46, 5783.
- T. E. Mallouk, G. L. Rosenthal, G. Mueller, R. Brusasco and N. Bartlett, *Inorg. Chem.*, 1984, 23, 3167.
- K. O. Christe, D. A. Dixon, D. McLemore, W. W. Wilson, J. A. Sheehy and J. A. Boatz, *J. Fluorine Chem.*, 2000, 101, 151.
- D. R. Lide, *CRC Handbook of Chemistry and Physics*, CRC Press/Taylor and Francis, Boca Raton, 90th edn, 2010.
- A. Klamt and G. Schuurmann, *J. Chem. Soc., Perkin Trans. 2*, 1993, 799.
- S. P. Hoffmann, T. Kato, F. S. Tham and C. A. Reed, *Chem. Commun.*, 2006, 767.
- A. Martens, P. Weis, M. C. Krummer, M. Kreuzer, A. Meierhöfer, S. C. Meier, J. Bohnenberger, H. Scherer, I. Riddlestone and I. Krossing, *Chem. Sci.*, 2018, 9, 7058.
- C. A. Reed, *Acc. Chem. Res.*, 1998, 31, 325.
- (a) V. G. Baonza, O. R. Montoro, M. Taravillo, M. Cáceres and J. Núñez, *J. Chem. Phys.*, 2004, 121, 11156; (b) C. Mediavilla, J. Tortajada and V. G. Baonza, *Chem. Phys. Lett.*, 2008, 454, 387.
- S. G. Wierschke, J. Chandrasekhar and W. L. Jorgensen, *J. Am. Chem. Soc.*, 1985, 107, 1496.
- E. Hey, F. Weller and K. Dehnicke, *Z. Anorg. Allg. Chem.*, 1983, 502, 45.





- 24 R. F. Bryan, *J. Am. Chem. Soc.*, 1964, **86**, 733.
- 25 H. H. Freedman and A. E. Young, *J. Am. Chem. Soc.*, 1964, **86**, 734.
- 26 G. Maier, R. Emrich, K.-D. Malsch, K.-A. Schneider, M. Nixdorf and H. Irngartinger, *Chem. Ber.*, 1985, **118**, 2798.
- 27 G. A. Olah, J. S. Staral, R. J. Spear and G. Liang, *J. Am. Chem. Soc.*, 1975, **97**, 5489.
- 28 J. B. Koster, G. J. Timmermans and H. van Bekkum, *Synthesis*, 1971, **1971**, 139.
- 29 P. B. J. Driessen and H. Hogeveen, *J. Am. Chem. Soc.*, 1978, **100**, 1193.
- 30 M. Simonetta and S. Winstein, *J. Am. Chem. Soc.*, 1954, **76**, 18.
- 31 C. Krüger, P. J. Roberts, Y.-H. Tsay and J. B. Koster, *J. Organomet. Chem.*, 1974, **78**, 69.
- 32 (a) Š. Janková, J. Schulz, S. Hybelbauerová, I. Císařová, P. Štěpnička and M. Katora, *Eur. J. Org. Chem.*, 2013, **2013**, 44; (b) P. B. J. Driessen and H. Hogeveen, *J. Organomet. Chem.*, 1978, **156**, 265; (c) M. J. Marsella, S. Estassi, L.-S. Wang and K. Yoon, *Synlett*, 2004, 192.
- 33 K. S. Fongers, H. Hogeveen and R. F. Kingma, *Tetrahedron Lett.*, 1983, **24**, 1423.
- 34 H. Hogeveen, H. Jorritsma, P. A. Wade, F. van Rantwijk, J. B. Koster, J. J. Prooi, A. Sinnema and H. van Bekkum, *Tetrahedron Lett.*, 1974, **15**, 3915.
- 35 T. Bally and S. Masamune, *Tetrahedron*, 1980, **36**, 343.
- 36 A. Fattahi, L. Lis, Z. Tian and S. R. Kass, *Angew. Chem., Int. Ed.*, 2006, **45**, 4984.
- 37 (a) H. Irngartinger and H. Rodewald, *Angew. Chem., Int. Ed.*, 1974, **13**; (b) H. Kimling and A. Krebs, *Angew. Chem., Int. Ed.*, 1972, **11**, 932.
- 38 (a) J. Possart, A. Martens, M. Schleep, A. Ripp, H. Scherer, D. Kratzert and I. Krossing, *Chem.-Eur. J.*, 2017, **23**, 12305; (b) M. Rohde, L. O. Müller, D. Himmel, H. Scherer and I. Krossing, *Chem.-Eur. J.*, 2014, **20**, 1218.
- 39 (a) R. J. Wehmschulte and L. Wojtas, *Inorg. Chem.*, 2011, **50**, 11300; (b) M. Kira, T. Hino and H. Sakurai, *Chem. Lett.*, 1992, **21**, 555.

