

Cite this: *Chem. Sci.*, 2019, 10, 2678

All publication charges for this article have been paid for by the Royal Society of Chemistry

Selective C–C bond formation from rhodium-catalyzed C–H activation reaction of 2-arylpyridines with 3-aryl-2H-azirines†

Yonghyeon Baek,^{‡a} Jinwoo Kim,^{‡cd} Hyunseok Kim,^a Seung Jin Jung,^a Ho Ryu,^{cd} Suyeon Kim,^{cd} Jeong-Yu Son,^a Kyusik Um,^a Sang Hoon Han,^a Hyung Jin Seo,^a Juyoung Heo,^a Kooyeon Lee,^b Mu-Hyun Baik^{id}*^{cd} and Phil Ho Lee^{id}*^a

A novel method for the synthesis of acylmethyl-substituted 2-arylpyridine derivatives using 3-aryl-2H-azirines was developed by exploring a prototype reaction using DFT-calculations and carrying out targeted experiments guided by the calculated mechanism. 2H-Azirine was initially hypothesized to ring-open at the metal center to furnish familiar metal nitrene complexes that may undergo C–N coupling. Computational studies quickly revealed and prototype experimental work confirmed that neither the formation of the expected metal nitrene complexes nor the C–N coupling were viable. Instead, azirine ring-opening followed by C–C coupling was found to be much more favorable to give imines that readily underwent hydrolysis in aqueous conditions to form acylmethyl-substituted products. This new method was highly versatile and selective toward a wide range of substrates with high functional group tolerance. The utility of the new method is demonstrated by a convenient one-pot synthesis of biologically relevant heterocycles such as pyridoisoindole and pyridoisoquinoline.

Received 17th November 2018

Accepted 5th January 2019

DOI: 10.1039/c8sc05142a

rsc.li/chemical-science

Introduction

Controlling the chemo-, regio- and stereoselectivities of reactions is of central importance in general, but particularly so in organic chemistry.¹ And C–H bond activation reactions² are among the most desirable to carry out with high degrees of selectivity and significant research efforts were made in the past to couple C–H activation to C–C and C–N bond forming transformations. Alkyl and aryl azides featured prominently as valuable reagents in this endeavor and many C–N bond forming reagents based on C–H activation reactions have been developed (Scheme 1, eqn (1)).³ Acyl azides are attractive components,⁴ as they are capable of forming C–C bonds through isocyanates generated *in situ* within the framework of well-studied Curtius rearrangement pathways.⁵ Most notably, Chang recently developed a method for forming C–N bonds

using acyl azides *via* C–H activation employing a ruthenium catalyst.⁶ Stable sulfonyl azides have also been used to form C–N bonds in C–H activation reactions⁷ and, most recently, dioxazolones were identified as highly versatile surrogates of azides for installing C–N bonds under mild reaction conditions.⁸ These studies highlight that the choice of an appropriate reagent that will mediate the formation of C–N and/or C–C bonds is critical. Thus, we sought to expand the scope of C–H activation reactions by identifying new reagents that may act as surrogates of azides in a C–H activation reaction. Given our previous experience with azirines,⁹ we wondered if they can be employed to furnish vinyl nitrenoids that may be exploited to form C–C and C–N bond selectively (Scheme 1, eqn (2)).

To increase the efficiency of the new reaction discovery process, we used density functional theory (DFT) based calculations¹⁰ to first examine the feasibility of putative reaction mechanisms and identify the most promising strategy for reaction design. As shown in eqn (3) (Scheme 1), we initially speculated that the 3-aryl-2H-azirine substrate may coordinate to a metallacycle intermediate that should be readily accessible *via* C–H activation using a well-known Ir, Rh or Co catalyst. Subsequent ring-opening of the azirine¹¹ was envisioned to give a vinyl metal-nitrene,¹² from which a migratory insertion may result in a C–N coupled product. Surprisingly, DFT-calculations showed that the familiar C–N bond formation is not likely in this case, but suggested that C–C bond formation can be furnished efficiently. Specifically, the barrier for the formation of the eight-membered metallacycle was computed to be low. In

^aNational Creative Research Initiative Center for Catalytic Organic Reactions, Department of Chemistry, Kangwon National University, Chuncheon 24341, Republic of Korea. E-mail: phlee@kangwon.ac.kr

^bDepartment of Bio-Health Technology, Kangwon National University, Chuncheon 24341, Republic of Korea


^cDepartment of Chemistry, Korea Advanced Institute of Science and Technology (KAIST), Daejeon 34141, Republic of Korea. E-mail: mbaik2805@kaist.ac.kr

^dCenter for Catalytic Hydrocarbon Functionalizations, Institute for Basic Science (IBS), Daejeon 34141, Republic of Korea

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c8sc05142a

‡ These authors contributed equally to this work.





Scheme 1 Selective C–C and C–N bond formation using azides and its surrogates in C–H activation.

good agreement with this DFT-based proposal, treatment of 2-phenylpyridine with 3-(4-methoxyphenyl)-2H-azirine in the presence of a Rh catalyst gave the acylmethylated product in 17% yield, whereas no evidence for C–N coupling was detectable.¹³ Based on these initial results, we developed an efficient and generally applicable new method that allows for the acylmethylation of 2-arylpyridine derivatives through selective C–C bond formation employing 2-arylpyridine and 3-aryl-2H-azirine using a Rh catalyst for C–H activation.

Results and discussion

Although our initial reaction design aimed to use azirines to form a nitrenoid intermediate that can be employed to facilitate C–N coupling, our DFT-calculations suggested that azirines cannot be used in this capacity. This result is surprising, as the steric energy stored in the three-membered ring is expected to lead to a rapid ring-opening during a reaction with metal catalysts capable of performing similar reactions with azides that are driven by the similarly favorable release of N₂. As a likely successful catalyst platform, we chose the ubiquitous Rh(III)-pentamethylcyclopentadienyl ([Cp*Rh]²⁺) platform carrying labile acetate ligands that is well-known to be an

effective catalyst for C–H activation, nitrenoid formation and C–N coupling reactions.

Fig. 1 shows the computed reaction energy profile of a plausible catalytic cycle employing the prototype substrates phenylpyridine and 2H-azirine. First, C–H activation *via* a concerted metalation deprotonation (CMD) step moderated by a Brønsted base such as acetate leads to the cyclometalation. Our calculations show in good agreement with previous work¹⁴ that this process is relatively easy with a barrier of only 17.6 kcal mol^{−1}. Ligand exchange transforms the cyclometalated intermediate **B** to the N-bound azirine complex **C**, which is predicted to ring-open the azirine readily traversing the transition state **C-TS** with a barrier of 22.8 kcal mol^{−1} to give the singlet intermediate **¹D**. It is at this juncture that the azirine substrate displays a decisive difference compared to the azides. In azides, there is a driving force to completely cleave the RN–N₂ bond and release a neutral nitrogen molecule and form a formally dianionic imido (R–N^{2−}) functionality upon oxidative coupling to the metal-center. In the case of azirine, ring-opening followed by rearrangement of the double bond can produce the vinyl-nitrene moiety that could undergo a similar oxidative coupling to the metal to produce a vinyl-imido complex, as envisioned (Scheme 1, eqn (2)). Our calculations show, however,






Fig. 1 Computed energy profile for the Cp*Rh(III)-catalyzed C–C coupling reaction using 2-phenylpyridine and 2*H*-azirine.

that azirine ring-opening and oxidative addition gives the four-membered rhodaheterocycle ${}^1\text{D}$ that is relatively high in energy and readily attacks the phenyl-carbon of the phenylpyridine ligand and rapidly inserts into the metal–carbon bond to afford the C–C coupled product **E**. The azirine ring-opening can be imagined to involve a homolytic C–N bond cleavage, which may lead to a triplet pathway after rapid spin-inversion promoted by the strong spin–orbit coupling provided by the Rh-center. We examined this possibility, but rejected it based on the much higher energy of the ${}^3\text{D}$ intermediate shown in red in Fig. 1.

As anticipated, the release of the strain energy renders this last step exergonic with the free energy of -20.9 kcal mol $^{-1}$ being assigned to **E**. Formerly an imido-complex, intermediate **E** can readily undergo proto-demetalation assisted by acetic acid, where the protonation of the imido functionality gives the corresponding imine product complex **G**, as highlighted in Fig. 1.

Fig. 2 illustrates the reaction energy profile of a putative C–N coupling mechanism. The initial phase of C–H activation of phenylpyridine is of course identical with what was shown in Fig. 1. To ring-open the azirine and facilitate the C–N bond formation with the phenylpyridine substrate, the insertion described above must be carried out by the *N*-atom of the metallaheterocycle, leading to the alternative insertion product **H**, which can be accomplished either on a singlet or a triplet pathway. Our calculations indicate that neither reaction trajectories afford a reasonable transition state, the migratory

insertion to form the C–N bond requires passage through the transition states ${}^1\text{D-TS}'$ and ${}^3\text{D-TS}$, located at 31.3 and 32.5 kcal mol $^{-1}$, giving rise to activation barriers of 36.3 and 37.5 kcal mol $^{-1}$, respectively. These energies are too high for the reaction to be viable under realistic conditions and considering the C–C bond forming reaction discussed above, the conclusion can be drawn that the azirine substrate will give exclusively C–C coupled products and C–N coupling is not possible. Thermodynamically, the C–N coupled product **H** at -21.4 kcal mol $^{-1}$ is slightly lower in energy than the C–C coupled intermediate **E**, which was found at -20.9 kcal mol $^{-1}$. And our calculations show that the final portion of the putative catalysis consisting of acetic acid promoted demetalation to form the final product is reasonable. Thus, it is the kinetic inhibition of inserting the imido functionality that makes the C–N coupling pathway impossible.

On the basis of these DFT-calculations, experimental studies were carried out using 2-phenylpyridine (**1a**) and 3-(4-methoxyphenyl)-2*H*-azirine (**2a**) in the presence of [Cp*RhCl $_2$] $_2$ with a number of additives and solvents, as summarized in Table 1. In good agreement with the mechanism discussed above, substrate **1a** (0.2 mmol, 1.0 equiv.) reacted with **2a** (1.2 equiv.) in the presence of [Cp*RhCl $_2$] $_2$ (4.0 mol%), AgSbF $_6$ (16.0 mol%), H $_2$ O (1.0 equiv.), and acetic acid (1.0 equiv.) in dichloroethane (DCE). In 3 hours at 80 °C and after workup, we obtained the acylmethylated product 1-(4-methoxyphenyl)-2-(2-(pyridin-2-yl)phenyl)ethan-1-one (**3a**) in 35% yield (entry 2), which we






Fig. 2 Computed energy profile for the Cp*Rh(III)-catalyzed C–N coupling reaction using 2-phenylpyridine and 2*H*-azirine.


propose is the hydrolyzed product of the computationally identified imine product. To test the role of acetate/acetic acid, which plays a prominent role in the computed mechanism, the experiments were repeated under identical conditions without adding acetic acid. As anticipated, no acylmethylated product was observed (entry 1).

Whereas computer models are powerful in suggesting, identifying and comparing possible reaction mechanisms, as we demonstrated above, predicting the impact of subtle environmental changes, such as the nature of the solvent or counter ions to the overall yield, is exceedingly difficult.¹⁵ Unsatisfied with the relatively low yield of only 35% in DCE, we complemented our initial findings by more classical screening efforts and identified trifluoroethanol (TFE) as the optimal solvent. Other solvents, such as DCE, MeOH, hexafluoroisopropanol (HFIP), and tetrahydrofuran (THF), gave inferior results (entries 2–6). Several counter ion additives, including AgSbF₆, AgNTf₂, and AgPF₆, were also examined and AgSbF₆ gave the desired product in 73% yield (entries 4, 7, and 8). Interestingly, when [Cp*IrCl₂]₂ (4.0 mol%) and [Cp*CoCl₂]₂ (4.0 mol%) were used as catalyst, the reaction was totally ineffective (entries 10 and 11). However, [Ru(*p*-cymene)Cl₂]₂ (4.0 mol%) provided **3a** in 65% yield (entry 12). The best result was obtained from the reaction of **1a** (2.0 equiv.) with **2a** (0.2 mmol, 1.0 equiv.) in the presence of [Cp*RhCl₂]₂ (4.0 mol%) with AgSbF₆ (16.0 mol%), water (1.0 equiv.), and AcOH (1.0 equiv.) in TFE at 80 °C for 3 h, affording **3a** in 86% yield (entry

9). Interestingly, attempts towards isolating the imine product under the anhydrous conditions were not successful (entry 13).

Next, we investigated the scope and limitation of Rh-catalyzed acylmethylation of 2-phenylpyridine (**1a**) with a wide range of 3-aryl-2*H*-azirines (**2**), as summarized in Table 2. When 3-phenyl-2*H*-azirine was treated with **1a** under the optimum reaction conditions, product **3b** was obtained in 84% yield. Electronic variations in the substituents on the aryl group of 3-aryl-2*H*-azirines (**2**) did not influence the reaction efficiency notably. Electron-donating groups, including 2-methyl, 3-methyl, and 4-methyl substituent, afforded the corresponding acylmethylated products (**3c**, **3d**, and **3e**) in high yields ranging from 88% to 91%. These results indicate that both steric and electronic effects were negligible in the case of 3-aryl-2*H*-azirines bearing electron-donating groups. Likewise, an electron-donating 4-*tert*-butyl group provided the desired pyridine (**3f**) in 73% yield. However, acylmethylation was slightly affected by electron-withdrawing groups on the aryl substituents of 3-aryl-2*H*-azirines (**2**). For example, a variety of electron-withdrawing groups, including 4-fluoro, 2-chloro, 3-chloro, 4-chloro, and 4-bromo groups, gave the corresponding acylmethylated 2-arylpyridines (**3g**, **3h**, **3i**, **3j**, and **3k**) in yields ranging from 57% to 77%. Strongly electron-withdrawing 4-trifluoromethyl, 4-nitro, and 4-ethoxycarbonyl-substituted 3-aryl-2*H*-azirines were less reactive and the products **3l**, **3m**, and **3n** were produced in moderated yields ranging from 42% to 54%. Biphenyl- and 1-naphthyl-substituted 2*H*-azirines were found to be compatible




Table 1 Experimental reaction optimization^a

Entry	Catalyst	Additive	Solvent	Yield ^b (%)
1	[Cp*RhCl ₂] ₂	AgSbF ₆	DCE	0 ^c
2	[Cp*RhCl ₂] ₂	AgSbF ₆	DCE	35
3	[Cp*RhCl ₂] ₂	AgSbF ₆	MeOH	20
4	[Cp*RhCl ₂] ₂	AgSbF ₆	TFE	73
5	[Cp*RhCl ₂] ₂	AgSbF ₆	HFIP	46
6	[Cp*RhCl ₂] ₂	AgSbF ₆	THF	28
7	[Cp*RhCl ₂] ₂	AgNTf ₂	TFE	37
8	[Cp*RhCl ₂] ₂	AgPF ₆	TFE	70
9	[Cp*RhCl ₂] ₂	AgSbF ₆	TFE	89(86) ^{d,e}
10	[Cp*IrCl ₂] ₂	AgSbF ₆	TFE	0
11	[Cp*CoCl ₂] ₂	AgSbF ₆	TFE	0
12	[Ru(<i>p</i> -cymene)Cl ₂] ₂	AgSbF ₆	TFE	65
13	[Cp*RhCl ₂] ₂	AgSbF ₆	TFE	17 ^f

^a Reactions were carried out with **1a** (0.2 mmol, 1.0 equiv.) and **2a** (1.2 equiv.) in the presence of catalyst (4.0 mol%), additive (16.0 mol%), acetic acid (1.0 equiv.), and H₂O (1.0 equiv.) in solvent (2.0 mL) at 80 °C for 3 h under a nitrogen atmosphere. ^b NMR yield using dibromomethane as an internal standard. ^c Acetic acid was not used. ^d Isolated yield. ^e **1a** (0.4 mmol, 2.0 equiv.) was used. ^f Water was not used.

with the reaction conditions, leading to the formation of **3o** and **3p** in 83% and 84% yields, respectively. The present method worked equally well with 2*H*-azirines possessing a thiophen-1-yl group to furnish **3q** in 71% yield.

Next, the substrate scope and the functional group tolerance with a variety of 2-arylpyridines (**1**) were investigated, as shown in Table 3. Modification of the substituents on the aryl ring at 2-position of 2-arylpyridines **1** did not influence the efficiency of the acylmethylation. Both electron-donating and electron-withdrawing groups on the 2-arylpyridines were compatible, thus equally affording the corresponding 2-acylmethyl-substituted arylpyridines in good yields. When 2-(*m*-tolyl)pyridine and 2-(*p*-tolyl)pyridine were treated with 3-(*p*-tolyl)-2*H*-azirine (**2e**), the desired acylmethylated products **5a** and **5b** were produced in 79% and 88% yields, respectively. The strongly electron-donating 4-methoxy group did not affect the efficiency of the reaction, and the desired product **5c** was obtained in 89% yield. The transformation of 2-arylpyridine having 3,4-methylenedioxy group was also highly facile, providing **5d** in 89% yield. Substrates with a halogen atom provided the desired compounds in good yields. Indeed, the 4-fluoro-substituted 2-phenylpyridine was reacted with 3-(*p*-tolyl)-2*H*-azirine (**2e**), furnishing the arylmethylated product **5e** in 83% yield. 2-Arylpyridines possessing 3-chloro, 4-chloro, 4-trifluoromethyl, and 4-acetyl groups were smoothly acylmethylated to produce **5f–5i** in good yields ranging from 61% to 74%. The tolerance of fluoro, chloro, and acetyl groups is very important, because these


Table 2 Scope of 3-aryl-2*H*-azirines^a

^a Reactions were carried out with **1a** (2.0 equiv.) and **2** (0.2 mmol, 1.0 equiv.) in the presence of catalyst (4.0 mol%), additive (16.0 mol%), acetic acid (1.0 equiv.), and H₂O (1.0 equiv.) in TFE (2.0 mL) at 80 °C for 3 h under a nitrogen atmosphere.

functional groups will allow further decoration and processing of the products. Pyridine substrates bearing 2-biphenyl and 2-naphthyl groups underwent the acylmethylation reaction, affording the corresponding products (**5j** and **5k**) in 74% and 83% yields, respectively. Moreover, the arene is not limited to a benzene skeleton. 2-(Thiophene-1-yl)pyridine was applied to the present Rh-catalyzed acylmethylation, resulting in the production of **5l** in 60% yield. When 2-arylpyridine possessing an estrone moiety was employed as the substrate, the acylmethylated product **5m** was produced in 80% yield.

Furthermore, tolerance of a wide range of substituents, including methyl, fluoro, ethoxycarbonyl, and acetyl, on the pyridine moiety was examined (Table 4). 5-Methyl-2-phenylpyridine underwent the acylmethylation with 3-(*p*-tolyl)-2*H*-azirine (**2e**), providing the desired product **5n** in 83% yield without notably affecting the catalytic effectiveness. Substrate bearing 4-fluoro group on the pyridine ring of 2-phenylpyridine gave rise to **5o** in acceptable yield. To our delight, the present acylmethylation proceeded despite the presence of an




Table 3 Scope of 2-arylpyridines^a

^a Reactions were carried out with **1** (2.0 equiv.) and **2e** (0.2 mmol, 1.0 equiv.) in the presence of catalyst (4.0 mol%), additive (16.0 mol%), acetic acid (1.0 equiv.), and H₂O (1.0 equiv.) in TFE (2.0 mL) at 80 °C for 3 h under a nitrogen atmosphere.

ethoxycarbonyl and acetyl group on the phenyl ring and afforded **5p** and **5q** in 67% and 68% yields, respectively.

Although the DFT-calculated mechanism discussed above serves as a useful guide for understanding our experimental results and offers a plausible concept for the reaction mechanism, there are a number of unresolved issues: (i) we assumed that hydrolysis affords the final acylmethylation product, which should be confirmed. (ii) Our DFT-calculations employ a model Rh-catalyst carrying acetate ligands for convenience to start the catalytic cycle. This putative resting state of the catalyst also serves as the reference state with a relative free energy of zero kcal mol⁻¹, which is used to evaluate the barriers of the initial C–H activation.¹⁶ This model reference state leads to a C–H activation barrier of only 17.6 kcal mol⁻¹, whereas the free energy of the azirine-bound intermediate **C** is –5.0 kcal mol⁻¹ and that affords a barrier for the concerted C–C coupling and azirine ring-opening to be 22.8 kcal mol⁻¹. It is not clear how representative and reliable these barrier estimates are. Given that the reaction typically requires 80 °C and 3 hours, these barriers seem underestimated.

To answer these mechanistic questions that are challenging to answer using computations, a series of experiments were


Table 4 Scope of 2-phenylpyridines^a

^a Reactions were carried out with **1** (2.0 equiv.) and **2e** (0.2 mmol, 1.0 equiv.) in the presence of catalyst (4.0 mol%), additive (16.0 mol%), acetic acid (1.0 equiv.), and H₂O (1.0 equiv.) in TFE (2.0 mL) at 80 °C for 3 h under a nitrogen atmosphere.

designed, as summarized in Scheme 2. First, isotopic labeling studies were performed to examine the source of oxygen in **3e**. When **1a** (2.0 equiv.) was treated with **2a** (0.2 mmol, 1.0 equiv.) in the presence of [Cp*RhCl₂]₂ (4.0 mol%) with AgSbF₆ (16.0 mol%), H₂¹⁸O (1.0 equiv.), and AcOH (1.0 equiv.) in TFE at 80 °C for 3 h, the corresponding ¹⁸O-inserted-ketone **3e**[¹⁸O] was produced in 83% yield (eqn (4)). These results confirm that the oxygen in the acylmethylation reaction originates from water, as we had assumed. Next, the kinetic isotope effect (KIE) was determined by utilizing the deuterated phenylpyridine substrate (eqn (5)). The KIE was found to be *k*_H/*k*_D = 3.13 from the intermolecular competition reaction¹⁷ between **1a** and **1a**[*d*₅]. These results indicate that the C–H cleavage at the *ortho*-position of 2-phenylpyridine is most likely involved in the rate-determining step. Although **1a**[*d*₅] was treated with **2e** under the optimum reaction conditions, the corresponding deuterium-inserted product at benzylic position was not detected (eqn (6)). This result implies that deuterium transfer through C–H activation did not occur.

The KIE of 3.13 provides significant insight that our DFT-calculations are unable to capture. As discussed above, we employ the Rh-acetate model for convenience and stoichiometric consistency during the computer simulation. For the KIE study, the chloride derivative of the Rh-catalyst is used in combination with a silver salt, which removes the chloride by forming poorly soluble silver chloride deposits. The counter anion of the silver salt has a notable impact on the yield, as we showed above. DFT-calculations are not capable of incorporating these effects accurately into the mechanistic model. One meaningful result that can be utilized is the DFT-predicted energy difference of 8.9 kcal mol⁻¹ between the product complex **G** and the recovered catalyst **A** upon release of the imine product **3a'**. Ignoring the aforementioned complications with the silver salt and the chloride ligands, the calculated energy difference between **G** and **A** suggests that after the first catalytic cycle, the product complex **G** becomes the resting state of the catalytic cycle and, thus, should be taken as the reference for evaluating the C–H activation barrier. And because the





Scheme 2 Experimental mechanistic studies.


energy of **G** is lower than that of the azirine-adduct **C**, the barrier of the C–C coupling reaction must also be referenced against **G**. The adjusted barriers for the C–H and C–C activations become 26.5 and 26.7 kcal mol⁻¹, respectively. We note that these numbers are much more consistent with the observed reaction conditions of 80 °C. And because the computed barriers are nearly identical and our DFT-calculations are fundamentally unable to offer information about the collision factor of the Arrhenius equation that would allow for computing the rate determining step, our DFT-model is unable to clearly identify the rate determining step. The KIE of 3.13 clarifies this point and strongly suggests that the C–H bond activation step is rate determining.

Scheme 3 summarizes the proposed mechanism incorporating the insights derived both from computations and experiments. We propose that the rate determining step is the C–H activation, giving rise to an experimental KIE value of 3.13 and turns the reactant complex **A** to the cyclometalated intermediate **B**. Coordination on 2*H*-azirine to Rh affords key intermediate **C**, which can ring-open the azirine *via* oxidative addition to form the rhodaheterocycle intermediate **D**. Subsequent insertion into the Rh–phenyl bond leads to the C–C coupling to give the product complex **E**. The alternative C–N coupling is found to be much less likely, as the migratory insertion for that process has a barrier that is ~15 kcal mol⁻¹ higher than the C–C coupling. The catalytic cycle closes *via* proto-demetalation with acetic acid to give the imine product **3a'**, which is readily hydrolyzed to give the acylmethylated product **3a**.

This new methodology is attractive for the synthesis of biologically relevant heterocycles such as pyridoisoindole and pyridoisoquinolinone through cyclization of the acylmethylated compounds (Scheme 4). Reactant **3j** was smoothly cyclized in


the presence of $\text{Cu}(\text{OAc})_2$ (50.0 mol%) to provide pyridoisoindole **6a** in 91% yield (eqn (7)).

Because 2-arylpyridine bearing an acyl diazo group can be easily obtained from the diazotization of the corresponding acylmethyl-substituted 2-arylpyridine, we envisioned that the intramolecular cyclization using *in situ* generated diazo intermediate should be possible in a convenient one-pot procedure. To test this idea, 1-(4-methoxyphenyl)-2-(2-(pyridin-2-yl)phenyl) ethan-1-one (**3a**), tosyl azide, DBU, and DCE were placed in a reaction vessel, and the reaction mixture was stirred at 40 °C.



Scheme 3 Proposed mechanism.





Scheme 4 Synthetic applications.

After 30 min, the reaction mixture was treated with 1.0 mol% $\text{Cu}(\text{OTf})_2$ and stirred at 80 °C for 1 h, leading to the formation of pyridoisoindole **6b** (45%) and pyridoisoquinolinone **7** (40%) through Curtius rearrangement (eqn (8)).

Conclusions

In conclusion, we developed a novel Rh-catalyzed synthetic method for a wide range of acylmethyl-substituted 2-arylpyridine derivatives using 3-aryl-2*H*-azirines as the reaction partner. The oxygen in the acylmethylation reaction was confirmed to originate from water, which hydrolyzes the initial imine product. C–H cleavage at the *ortho*-position of 2-phenylpyridine is most likely the rate-limiting step. Initially, we had aimed at developing a C–N coupling reaction, but DFT-calculations and screening results clearly suggest that C–C coupling is much faster and preferable after the azirine substrate is ring-opened at the metal center of the catalyst. The present method is highly efficient and selective, displays a broad substrate scope and a high tolerance of various functional groups, including fluoro, chloro, bromo, ketone, ester, nitro, alkyl, alkoxy, trifluoromethyl, 2-naphthyl-, and thiophen-1-yl. The mechanism was elucidated by combining computational and experimental studies, where the weaknesses and strengths of the two complementary methods of mechanistic inquiries were combined to obtain a precise and convincing mechanism for this novel reaction.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This paper is dedicated to Professor Chul-Ho Jun (Yonsei University) for his honorable retirement. This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIP) (2011-0018355 and 2017R1A4A1015405). We thank the Institute for Basic Science in Korea for financial support (IBS-R10-A1).

Notes and references

- (a) J. Royer, *Asymmetric Synthesis of Nitrogen Heterocycles*, Wiley-VCH Verlag, Weinheim, 2009; (b) X.-F. Wu, *Transition Metal-Catalyzed Heterocycle Synthesis via C–H Activation*, John Wiley & Sons, Weinheim, 2015.
- (a) D. A. Colby, R. G. Bergman and J. A. Ellman, *Chem. Rev.*, 2010, **110**, 624; (b) T. W. Lyons and M. S. Sanford, *Chem. Rev.*, 2010, **110**, 1147; (c) Z. Chen, B. Wang, J. Zhang, W. Yu, Z. Liu and Y. Zhang, *Org. Chem. Front.*, 2015, **2**, 1107.
- Y. Park, Y. Kim and S. Chang, *Chem. Rev.*, 2017, **117**, 9247.
- For selected literatures on the usage of acyl azide, see: (a) J. Ryu, J. Kwak, K. Shin, D. Lee and S. Chang, *J. Am. Chem. Soc.*, 2013, **135**, 12861; (b) J. Peng, Z. Xie, M. Chen, J. Wang and Q. Zhu, *Org. Lett.*, 2014, **16**, 4702.
- (a) T. Curtius, *Ber. Dtsch. Chem. Ges.*, 1890, **23**, 3023; (b) E. F. V. Scriven and K. Turnbull, *Chem. Rev.*, 1988, **88**, 297; (c) D. Li, T. Wu, K. Liang and C. Xia, *Org. Lett.*, 2016, **18**, 2228; (d) H. Lebel and O. Leogane, *Org. Lett.*, 2005, **7**, 4107.
- K. Shin, J. Ryu and S. Chang, *Org. Lett.*, 2014, **16**, 2022.
- For selected examples on the usage of sulfonyl azide, see: (a) E. D. Goddard-Borger and R. V. Stick, *Org. Lett.*, 2007, **9**, 3797; (b) J. Y. Kim, S. H. Park, J. Ryu, S. H. Cho, S. H. Kim and S. Chang, *J. Am. Chem. Soc.*, 2012, **134**, 9110; (c) B. Sun, T. Yoshino, S. Matsunaga and M. Kanai, *Adv. Synth. Catal.*, 2014, **356**, 1491; (d) D. Lee, Y. Kim and S. Chang, *J. Org. Chem.*, 2013, **78**, 11102; (e) B. Zhu, X. Cui, C. Pi, D. Chen and Y. Wu, *Adv. Synth. Catal.*, 2016, **358**, 326.
- For selected literature references on the usage of dioxazolone, see: (a) Y. Park, K. T. Park, J. G. Kim and S. Chang, *J. Am. Chem. Soc.*, 2015, **137**, 4534; (b) Y. Park, J. Heo, M.-H. Baik and S. Chang, *J. Am. Chem. Soc.*, 2016, **138**, 14020; (c) H. Wang, G. Tang and X. Li, *Angew. Chem., Int. Ed.*, 2015, **54**, 13049; (d) F. Wang, H. Wang, Q. Wang, S. Yu and X. Li, *Org. Lett.*, 2016, **18**, 1306; (e) Y. Liang, Y.-F. Liang, C. Tang, Y. Yuan and N. Jiao, *Chem.–Eur. J.*, 2015, **21**, 16395; (f) R. Mei, J. Loup and L. Ackermann, *ACS Catal.*, 2016, **6**, 793; (g) H. Wang, M. M. Lorion and L. Ackermann, *Angew. Chem., Int. Ed.*, 2016, **55**, 10386; (h)



- X. Wang, A. Lerchen and F. Glorius, *Org. Lett.*, 2016, **18**, 2090; (i) N. Barsu, M. A. Rahman, M. Sen and B. Sundararaju, *Chem.–Eur. J.*, 2016, **22**, 9135; (j) J. Wang, S. Zha, K. Chen, F. Zhang, C. Song and J. Zhu, *Org. Lett.*, 2016, **18**, 2062; (k) B. Jeon, U. Yeon, J.-Y. Son and P. H. Lee, *Org. Lett.*, 2016, **18**, 4610.
- 9 (a) T. Ryu, Y. Baek and P. H. Lee, *J. Org. Chem.*, 2015, **80**, 2376; (b) Y. Baek, C. Maeng, H. Kim and P. H. Lee, *J. Org. Chem.*, 2018, **83**, 2349.
- 10 See ESI for details.†
- 11 (a) A. Padwa and T. Stengel, *Tetrahedron Lett.*, 2004, **45**, 5991; (b) F. Palacios, A. M. O. de Retana, E. M. de Marigorta and J. M. de los Santos, *Eur. J. Org. Chem.*, 2001, **2001**, 2401; (c) S. Chiba, G. Hattori and K. Narasaka, *Chem. Lett.*, 2007, **36**, 52; (d) D. A. Candito and M. Lautens, *Org. Lett.*, 2010, **12**, 3312; (e) T. Izumi and H. Alper, *Organometallics*, 1982, **1**, 322.
- 12 (a) H. M. L. Davies and J. R. Manning, *Nature*, 2008, **451**, 417; (b) J. L. Roizen, M. E. Harvey and J. D. Bois, *Acc. Chem. Res.*, 2012, **45**, 911; (c) S. Jana, M. D. Clements, B. K. Sharp and N. Zheng, *Org. Lett.*, 2010, **12**, 3736.
- 13 (a) M. Barday, C. Janot, N. H. Halcovitch, J. Muir and C. Aïssa, *Angew. Chem., Int. Ed.*, 2017, **56**, 13117; (b) Y. Xu, X. Zhou, G. Zheng and X. Li, *Org. Lett.*, 2017, **19**, 5256; (c) H. Oh, S. Han, A. K. Pandey, S. H. Han, N. K. Mishra, S. Kim, R. Chun, H. S. Kim, J. Park and I. S. Kim, *J. Org. Chem.*, 2018, **83**, 4641; (d) G. Zheng, M. Tian, Y. Xu, X. Chen and X. Li, *Org. Chem. Front.*, 2018, **5**, 998; (e) S. Ji, K. Yan, B. Li and B. Wang, *Org. Lett.*, 2018, **20**, 5981.
- 14 (a) S. H. Park, J. Kwak, K. Shin, J. Ryu, Y. Park and S. Chang, *J. Am. Chem. Soc.*, 2014, **136**, 2492; (b) D. L. Davies, S. M. A. Donald, O. Al-Duaij, S. A. Macgregor and M. Pölleth, *J. Am. Chem. Soc.*, 2006, **128**, 4210; (c) D. L. Davies, S. M. A. Donald, O. Al-Duaij, J. Fawcett, C. Little and S. A. Macgregor, *Organometallics*, 2006, **25**, 5976; (d) H. M. L. Davies and D. Morton, *Chem. Soc. Rev.*, 2011, **40**, 1857; (e) S. H. Cho, J. Y. Kim, J. Kwak and S. Chang, *Chem. Soc. Rev.*, 2011, **40**, 5068; (f) L. Li, W. W. Brennessel and W. D. Jones, *Organometallics*, 2009, **28**, 3492; (g) D. Balcells, E. Clot and O. Eisenstein, *Chem. Rev.*, 2010, **110**, 749; (h) H. M. L. Davies and D. Morton, *Chem. Soc. Rev.*, 2011, **40**, 1857; (i) G. Rouquet and N. Chatani, *Angew. Chem., Int. Ed.*, 2013, **52**, 11726.
- 15 (a) D. J. Tantillo, *Applied Theoretical Organic Chemistry*, World Scientific Publishing Europe Ltd, London, 2018; (b) S. M. Bachrach, *Computational Organic Chemistry*, John Wiley & Sons, Inc, Hoboken, New Jersey, 2014.
- 16 S. Kozuch and S. Shaik, *Acc. Chem. Res.*, 2011, **44**, 101.
- 17 (a) W. D. Jones, *Acc. Chem. Res.*, 2003, **36**, 140; (b) F. Kakiuchi, Y. Matsuura, S. Kan and N. Chatani, *J. Am. Chem. Soc.*, 2005, **127**, 5936.

