Chemical Science

EDGE ARTICLE

Check for updates

Cite this: Chem. Sci., 2019, 10, 3987

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 30th January 2019 Accepted 21st February 2019

DOI: 10.1039/c9sc00545e

rsc.li/chemical-science

Introduction

Nitrogen-containing heterocyclic compounds are prevalent in natural products and bioactive molecules. They are also found in a large number of drugs used to combat a broad range of diseases (Scheme 1a). Advances in transition-metal catalyzed C-H functionalization have significantly streamlined the synthesis of a plethora of heterocyclic molecules.¹ In many cases, the heteroatom directing group participated in the annulation reaction besides offering chelation assistance, which calls for sufficient interactions between the metal center and the directing group. In this context, C-H bond activation catalyzed by high valent Rh(m) complexes has stood out as a powerful strategy for step-economical construction of C-C bonds and has become one of the increasingly important

‡ D. B. and J. X. contributed equally to this work.

Rhodium(\mathfrak{m})-catalyzed diverse [4 + 1] annulation of arenes with 1,3-enynes *via* sp³/sp² C–H activation and 1,4-rhodium migration[†]

Dachang Bai, ^[10] ‡^a Jintao Xia, ‡^{bd} Fangfang Song,^a Xueyan Li,^a Bingxian Liu, ^[10] ^a Lihong Liu,^a Guangfan Zheng,^c Xifa Yang,^b Jiaqiong Sun*^c and Xingwei Li

Nitrogen-rich heterocyclic compounds have a profound impact on human health. Despite the numerous synthetic methods, diversified, step-economic, and general synthesis of heterocycles remains limited. C-H bond functionalization catalyzed by rhodium(iii) cyclopentadienyls has proven to be a powerful strategy in the synthesis of diversified heterocycles. Herein we describe rhodium(iii)-catalyzed sp² and sp³ C-H activation-oxidative annulations between aromatic substrates and 1,3-enynes, where alkenyl-to-allyl 1,4-rhodium(iii) migration enabled the generation of electrophilic rhodium(iii) π -allyls *via* remote C-H functionalization. Subsequent nucleophilic trapping of these species by various sp²-hybridized N-nucleophiles delivered three classes (external salts, inner salts, and neutral azacycles) of five-membered azacycles bearing a tetrasubstituted saturated carbon center, as a result of [4 + 1] annulation with the alkyne being a one-carbon synthon. All the reactions proceeded under relatively mild conditions with broad substrate scope, high efficiency, and excellent regioselectivity. The synthetic applications of this protocol have also been demonstrated, and experimental studies have been performed to support the proposed mechanism.

strategies in organic syntheses.² In these systems, a large scope of directing groups can be accommodated to ensure high activity and selectivity of the C-H bond. In annulation reactions, development of mild and straightforward C-H activation processes assisted by a multifunctional directing group is especially attractive, allowing realization of molecular diversity and synthetic versatility.3 On the other hand, alkynes are widely used as a typical unsaturated coupling partner in numerous annulation reactions, and in most cases they react as a twocarbon synthon.4 To move beyond this limitation, the Lam group pioneered excellent work in oxidative coupling of arenes bearing a nucleophilic directing group and 1,3-envnes,^{5,6} which occurred via 1,4-rhodium migration and Rh(III) allyl intermediates, leading to [n + 3] and [n + 1] annulation in most cases. Still, the [n + 1] type annulation with the alkyne being a C1 synthon is still rare,^{5a,c} where the nucleophilic directing group has been limited to 1,3-dione, acidic OH groups, and imide groups, and the corresponding $C(sp^3)$ -H activation has not been realized.

On the other hand, quaternary ammonium salts are versatile building blocks for a number of naturally occurring products and functional materials.⁷Although a number of methods for the synthesis of such ionic compounds are available, many are limited by the lack of generality, limited functional group tolerance, and lengthy synthetic procedures.⁸ Recently, quaternary ammonium salts have been synthesized through sp² C–H activation of arenes and annulations with alkynes or diazo substrates in the presence of stoichiometric amounts of metal

This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.

Open Access Article. Published on 26 February 2019. Downloaded on 8/10/2025 11:32:58 PM.



View Article Online

View Journal | View Issue

^aHenan Key Laboratory of Organic Functional Molecule and Drug Innovation, School of Chemistry and Chemical Engineering, Henan Normal University, Xinxiang 453007, China

^bDalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, China. E-mail: xwli@dicp.ac.cn

^cSchool of Chemistry and Chemical Engineering, Shaanxi Normal University (SNNU), Xi'an 710062, China. E-mail: sunjq295@nenu.edu.cn

^dUniversity of Chinese Academy of Sciences, Beijing 100049, China

[†] Electronic supplementary information (ESI) available. CCDC 1861504 and 1457168. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c9sc00545e



Scheme 1 Rh(III)-catalyzed oxidative annulation of 1,3-enynes or alkynes. (a) Some drugs with isoindoline or the quaternary ammonium motif. (b) Quaternary ammonium salts formed *via* coupling with alkynes. (c) Synthesis of diversified heterocycles bearing a quaternary carbon center *via* coupling of arenes and 1,3-enynes.

oxidants/additives, as has been pioneered by Cheng and others. While quaternary ammonium salts could be synthesized through the [n + 2] reaction with alkynes, the synthesis of quaternary ammonium salts9 bearing a quaternary carbon center is limited because of the challenge of C(tertiary)-N reductive elimination,4,10 an inner-sphere reaction pathway (Scheme 1b). However, it is challenging to construct a quaternary carbon center through an outer-sphere mechanism. Herein, we report diverse synthesis of four classes of nitrogencontaining heterocycles via C-H activation, 1,4-rhodium(III) migration, and allyl-to-allyl rearrangement. Importantly, quaternary ammonium salts can be readily accessed via Rh(III)catalyzed sp² and sp³ C-H functionalization with Cu(OAc)₂/air as the oxidant. Besides ionic products, zwitterionic and neutral N-containing heterocycles bearing a quaternary carbon can also be accessed when oximes and ketimines are employed as the arene source. All the coupling systems have been realized with high regio- and chemoselectivity (Scheme 1c).

Results and discussion

We commenced our studies by exploring the reaction parameters of the coupling of 2-phenylpyridine (1a) and 1,3-enyne (3a, see ESI Table 1[†]). An excellent yield of quaternary ammonium salt 4aa was obtained with $Cp*Rh(OAc)_2$ as a catalyst, Cu(II)/airas the oxidant, and KPF_6 as the anion source under mild conditions. With the optimized conditions in hand, we next investigated the scope and generality of this coupling system (Scheme 2). Various 1,3-enynes were tolerated (4aa-4ai, 61– 98%), and the identity of product 4ae has been confirmed by



Scheme 2 Scope of the coupling of 2-phenylpyridines with 1,3-enynes. Reaction conditions: arene 1 or 2 (0.2 mmol), enyne 3 (0.22 mmol), [Cp*Rh(OAc)₂] (8 mol%), Cu(OAc)₂ (50 mol%), and KPF₆ (0.5 mmol) in CF₃CH₂OH (2.0 mL) under air, 50 °C, monitored by TLC, isolated yields. See general procedure A.

X-ray crystallography (CCDC 1861504[†]). However, 1,3-enyne 3h containing an alkylamino moiety showed poor reactivity. The phenyl-substituted 1,3-enyne 3r reacted with poor regio- and chemoselectivity, and the [4 + 2] annulation side product was isolated in 45% yield.4m 2-Phenylpyridines bearing Me, electrondonating, electron-withdrawing, and bromo groups were fully tolerated (4ba-4ia, 59-95%). We also investigated preliminarily the enantioselectivity of the reactions of 1a and 3a using chiral (R)-Rh1, and so far, only <5% of 4aa was obtained under standard conditions. The reaction also worked well for benzo[h]quinoline, and the product 4ja was isolated in 66% yield. The arene was not limited to 2-arylpyridines. Thus, the coupling for N-phenylpyrimidin-2-amine and N-phenylpyridin-2-amine afforded the ionic products 4la and 4ka in good yields, respectively. Moreover, several N-pyridylisoquinolones were also viable substrates, providing the desired products (5aa-5ca) in good to excellent yields (75-95%). In all cases, the well-explored [n+2] annulation side products were not detected.

To better define the scope of this reaction system, we next extended the reaction system to 8-methylquinolines (Scheme 3). To our delight, the quaternary ammonium salt **7aa** was obtained in 82% yield *via* benzylic C–H activation, and AgOAc and



Scheme 3 Scope of the coupling of 8-methylquinolines with 1,3enynes. Reaction conditions: 6 (0.2 mmol), 3 (0.3 mmol), $[Cp*Rh(OAc)_2]$ (8 mol%), AgSbF₆ (0.2 mmol), AgOAc (0.3 mmol), DCE (2.0 mL), monitored by TLC, isolated yields. see general procedure B.

AgSbF₆ were identified as the optimal oxidant and anion source (see ESI Table 2†). The scope of this sp³ C–H coupling system was examined under the optimal conditions. The scope of the 1,3-enynes is reasonably broad (**7aa-7af**, 52–82%) although several heteroatom-functionalized enynes (**3g** and **3h**) were inactive. Various substituted 8-methylquinolines bearing methyl, halogen, and methoxy groups at the 5- and 6-positions all reacted with moderate to good yields (**7ba-7ka**, 53–83%). 7-Substituted and 3,5-disubstituted quinolines also reacted smoothly but with slightly lower yields (**7la-7ma**, 46–52%)

To our delight, the arene substrates of this [4 + 1] coupling reaction system could be smoothly extended to oximes (Scheme 4) and ketimines (Scheme 5, see ESI Table 3[†]).¹¹ The coupling of readily available ketoximes afforded zwitterionic isoindole 2-oxides. The generality of this system was examined for oximes with different substituents at the para position using enyne 3b as a coupling reagent. Alkyl-substituted oximes gave good to excellent yields (57-96%, 9ab-9db) while relatively lower efficiency (9ib) was found when an electron-donating group (OMe) was present. Furthermore, an isoquinoline 9ib', resulting from redox-neutral [4 + 2] annulation, was also detected as a byproduct.4g Introduction of electron-withdrawing groups, such as -OCF₃ (9fb), -CF₃ (9gb), and -NO₂ (9hb) all resulted in high efficiency, indicative of tolerance of electronic effect. We next evaluated the compatibility of halogen groups, and 9jb-9mb were delivered in good yields. Moreover, the reaction also proceeded smoothly when various meta substituents were present (9nb-9pb, 46–75%). High regioselectivity at the less hindered site was obtained for meta Cl- and Br-substituted oximes, while nearly equal amounts of two regioisomeric products were



Scheme 4 Scope of the coupling of oximes with 1,3-enynes.^{a,b} ^aReaction conditions A: **8** (0.20 mmol), **3** (0.22 mmol), $[Rh*CpCl_2]_2$ (4.0 mol%), Cu(OAc)₂ (2.1 equiv.), MeOH (2 mL) at 40 °C under N₂ for 10 h. ^bIsolated yields. ^cReaction was performed with 4 mmol of **8** at 2.5 mol% catalyst loading. ^dIsoquinoline **9ib**', a [4 + 2] annulation product, was also isolated in 10% yield (see ESI†). ^eregioselectivity. See general procedure C.

obtained for the meta MeO-substituent. Besides, introduction of ortho Me (9qb) was also tolerated. Doubly substituted arenes at the meta and para positions were also tolerated, providing products 9rb-9ub in 55-85% yields. Other oximes 8v, 8w and 8x also worked well to yield 9vb-9xb in good yields (50-85%). The ketoxime (8a) was next evaluated in reaction with a range of 1,3envnes. 1,3-Envnes with the *n*-butyl substituent at the alkyne terminus gave an inferior result (9aa, 12%), probably due to the poor reactivity of such substrates, while unprotected or silvlprotected alcohol substituents were tolerated, affording the Noxide in high yields (9aj-9al, 82-89%). However, alkylaminosubstituted 1,3-envne 3h showed poor reactivity. Variation of the group *trans* to the alkyne from phenyl (3m) to methyl (3a) was also successful. We also examined 1,3-enyne 3n that contains a *trans* butenyl group, giving high E selectivity albeit with lower yield (9an, 14%). Furthermore, 1,3-enyne 3o containing a cyclohexyl group was also an effective coupling partner with 94% yield (9ao). To confirm whether the allylic hydrogen atom cis to the alkyne is necessary, ketoxime 8a was allowed to react with 1,3-envnes 3p and 3q. It was found that reaction with **3p** gave the [4 + 1] annulation product in 46% yield. However, no product was obtained when 3q was employed as a coupling partner. Those results demonstrated that allylic hydrogen cis to the alkyne in the 1,3-enyne is crucial for this reaction, which is consistent with a 1,4-Rh migration pathway.

Extension of the oxime to ketimine substrate was next explored. We reasoned that the oxidative [4 + 1] annulation may



Scheme 5 Oxidative annulation of ketimines with 1,3-enynes and sequential hydrogenation. Reaction conditions: imine **10** (0.20 mmol), enyne **3** (0.22 mmol), [RhCp*Cl₂]₂ (4.0 mol%), AgOAc (2.1 equiv.), HFIP (2 mL) at 100 °C under N₂ for 10 h; NaBH(OAc)₃ (2.5 equiv.)/HOAc (20 equiv.) was sequentially added and was kept for 30 min at room temperature, followed by quenching with NaOH solution. Isolated yields. See general procedure D.

afford an iminium salt. However, using a ketimine of acetophenone in the presence of a base may lead to further deprotonation of the putative iminium species, generating an enamine that is probably not air-stable. Indeed, reaction of ketoimine 10a and 1,3-envne 3b under streamlined conditions delivered 11ab' in excellent yield as an air-sensitive oil (see ESI Fig. 9[†]). To facilitate product isolation, intermediate 11ab' was hydrogenated using NaBH(OAc)₃,¹² which provided stable amine 11ab in 74% total yield albeit with low diastereoselectivity (Scheme 5). Next, various 1,3-enynes were tested through this two-step process. It was found that 1,3-enynes terminated with an *n*-butyl or alkylamino group failed to react (11aa and 11ah), while several other 1,3-enynes coupled in good yields but with low diastereoselectivity (11aj-11am, 11ao). Delightfully, the coupling with 1,3-envne containing a trans butenyl group (3n) provided excellent diastereoselectivity and high yield (11an). Therefore, the introduction of a methyl substituent at the alkenyl carbon had a positive effect on the diastereoselectivity in the hydrogenation step. Meanwhile, the identity of 11an was unambiguously confirmed by X-ray crystallography (CCDC 1457168[†]). Next, various electron-donating, electron-withdrawing, and halogen substituents at the para position of phenyl were examined by coupling with 3n, from which >20:1 dr and moderate to good yields were obtained (11bn-11jn). In addition, the arene substrates have been extended to disubstituted and heteroaryl ketimines, leading to 11kn-11mn in high diastereoselectivity and good yields.

Besides, a trimethyl-substituted 1,3-enyne was also evaluated in the reaction with ketamine **10a**, affording **11br** with good yield and diastereoselectivity.

Synthetic applications

Several derivatization reactions of an N-oxide product **9ab** have been performed to briefly demonstrate the synthetic utility of the coupling system. First, a gram-scale synthesis of **9ab** has been realized in 59% yield under reduced catalyst loading (Scheme 6a). Reduction of **9ab** by Zn delivered the corresponding isoindole **12** in 59% yield (Scheme 6b). Furthermore, hydrogenation of conjugated diene using Pd/C as a catalyst gave **13** in 95% yield with retention of the N–O group (Scheme 6b). Diels–Alder reaction of **1**,3-diene **9ab** and *N*-phenylmaleimide in toluene at 80 °C proceeded smoothly to give adduct **14** in 77% yield with >95 : 5 endo : exo selectivity. In addition, **9ab** underwent smooth oxidative cleavage when treated with NaIO₄ to deliver α , β -unsaturated ketone **15** in 45% yield (Scheme 6b).

Mechanistic investigations

A series of experiments have been conducted to probe the reaction mechanism. Rhodacyclic complex **16** has been prepared from cyclometalation of oxime **8a**, and it proved to be an active catalyst for the coupling of **8a** and **3b** (Scheme 7a), indicating relevancy of C–H activation. A kinetic isotope effect (KIE) value of 9.0 was then obtained in the competitive coupling of **8a** and **8a**-d₅ with **3b** under the standard conditions at a low conversion (Scheme 7b). A large KIE value was also obtained in the competitive coupling between **6a** and **6a**-d₃ with **3a** (see ESI Fig. 5†). These results indicated that cleavage of the C–H bond is likely involved in the turnover-limiting step of all the above coupling systems. When an equimolar mixture of **8d** and **8g** was allowed to competitively couple with **3b**, products **8dn** and **8gb** were obtained in a 1 : 2.2 ratio, indicating that an electron-poor oxime reacted at a slightly higher rate (Scheme 7c).

To understand the 1,4-Rh migration process, a reaction of **8a** and hexadeuterated 1,3-enyne $3\mathbf{b}$ -d₆ was then conducted (Scheme 7d). Observation of partial deuteration (83% D) at the



Scheme 6 Applications of the coupling reaction. (a) Reaction on a gram scale. (b) Derivations of a coupled product through chemo-selective reduction, Diels-Alder reaction and oxidative cleavage.

(a) Catalytic reactivity of rhodacyclic complex 16 16 (8 mol %) ndard conditio 76% (b) Kinetic isotope effect (competition experiment OF KIF = 9.010 (c) Comp conditions = 1 : 2.2 (d) Reaction with deuterio-enyne [D]6-3 standard condition: 23 min > 95% D D₃C > 95% D 77% 8a. 0.2 mmol [D]_-3b. 0.22 mmo 83% D 95% D (e) Control experimer [Cp*RhCl_]_/AqOAq not detected

Scheme 7 Mechanistic studies. (a) Catalytic reactivity of a rhodacyclic intermediate. (b) Kinetic isotope effect. (c) Competition experiment. (d) Reaction with deuterio-enyne [D]6-3b. (e) Control experiment.

alkenyl carbon close to the quaternary center and slight loss of deuterium at the alkene terminus positions (95% D) suggested relevancy of reversible 1,4-Rh migration and this migration may occur under acetate-assistance (H_a), and this result is consistent with those in Lam's work.^{5,6} The partial deuteration (61% D) of H_b may suggest reversible protonolysis at the allylic position in our system (see ESI Fig. 10† for another proposed mechanism).¹³ Interestingly, the total number of deuterium atoms in $[D]_n$ -**9ab** is greater than that in **3b**, and this may result from additional H/D exchange with the $[D]_6$ -**3b**. Additionally, a control experiment (Scheme 7e) revealed that an NH benzyl amine failed to react under similar conditions, indicating that the reaction had to be performed by a two-step process.

On the basis of mechanistic studies and previous reports,¹⁴ a plausible catalytic cycle is proposed in Scheme 8 for the representative coupling of **8a** and **3b**. Starting from an active $[Cp*Rh(OAc)_2]$ species (A), cyclorhodation of oxime **8a** gives a five-membered rhodacycle **B**. Coordination of the incoming 1,3-enyne **3b** and regioselective migratory insertion of Rh-C(aryl) afford a Rh(m) alkenyl intermediate C, and the regioselective insertion of this step is largely controlled by electronic effects of the biased alkyne.^{4m} Subsequently, reversible 1,4-Rh(m) migration occurs under acetate-assisted concerted metalation-deprotonation to generate intermediate **D**, and subsequent protonolysis may produce **E**. Allyl-to-allyl rearrangement of **F** generates a π -allylrhodium species **G**, and nucleophilic attack of the directing group at π -allyl carbon provides the



Scheme 8 Proposed mechanism.

conjugate acid of the [4 + 1] annulation product, deprotonation of which delivers the final product **9ab**. In the case of 2-phenylpyridine and 8-methylquinoline, the resulting ionic product is stable enough for isolation. The Rh(I) intermediate generated from the nucleophilic attack is reoxidized by Cu(II)/O₂ to regenerate the active rhodium(III) species for the next catalytic cycle.

Conclusions

In summary, we have realized Rh(m)-catalyzed diversified oxidative annulation of 2-arylpyridines, 8-methylquinolines, oximes, and ketimines with 1,3-enynes through alkenyl-to-allyl 1,4-rhodium(m) migration. The C–N coupling/annulation occurred *via* the nucleophilic attack of the sp²-hybridized Nnucleophiles. Various ionic, zwitterionic, and neutral Ncontaining five-membered azacycles bearing a quaternary carbon center have been efficiently synthesized. Mechanistic studies including KIE and competition experiments have been performed and supported the proposed mechanism. Further studies on the C–H activation of other arenes that highlight the unique role of 1,3-enynes are currently underway in our laboratories.

Conflicts of interest

The authors declare no competing financial interests.

Acknowledgements

The NSFC (No. 21525208 and 21801067), research fund from Henan Normal University (5101034011009), the Education Department of Henan Province Natural Science Research Program (18A150010), and start-up fund from Henan Normal University (qd17108) are gratefully acknowledged.

Notes and references

- 1 (a) V. Ritleng, C. Sirlin and M. Pfeffer, Chem. Rev., 2002, 102, 1731-1770; (b) O. Daugulis, H.-Q. Do and D. Shabashov, Acc. Chem. Res., 2009, 42, 1074-1086; (c) X. Chen, K. M. Engle, D. H. Wang and J. Q. Yu, Angew. Chem., 2009, 48, 5094-5115; (d) D. A. Colby, R. G. Bergman and J. A. Ellman, Chem. Rev., 2010, 110, 624-655; (e) T. W. Lyons and M. S. Sanford, Chem. Rev., 2011, 110, 1147-1169; (f) C. S. Yeung and V. M. Dong, Chem. Rev., 2011, 111, 1215-1292; (g) S. H. Cho, J. Y. Kim, J. Kwak and S. Chang, Chem. Soc. Rev., 2011, 40, 5068-5083; (h) L. Ackermann, Chem. Rev., 2011, 111, 1315-1345; (i) J. Wencel-Delord, T. Droge, F. Liu and F. Glorius, Chem. Soc. Rev., 2011, 40, 4740-4761; (j) W. R. Gutekunst and P. S. Baran, Chem. Soc. Rev., 2011, 40, 1976–1991; (k) L. McMurray, F. O'Hara and M. J. Gaunt, Chem. Soc. Rev., 2011, 40, 1885-1898; (l) J. Yamaguchi, A. D. Yamaguchi and K. Itami, Angew. Chem., Int. Ed., 2012, 51, 8960-9009; (m) B.-J. Li and Z.-J. Shi, Chem. Soc. Rev., 2014, 41, 5588-5598; (n) C. Zheng and S.-L. You, RSC Adv., 2014, 4, 6173-6214; (o) X.-X. Guo, D.-W. Gu, Z.-X. Wu and W.-B. Zhang, Chem. Rev., 2015, 115, 1622-1651; (p) L. Yang and H. M. Huang, Chem. Rev., 2015, 115, 3468-3517; (q) T. Gensch, M. N. Hopkinson, F. Glorius and J. Wencel-Delord, Chem. Soc. Rev., 2016, 45, 2900-2936.
- 2 (a) D. A. Colby, A. S. Tsai, R. G. Bergman and J. A. Ellman, Acc. Chem. Res., 2012, 45, 814–825; (b) N. Kuhl, N. Schroder and F. Glorius, Adv. Synth. Catal., 2014, 356, 1443–1460; (c) G.-Y. Song and X.-W. Li, Acc. Chem. Res., 2015, 48, 1007–1020.
- 3 (a) Y. Yang, K. Li, Y. Cheng, D. Wan, M. Li and J. You, Chem. Commun., 2016, 52, 2872–2884; (b) T. Piou and T. Rovis, Acc. Chem. Res., 2018, 51, 170–180; (c) N. Guimond, C. Gouliaras and K. Fagnou, J. Am. Chem. Soc., 2010, 132, 6908–6909; (d) F. W. Patureau, T. Besset, N. Kuhl and F. Glorius, J. Am. Chem. Soc., 2011, 133, 2154–2156; (e) B. J. Li, H. Y. Wang, Q. L. Zhu and Z. J. Shi, Angew. Chem., Int. Ed., 2012, 51, 3948–3952; (f) M.-B. Zhou, R. Pi, M. Hu, Y. Yang, R. J. Song, Y. Z. Xia and J. H. Li, Angew. Chem., Int. Ed., 2014, 53, 11338–11341; (g) T. Zhou, Y. W. Wang, B. Li and B. Q. Wang, Org. Lett., 2016, 18, 5066–5069; (h) R. T. Stemmler and C. Bolm, Adv. Synth. Catal., 2007, 349, 1185–1198.
- 4 (a) T. Satoh and M. Miura, Chem.-Eur. J., 2010, 16, 11212-11222; (b) T. Satoh and M. Miura, Org. Lett., 2007, 9, 1407-1409; (c) D. A. Colby, R. G. Bergman and J. A. Ellman, J. Am. Chem. Soc., 2008, 130, 3645-3651; (d) L. Li, W. W. Brennessel and W. D. Jones, J. Am. Chem. Soc., 2008, 130, 12414-12419; (e) D. R. Stuart, M. Bertrand-Laperle, K. M. N. Burgess and K. Fagnou, J. Am. Chem. Soc., 2008, 130, 16474-16475; (f) N. Umeda, H. Tsurugi, T. Satoh and M. Miura, Angew. Chem., Int. Ed., 2008, 47, 4019-4022; (g) N. Guimond and K. Fagnou, J. Am. Chem. Soc., 2009, 131, 12050-12051; (h) P. C. Too, Y.-F. Wang and S. Chiba, Org. Lett., 2010, 12, 5688-5691; (i) D. R. Stuart, P. Alsabeh, M. Kuhn and K. Fagnou, J. Am. Chem. Soc., 2010, 132, 18326-18339; (j) K. Muralirajan, K. Parthasarathy and

C.-H. Cheng, Angew. Chem., Int. Ed., 2011, 50, 4169-4172; (k) N. Umeda, K. Hirano, T. Satoh, N. Shibata, H. Sato and M. Miura, J. Org. Chem., 2011, 76, 13-24; (l) H. Yan, H.-L. Wang, X.-C. Li, X.-Y. Xin, C.-X. Wang and B.-S. Wan, Angew. Chem., Int. Ed., 2015, 54, 10613-10617; (m) M. P. Huestis, L. Chan, D. R. Stuart and K. Fagnou, Angew. Chem., Int. Ed., 2011, 50, 1338-1341; (n) D. Ghoraia and J. Choudhury, Chem. Commun., 2014, 50, 15159-15162; (o) D. Ghorai and J. Choudhury, ACS Catal., 2015, 5, 2692-2696; (p) D. L. Davies, C. E. Ellul, S. A. Macgregor, C. L. McMullin and K. Singh, J. Am. Chem. Soc., 2015, 137, 9659-9669; (q) P. Gandeepan and C. -H. Cheng, Chem.-Asian J., 2016, 11, 448-546; (r) J. L. Mascareñas, Angew. Chem., Int. Ed., 2016, 55, 11000-11019; (s) C.-Q. Wang, L. Ye, C. Feng and T.-P. Loh, J. Am. Chem. Soc., 2017, 139, 1762-1765.

- 5 (a) D. J. Burns and H. W. Lam, Angew. Chem., Int. Ed., 2014,
 53, 9931–9935; (b) D. J. Burns, D. Best, M. D. Wieczysty and
 H. W. Lam, Angew. Chem., Int. Ed., 2015, 54, 9958–9962; (c)
 J. D. Dooley and H. W. Lam, Chem.-Eur. J., 2018, 24, 4050–4054; (d) S. E. Korkis, D. J. Burns and H. W. Lam, J. Am. Chem. Soc., 2016, 138, 12252–12257.
- 6 (a) H. B. Hepburn and H. W. Lam, Angew. Chem., Int. Ed., 2014, 53, 11605–11610; (b) M. Callingham, B. M. Partridge, W. Lewis and H. W. Lam, Angew. Chem., Int. Ed., 2017, 56, 16352–16356; (c) B. M. Partridge, M. Callingham, W. Lewis and H. W. Lam, Angew. Chem., Int. Ed., 2017, 56, 7227–7232.
- 7 (a) J. E. Anthony, Angew. Chem., Int. Ed., 2008, 47, 452–483;
 (b) J. Fortage, C. Peltier, F. Nastasi, F. Puntoriero, F. Tuyèras, S. Griveau, F. Bedioui, C. Adamo, I. Ciofini and S. Campagna, J. Am. Chem. Soc., 2010, 132, 16700–16713.
- 8 (a) A. Núñez, A. M. Cuadro, J. Alvarez-Builla and J. J. Vaquero, Org. Lett., 2009, 9, 2977–2980; (b) T. Fukutani, N. Umeda, K. Hirano, T. Satoh and M. Miura, Chem. Commun., 2009, 5141–5143; (c) K. S. Stowers, C. Kevin, K. C. Fortner and M. S. Sanford, J. Am. Chem. Soc., 2011, 133, 6541–6544.
- 9 (a) J. Jayakumar, K. Parthasarathy and C. H. Cheng, Angew. Chem., Int. Ed., 2012, 51, 197-220; (b) D. Zhao, Q. Wu, X. Huang, F. Song, T. Lv and J. You, Chem.-Eur. J., 2013, 19, 6239–6244; (c) N. Senthilkumar, P. Gandeepan, J. Jayakumar and C. H. Cheng, Chem. Commun., 2014, 50, 3106-3108; (d) N. S. Upadhyay, J. Jayakumar and C. H. Cheng, Chem. Commun., 2017, 53, 2491-2494; (e) Y. R. Han, S.-H. Shim, D.-S. Kim and C.-H. Jun, Org. Lett., 2018, 20, 264-267; (f) J. Jayakumar and C.-H. Cheng, Chem.-Eur. J., 2016, 22, 1800-1804; (g) D. Sucunza, A. M. Cuadro, J. Alvarez-Builla and J. J. Vaquero, J. Org. Chem., 2016, 81, 10126–10135; (h) Y. Zhao, S. Li, X. Zheng, J. Tang, Z. She, G. Gao and J. You, Angew. Chem., Int. Ed., 2017, 56, 4286-4289; (i) G.-Y. Zhang, L. Yang, Y.-Y. Wang, Y.-J. Xie and H.-M. Huang, J. Am. Chem. Soc., 2013, 135, 8850-8853.
- 10 X. Yang, S. Liu, S. Yu, L. Kong, Y. Lan and X. Li, *Org. Lett.*, 2018, **20**, 2698–2701.
- 11 (a) Z. Shi, D. C. Koester, M. Boultadakis-Arapinis and F. Glorius, J. Am. Chem. Soc., 2013, 135, 12204–12207; (b)

Q. Wang, F. Wang, X. Yang, X. Zhou and X. Li, *Org. Lett.*, 2016, **18**, 6144–6147.

- 12 W. Zhang, P. Chen and G. Liu, *Angew. Chem., Int. Ed.*, 2017, **56**, 5336–5340.
- 13 (*a*) A. Lerchen, T. Knecht, M. Koy, J. B. Ernst, K. Bergander, C. G. Daniliuc and F. Glorius, *Angew. Chem., Int. Ed.*, 2018,

57, 15248–15252; (b) T. A. F. Nelson and S. B. Blakey, Angew. Chem., Int. Ed., 2018, 57, 14911–14915.

14 (a) S. Ma and Z. Gu, Angew. Chem., Int. Ed., 2005, 44, 7512–7517; (b) Y. Ikeda, K. Takano, S. Kodama and Y. Ishii, Chem. Commun., 2013, 49, 11104–11106; (c) Y. Ikeda, K. Takano, M. Waragai, S. Kodama, N. Tsuchida, K. Takano and Y. Ishii, Organometallics, 2014, 33, 2142–2145.

Edge Article