

Cite this: *Chem. Sci.*, 2019, 10, 5787

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 5th February 2019

Accepted 4th May 2019

DOI: 10.1039/c9sc00640k

rsc.li/chemical-science


# Cyano-borrowing reaction: nickel-catalyzed direct conversion of cyanohydrins and aldehydes/ketones to $\beta$ -cyano ketone†

Zhao-Feng Li, Qian Li, Li-Qing Ren, Qing-Hua Li, Yun-Gui Peng\* and Tang-Lin Liu \*

A direct nickel-catalyzed, high atom- and step-economical reaction of cyanohydrins with aldehydes or ketones *via* an unprecedented “cyano-borrowing reaction” has been developed. Cleavage of the C–CN bond of cyanohydrins followed by aldol condensation and conjugate addition of cyanide to  $\alpha,\beta$ -unsaturated ketones proceeded to deliver a range of racemic  $\beta$ -cyano ketones with good to high yields. The practical procedure with the use of a commercial and less-toxic CN source bodes well for wide application of this protocol.

## Introduction

Nitriles are important synthesis intermediates in transformation<sup>1</sup> and are key components in various natural products, medicinal pharmacophores and drugs,<sup>2</sup> and in organic synthesis, the cyano group is equivalent to an amine or carbonyl group. The catalytic addition of the cyano group to the C=C bond has been established as one of the most direct pathways for the synthesis of nitriles.<sup>3,4</sup> Among the important nitriles,  $\beta$ -cyano-ketones are commonly utilized in organic synthesis.<sup>5</sup> One of the classical approaches to deliver these compounds is the catalyzed conjugate addition of cyanide to  $\alpha,\beta$ -unsaturated carbonyl compounds (hydrocyanation procedure, Scheme 1a),<sup>6</sup> which utilizes the highly toxic and explosive HCN gas as the cyano source. An alternative strategy is transfer hydrocyanation, which involves the commercially available, less-toxic and less-explosive cyanohydrin to deliver nitriles (Scheme 1b), but with low atom economy.<sup>7</sup> Recently, Morandi developed a nickel-catalyzed transfer hydrocyanation reaction between alkyl nitriles and alkenes or aryl chlorides, which utilizes non-toxic alkyl nitriles as the cyanide source.<sup>8</sup> Although the catalyzed hydrocyanation and transfer hydrocyanation reactions have been well developed,<sup>6,7</sup> it remains an important challenge to bypass the usage of toxic HCN gas as the cyano source and overcome the issue of atom-economy. To mitigate these concerns and inspired by the atom- and step-economical procedure of  $\beta$ -alkylation of secondary alcohols with aldehydes *via* borrowing hydrogen reactions,<sup>9,10</sup> we postulated that the cyano group could be tolerated in the reaction with a mechanism analogous to the borrowing hydrogen reactions.



High atom-economy:  $\text{H}_2\text{O}$  as the sole side product  
 C-CN bond cleavage and formation Commercial, non-toxic CN source

Scheme 1 Development of cyano-borrowing.

School of Chemistry and Chemical Engineering, Southwest University, Chongqing 400715, China. E-mail: liuschop@swu.edu.cn; pyg@swu.edu.cn

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c9sc00640k


Guided by these considerations, we envisioned that, as shown in Scheme 1c, under the catalysis of a transition metal, there is cleavage of the C–CN bond<sup>11,12</sup> of the cyanohydrin and delivery of the corresponding ketones and metal–cyano intermediate ( $[M]^n-CN$ ), followed by aldol condensation of ketones with aldehydes, and subsequent conjugate addition of cyanide to chalcones and utilization of the  $[M]^{n+1}-CN$  as the cyano donor to deliver the desired products (Scheme 1c). This hypothesis is notable in that cyanohydrin plays a dual role both as the source of ketone and the cyanide donor with high atom- and step-economy. Herein, we report the first catalytic process of the direct transformation of cyanohydrins with aldehydes to deliver  $\beta$ -cyano ketones as the sole product *via* nickel catalyzed cyano-borrowing reactions.<sup>13</sup>

## Results and discussion

To test our hypothesis, we initiated the cyano-borrowing reaction using commercially available acetophenone cyanohydrin **1a** which could be prepared from acetophenone and TMSCN and benzaldehyde **2a** for the optimization of the reaction conditions. After screening an array of transition metal catalysts, we found that nickel complexes showed better performance. To our delight, a cocktail consisting of  $NiBr_2$ ,  $PPh_3$ , and  $LiOH$  as the base in dioxane at 100 °C could deliver the desired racemic  $\beta$ -cyano ketone **3aa** in 52% yield (Table 1, entry 1). Notably, determined by the crude  $^1H$  NMR of the reaction mixture, not even a trace of 1,2-addition products was observed.<sup>14</sup> As shown in Table 1, we then examined an extensive array of parameters. By varying the anion of the nickel salt (*i.e.*,  $NiCl_2$ ,  $Ni(OAc)_2$ ,  $Ni(OTf)_2$  and  $Ni(acac)_2$ ) as precatalysts,  $NiCl_2$  exhibited the best yield (Table 1, entries 3–5 *vs.* 2). The screening of different phosphines revealed that this transformation requires a bulky, electron rich ligand, and  $^tBuPAD_2$  showed the highest reactivity with 82% yield (Table 1, entry 11 *vs.* 2 and 6–10). Stronger bases such as  $NaOH$  and  $KO^tBu$  show low reactivity. Meanwhile,  $CS_2CO_3$  and organic bases (*i.e.*, DMAP and DBU) failed to provide the target products (Table 1, entries 14–16). Optimization of the solvent led to no improvement (Table 1, entries 17–19). Lowering the temperature to 80 °C gives a lower yield (Table 1, entry 20). Control experiments verified that the presence of a Ni-complex was necessary to achieve high yield in the cyano-borrowing reaction (Table 1, entry 21). No reaction occurred in the absence of  $LiOH$  (Table 1, entry 22). After the screening of the reaction parameters, we found that  $NiCl_2$  and  $^tBuPAD_2$  as the precatalyst and  $LiOH$  as the base in dioxane at 100 °C for 20 h (82% yield) were the optimal conditions.

Having identified the optimized reaction conditions, we further explored the substrate scope of this reaction. Various commercially available ketone cyanohydrins were examined with benzaldehyde (**2a**) and the results are summarized in Scheme 2. The electron-deficient substrate with varying substituent patterns at the *p*-position did not dramatically influence yields (**3ba–3da**). Substituting the cyanohydrin with *p*-methyl also shows high reactivity and 72% of the desired product was obtained (Scheme 2, **3ea**). However, substituting

Table 1 Screening studies of  $\beta$ -alkylation of cyanohydrin **1a** with benzaldehyde **2a**<sup>a</sup>




Entry	[Ni]	L	Base	Solvent	Yield <sup>b</sup> (%)
1	$NiBr_2$	$PPh_3$	$LiOH$	Dioxane	52
2	$NiCl_2$	$PPh_3$	$LiOH$	Dioxane	76
3	$Ni(OAc)_2$	$PPh_3$	$LiOH$	Dioxane	66
4	$Ni(OTf)_2$	$PPh_3$	$LiOH$	Dioxane	62
5	$Ni(acac)_2$	$PPh_3$	$LiOH$	Dioxane	71
6	$NiCl_2$	$PCy_3$	$LiOH$	Dioxane	76
7	$NiCl_2$	dppe	$LiOH$	Dioxane	72
8	$NiCl_2$	dppp	$LiOH$	Dioxane	78
9	$NiCl_2$	dppb	$LiOH$	Dioxane	37
10	$NiCl_2$	BINAP	$LiOH$	Dioxane	44
11	$NiCl_2$	$^tBuPAD_2$	$LiOH$	Dioxane	82
12	$NiCl_2$	$^tBuPAD_2$	$NaOH$	Dioxane	28
13	$NiCl_2$	$^tBuPAD_2$	$KO^tBu$	Dioxane	Trace
14	$NiCl_2$	$^tBuPAD_2$	$CS_2CO_3$	Dioxane	0
15	$NiCl_2$	$^tBuPAD_2$	DMAP	Dioxane	0
16	$NiCl_2$	$^tBuPAD_2$	DBU	Dioxane	0
17	$NiCl_2$	$^tBuPAD_2$	$LiOH$	Toluene	22
18	$NiCl_2$	$^tBuPAD_2$	$LiOH$	TBME	23
19	$NiCl_2$	$^tBuPAD_2$	$LiOH$	THF	72
20 <sup>c</sup>	$NiCl_2$	$^tBuPAD_2$	$LiOH$	Dioxane	59
21	—	—	$LiOH$	Dioxane	<10
22	$NiCl_2$	$^tBuPAD_2$	—	Dioxane	0

<sup>a</sup> The reaction was carried out with 0.4 mmol of **1a**, 0.4 mmol of **2a**, 5 mol%  $[Ni]$ , 10 mol% ligand (**L**) and 300 mol% base in 0.5 mL of solvent at 100 °C for 18 h. <sup>b</sup> Isolated yield. <sup>c</sup> The reaction was carried out at 80 °C. dppm = bis(diphenylphosphino)methane; dppe = bis(diphenylphosphino)ethane; dppp = bis(diphenylphosphino)propane; dppb = bis(diphenylphosphino)butane;  $^tBuPAD_2$  = di(1-adamantyl)- $^n$ -butylphosphine; TBME = methyl *tert*-butyl ether.


the *para* position with a more electron-rich functionality (*i.e.*, *p*-MeO) led to diminished yields, but the desired product could be delivered with 86% yield while increasing the reaction temperature to 120 °C (Scheme 2, **3fa**). Reactions of cyanohydrin with *ortho*- and *meta*-substitution on aryl groups gave excellent yields (Scheme 2, **3ga–3ja**). The reaction proceeds smoothly in the case of the cyanohydrin bearing a thienyl group, affording the corresponding product **3la** in 82% isolated yield. In terms of the alkyl substituents, cyanohydrin derived from butanone proceeded smoothly to afford the desired product with 40% yield (Scheme 2, **3ma**). The substrate bearing a cyclopropyl group was well tolerated, leading to the cyclopropyl substituted product with 57% yield (Scheme 2, **3na**). Importantly, the cyclopropyl group remains untouched, which indicates that this nickel-catalyzed protocol does not proceed *via* a radical pathway.

Next, various aldehydes were investigated with acetophenone cyanohydrin (**1a**) using the optimized reaction conditions, and representative results are summarized in Scheme 3. Benzaldehydes bearing various electron-deficient (**3ab–3ad** and **3ag–3ah**), electron-neutral (**3al** and **3am**) and electron-rich (**3ae–3af**





Scheme 2 Reaction scope of ketone cyanohydrins. <sup>a</sup>The reaction was carried out at  $120^\circ\text{C}$ .




Scheme 3 Reaction scope of aldehydes. <sup>a</sup>The reaction was carried out at  $120^\circ\text{C}$ .

and **3ai–3ak**) substituents reacted with **1a** to deliver the desired products in moderate to excellent yields (58–90%). Furthermore, the reaction is also compatible with heteroaryl rings, such as 2-furanyl (**3an**), 2-thiophenyl (**3ao**) and unprotected 3-indolyl (**3ap**), providing diverse  $\beta$ -cyano ketones in 51–77% yield. Remarkably, compared with the aryl aldehydes, the alkyl substituted aldehydes (e.g., 2-phenylacetaldehyde and cyclohexanecarbaldehyde) reacted with **1a** to afford the corresponding products **3aq** and **3ar** in 64% and 80% yields, respectively.

The aldehyde containing a sulfur atom is also tolerated under the nickel-catalyzed protocol and delivers the desired product **3as** with a slightly lower yield. To demonstrate the practicality and scalability of our protocol, we proceeded to carry out a gram-scale reaction with 5.0 mmol **2i** reacted with 10.0 mmol **1a** catalyzed by 5 mmol% of  $\text{NiCl}_2$ , affording 1.12 g **3ai** in 85% yield, suggesting that this procedure is quite reliable and practically applicable.


Inspired by the success with the nickel-catalyzed cyano-borrowing reaction of cyanohydrins derived from ketones **1** with aldehydes, we further demonstrate the cyano-borrowing protocol with more challenging substrates such as aldehyde cyanohydrin **5**. To our delight, benzaldehyde cyanohydrin **5a** reacted with acetophenone **4a** smoothly under standard conditions, yielding the corresponding product **3aa** with full conversion and 83% isolated yield. To indicate the generality of this protocol, we then examined the scope of benzaldehyde cyanohydrins and ketones. In addition to the phenyl group, it was found that substrates bearing electron-rich or electron-deficient substituents on the benzene ring were well tolerated to give  $\beta$ -cyano ketones in moderate to good yields (**3ba**, **3ea**, **3ad** and **3af** in Scheme 4). Moreover, heteroaryl and alkyl substituents also participated in this protocol very well (Scheme 4, **3la**, **3ma**, **3ao** and **3ar**).

To illustrate the scope and limitations of the new transformation, challenging substrates beyond methyl ketones were further examined, shown in Scheme 5. The propiophenone cyanohydrin **6** was selected to react with benzaldehyde **2a** under standard reaction conditions, and the desired product **8** was obtained in 31% isolated yield and with high diastereoselectivity ( $\text{dr} > 10 : 1$ ) (Scheme 5a, left). Furthermore, we also examined benzaldehyde cyanohydrin **5a** and propiophenone **7**, with a nickel catalyst, and the corresponding product **8** was obtained in 25% yield (Scheme 5a, right). Notably, the bioactive



Scheme 4 Examples of aldehyde cyanohydrin with ketone. <sup>a</sup>The reaction was carried out at  $120^\circ\text{C}$ .






Scheme 5 Cyano-borrowing beyond methyl ketones.

ketone, epiandrosterone was tested in this reaction with the partner of benzaldehyde cyanohydrin **5a**, and the corresponding product **10** was delivered in 57% yield with 5 : 1 dr (Scheme 5b), which shows the potential of this nickel-catalyzed cyano borrowing process for the selective modification of bioactive ketones.

To shed light on the mechanism for the nickel-catalyzed cyano-borrowing protocol, a series of control experiments were conducted. As shown in Scheme 6, acetophenone cyanohydrin **1a** reacted smoothly with chalcone **11** and give the corresponding product **3aa** in excellent yield, which indicated that cyanohydrin is the source of the cyano group in transfer hydrocyanation under standard reaction conditions. Benzaldehyde cyanohydrin **5a** could also react with chalcone **11** efficiently, delivering **3aa** with 89% yield, and the hydrogen-borrowing product **12** was not observed, showing that cleavage of the C–CN bond is easier than that of the C–H bond in cyanohydrins. Meanwhile, in the crossover reaction of **1a**, **2e** and **11** under standard conditions, we got the corresponding products **3aa** and **3ae** with the ratio of 1.05 : 1, which shows that the cyano group from the cleavage of the C–CN bond of cyanohydrin was a free anion in this nickel-catalyzed protocol



Scheme 6 Mechanism studies.

and has the same opportunity to conjugate to each chalcone. Together, these experimental results support our hypothesis on nickel-catalyzed step- and atom-economical cyano-borrowing reaction of cyanohydrin with aldehydes or ketones (Scheme 1c) (for more details of the mechanism studies, please see the ESI†).

## Experimental

### General procedure

**Method A.** To a vial equipped with a dried stir bar was added aldehydes (0.2 mmol), ketone cyanohydrins (0.4 mmol),  $\text{NiCl}_2$  (5 mol%),  $n\text{-BuPAD}_2$  (5 mol%),  $\text{LiOH}$  (0.6 mmol), 100 mg 4Å MS and anhydrous dioxane (1 mL) in a glovebox. The reaction mixture was taken outside the glovebox and allowed to stir at room temperature for 30 min. After that, the reaction mixture was allowed to stir at 100 °C for 18 hours. The crude reaction mixture was concentrated under reduced pressure and directly purified by silica gel chromatography to give pure products.

**Method B.** To a vial equipped with a dried stir bar was added ketones (0.2 mmol), aldehyde cyanohydrins (0.4 mmol),  $\text{NiCl}_2$  (5 mol%),  $n\text{-BuPAD}_2$  (5 mol%),  $\text{LiOH}$  (0.6 mmol), 100 mg 4Å MS and anhydrous dioxane (1 mL) in a glovebox. The procedure was the same as Method A.

## Conclusions

In conclusion, we have developed an unprecedented nickel-catalyzed protocol for the direct conversion of cyanohydrins and aldehydes or ketones into racemic  $\beta$ -cyano ketones *via* a nickel-catalyzed cyano-borrowing reaction. A range of cyanohydrins derived from aldehydes or ketones could be tolerated and delivered products with high regioselectivity and good to excellent yields. To the best of our knowledge, catalytic conversion of cyanohydrins into  $\beta$ -cyano ketones by reaction with aldehydes/ketones has not been reported. Further studies of catalytic cyano-borrowing reaction of cyanohydrins are in progress in our research lab and will be reported in due course.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

We are grateful for the financial support provided by the National Natural Science Foundation of China (21801209 and 21801210), the Fundamental Research Funds for the Central Universities (SWU118117, XDJK2018C044 and SWU118028) and the Venture & Innovation Support Program for Chongqing Overseas Returnees (cx2018085).

## Notes and references

- (a) R. J. H. Gregory, *Chem. Rev.*, 1999, **99**, 3649–3682; (b) D. Enders and J. P. Shilcock, *Chem. Soc. Rev.*, 2000, **29**, 359–373; (c) M.-X. Wang, *Top. Catal.*, 2005, **35**, 117–130; (d)





- M. B. Smith, *March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*, Wiley-Interscience, 6th edn, 2007; (e) R. Wang and J. R. Falck, *Catal. Rev.: Sci. Eng.*, 2014, **56**, 288–331; (f) M. Tobisu, R. Nakamura, Y. Kita and N. Chatani, *J. Am. Chem. Soc.*, 2009, **131**, 3174–3175; (g) G. A. Chotana, M. A. Rak and M. R. Smith, *J. Am. Chem. Soc.*, 2005, **127**, 10539–10544; (h) T. Patra, S. Agasti, A. Modak and D. Maiti, *Chem. Commun.*, 2013, **49**, 8362–8364; (i) T. Patra, S. Agasti, Akanksha and D. Maiti, *Chem. Commun.*, 2013, **49**, 69–71; (j) M. Tobisu, H. Kinuta, Y. Kita, E. Rémond and N. Chatani, *J. Am. Chem. Soc.*, 2012, **134**, 115–118; (k) M. Tobisu, Y. Kita and N. Chatani, *J. Am. Chem. Soc.*, 2006, **128**, 8152–8153; (l) E. C. Keske, T. H. West and G. C. Lloyd-Jones, *ACS Catal.*, 2018, 8932–8940.
- 2 (a) P. Pollak, G. Romeder, F. Hagedorn and H. P. Gelbke, *Nitriles, Ullmann's Encyclopedia of Industrial Chemistry*, Wiley-VCH, June 15, 2000; (b) J. S. Miller and J. L. Manson, *Acc. Chem. Res.*, 2001, **34**, 563–570; (c) A. Kleemann, J. Engel, B. Kutscher and D. Reichert, *Pharmaceutical substances: Syntheses, Patents, Applications of the most relevant APIs*, Georg Thieme Verlag, Stuttgart, New York, 2001; (d) L. J. Goujon, A. Khaldi, A. Maziz, C. Plesse, G. T. M. Nguyen, P.-H. Aubert, F. Vidal, C. Chevrot and D. Teyssié, *Macromolecules*, 2011, **44**, 9683–9691; (e) F. F. Fleming, *Nat. Prod. Rep.*, 1999, **16**, 597–606; (f) F. F. Fleming, L. Yao, P. C. Ravikumar, L. Funk and B. C. Shook, *J. Med. Chem.*, 2010, **53**, 7902–7917.
- 3 For reviews, see: (a) T. V. Rajanbabu, *Hydrocyanation of Alkenes and Alkynes*, in *Organic Reactions*, John Wiley & Sons, Inc., Hoboken, NJ, 2011, ch. 1, vol. 75, pp. 1–74; (b) M. Beller, J. Seayad, A. Tillack and H. Jiao, *Angew. Chem., Int. Ed.*, 2004, **43**, 3368–3398 and the references therein; (c) T. V. Rajanbabu and A. L. Casalnuovo, in *Comprehensive Asymmetric Catalysis*, ed. E. N. Jacobsen, A. Pfaltz and H. Yamamoto, Springer, Berlin, 1999, vol. 1, pp. 367–378; (d) J. Wilting and D. Vogt, in *Handbook of C-H Transformations*, ed. G. Dyker, Wiley-VCH, Weinheim, 1st edn, 2005, vol. 1, pp. 87–96; (e) L. Bini, C. Müller and D. Vogt, *ChemCatChem*, 2010, **2**, 590–608; (f) P. W. N. M. van Leeuwen, in *Science of Synthesis: Stereoselective Synthesis 1*, ed. J. G. de Vries, Thieme, Stuttgart, 2011, pp. 409–475.
- 4 For reviews on the conjugate reactions, see: (a) A. Córdova, *Catalytic Asymmetric Conjugate Reactions*, WileyVCH, Weinheim, Germany, 2014, pp. 1–439; (b) M. Mauduit, O. Baslé, H. Clavier, C. Crévisy and A. DenicourtNowicki, *Metal-Catalyzed Asymmetric Nucleophilic Addition to Electron-Deficient Alkenes*, Wiley-VCH, Weinheim, Germany, 2014, p. 189; (c) J. L. Vicario, E. Reyes, L. Carrillo and U. Uria, *Organocatalytic Asymmetric Nucleophilic Addition to Electron-Deficient Alkenes*, WileyVCH, Weinheim, Germany, 2014, p. 119; (d) K. Zheng, X. Liu and X. Feng, *Chem. Rev.*, 2018, **118**, 7586–7656.
- 5 (a) G. M. Sammis and E. N. Jacobsen, *J. Am. Chem. Soc.*, 2003, **125**, 4442–4443; (b) T. Mita, K. Sasaki, M. Kanai and M. Shibasaki, *J. Am. Chem. Soc.*, 2005, **127**, 514–515; (c) I. Fujimori, T. Mita, K. Maki, M. Shiro, A. Sato, S. Furusho, M. Kanai and M. Shibasaki, *Tetrahedron*, 2007, **63**, 5820–5831.
- 6 (a) G. M. Sammis, H. Danjo and E. N. Jacobsen, *J. Am. Chem. Soc.*, 2004, **126**, 9928–9929; (b) C. Mazet and E. N. Jacobsen, *Angew. Chem., Int. Ed.*, 2008, **47**, 1762–1765; (c) Y. Tanaka, M. Kanai and M. Shibasaki, *J. Am. Chem. Soc.*, 2008, **130**, 6072–6073; (d) Y. Tanaka, M. Kanai and M. Shibasaki, *J. Am. Chem. Soc.*, 2010, **132**, 8862–8863; (e) N. Kurono, N. Nii, Y. Sakaguchi, M. Uemura and T. Ohkuma, *Angew. Chem., Int. Ed.*, 2011, **50**, 5541–5544.
- 7 (a) B. A. Provencher, K. J. Bartelson, Y. Liu, B. M. Foxman and L. Deng, *Angew. Chem., Int. Ed.*, 2011, **50**, 10565–10569; (b) J. Wang, W. Li, Y. Liu, Y. Chu, L. Lin, X. Liu and X. Feng, *Org. Lett.*, 2010, **12**, 1280–1283; (c) Y.-F. Wang, W. Zeng, M. Sohail, J. Guo, S. Wu and F.-X. Chen, *Eur. J. Org. Chem.*, 2013, **2013**, 4624–4633.
- 8 (a) X. Fang, P. Yu and B. Morandi, *Science*, 2016, **351**, 832–836; (b) P. Yu and B. Morandi, *Angew. Chem., Int. Ed.*, 2017, **56**, 15693–15697.
- 9 For selected reviews on the borrowing-hydrogen methodology, see: (a) Q. Yang, Q. Wang and Z. Yu, *Chem. Soc. Rev.*, 2015, **44**, 2305–2329; (b) M. H. S. A. Hamid, P. A. Slatford and J. M. J. Williams, *Adv. Synth. Catal.*, 2007, **349**, 1555–1575; (c) T. D. Nixon, M. K. Whittlesey and J. M. J. Williams, *Dalton Trans.*, 2009, 753–762; (d) G. E. Dobereiner and R. H. Crabtree, *Chem. Rev.*, 2010, **110**, 681–703; (e) G. Guillena, D. J. Ramón and M. Yus, *Chem. Rev.*, 2010, **110**, 1611–1641; (f) A. J. A. Watson and J. M. J. Williams, *Science*, 2010, **329**, 635–636; (g) S. Bähn, S. Imm, L. Neubert, M. Zhang, H. Neumann and M. Beller, *ChemCatChem*, 2011, **3**, 1853–1864; (h) S. Pan and T. Shibata, *ACS Catal.*, 2013, **3**, 704–712; (i) A. Corma, J. Navas and M. J. Sabater, *Chem. Rev.*, 2018, **118**, 1410–1459; (j) C. Gunanathan and D. Milstein, *Science*, 2013, **341**, 1229712; (k) T. Irrgang and R. Kempe, *Chem. Rev.*, 2018, **118**, 7586–7656. For a recent review on enantioselective borrowing hydrogen, see: (l) A. Quintard and J. Rodriguez, *Chem. Commun.*, 2016, **52**, 10456–10473.
- 10 For selected reviews on  $\alpha$ -alkylation reactions using alcohols with hydrogen borrowing methodologies, see: (a) F. Huang, Z. Liu and Z. Yu, *Angew. Chem., Int. Ed.*, 2016, **55**, 862–875; (b) Y. Obora, *Top. Curr. Chem.*, 2016, **374**, 11; (c) Y. Obora, *ACS Catal.*, 2014, **4**, 3972–3981; for selected examples of  $\alpha$ -alkylation reactions using alcohols with hydrogen borrowing methodologies, see: (d) M. Peña-López, P. Piehl, S. Elangovan, H. Neumann and M. Beller, *Angew. Chem., Int. Ed.*, 2016, **55**, 14967–14971; (e) C. Seck, M. D. Mbaye, S. Coufourier, A. Lator, J. F. Lohier, A. Poater, T. R. Ward, S. Gaillard and J. L. Renaud, *ChemCatChem*, 2017, **9**, 4410–4416; (f) F. Freitag, T. Irrgang and R. Kempe, *Chem.-Eur. J.*, 2017, **23**, 12110–12113; (g) G. Zhang, J. Wu, H. Zeng, S. Zhang, Z. Yin and S. Zheng, *Org. Lett.*, 2017, **19**, 1080–1083; (h) B. C. Roy, S. Debnath, K. Chakrabarti, B. Paul, M. Maji and S. Kundu, *Org. Chem. Front.*, 2018, **5**, 1008–1018.
- 11 F. Chen, T. Wang and N. Jiao, *Chem. Rev.*, 2014, **114**, 8613–8661.



- 12 For selected examples of acetone cyanohydrin in hydrocyanation addition reactions, see: (a) A. Siby, O. Loreau and F. Taran, *Synthesis*, 2009, 2365–2370; (b) M. de Greef and B. Breit, *Angew. Chem., Int. Ed.*, 2009, **48**, 551–554; (c) A. Falk, A.-L. Göderz and H.-G. Schmalz, *Angew. Chem., Int. Ed.*, 2013, **52**, 1576–1580; (d) S. Arai, Y. Amako, X. Yang and A. Nishida, *Angew. Chem., Int. Ed.*, 2013, **52**, 8147–8150; (e) K. Nemoto, T. Nagafuchi, K.-i. Tominaga and K. Sato, *Tetrahedron Lett.*, 2016, **57**, 3199–3203; (f) F. Ye, J. Chen and T. Ritter, *J. Am. Chem. Soc.*, 2017, **139**, 7184–7187.
- 13 (a) B. N. Bhawal and B. Morandi, *ACS Catal.*, 2016, **6**, 7528–7535; (b) B. N. Bhawal and B. Morandi, *Chem.–Eur. J.*, 2017, **23**, 12004–12013.
- 14 For selected examples of 1,2-addition of cyanide to chalcones, see: (a) D. E. Fuerst and E. N. Jacobsen, *J. Am. Chem. Soc.*, 2005, **127**, 8964–8965; (b) X. Liu, B. Qin, X. Zhou, B. He and X. Feng, *J. Am. Chem. Soc.*, 2005, **127**, 12224–12225; (c) Q.-W. Yu, L.-P. Wu, T.-C. Kang, J. Xie, F. Sha and X.-Y. Wu, *Eur. J. Org. Chem.*, 2018, **2018**, 3992–3996; (d) X.-P. Zeng, Z.-Y. Cao, X. Wang, L. Chen, F. Zhou, F. Zhu, C.-H. Wang and J. Zhou, *J. Am. Chem. Soc.*, 2016, **138**, 416–425.

