


Chemical Science

rsc.li/chemical-science



ISSN 2041-6539



ROYAL SOCIETY
OF CHEMISTRY

Celebrating
IYPT 2019

EDGE ARTICLE

Janusz Lewiński *et al.*

Mechanochemical and slow-chemistry radical transformations: a case of diorganozinc compounds and TEMPO

Cite this: *Chem. Sci.*, 2019, 10, 7149

All publication charges for this article have been paid for by the Royal Society of Chemistry

Mechanochemical and slow-chemistry radical transformations: a case of diorganozinc compounds and TEMPO†

Krzysztof Budny-Godlewski,^a Iwona Justyniak,^b Michał K. Leszczyński^b and Janusz Lewiński^{*ab}

From the green chemistry perspective, molecular solid-state transformations conducted under mild conditions are of great interest and desirability. However, research in this area lacked popularity in the previous century, and thus progressed slowly. In particular, the application of radical reactions in solid-state chemistry has been hampered by several long-standing challenges that are intrinsically associated with the apparent unpredictable nature of radical chemistry. We present a comparative study of model mechanochemical, slow-chemistry and solution radical reactions between TEMPO and homoleptic organozinc compounds (*i.e.*, di-*tert*-butylzinc and diphenylzinc). In the case of the *t*Bu₂Zn/TEMPO reaction system only a dimeric diamagnetic complex [tBuZn(μ-TEMPO*)]₂ is obtained in yields slightly varying with the method chosen. In contrast, when TEMPO is mixed with diphenylzinc in a 2 : 1 molar ratio a novel paramagnetic Lewis acid–base adduct [(Ph₂Zn(η¹-TEMPO)]·TEMPO is isolated in high yields regardless of the applied methodology. This adduct is also formed in the slow-chemistry process when TEMPO is gently mixed with Ph₂Zn in a 1 : 1 molar ratio and left for two weeks at ambient temperature. Within the next week the reaction mixture gives in high yield a diamagnetic dinuclear compound [PhZn(μ-TEMPO*)][PhZn(μ₂-η¹:η¹-TEMPO*)] and biphenyl. The analogous reaction conducted in toluene results in a much lower conversion rate. The reported results open up a new horizon in molecular solid-state radical transformations.

Received 21st March 2019

Accepted 16th June 2019

DOI: 10.1039/c9sc01396b

rsc.li/chemical-science

Introduction

The rational development of solid-state reactions is one of the objectives of the green chemistry movement. Notably, in the previous century research on molecular solid-state reactions mediated by mechanical forces progressed slowly which was aptly annotated by Braga and Grepioni: “Although research work in this area has never stopped, it has also lacked popularity and has remained confined to a few connoisseurs”.¹ Only the last decade has witnessed a true renaissance of mechanochemistry^{2–5} applications of which include organic^{6–10} and organometallic synthesis,^{11–14} polymer chemistry,^{15,16} and materials science (*e.g.* in the preparation of MOFs,^{17,18} perovskites for solar cells^{19–22} or nanomaterials).^{3,23–26} In some cases mechanical stress can support electron transfer²⁷ or cause

homolytic bond cleavage and in consequence the formation of radicals. A textbook example of a radical solid-state chemistry phenomenon, which was already noted in ancient Greece, is the attraction between rubbed amber and some small objects.²⁸ Only in 1967 Urbański revealed the formation of radicals on the surface of ground amber.²⁹ The formation of radicals has also been observed upon mechanically induced reactions in polymeric materials.^{30,31} Grzybowski and co-workers demonstrated that these mechanoradicals can be responsible for static charge stabilization, and thus, polymers intentionally doped with radical scavengers (*inter alia* free stable radicals) are less vulnerable to static charging.^{32–34} Despite these seminal pioneering discoveries solid-state radical reactions have remained an underdeveloped area of research.

Even though mechanochemistry is slowly becoming a routine solid-state synthesis tool, knowledge on the intimate mechanistic details is very limited.^{26,35,36} Remarkably, an emerging field of slow-chemistry (mild condition reactions occurring in the solid-state without an external stimuli)³⁷ or accelerated aging reactions (*i.e.*, solid-state transformations where reaction rates are enhanced by the presence of solvent vapour)³⁸ nicely complements and enriches solid-state processes, and may turn out to be an area to help better understand mechanochemical reactions. Slow-chemistry

^aFaculty of Chemistry, Warsaw University of Technology, Noakowskiego 3, 00-664 Warsaw, Poland. E-mail: lewin@ch.pw.edu.pl; Fax: +48 22 6282741; Tel: +48 22 2347315

^bInstitute of Physical Chemistry, Polish Academy of Sciences, Kasprzaka 44/52, 01-224 Warsaw, Poland

† Electronic supplementary information (ESI) available: Characterization data of new products: the NMR, EPR, IR spectra, extended PXRD diffractograms and crystallographic data. CCDC 1550886–1550888. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c9sc01396b



reactions are widely observed in nature, *e.g.* biomineralization, pearl formation or mineral neogenesis, and only recently have these natural processes been mimicked in synthetic procedures.^{39–42}

Herein we present unprecedented solvent-free slow-chemistry and mechanical force triggered radical transformations involving solid homoleptic organozinc compounds and an intentionally added free stable nitroxyl radical, (2,2,6,6-tetramethylpiperidin-1-yl)oxyl (TEMPO). Very recently, in contrast to common beliefs,⁴³ our group and independently Stephan's group demonstrated that in solution TEMPO is non-innocent towards Et_2Zn ^{44,45} and $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Zn}$,⁴⁶ respectively, and smoothly induces radical transformations triggered by single-electron transfer (SET)⁴⁷ from the Zn–C bond to TEMPO, to afford organozinc complexes incorporating TEMPO anions. We were wondering whether TEMPO will also remain non-innocent towards homoleptic diorganozinc compounds in the solid-state and to this end we chose di-*tert*-butylzinc ($t\text{Bu}_2\text{Zn}$) and diphenylzinc (Ph_2Zn) solid homoleptic organometallics with versatile properties and reactivity.^{48–50} The solid-state reactions were commenced with comparative solution phase investigations. Remarkably, while in the case of the $t\text{Bu}_2\text{Zn}/\text{TEMPO}$ system either wet chemistry or mechanochemical and slow-chemistry transformations provided essentially the same outcome, the character of the products of the $\text{Ph}_2\text{Zn}/\text{TEMPO}$ system depends significantly on the stoichiometry of the reaction and the selected synthetic methodology.

Results and discussion

Comparative investigations in solution

Initially, when a toluene solution of $t\text{Bu}_2\text{Zn}$ was treated with an orange toluene solution of TEMPO (2 equiv.) at -78°C , an instantaneous decolourization took place. Standard work-up of the post-reaction mixture afforded colourless crystals of a diamagnetic dimeric complex $[\text{tBuZn}(\mu\text{-TEMPO}^*)]_2$ (**1**₂) (Fig. 1a, TEMPO* stands for the TEMPO anion) and TEMPO-*t*Bu as the primary products. The formation of **1**₂ is likely triggered by an incipient $t\text{Bu}_2\text{Zn}/\text{TEMPO}$ adduct followed by an intramolecular SET from the Zn-*t*Bu bond to TEMPO, and TEMPO-*t*Bu results from the subsequent release of a *t*Bu[•] radical and its recombination with the second TEMPO molecule.⁵¹ In this case we were not able to isolate the anticipated paramagnetic Lewis acid–base $\text{R}_2\text{Zn}(\text{TEMPO})$ adduct due to the high reactivity of homoleptic zinc alkyls towards TEMPO. Remarkably, the respective paramagnetic adduct was isolated from a similar reaction with Ph_2Zn . Clearly, Ph_2Zn is less prone to react as a SET reagent, and moreover the $\text{Ph}_2\text{Zn}/\text{TEMPO}$ system appeared also sensitive to the molar ratio of the reagents and reaction time. A 1 : 1 mixture of the starting materials was dissolved in toluene and stirred at room temperature for an hour and then cooled to -26°C . Within 24 hours yellow crystals of a supramolecular complex $[[\text{Ph}_2\text{Zn}(\eta^1\text{-TEMPO})]\cdot\text{TEMPO}]$ (**2**·TEMPO) deposited in moderate yield. Compound **2**·TEMPO is composed of a paramagnetic Lewis acid–base adduct $[\text{Ph}_2\text{Zn}(\eta^1\text{-TEMPO})]$ (**2**) with a hydrogen-bonded TEMPO molecule (Fig. 1b). Interestingly, a prolonged (two weeks) reaction

between Ph_2Zn and 1 equiv. of TEMPO conducted at ambient temperature in toluene generated a new diamagnetic dinuclear complex $[\text{PhZn}(\mu\text{-TEMPO}^*)][\text{PhZn}(\mu_2\text{-}\eta^1\text{-}\eta^1\text{-TEMPO}^*)]$ (**3**, Fig. 1c) (yield *ca.* 35%) as well as biphenyl (PhPh) resulting from homocoupling of the Ph[•] radicals. Remarkably, the Ph[•] radicals do not recombine with TEMPO molecules on this occasion as we did not observe the signals characteristic of TEMPO–Ph in the ¹H NMR spectrum of the post-reaction mixture.

To clarify the factors determining the formation of the paramagnetic complex **2**·TEMPO and the diamagnetic complex **3**, we performed the reaction of Ph_2Zn with 2 equiv. of TEMPO in a manner analogous to that of the equimolar system (for details, see the Experimental procedures). In this case the SET process was suppressed and **2**·TEMPO was isolated essentially quantitatively. Undoubtedly, a key aspect of the $\text{Ph}_2\text{Zn}/\text{TEMPO}$ reaction system is the reduction of TEMPO to its anionic form by Ph_2Zn (with the subsequent coupling of the formed Ph[•] radicals to yield PhPh) only for the 1 : 1 mixture while a higher amount of TEMPO inhibits the SET step (for details see Fig. S22 in the ESI†).

Crystal structure analysis

Compounds **1**₂, **2**·TEMPO and **3** were characterized by NMR spectroscopy and single-crystal X-ray diffraction, and additionally the paramagnetic character of **2**·TEMPO was confirmed by EPR spectroscopy. Compounds **1**₂ and **3** show sharp ¹H NMR resonances characteristic of the zinc bonded organic groups and the TEMPO* anion, while the NMR spectrum of **2**·TEMPO confirms its paramagnetic character (Fig. S1–S8 in the ESI†). The diffusion ordered NMR spectroscopy (DOSY) experiments show that **1**₂ and **3** most likely exist as monomers in solution (for details, see Table S1 in the ESI†). Remarkably, the solution and solid-state EPR spectra of **2**·TEMPO show one set of signals analogous to that observed for the free TEMPO radical (see Fig. S13 and S14†).

X-ray crystal structure analysis showed that the molecular structure of the *tert*-butyl–zinc derivative **1**₂ (Fig. 1a) is essentially analogous to the previously reported diamagnetic ethylzinc derivative $[\text{EtZn}(\mu\text{-TEMPO}^*)]_2$,⁴⁶ and comprises bridging $\mu\text{-TEMPO}^*$ ligands and a planar Zn_2O_2 ring (Zn1–O1 = 1.939(3) Å and Zn1–O1' = 1.964(3) Å). In turn, the relevant phenyl–zinc derivative **3** crystallizes as an asymmetric dimer that contains a rarely encountered mix of coordination $\mu\text{-O-}$ and $\mu\text{-}\eta^1\text{:}\eta^1\text{-NO-}$ modes of the anionic TEMPO* ligands (Fig. 1c).⁵² Two three-coordinate zinc atoms are bridged by $\mu\text{-O-TEMPO}^*$ and $\mu\text{-}\eta^1\text{:}\eta^1\text{-NO-TEMPO}^*$ units, thus forming a slightly bent five-membered $[\text{ZnOZnNO}]$ ring (Zn1–O1 = 1.916(3) Å, O1–Zn2 = 1.943(3) Å, Zn2–N2 = 2.078(3) Å, N2–O2 = 1.441(4) Å, and O2–Zn1 = 1.898(2) Å).

In the paramagnetic adduct **2**·TEMPO, one TEMPO radical coordinates to the zinc centre in an η^1 -mode (Zn1–O1 = 2.0841(18) Å, O1–N1 = 1.294(3) Å, and Zn1–O1–N1 = 143.50(17)°) (Fig. 1b). The second TEMPO molecule interacts by hydrogen bonding involving the oxygen atom and the C–H bond of one Zn-bonded Ph group (H5–O2 = 2.323(9) Å, O2–N2 = 1.288(3) Å, and H5–O2–N2 = 86.517(13)°) and the observed






Fig. 1 Unprecedented variety of outcomes of the solid-state and wet reactions of TEMPO with $t\text{Bu}_2\text{Zn}$ and Ph_2Zn : (a) $[\text{tBu}_2\text{Zn}(\mu\text{-TEMPO}^*)]_2$ (1_2), (b) $[\text{Ph}_2\text{Zn}(\eta^1\text{-TEMPO}^*)\cdot\text{TEMPO}]$ ($2\cdot\text{TEMPO}$), and (c) $[\text{PhZn}(\mu\text{-TEMPO}^*)][\text{PhZn}(\mu_2\text{-}\eta^1:\eta^1\text{-TEMPO}^*)]$ (3).

hydrogen-bonding interaction torsion angle $\text{N2-O2}\cdots\text{H5-C5} = 45.227(20)^\circ$ is similar to the calculated $\text{N-O}\cdots\text{H-C}$ torsion angle (45°) for the $\text{CH}_3\text{CN}/\text{TEMPO}$ system.⁵³ The zinc atom occupies a site within a trigonal-planar arrangement of the oxygen and carbon atoms ($\text{C1-Zn1-C7} = 145.61(12)$, $\text{C1-Zn1-O1} = 112.8(1)^\circ$, and $\text{O1-Zn1-O7} = 101.4(1)$) with the Zn1-O1 bond length markedly longer than that observed for the Zn-O bond in zinc complexes incorporating an external anionic $\eta^1\text{-TEMPO}^*$ ligand (*cf.* the Zn-O of 1.854(6) in $[\text{Zn}(\text{TEMPO}^*)_2]$).⁴⁶ To the best of our knowledge, the adduct **2** is the first isolated and structurally characterized simple Lewis acid adduct of TEMPO with an organometallic molecule.⁵⁴ The revealed hydrogen-bond interaction between the free TEMPO molecule and the Zn-bonded Ph group as well as a vast number of other non-covalent interactions observed in the solid-state structure of $2\cdot\text{TEMPO}$

(see Fig. S27[†]) are likely responsible for an increased stability of $2\cdot\text{TEMPO}$ in the solid-state (note that $2\cdot\text{TEMPO}$ does not transform into **3** in the solid-state, *vide infra*).

Reactions in the solid-state

Equipped with the above knowledge about the reactivity of TEMPO towards $t\text{Bu}_2\text{Zn}$ and Ph_2Zn in solution, we performed analogous reactions with these reagents in the solid-state. The progress of both slow-chemistry and mechanochemical reactions was monitored using PXRD analysis (for extended diffractograms see Fig. S9–S12[†]) and additionally confirmed by ^1H NMR spectroscopy; all operations were conducted under an argon inert gas atmosphere.

We revealed that both slow- and mechanochemical reactions take place in the case of the $t\text{Bu}_2\text{Zn}/\text{TEMPO}$ system, and



selectively afford **1**₂ and TEMPO-*t*Bu in high yield (Fig. 2). The mechanochemical reaction involving grinding *t*Bu₂Zn with 2 equiv. of TEMPO with a glass rod at 0 °C occurs within circa 20 minutes (Fig. 2b). In the slow-chemistry reaction, *t*Bu₂Zn was gently mixed with 2 equiv. of TEMPO and then stored at 5 °C under an inert gas atmosphere and in this case the reaction required circa 24 hours for completion (Fig. 2c).




Fig. 2 The PXRD patterns for the reactions of TEMPO with *t*Bu₂Zn: (a) the simulated PXRD spectrum of **1**₂, (b) the mixture of *t*Bu₂Zn and 2 equiv. of TEMPO ground with a glass rod for 20 minutes at 0 °C and (c) the mixture of *t*Bu₂Zn and 2 equiv. of TEMPO gently mixed and stored at 5 °C for 24 h.




Fig. 3 The PXRD patterns of various reaction systems involving TEMPO and Ph₂Zn: (a) the simulated PXRD spectrum of **2**·TEMPO, (b) Ph₂Zn ground with 2 equiv. of TEMPO, (c) Ph₂Zn gently mixed with 2 equiv. of TEMPO and stored under an inert gas atmosphere at 20 °C for 5 days, (d) Ph₂Zn ground with 1 equiv. of TEMPO, and (e) Ph₂Zn. (f) The simulated PXRD spectrum of **3** and (g) Ph₂Zn gently mixed with 1 equiv. of TEMPO stored at 20 °C for 21 days.

Strikingly, transformations involving Ph₂Zn and TEMPO proceed entirely differently than in the case of the *t*Bu₂Zn/TEMPO system (Fig. 3). Grinding Ph₂Zn with 1 equiv. of TEMPO at ambient temperature for over 15 minutes gives a mixture of the paramagnetic adduct **2**·TEMPO and unreacted Ph₂Zn as judged by the PXRD analysis (Fig. 3b). However, when Ph₂Zn is ground with 2 equiv. of TEMPO, then **2**·TEMPO is obtained selectively as the only product (Fig. 3c). Regardless of the grinding conditions (molar ratio of substrates, prolonged time, and higher frequency of milling), no trace of the **3** phase was observed in the PXRD pattern of the postreaction powder.‡ What is more, the pure compound **2**·TEMPO does not transform into the diamagnetic compound **3** even after over 6 months of aging (20 °C and inert gas atmosphere).

The slow chemistry reactions involving Ph₂Zn and TEMPO in various molar ratios are more subtle. The paramagnetic adduct **2**·TEMPO is obtained selectively within *ca.* 5 days of aging (20 °C) of a mixture consisting of Ph₂Zn gently mixed with 2 equiv. of TEMPO (Fig. 3d). No further transformation of **2**·TEMPO was observed in the solid state even after over 6 months of aging. In the case of the equimolar mixture of Ph₂Zn and TEMPO the slow-chemistry reaction proceeds differently and consists of two distinguishable steps. In the first step, the reaction yields an intermediate mixture of **2**·TEMPO and Ph₂Zn, which in the second step quantitatively transforms into an equimolar mixture of **3** and PhPh (Fig. 4). Moreover, the evolution of PXRD diffractograms of the equimolar mixture of Ph₂Zn and TEMPO over time depends on the way the substrates were initially mixed (ground or gently mixed). The PXRD analysis of the initially ground substrate mixture reveals patterns characteristic of **2**·TEMPO and Ph₂Zn. Then after 21 days of aging the PXRD patterns evolve into that characteristic of **3** (patterns characteristic of PhPh are not observed but its formation was




Fig. 4 Proposed mechanism of the reaction of TEMPO with Ph₂Zn (1 : 1 molar ratio) in the solid-state.



confirmed by ^1H NMR spectroscopy). In contrast, the gently mixed substrate mixture affords an intermediate mixture of $2\cdot\text{TEMPO}$ and Ph_2Zn within 7 days of aging and then within the next 14 days transforms into **3** and PhPh .

The above results indicate that the solid-state equimolar mixture of Ph_2Zn and TEMPO initially smoothly transforms into a $2\cdot\text{TEMPO}$ supramolecular complex while half the amount of Ph_2Zn remains intact. Subsequently, the mixture of $2\cdot\text{TEMPO}$ and Ph_2Zn transforms into **3** in the slow-chemistry process. Remarkably, this transformation is not observed for the pure phase $2\cdot\text{TEMPO}$ complex under either mechanochemical or slow-chemistry conditions. Thus, at this stage of studies it is reasonable to propose that $2\cdot\text{TEMPO}$ and Ph_2Zn equilibrate in the solid state with the simple Lewis acid–base adduct **2** that is active in the SET process leading to **3** (Fig. 4).

Conclusions

Investigations of solid-state chemistry reactions under mild conditions have remained both one of the greatest mysteries and one of the most challenging tasks for chemists and materials scientists. We demonstrated that mechanochemical and slow-chemistry processes can be used as efficient tools to control the outcome of organometallic radical reactions. Moreover, in our opinion, the reported studies are a significant advance in molecular solid-state chemistry and should aid in unlocking the potential of radical slow-chemistry reactions as a novel tool for the preparation of desired products. At this stage with little knowledge of the mechanisms of the titular solid-state transformations, the best approach to extending this idea would seemingly be through trial and error. Further work in this regard is currently underway in our laboratory.

Experimental

$t\text{Bu}_2\text{Zn}$ was prepared according to literature procedures⁴⁸ under an argon atmosphere using standard Schlenk techniques. Ph_2Zn and all other reagents were of commercial grade obtained from Sigma-Aldrich Co. The solvents were carefully dried and distilled under a nitrogen atmosphere prior to use. TEMPO was sublimed and stored under an argon atmosphere.

$[\text{tBuZn}(\mu\text{-TEMPO}^*)]_2$ (**1**)

Solution. An orange solution of TEMPO (2 mmol, 0.312 g) in toluene (2.7 mL) was added dropwise to a stirred solution of $t\text{Bu}_2\text{Zn}$ (1 mmol, 0.180 g) in toluene (8 mL) at -78°C . A suspension of white solids in colourless solution was formed. The mixture was warmed to RT resulting in a colourless clear solution. After 24 h at -15°C colourless crystals of **1** deposited from this solution (78%). ^1H NMR (benzene- d_6 , 400 MHz, 25°C , ppm): δ 1.50 (s, 9H; $\text{ZnC}(\text{CH}_3)_3$), 1.42–1.37 (br. m, 4H; $\beta\text{-CH}_2$), 1.30–1.23 (br. m, 2H; $\gamma\text{-CH}_2$), 1.18 (s, 12H; $\text{C}(\text{CH}_3)_2$). $^{13}\text{C}\{^1\text{H}\}$ NMR (benzene- d_6 , 125 MHz, 25°C , ppm): δ 58.87 ($\text{C}(\text{CH}_3)_2$), 40.49 ($\beta\text{-CH}_2$), 34.02 ($\text{C}(\text{CH}_3)_2$), 25.50 ($\text{C}(\text{CH}_3)_3$), 17.76 ($\gamma\text{-CH}_2$). C,H,N analysis (%) calcd for $\text{C}_{22}\text{H}_{46}\text{N}_2\text{O}_2\text{Zn}_2$: C 52.70, H 9.25, N 5.59; found: C 52.73, H 9.22, N 5.55.

Mechanochemistry. TEMPO (2.4 mmol, 0.376 g) and $t\text{Bu}_2\text{Zn}$ (1.2 mmol, 0.448 g) were ground with a glass rod for 20 minutes at 0°C in a Schlenk tube until the product mixture became light-beige in colour and obtained homogeneous paste-like consistency. Then the volatiles were removed *in vacuo* at 0°C . The crude product mixture was analysed by PXRD and ^1H NMR. The mixture was dissolved in toluene affording colourless crystals in 81% yield after 24 h at -5°C .

Slow-chemistry. TEMPO (0.88 mmol, 0.137 g) and $t\text{Bu}_2\text{Zn}$ (0.44 mmol, 0.079 g) were gently mixed with a needle in a Schlenk tube at -30°C and then stored at 5°C for 24 hours. The colour of the mixture turned from white-orange to white. The volatiles were removed *in vacuo* at 5°C . The crude product mixture was analysed by PXRD and ^1H NMR spectroscopy. The mixture was dissolved in toluene affording colourless crystals in 77% yield after 24 h at -15°C .

$[\text{Ph}_2\text{Zn}(\eta^1\text{-TEMPO})\cdot\text{TEMPO}]$ ($2\cdot\text{TEMPO}$)

Solution. An orange solution of TEMPO (2 mmol, 0.312 g) in toluene (2.1 mL) was added to a suspension of Ph_2Zn (1 mmol, 0.180 g) in toluene (8 mL) at RT. The clear yellow reaction mixture was stirred at RT for 1 hour and then placed in a freezer. After 24 h at -26°C yellow crystals suitable for X-ray analysis were obtained in 69% yield. The isolated yield was lower for a similar reaction conducted for the 1 : 1 molar ratio (*ca.* 39%). The paramagnetic nature of the product was confirmed by EPR spectroscopy. Pure compound $2\cdot\text{TEMPO}$ left for over 6 months at 20°C under an inert gas atmosphere does not transform into the diamagnetic compound **3**. C,H,N analysis (%) calcd for $\text{C}_{30}\text{H}_{46}\text{N}_2\text{O}_2\text{Zn}_1$: C 67.72, H 8.71, N 5.26; found: C 67.75, H 8.68, N 5.24.

Mechanochemistry. Ph_2Zn (0.5 mmol, 0.110 g) was gently mixed with TEMPO (1.0 mmol, 0.156 g) at 20°C and transferred into an agate ball mill. The mixture was milled for a total of 15 minutes (25 Hz, 1 agate ball) in three intervals (with 3 minute breaks, to prevent excessive heating). The colour of the mixture turned yellow. The crude product was analysed by PXRD. No trace of the **3** phase was observed in the PXRD pattern of the postreaction mixture regardless of the process conditions (prolonged time (over 1 hour in total) and higher frequency of milling).

Slow-chemistry. Ph_2Zn (0.5 mmol, 0.110 g) was gently mixed with TEMPO (1.0 mmol, 0.156 g) with a needle and stored under an inert gas atmosphere at 20°C for 5 days. The colour of the mixture turned yellow. The crude product was analysed by PXRD.

$[\text{PhZn}(\mu_2\text{-TEMPO}^*)][\text{PhZn}(\mu_2\text{-}\eta^1\text{-}\eta^1\text{-TEMPO}^*)]$ (**3**)

Solution. An orange solution of TEMPO (0.8 mmol, 0.125 g) in toluene (0.8 mL) was added to a suspension of Ph_2Zn (0.8 mmol, 0.144 g) in toluene (5 mL) at RT. The clear yellow reaction mixture was stirred at RT for 21 days. After 24 h at -26°C colourless crystals suitable for X-ray analysis deposited (isolated yield 24%). ^1H NMR (benzene- d_6 , 500 MHz, 25°C , ppm): δ 8.01 (dd, $^3J = 7.82$ Hz, $^4J = 1.35$ Hz, 2H, *o*- $\text{CH}_{(\text{ar})}$), 7.42 (t, 2H, $J = 7.4$ Hz, *m*- $\text{CH}_{(\text{ar})}$), 7.30 (tt, 1H, $^3J = 7.4$ Hz, $^4J = 1.34$



Hz, p -CH_(ar)), 1.47 (br m, 4H, β -CH₂ (TEMPO-ring)), 1.28 (s, 12H, C(CH₃)₂), 1.22 (m, 2H, γ -CH₂ (TEMPO-ring)). ¹³C{¹H} NMR (benzene-*d*₆, 125 MHz, 20 °C, ppm): δ 18.2 (C(CH₃)₂), 40.5 (TEMPO-ring), 41.0 (C(CH₃)₂), 59.2 (TEMPO-ring). C,H,N analysis (%) calcd for C₃₆H₇₂N₄O₄Zn₂: C 57.21, H 9.60, N 7.41; found: C 57.28, H 9.55, N 7.40.

Slow-chemistry. TEMPO (1.22 mmol, 0.190 g) and Ph₂Zn (1.22 mmol, 0.267 g) were gently mixed with a glass rod in a Schlenk tube under an inert gas atmosphere at 20 °C. The mixture was stored for 21 days at 20 °C and the progress of the process was monitored by PXRD. The colour of the mixture turned from yellowish to white. Then the mixture was dissolved in toluene. After 24 h at -26 °C colourless crystals of 3 deposited (isolate yield 92%). The nature of the product was confirmed by ¹H NMR spectroscopy. We note that the slow-chemistry process occurs regardless of whether the substrates were initially gently mixed or ground. However, the initial treatment of the mixture drastically changes the evolution of its PXRD patterns over time (see Fig. S11 and S12† to compare the PXRD patterns of the initially ground and mixed mixtures). Prolonged grinding of the reaction mixture (conducted with 3 minute breaks to prevent excessive heating) results in a mixture of Ph₂Zn and 2·TEMPO. It is noteworthy that prolonged uninterrupted grinding results in intensive heating of the reaction mixture.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors gratefully acknowledge the financial support of the Foundation for Polish Science TEAM Program co-financed by the European Union under the European Regional Development Fund TEAM/2016-2/14 and the support of the National Science Centre 2015/19/N/ST5/02194. We are grateful to Dr Krzysztof Kruczała and Prof. Zbigniew Sojka for the EPR measurements and Prof. Zbigniew Ochal for GC measurements.

Notes and references

‡ Bearing in mind that TEMPO melts at approximately 40 °C, we conducted an experiment between an equimolar mixture of Ph₂Zn and TEMPO at 40 °C. According to PXRD analysis, the reaction gives a mixture of 2·TEMPO and Ph₂Zn within 30 minutes, which then after 4 days at 40 °C gives 3 and PhPh in excellent yield (for details see Fig. S24). We assume that prolonged (days) grinding will lead to the same results as the simple thermal experiment described above, although as a reaction in the melted phase it should no longer be considered as mechanochemistry.

- D. Braga and F. Grepioni, *Angew. Chem., Int. Ed.*, 2004, **43**, 4002–4011.
- V. Boldyrev and K. Tkáčová, *J. Mater. Synth. Process.*, 2000, **8**, 121–132.
- S. L. James, C. J. Adams, C. Bolm, D. Braga, P. Collier, T. Frišćić, F. Grepioni, K. D. M. Harris, G. Hyett, W. Jones, A. Krebs, J. Mack, L. Maini, A. G. Orpen, I. P. Parkin, W. C. Shearouse, J. W. Steed and D. C. Waddell, *Chem. Soc. Rev.*, 2011, **41**, 413–447.
- Themed Issue: Mechanochemistry, *Chem. Soc. Rev.*, 2013, **42**, 7487–7740.
- J.-L. Do and T. Frišćić, *ACS Cent. Sci.*, 2017, **3**, 13–19.
- A. Stolle, T. Szuppa, S. E. S. Leonhardt and B. Ondruschka, *Chem. Soc. Rev.*, 2011, **40**, 2317–2329.
- J. L. Howard, Q. Cao and D. L. Browne, *Chem. Sci.*, 2018, **9**, 3080–3094.
- G.-W. Wang, *Chem. Soc. Rev.*, 2013, **42**, 7668–7700.
- D. Tan and T. Frišćić, *Eur. J. Org. Chem.*, 2018, **2018**, 18–33.
- J. Andersen and J. Mack, *Green Chem.*, 2018, **20**, 1435–1443.
- D. Tan and F. Garcia, *Chem. Soc. Rev.*, 2019, **48**, 2274–2292.
- N. R. Rightmire and T. P. Hanusa, *Dalton Trans.*, 2016, **45**, 2352–2362.
- J. Lewiński, M. Dutkiewicz, M. Lesiuk, W. Śliwiński, K. Zelga, I. Justyniak and J. Lipkowski, *Angew. Chem., Int. Ed.*, 2010, **49**, 8266–8269.
- J. G. Hernández, I. S. Butler and T. Frišćić, *Chem. Sci.*, 2014, **5**, 3576–3582.
- J. Li, C. Nagamani and J. S. Moore, *Acc. Chem. Res.*, 2015, **48**, 2181–2190.
- N. Willis-Fox, E. Rognin, T. A. Aljohani and R. Daly, *Chem.*, 2018, **4**, 2499–2537.
- D. Prochowicz, K. Sokołowski, I. Justyniak, A. Kornowicz, D. Fairen-Jimenez, T. Frišćić and J. Lewiński, *Chem. Commun.*, 2015, **51**, 4032–4035.
- K. Užarević, T. C. Wang, S.-Y. Moon, A. M. Fidelli, J. T. Hupp, O. K. Farha and T. Frišćić, *Chem. Commun.*, 2016, **52**, 2133–2136.
- D. Prochowicz, M. Franckevičius, A. M. Cieślak, S. M. Zakeeruddin, M. Grätzel and J. Lewiński, *J. Mater. Chem. A*, 2015, **3**, 20772–20777.
- P. Pal, S. Saha, A. Banik, A. Sarkar and K. Biswas, *Chem.–Eur. J.*, 2018, **24**, 1811–1815.
- D. J. Kubicki, D. Prochowicz, A. Hofstetter, M. Sasaki, P. Yadav, D. Bi, N. Pellet, J. Lewiński, S. M. Zakeeruddin, M. Grätzel and L. Emsley, *J. Am. Chem. Soc.*, 2018, **140**, 3345–3351.
- M. Sasaki, D. Prochowicz, W. Marynowski and J. Lewiński, *Eur. J. Inorg. Chem.*, 2019, 2680–2684.
- M. J. Rak, T. Frišćić and A. Moores, *Faraday Discuss.*, 2014, **170**, 155–167.
- M. Y. Malca, H. Bao, T. Bastaille, N. K. Saadé, J. M. Kinsella, T. Frišćić and A. Moores, *Chem. Mater.*, 2017, **29**, 7766–7773.
- A. Moores, *Current Opinion in Green and Sustainable Chemistry*, 2018, **12**, 33–37.
- P. Krupiński, A. Kornowicz, K. Sokołowski, A. M. Cieślak and J. Lewiński, *Chem.–Eur. J.*, 2016, **22**, 7817–7823.
- Y. Li, N. L. Haworth, L. Xiang, S. Ciampi, M. L. Coote and N. Tao, *J. Am. Chem. Soc.*, 2017, **139**, 14699–14706.
- P. Iversen and D. J. Lacks, *J. Electroanal. Chem.*, 2012, **70**, 309–311.
- T. Urbański, *Nature*, 1967, **216**, 577–578.
- M. K. Beyer and H. Clausen-Schaumann, *Chem. Rev.*, 2005, **105**, 2921–2948.
- M. M. Caruso, D. A. Davis, Q. Shen, S. A. Odom, N. R. Sottos, S. R. White and J. S. Moore, *Chem. Rev.*, 2009, **109**, 5755–5798.



- 32 B. Baytekin, H. T. Baytekin and B. A. Grzybowski, *Angew. Chem., Int. Ed.*, 2014, **53**, 6946–6950.
- 33 H. T. Baytekin, B. Baytekin and B. A. Grzybowski, *Angew. Chem., Int. Ed.*, 2012, **51**, 3596–3600.
- 34 H. T. Baytekin, B. Baytekin, T. M. Hermans, B. Kowalczyk and B. A. Grzybowski, *Science*, 2013, **341**, 1368–1371.
- 35 A. M. Belenguer, G. I. Lampronti, D. J. Wales and J. K. M. Sanders, *J. Am. Chem. Soc.*, 2014, **136**, 16156–16166.
- 36 K. Užarević, I. Halasz and T. Friščić, *J. Phys. Chem. Lett.*, 2015, **6**, 4129–4140.
- 37 X. Lim, *Nature*, 2015, **524**, 20–21.
- 38 M. J. Cliffe, C. Mottillo, R. S. Stein, D.-K. Bučar and T. Friščić, *Chem. Sci.*, 2012, **3**, 2495–2500.
- 39 F. Nudelman and N. A. J. M. Sommerdijk, *Angew. Chem., Int. Ed.*, 2012, **51**, 6582–6596.
- 40 C. Mottillo, Y. Lu, M.-H. Pham, M. J. Cliffe, T.-O. Do and T. Friščić, *Green Chem.*, 2013, **15**, 2121–2131.
- 41 F. Qi, R. S. Stein and T. Friščić, *Green Chem.*, 2014, **16**, 121–132.
- 42 C. Mottillo and T. Friščić, *Molecules*, 2017, **22**, 144–181.
- 43 A. Gorrane, I. Resa, A. Rodriguez, E. Carmona, E. Alvarez, E. Gutierrez-Puebla, A. Monge, A. Galindo, D. del Río and R. A. Andersen, *J. Am. Chem. Soc.*, 2007, **129**, 693–703.
- 44 K. Budny-Godlewski, D. Kubicki, I. Justyniak and J. Lewiński, *Organometallics*, 2014, **33**, 5093–5096.
- 45 M. Kubisiak, K. Zelga, W. Bury, I. Justyniak, K. Budny-Godlewski, Z. Ochal and J. Lewiński, *Chem. Sci.*, 2015, **6**, 3102–3108.
- 46 P. Jochmann and D. W. Stephan, *Chem. Commun.*, 2014, **50**, 8395–8397.
- 47 For selected examples of the role of SET in organozinc chemistry, see: (a) E. Wissing, S. van der Linden, E. Rijnberg, J. Boersma, W. J. J. Smeets, A. L. Spek and G. van Koten, *Organometallics*, 1994, **13**, 2602–2608; (b) J. Lewiński, W. Śliwiński, M. Dranka, I. Justyniak and J. Lipkowski, *Angew. Chem., Int. Ed.*, 2006, **45**, 4826–4829; (c) J. Lewiński, K. Suwała, M. Kubisiak, Z. Ochal, I. Justyniak and J. Lipkowski, *Angew. Chem., Int. Ed.*, 2008, **47**, 7888–7891; (d) I. Dranka, M. Kubisiak, I. Justyniak, M. Lesiuk, D. Kubicki and J. Lewiński, *Chem.-Eur. J.*, 2011, **17**, 12713–12721; (e) E. Folkertsma, S. H. Benthem, J. T. B. H. Jastrzebski, M. Lutz, M.-E. Moret and R. J. M. Klein Gebbink, *Eur. J. Inorg. Chem.*, 2018, 1167–1175.
- 48 J. Lewiński, M. Dranka, W. Bury, W. Śliwiński, I. Justyniak and J. Lipkowski, *J. Am. Chem. Soc.*, 2007, **129**, 3096–3098.
- 49 G. Wittig, F. J. Meyer and G. Lange, *Justus Liebigs Ann. Chem.*, 1951, **571**, 167–201.
- 50 P. R. Markies, G. Schat, O. S. Akkerman, F. Bickelhaupt, W. J. J. Smeets and A. L. Spek, *Organometallics*, 1990, **9**, 2243–2247.
- 51 D. R. Armstrong, L. Balloch, J. J. Crawford, B. J. Fleming, L. M. Hogg, A. R. Kennedy, J. Klett, R. E. Mulvey, C. T. O'Hara, S. A. Orr and S. D. Robertson, *Chem. Commun.*, 2012, **48**, 1541–1543.
- 52 For selected examples of TEMPO* anion coordination modes in organozinc chemistry, see: ref. 44 and 46, and (a) G. C. Forbes, A. R. Kennedy, R. E. Mulvey and P. J. A. Rodger, *Chem. Commun.*, 2001, 1400–1401; (b) Z. Zhu, J. C. Fettinger, M. M. Olmstead and P. P. Power, *Organometallics*, 2009, **28**, 2091–2095; (c) J. Francos, A. R. Kennedy and C. T. O'Hara, *Dalton Trans.*, 2016, **45**, 6222–6233.
- 53 J. L. Russ, J. Gu, K.-H. Tsai, T. Glass, J. C. Duchamp and H. C. Dorn, *J. Am. Chem. Soc.*, 2007, **129**, 7018–7027.
- 54 M. R. Haneline and F. P. Gabbaï, *Inorg. Chem.*, 2005, **44**, 6248–6255.

