



Cite this: *Chem. Sci.*, 2019, 10, 5719

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 27th March 2019
Accepted 1st May 2019

DOI: 10.1039/c9sc01502g

rscl.li/chemical-science

Synthesis of aminyl biradicals by base-induced Csp³–Csp³ coupling of cationic azo dyes†

Yizhu Liu,^a Paul Varava,^a Alberto Fabrizio,^a Léonard Y. M. Eymann,^a Alexander G. Tskhovrebov,^a Ophélie Marie Planes,^a Euro Solari,^a Farzaneh Fadaei-Tirani,^a Rosario Scopelliti,^a Andrzej Sienkiewicz,^b Clémence Corminboeuf^a and Kay Severin^{*,a}

The synthesis of the industrially important polymer parylene is achieved by polymerization of *p*-quinodimethane (*p*-QDM). The polymerization is thought to proceed *via* a biradical *p*-QDM dimer, but isolation or characterization of such a biradical has remained elusive. Here, we describe the synthesis of an aza-analogue of this *p*-QDM dimer. The biradical is formed by base-induced dimerization of an azoimidazolium dye. Due to the presence of sterically shielded aminyl radicals instead of terminal H₂C groups, the stability of this dimer is sufficient for analyses by ESR spectroscopy and X-ray crystallography. A similar Csp³–Csp³ coupling was observed for an azotriazolium dye, suggesting that base-induced C–C coupling reactions can be realized for different types of azo dyes.


Introduction

p-Quinodimethane (*p*-QDM, or *p*-xylylene) is an important hydrocarbon, from both a fundamental and application point of view.¹ *p*-QDM was first obtained through pyrolysis of *p*-xylene.² Later, it was found that [2.2]-*p*-cyclophane is a more convenient starting material.^{3,4} *p*-QDM readily polymerizes in the condensed phase generating poly(*p*-xylylene), also known as parylene (Scheme 1a).⁵ This polymer has interesting materials properties, and it is used for numerous applications.^{4,5}

Mechanistically, the polymerization of *p*-QDM is intriguing. Despite its radical-type reactivity,⁶ *p*-QDM is characterized by a closed-shell quinoidal electronic ground state.⁷ Recent theoretical investigations suggest that the triplet open-shell electronic state lies as high as 38 kcal mol^{−1} above the singlet closed-shell ground state.⁸ However, the initial C–C coupling step involving a singlet-to-triplet spin transition requires a lower activation energy,⁹ and it is assumed that the polymerization of *p*-QDM starts with the formation of a dimeric biradical (Scheme 1a).^{5,6,9} This proposition is supported by the fact that solutions of freshly prepared *p*-QDM give [2.2]-*p*-cyclophane and larger macrocycles, along with oligomers.¹⁰ Further evidence for a stepwise oligomerization mechanism

involving biradicals was provided by studying structurally related compounds.¹¹

The postulated biradical dimer of *p*-QDM is highly reactive,⁹ and its isolation or direct characterization has remained elusive. Here we describe the synthesis of a structural analogue of the *p*-QDM dimer, in which the terminal H₂C groups are replaced by aminyl radicals (Scheme 1b). The stability of this aminyl biradical¹² is sufficient for analyses by ESR spectroscopy and X-ray crystallography.



Scheme 1 The chemistry of *p*-QDM (a), and synthesis of an aminyl biradical by base-induced dimerization of an azo dye (b).

^aInstitut des Sciences et Ingénierie Chimiques, Ecole Polytechnique Fédérale de Lausanne (EPFL), 1015 Lausanne, Switzerland. E-mail: kay.severin@epfl.ch

^bInstitute of Physics, EPFL, Switzerland

† Electronic supplementary information (ESI) available. CCDC 1903893–1903897. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c9sc01502g




Results and discussion


Azoimidazolium salts are industrially important dyes, which are used for the dyeing of synthetic and natural fibres.¹³ We have recently reported a new route for preparing these dyes.¹⁴ Importantly, this methodology can be used for the synthesis of dyes with aryl substituents attached to the heterocycle. Subsequent studies have shown that azoimidazolium dyes with *N*-aryl groups can be reduced to give stable aminyl radicals (Scheme 2a).¹⁵ Deprotonation of these dyes with potassium bis(trimethylsilyl)amide (KHMDS) results in the formation of mesoionic carbenes ('Azo-MICs'), which can be captured with metal complexes (Scheme 2b).¹⁶

While studying the formation of mesoionic carbenes, we made an unexpected observation: when one equivalent of KHMDS was added to a solution of dye **1** (Scheme 3) in THF-*d*₈, the color of the solution changed immediately from bright orange to dark rose-brown, and the ¹H NMR signals vanished except for the solvent and HMDS peaks (Fig. S1, ESI†).

When the reaction was performed with temperature control $-78\text{ }^{\circ}\text{C} \rightarrow \text{RT}$ on a preparative scale, black crystals of **3** were isolated in 42% yield (Scheme 3). The molecular structure of **3** was unambiguously established by single crystal X-ray crystallography (Fig. 1). Compound **3** can be regarded as a dimer of the hypothetical intermediate **2**, which is formed upon deprotonation of **1**. The two parts are linked *via* an ethylene bridge. The average length of this C–C bond is 1.484 Å, which is comparable to what has been reported for *para*-*N*-containing 1,2-diphenylethanes.¹⁷ Given that compound **3** has an overall charge of zero, each side of the molecule is expected to feature one unpaired electron. In fact, compound **3** is structurally similar to our previously reported aminyl radicals (Scheme 2a).¹⁵ For example, the bond length alternation (BLA)¹⁸ of **3** as a measure of the aromaticity is 0.014 Å, and comparable values were observed for the reduced dyes.¹⁵ The N–C_{imidazole} moieties are coplanar with the heterocyclic rings, but dihedral angles of *ca.* 27° are observed between the N₂C groups and the arene rings. Moreover, like the aminyl radicals reported earlier,¹⁵ **3** is characterized with short N–C_{imidazole} (avg. 1.317 Å) bonds, long N–C_{arene} (avg. 1.386 Å) bonds, and moderate N–N (avg. 1.350 Å) bonds.



Scheme 2 Azoimidazolium dyes can be used as precursors for stable aminyl radicals (a), and they can be converted into mesoionic carbene ligands (b).



Scheme 3 The synthesis of the biradical **3** and mesomeric forms of the proposed intermediate **2**.

These structural data suggest that **3** can be described as an aminyl biradical, as depicted in Scheme 3.

The biradical nature of **3** is further supported by ESR spectroscopy. Compound **3** displays limited stability in solution, and the ESR spectrum was thus recorded in the solid state (for




Fig. 1 The molecular structure of **3** in the crystal. The ellipsoids are shown at 30% probability. The unit cell contains two independent molecules, and only one of them is shown here. Hydrogen atoms, except those on the ethylene bridge, and solvent molecules are omitted for clarity. Selected bond lengths (Å) and bond angles (°), averaged between the two asymmetric molecules in the unit cell: C30–C31 1.458(6), C25–C30 1.512(6), C24–C25 1.375(8), C23–C24 1.395(6), C22–C23 1.417(6), C22–C27 1.408(7), C26–C27 1.393(6), C25–C26 1.394(7), C22–N4 1.386(5), N3–N4 1.343(5), N3–C1 1.331(5), C1–N1 1.365(5), C1–N2 1.371(6), C31–C32 1.522(6), C32–C33 1.383(8), C33–C34 1.382(5), C34–C35 1.415(7), C35–C36 1.426(7), C36–C37 1.398(6), C32–C37 1.395(8), C35–N5 1.386(5), N5–N6 1.338(5), N6–C40 1.311(5), C40–N7 1.365(6), C40–N8 1.378(6), C25–C30–C31 113.8(4), C22–N4–N3 114.6(4), N4–N3–C1 113.7(4), C30–C31–C32 113.6(3), C35–N5–N6 115.7(4), N5–N6–C40 114.3(4). Note that averaged values are given in the main text.




details, see ESI†). The spectrum pointed to a spin doublet with the g -factor of 2.0037 (ESI, Fig. S18†). This value is again very similar to what was found for monomeric aminyl radicals prepared by reduction of azoimidazolium dyes.¹⁵

The generation of the biradical **3** by reaction of **1** with base is an unexpected finding, considering that other azoimidazolium dyes are deprotonated at the heterocycle to give mesoionic carbenes (Scheme 2b).¹⁶ The apparent $\text{Csp}^3\text{-Csp}^3$ coupling reaction and the generation of a biradical suggests a radical pathway, notwithstanding the fact that both the starting material **1** and the base are closed-shell compounds. Therefore, we propose that the observed reactivity resembles that of p -QDM. Deprotonation of **1** at the benzylic position gives zwitterionic **2A**, which is in resonance with the quinoidal form **2B** (Scheme 3). Furthermore, a diradical¹² structure is conceivable (**2C**). According to density functional theory (DFT) computations at (U)TPSS-D3/cc-pVDZ level, the open-shell triplet form of **2C** is energetically disfavored by 25 kcal mol⁻¹ compared to the closed-shell quinoidal structure **2B** (see ESI†). As discussed in the introduction, a similar situation is found for p -QDM.⁸ Moreover, the optimized structure of **2** shows very similar structural parameters for the central benzene ring and the terminal methylene group as p -QDM (ESI, Table S1†).^{7a} All these results suggest that the electronic structures of **2** and p -QDM are similar. The formation of the biradical **3** can thus be regarded as an aza-version of the p -QDM dimerization. Further oligomerization is prohibited by steric hindrance of the aminyl radical.

The biradical **3** is labile in solution and spontaneously converts into compound **4** at a slow rate. In fact, crystallization of **3** was only achievable at low temperature (−40 °C), whereas attempts performed at RT gave crystals of **4**. The conversion of **1** into **4** can be facilitated with an external hydrogen atom abstractor such as TEMPO, and the best yield was obtained when using KO t Bu as base (Scheme 4). In comparison to the dark rose brown parental **3**, **4** displays a turquoise color in solution.

Two resonance structures have to be considered for **4**: a closed-shell, quinoidal form **4A**, and an open-shell diradical¹² form **4B**, which would benefit from the formation of two



Scheme 4 The synthesis of compound **4**.

aromatic rings.²⁰ Experimental (X-ray, NMR) and computational results suggest that **4** is best described as a closed shell system with low diradicaloid character.²⁰

The molecular structure of compound **4** in the crystal is depicted in Fig. 2. Comparison with **3** in some structural parameters is given in Table S3 (ESI†). In contrast to **3**, the central hydrocarbon fragment of **4** is essentially co-planar, consistent with a more conjugated framework. Significant changes for the bridging moieties are observed, including an increased “zigzag” angle (avg. 112.8° in **3** vs. 124.6° in **4**), and a shortened distance of the central C–C bond (1.484 Å in **3** vs. 1.408 Å in **4**). The bridging hydrocarbon fragment undergoes considerable quinoidization, with C25–C29, C23–C24 (and C26–C27), and C22–41 bonds shortened substantially, and the BLA increased to 0.07 Å (compared to 0.014 Å for **3**). It is worth noting that the BLA of **4** is larger than that of ‘Tschitschibabin’s hydrocarbon’ (Ph₂C(C₆H₄)₂CPh₂, BLA = 0.052 Å, see Table S3, ESI,† for more details).²¹ This compound has been studied extensively, and it was concluded to have a closed-shell quinoidal structure in the ground state.^{20,21}

DFT computations ((U)TPSS-D3/cc-pVDZ) suggest that **4** is characterized by a closed-shell ground state, stabilized relative to the open-shell triplet and singlet states by 8.6 and 38.8 kcal mol⁻¹, respectively.

In line with these results, a solution of **4** in THF-*d*₈ gave rise to a well-resolved ¹H NMR spectrum. At room temperature, only one set of signals was observed for the four mesityl groups. Apparently, rotation around the C_{imidazole}–N bond is fast on the NMR time scale. At lower temperatures, however, the ¹H NMR spectra are more complex, indicating a reduced rotational freedom.

The ESR characterization of **4** was also performed. Surprisingly, the ESR spectrum suggested a spin doublet. However, the intensity of the signal was weak, and its percentage varied from




Fig. 2 The molecular structure of **4** in the crystal. The unit cell contains two independent molecules, and only one of them is shown here. The ellipsoids are shown at 30% probability. Hydrogen atoms, except those on the vinyl bridge, and solvent molecules, are omitted for clarity. Selected bond lengths (Å) and bond angles (°): C29–C29′ 1.403(4), C29–C25 1.415(3), C24–C25 1.421(3), C23–C24 1.371(2), C22–C23 1.452(2), C22–C27 1.452(2), C26–C27 1.367(2), C25–C26 1.421(3), C22–N4 1.339(2), N3–N4 1.357(2), N3–C1 1.321(2), C1–N2 1.366(2), C1–N1 1.366(2), C25–C29–C29′ 124.9(2), C22–N4–N3 116.2(2), N4–N3–C1 112.4(2). Note that averaged values are given in the main text.




batch to batch. We assume that the signal is due to small impurities of a radical cation, which is formed from **4** by oxidation. In fact, compound **4** is a strong reducing agent with a reversible redox transition at -0.76 V vs. ferrocenium/ferrocene (Fc^+/Fc) as revealed by cyclic voltammetry (for details see ESI[†]). It should be noted that ESR-active impurities have been observed for structurally related compounds such as 'Tschitschibabin's hydrocarbon'.²¹

In order to examine if similar $\text{Csp}^3\text{-Csp}^3$ coupling reactions could be observed for other cationic azo dyes, we have investigated the reaction of the azotriazolium dye **5** (ref. 14a) with KHMDS. When base was added to a solution of **5** in THF- d_8 , vanishing of the ^1H NMR signals was observed, as in the case of **1** (ESI, Fig. S8[†]). Attempts to isolate a defined product from reactions on preparative scale were first not successful. However, when two equivalents of KHMDS were employed, we were able to isolate the *N*-demethylated dimer **6** in 36% yield (Scheme 5). Dark brown-green crystals, suitable for X-ray crystallography, were obtained by slow diffusion of pentane into a THF solution of **6** at -40 °C.

Fig. 3a shows the structure of **6** in the crystal. As observed for **1**, the reaction with KHMDS had resulted in a C-C coupling reaction involving the *p*- CH_3 groups of the mesityl substituent. The dimer has an overall charge of minus two, and charge compensation is achieved by two potassium ions. The latter are coordinated in a κ^2 -type fashion to the heterocycle, with the remaining coordination sites being occupied by THF molecules (not shown). The shorter K1-N2 distance (2.810(4) Å) compared to the K1-N3 distance (2.969(5) Å) suggests a higher charge density at the N2 atom. Both N-K distances fall in the range reported for compounds with a side-on $\kappa^2\text{-K}(\text{N}=\text{N})^-$ motif.²² Compared to the structure of the starting material **5**,^{14a} a comprehensive expansion of the triazole rings is observed, with a lengthening of all five bonds. This change is consistent with the conversion of a cationic into an anionic ring system.

Like **3**, compound **6** was expected to feature two unpaired electrons. This assumption is in line with the observed



Scheme 5 Synthesis of **6** and identification of **7** by X-ray diffraction crystallography.





Fig. 3 (a) The molecular structure of **6** in the crystal. The ellipsoids are shown at 30% probability. Hydrogen atoms, except those on the ethylene bridge, and THF molecules are omitted for clarity. Selected bond lengths (Å) and bond angles (°): C30–C30' 1.453(9), C30–C26 1.483(7), C25–C26 1.389(7), C24–C25 1.381(7), C23–C24 1.445(6), C23–C28 1.434(6), C27–C28 1.368(7), C26–C27 1.436(7), C23–N5 1.365(6), N4–N5 1.348(5), N4–C2 1.340(6), C1–C2 1.411(6), C1–N3 1.358(7), N2–N3 1.327(5), N1–N2 1.383(6), N1–C2 1.385(6), K1–N2 2.810(4), K1–N3 2.969(5), C26–C30–C30' 119.5(6), C23–N5–N4 115.2(4), N5–N4–C2 113.3(4). (b) Total spin density (isovalue: ± 0.01) of the open-shell singlet ground state of **6**. Blue positive spin density (excess α -spin), orange negative spin density (excess β -spin). K^+ cations and coordinating THF molecules were omitted in the computations.

vanishing of the ^1H NMR signals of **5** upon addition of base. Further evidence of its biradical nature was obtained through ESR measurements. The ESR spectrum acquired at room temperature for **6** in the solid state pointed to a spin doublet with the *g*-factor of 2.0036 (ESI, Fig. S19[†]). The electronic structure computations ((U)TPSS-D3/cc-pVDZ) shows that dimer **6** favors an open-shell singlet ground state, with the triplet state almost isoenergetic. The largest spin density is found on the azo bridge, with the nitrogen atoms accounting symmetrically and in pairs for 21% (N5 and N10) and for 12% (N4 and N9) of the α - and β -spin densities (Fig. 3b).

Crystallization of a THF solution of **6** under air gave bright red crystals of the neutral dimer **7**, the structure of which was analyzed by X-ray crystallography (ESI, Fig. S20[†]). The ethylene bridge is well preserved, while the remaining parts of the molecule restore complete aromaticity, with small BLA values (0.019 Å) for the benzene rings, well-defined azo ($\text{N}=\text{N}$) bonds, and characteristic neutral triazole rings.

The generation of the biradical **6** from the cationic monomer **5** requires two chemical transformations, namely *N*-demethylation of the heterocycle,²³ and base-induced C-C coupling. *A priori*, two reaction pathways can be envisaged, depending on the sequence of these two transformations (Scheme 6). In order to address this point, we have synthesized the neutral azo dye





Scheme 6 Possible pathways for the formation of the biradical 6.

12, which represents the expected intermediate if *N*-demethylation proceeds before C–C coupling (Pathway II). The synthesis of 12 was accomplished by demethylation of the known diazotate 10 (ref. 14a) with LiHMDS, followed by azo coupling with mesitylene (Scheme 6, for a crystallographic characterization of the intermediate lithium salt 11, see ESI†). Compound 12 turned out to be inert towards KHMDS. Consequently, we propose that the formation of 6 proceeds *via* Pathway I, involving the initial formation of 8, which displays a *p*-QDM-like reactivity. Dimerization gives the biradical 9, which undergoes a two-fold *N*-demethylation to afford 6.

Conclusions

In summary, we have shown that cationic azo dyes with diazenylmesityl groups can undergo a base-induced Csp³–Csp³ coupling reaction to give biradicals. The presence of a CH₃ group in the *para* position to the azo group is essential for this coupling reaction, because dyes with CH₃ groups in *ortho* or *meta* position are deprotonated at imidazolium heterocycle.¹⁶ The dimerization of the azo dyes is proposed to

involve the formation of the neutral monomers 2 and 8, which display *p*-QDM-like structures and reactivity.

The polymerization of *p*-QDM is an intriguing reaction, because it appears to proceed *via* a radical mechanism, even though *p*-QDM has a closed-shell structure. It is generally assumed that the polymerization is initiated by the formation of a biradical *p*-QDM dimer, but the characterization of this highly reactive intermediate has remained elusive. Our results show that it is possible to intercept dimers with a biradical structure if the terminal radical sites are stabilized. In the case of 3 and 6, this stabilization is achieved by replacing the H₂C groups with sterically shielded aminyl radicals.

The dimerization reactions described in this contribution are also interesting from a synthetic chemistry point of view, because the homo-coupling of azo dyes represents a new way to synthesize more complex dye structures. Polyconjugated compounds such as 4, for example, could also be of interest for optoelectronic applications.^{19a,20a}

Conflicts of interest

There are no conflicts to declare.

Author contributions

Y. L., P. V., L. Y. M. E. and A. G. T. synthesized and analyzed the compounds, A. F. and C. C. performed the computational study, O. M. P. performed the CV experiments, E. S., F. F.-T., and R. S. carried out the crystallographic analyses, A. S. performed the ESR analyses, K. S. initiated and coordinated the study, and all authors except E. S. and A. G. T. contributed to writing of the manuscript.

Acknowledgements

Y. L. was an EPFL Fellow co-funded by the Marie Skłodowska-Curie program (grant No. 665667). The work was supported by the Swiss National Science Foundation, and by the Ecole Polytechnique Fédérale de Lausanne (EPFL). We thank Abdusalom Suleymanov for help with the ESR measurements, and Emilie Baudat for NMR measurements.

References

- (a) A. Konishi and T. Kubo, *Top. Curr. Chem.*, 2017, **375**, 83; (b) J. Casado, *Top. Curr. Chem.*, 2017, **375**, 73.
- (a) L. A. Errede and M. Szwarc, *Q. Rev., Chem. Soc.*, 1958, **12**, 301; (b) M. Szwarc, *J. Chem. Phys.*, 1948, **16**, 128; (c) M. Szwarc, *Nature*, 1947, **160**, 403; (d) M. Szwarc, *Discuss. Faraday Soc.*, 1947, **2**, 46–49.
- W. F. Gorham, *J. Polym. Sci., Part A-1: Polym. Chem.*, 1966, **4**, 3027–3039.
- H. Hopf, *Angew. Chem., Int. Ed.*, 2008, **47**, 9808.
- (a) C. P. Tan and H. C. Craighead, *Materials*, 2010, **3**, 1803; (b) J. B. Fortin and T.-M. Lu, *Chemical vapor deposition polymerization: the growth and properties of parylene thin*



- films*, Kluwer, Boston, 2004; (c) M. Szwarc, *Polym. Eng. Sci.*, 1976, **16**, 473.
- 6 L. A. Errede and J. M. Hoyt, *J. Am. Chem. Soc.*, 1960, **82**, 436.
- 7 For the spectroscopic characterization of p-DQM, see: (a) P. G. Mahaffy, J. D. Wieser and L. K. Montgomery, *J. Am. Chem. Soc.*, 1977, **99**, 4514; (b) T. Koenig, R. Wielesek, W. Snell and T. Balle, *J. Am. Chem. Soc.*, 1975, **97**, 3225; (c) Y. Yamakita, Y. Furukawa and M. Tasumi, *Chem. Lett.*, 1993, **22**, 311; (d) J. M. Pearson, H. A. Six, D. J. Williams and M. Levy, *J. Am. Chem. Soc.*, 1971, **93**, 5034; (e) D. J. Williams, J. M. Pearson and M. Levy, *J. Am. Chem. Soc.*, 1970, **92**, 1436.
- 8 M. Bobrowski, P. Skurski and S. Freza, *Chem. Phys.*, 2011, **382**, 20.
- 9 K. Smalara, A. Gieldoń, M. Bobrowski, J. Rybicki and C. Czaplewski, *J. Phys. Chem. A*, 2010, **114**, 4296.
- 10 (a) W. S. Trahanovsky and S. P. Lorimor, *J. Org. Chem.*, 2006, **71**, 1784; (b) L. A. Errede, R. S. Gregorian and J. M. Hoyt, *J. Am. Chem. Soc.*, 1960, **82**, 5218.
- 11 (a) D. A. Klumpp, T. M. Gilbert and W. S. Trahanovsky, *Eur. J. Org. Chem.*, 2016, 5559; (b) W. S. Trahanovsky and D. A. Klumpp, *Tetrahedron Lett.*, 2016, **57**, 2386; (c) X. Fu and D. Zhao, *Org. Lett.*, 2015, **17**, 5694; (d) W. S. Trahanovsky, D. L. Miller and Y. Wang, *J. Org. Chem.*, 1997, **62**, 8980.
- 12 The term 'biradical' is used for molecules in which the two unpaired electrons are largely independent. Electronically coupled systems are referred to as 'diradicals' or 'diradicaloids'. For a more detailed discussion see: M. Abe, *Chem. Rev.*, 2013, **113**, 7011.
- 13 R. Raue, *Methine Dyes and Pigments*. Ullmann's Encyclopedia of Industrial Chemistry, 2000, pp. 1–55.
- 14 (a) L. Y. M. Eymann, R. Scopelliti, F. Fadaei Tirani and K. Severin, *Chem.–Eur. J.*, 2018, **24**, 7957; (b) A. G. Tskhovrebov, L. C. E. Naested, E. Solari, R. Scopelliti and K. Severin, *Angew. Chem., Int. Ed.*, 2015, **54**, 1289.
- 15 L. Y. M. Eymann, A. G. Tskhovrebov, A. Sienkiewicz, J. L. Bila, I. Živković, H. M. Rønnow, M. D. Wodrich, L. Vannay, C. Corminboeuf, P. Pattison, E. Solari, R. Scopelliti and K. Severin, *J. Am. Chem. Soc.*, 2016, **138**, 15126.
- 16 F. M. Chadwick, B. F. E. Curchod, R. Scopelliti, F. Fadaei Tirani, E. Solari and K. Severin, *Angew. Chem., Int. Ed.*, 2019, **58**, 1764.
- 17 For selected examples, see: (a) B. Manna, B. Joarder, A. V. Desai, A. Karmakar and S. K. Ghosh, *Chem.–Eur. J.*, 2014, **20**, 12399; (b) N. Wang, J.-G. Ma, W. Shi and P. Cheng, *CrystEngComm*, 2012, **14**, 5634; (c) B. Manna, A. K. Chaudhari, B. Joarder, A. Karmakar and S. K. Ghosh, *Angew. Chem., Int. Ed.*, 2012, **52**, 998; (d) A. Ghames, T. Douadi, D. Haffar, S. Chafaa, M. Allain, M. A. Khan and G. M. Bouet, *Polyhedron*, 2006, **25**, 3201; (e) M. Albrecht, I. Janser, H. Houjou and R. Fröhlich, *Chem.–Eur. J.*, 2004, **10**, 2839; (f) Z. Feng, A. Fan, S. Valiyaveetil and J. J. Vittal, *Cryst. Growth Des.*, 2003, **3**, 555; (g) P. Mandal, S. Paul, K. Goubitz and H. Schenik, *Mol. Cryst. Liq. Cryst. Sci. Technol., Sect. A*, 1995, **258**, 209.
- 18 M. Kertesz, C. H. Choi and S. Yang, *Chem. Rev.*, 2005, **105**, 3448.
- 19 (a) J. Messelberger, A. Grünwald, P. Pinter, M. M. Mansmann and D. Munz, *Chem. Sci.*, 2018, **9**, 6107; (b) S. Grimme and A. Hansen, *Angew. Chem., Int. Ed.*, 2015, **54**, 12308.
- 20 For selected review articles about diradicaloids, see: (a) X. Hu, W. Wang, D. Wang and Y. Zheng, *J. Mater. Chem. C*, 2018, **6**, 11232; (b) Y. Huang and E. Egap, *Polym. J.*, 2018, **50**, 603–614; (c) T. Y. Gopalakrishna, W. Zeng, X. Lu and J. Wu, *Chem. Commun.*, 2018, **54**, 2186; (d) G. Tan and X. Wang, *Acc. Chem. Res.*, 2017, **50**, 1997–2006; (e) Y. Ni and J. Wu, *Tetrahedron Lett.*, 2016, 557, 5426; (f) G. Gryn'ova, M. L. Coote and C. Corminboeuf, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2015, **5**, 440; (g) Z. Zeng, X. Shi, C. Chi, J. T. Navarette, J. Casado and J. Wu, *Chem. Soc. Rev.*, 2015, **44**, 6578.
- 21 L. K. Montgomery, J. C. Huffman, E. A. Jurczak and M. P. Grendze, *J. Am. Chem. Soc.*, 1986, **108**, 6004.
- 22 (a) C. Pi, Y. Wang, W. Zheng, L. Wan, H. Wu, L. Weng, L. Wu, Q. Li and P. von Ragué Schleyer, *Angew. Chem., Int. Ed.*, 2010, **49**, 1842; (b) K. Kunz, H.-W. Lerner and M. Bolte, *Acta Crystallogr.*, 2009, **E65**, m171; (c) M.-Á. Velázquez-Carmona, A.-J. Metta-Magaña, S.-A. Cortés-Llamas, V. Montiel-Palma and M.-Á. Muñoz-Hernández, *Polyhedron*, 2009, **28**, 205; (d) L. Wan, C. Pi, L. Zhang, W. Zheng, L. Weng, Z. Chen and Y. Zhang, *Chem. Commun.*, 2008, 2266; (e) T. J. Hebden, W. W. Brennessel, C. J. Flaschenriem and P. L. Holland, *Dalton Trans.*, 2006, 3855; (f) S.-Á. Cortés-Llamas, R. Hernández-Lamonedá, M.-Á. Velázquez-Carmona, M.-A. Muñoz-Hernández and R. A. Toscano, *Inorg. Chem.*, 2006, **45**, 286; (g) X.-R. Hu, Y.-W. Wang and J.-M. Gu, *Acta Crystallogr.*, 2005, **E61**, m1686.
- 23 For examples of N-dealkylation of 1,2,3-triazolium salts see: (a) Z. Monasterio, A. Irastorza, J. I. Miranda and J. M. Aizpurua, *Org. Lett.*, 2016, **18**, 2511; (b) R. Maity, T. Tichter, M. van der Meer and B. Sarkar, *Dalton Trans.*, 2015, **44**, 18311; (c) S. Koguchi and K. Izawa, *Synthesis*, 2012, **44**, 3603; (d) J. Bouffard, B. K. Keitz, R. Tonner, G. G. Barrios, G. Frenking, R. H. Grubbs and G. Bertrand, *Organometallics*, 2011, **30**, 2617.

