

Cite this: *Chem. Sci.*, 2019, 10, 9483

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 18th July 2019
Accepted 15th August 2019

DOI: 10.1039/c9sc03565f

rsc.li/chemical-science

Design of non-ionic carbon superbases: second generation carbodiphosphoranes†

Sebastian Ullrich,^a Borislav Kovačević,^b Björn Koch,^a Klaus Harms^{ID}^a and Jörg Sundermeyer^{ID}^{*a}

A new generation of carbodiphosphoranes (CDPs), incorporating pyrrolidine, tetramethylguanidine, or tris(dimethylamino)phosphazene as substituents is introduced as the most powerful class of non-ionic carbon superbases on the basicity scale to date. The synthetic approach as well as NMR spectroscopic and structural characteristics in the free and protonated form are described. Investigation of basicity in solution and in the gas phase by experimental and theoretical means provides the to our knowledge first reported pK_{BH}^+ values for CDPs in the literature and suggest them as upper tier superbases.

Introduction

Much theoretical and synthetic effort has been devoted to lift non-ionic organic bases to the basicity level of common inorganic or metalorganic bases.^{1,2} With his famous phosphazenes Schwesinger established a widely used and commercially available class of (organo-)superbases.^{3,4} His homologization concept, the stepwise expansion of the molecular scaffold in order to better delocalize the positive charge formed upon protonation, was also applied to synthesize higher-order N-superbases of guanidines,^{5,6} imidazolidine amines⁷ and cyclopropeneimines.^{8,9} However, such basicity enhancement is accompanied by an unwanted growth of the bases' molecular weight. Therefore, other strategies for augmenting the intrinsic proton affinity have been investigated: in proton sponges, a second nitrogen basicity centre in close proximity to the first one increases the pK_{BH}^+ value up to 16 orders of magnitude by intramolecular hydrogen bonding compared to corresponding non chelating bases.¹⁰ Additional thermodynamic driving force comes from relief of strain of the aromatic backbone.¹¹ Many derivatives of such proton sponges were designed by combining aforementioned superbasic functionalities with the 1,8-diaminonaphthalene structural motif¹² or as proton pincers with different backbones.¹³

Atoms other than nitrogen as basicity centre were also applied, such as phosphorus.^{14,15} Recently, we demonstrated, that N-phosphazenylyl substituted phosphines (PAPs) possess higher pK_{BH}^+ values as P^{III} bases than their corresponding

phosphazene P^VNzBu counterparts as N bases.¹⁶ So far the limit of homologization is reached at the P₇ level both in phosphazenylyl phosphazenes and phosphazenylyl phosphines as both P₇ benchmark bases have only been isolated in their protonated form.^{16,17}

Non-ionic carbon is another contender to extend the basicity ladder to unmatched regions.¹⁸ In this respect phosphorus (mono-)ylides^{19,20} as well as bisylidic proton sponges²¹ were investigated on theoretical and experimental level. Although identified as potential superbases, the application of N-heterocyclic carbenes (NHCs),²² cyclic alkyl amino carbenes (CAACs),²³ carbodicarbenes (CDCs),²⁴ and carbodiphosphoranes (CDPs)²⁵ has been exploited predominantly as strong Lewis bases towards transition and main group elements other than the proton.²⁶

The prototypic hexaphenyl carbodiphosphorane ((Ph)₆-CDP) was first synthesized 1961 by Ramirez *et al.*²⁷ Further compounds like the hexamethyl carbodiphosphorane ((Me)₆-CDP),²⁸ hexakis(dimethylamino) carbodiphosphorane ((dma)₆-CDP),²⁹ and mixed representatives followed.^{30–32}

Herein we promote carbodiphosphoranes with their electron-rich R₃P–C–PR₃ functionality as exceptionally strong carbon Brønsted bases. As bisylides with a π -symmetric HOMO and σ -symmetric HOMO–1, both mainly located as lone pairs at the carbon, only slightly stabilized by back-bonding *via* negative hyperconjugation,³³ they provide outstanding pK_{BH}^+ values in particular for the first of two protonation steps. We present a synthesis for hexa(pyrrolidino) carbodiphosphorane ((pyrr)₆-CDP) with its calculated first and second proton affinity (PA) of 287.6 and 188.9 kcal mol^{–1},³⁴ which exceeds the PAs of (Ph)₆-CDP (280.0 and 185.6 kcal mol^{–1})³⁴ and (dma)₆-CDP (279.9 and 174.9 kcal mol^{–1}).³⁴ Furthermore we apply the homologization concept to CDPs by introducing PR₂R' units bearing one intrinsically superbasic substituent R' to access CDP

^aFachbereich Chemie, Philipps-University Marburg, Hans-Meerwein-Straße, 35032 Marburg, Germany. E-mail: jsu@staff.uni-marburg.de

^bThe Group for Computational Life Sciences, Rudjer Bošković Institute, Bijenička c. 54, HR-10000 Zagreb, Croatia

† Electronic supplementary information (ESI) available. CCDC 1903830, 1903833, 1903838, 1903840, 1903841 and 1903843. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c9sc03565f



superbases of second-order.⁸ We thereby focused on *N*-tetramethylguanidynyl (tmg) and *N*-tris(dimethylamino)phosphazanyl (dmaP₁) substituents targeting new carbodiphosphoranes *sym*-(tmg)₂(dma)₄-CDP and *sym*-(dmaP₁)₂(dma)₄-CDP.

Results and discussion

Synthesis

We experienced, that the established synthesis routes to CDPs are inappropriate for phosphines more electron-rich than P(NMe₂)₃: reactions between such phosphines P(NR₂)₂R' and CCl₄ did not follow the pattern outlined in ref. 32 and 35 but exclusively led to chlorination of the phosphine, whilst reactions with methylene bromide did not selectively follow the path outlined in ref. 30 and 36, but led to a 1 : 1-mixture of the methylated phosphonium bromide [R'(NR₂)₂P-Me]Br and the brominated species [R'(NR₂)₂P-Br]Br. Therefore we further developed an alternative strategy laid out by Appel *et al.* for the synthesis of (dma)₆-CDP.²⁹ The doubly protonated precursors of the second-order carbodiphosphorane superbases, *sym*-(tmg)₂(dma)₄-CDP (**1**) and *sym*-(dmaP₁)₂(dma)₄-CDP (**2**), were obtained in an oxidative imination sequence as shown in Scheme 1. Bis[bis(dimethylamino)phosphino]methane (**3**) was oxidized by CCl₄ in presence of tetramethylguanidine (Htmg) or tris(dimethylamino)phosphazene ((dma)P₁-H) instead of dimethylamine as nucleophile and auxiliary base. This reaction offers the advantage of preformed C–P bonds avoiding the preparation of respective P^{III} nucleophiles.^{15,20,37} **3** is readily synthesized in two steps on a large scale³⁸ and the selected superbasic building blocks oxidatively introduced as nucleophiles are either commercially available or easily accessible in few steps.⁴


The synthesis of **4**·2HBF₄, the precursor for (pyrr)₆-CDP **4**, was accomplished in a one-pot synthesis (Scheme 2), since the intermediate bis[di(pyrrolidino)phosphino]methane (**5**) turned

out to decompose upon vacuum distillation. Starting from bis(dichlorophosphino)methane³⁸ (**6**), **5** was prepared *in situ* with an excess of pyrrolidine (Fig. S1 in the ESI†) and directly oxidized with CCl₄.

In all three reactions the respective monoprotonated hydrochloride adducts were identified as products *via* ³¹P NMR spectroscopy. Therefore the second p*K*_{BH}⁺ values in THF of these new CDPs are obviously lower than that of the auxiliary base pyrrolidine (13.5),³⁹ tetramethylguanidine (15.5),⁴⁰ or tris(dimethylamino)phosphazene **2a** (19.7),⁴⁰ respectively. For purification, the crude products were precipitated with NaBF₄ from aqueous solution. These conditions lead to second protonation at the central carbon atom and a strongly alkaline solution. Therefore, even the monoprotonated CDPs can be considered as strong cationic bases in aqueous medium. Similar behaviour was found for (Ph)₆-CDP in water, although the latter is slowly hydrolysed under ambient conditions,²⁷ which is not the case for peraminated CDPs **1**, **2** and **4** reported here.


The bis(tetrafluoridoborate) salts of **1**, **2** and **4** were obtained in 50–60% yield as water and air stable, colourless solids, indefinitely storable. They are well soluble in polar organic solvents like methanol, acetonitrile or DMSO but insoluble in less polar solvents such as ethers and hydrocarbons.

For the liberation of the free CDPs different suitable bases were identified: for **4** potassium bis(trimethylsilyl)amide (KHMSD) is of sufficient basicity, whilst for **1** the more basic sodium amide (NaNH₂) is necessary for full deprotonation. Both new bases **1** and **4** could be isolated in 70% and 60% yield, respectively, from *n*-hexane as pure colourless crystalline solids, indefinitely storable at room temperature under inert conditions. Contrastingly we were not able to isolate **2** as free CDP base form. Sodium amide in liquid ammonia or suspended in THF at room temperature selectively abstracts the first proton under formation of **2**·HBF₄ as colourless solid in 69% yield. At elevated temperature the central carbon atom is not further



Scheme 1 Preparation of CDP precursors **1**·2HBF₄ and **2**·2HBF₄ together with subsequent deprotonation to **1** (one exemplary mesomeric structure displayed) and **7**, respectively. Numbering schemes refer to assigned NMR signals in the experimental section.





Scheme 2 *In situ* preparation of 5 with subsequent oxidation by CCl₄ in presence of excess of pyrrolidine (Hpyrr) to 4·2HBF₄. Deprotonation with KHMDS lead to the free CDP 4 (displayed in exemplarily bisylidic notation). The numbering scheme refers to assigned NMR signals in the experimental section.

deprotonated, even though it is the thermodynamically most acidic site (see Theoretical Calculations). Instead NaNH₂ deprotonates selectively one of the dimethylamino groups at the terminal phosphazene moiety which results in the irreversible elimination of *N*-methylmethanimine and reduction of the phosphazene to a phosphine (Scheme 1). A related deprotonation and reduction of tetrakis(dimethylamino)phosphonium bromide under the action of NaNH₂ was described by Pinchuk *et al.*⁴¹ In case of 2 this reaction is slow but highly selective and 7 could be obtained as sole product as pale yellow highly viscous oil. The proposed configuration was confirmed *via* ¹H, ¹³C, and ³¹P NMR spectroscopy and by HR mass spectrometry. 7 can be considered as a hybrid between mixed valence phosphazeny phosphines^{15,16} and ylidic P^{III}/P^V compounds of the type (Me₂N)₃P=C(H)-PR₂ (ref. 42) or other ylide-functionalized phosphines.⁴³ Further attempts to deprotonate 2·2HBF₄ with other bases or reducing agents resulted either in only single deprotonation (benzyl potassium in THF), in an unselective disintegration (*n*BuLi) or in the same deprotonation of the P-NMe₂ group (potassium in liquid ammonia, ethylene diamine, THF, or DME or an excess of benzyl potassium in THF). The reaction of potassium hydride in THF gave a mixture of 7 as minor component and presumably free CDP 2 as major product by means of ³¹P NMR spectroscopy (Fig. S29 in the ESI[†]). Clearly the acidity of P^V-attached NMe₂ groups limits the accessibility of 2. Under the action of excess of strong inorganic bases at elevated temperatures the stability limit of these phosphazene moieties seems to have been reached.

For analytical reasons the monoprotonated forms of 1 and 4 were prepared on NMR scale either *via* commutation between the free CDP and its bisprotonated form or by protonating the free CDPs with one equivalent triflimidic acid (HTFSI).

Structural features

For X-ray structure determination suitable single crystals were obtained from *n*-hexane for both presented CDPs 4 and 1. They crystallize solvent-free in space group *P2₁/c* or *Pbca*, respectively, with one complete molecule per asymmetric unit (Fig. 1). Contrary to the parent compound (dma)₆-CDP, one of the hitherto two reported linear CDPs,^{29,44} a bent structure with P-C-P angles of 155.9(2)° and 147.30(9)°, respectively is found. Since the potential for bending at the central P-C-P carbon atom in polymorphic (Ph)₆-CDP is very flat⁴⁴ and reveals high dependence of the crystallization method,⁴⁵ the obtained

crystals of (dma)₆-CDP from the melt are maybe the reason for its linearity.²⁹ The P-C_{central} distances are with 1.606 Å (4) and 1.618 Å (1) in the for CDPs reported range: (dma)₆-CDP: 1.584(1) Å,²⁹ (Me)₆-CDP: 1.594(3) Å,⁴⁶ (Ph)₆-CDP: 1.601–1.635 Å.^{44,47} On average, pyrrolidine N-P distances in 4 are 1.68 Å while those of dma and tmg groups in 1 are 1.70 Å and 1.66 Å respectively.

Single crystals obtained from reaction control samples during the synthesis of 4·2HBF₄ turned out to be a cocrystallizate of 4·2HCl and pyrrolidinium chloride (Fig. 2). Cations and anions form a C-H...Cl...H-N hydrogen bond network with C...Cl distances of 3.600(2) Å and N...Cl distances of 3.018(2) Å and




Fig. 1 Molecular structure of 4 (top) and 1 (bottom). Hydrogen atoms omitted for clarity, ellipsoids at 50% probability. Selected bond length/Å and angles/°: 4 P1-C1 1.605(2), P1-N1 1.672(2), P1-N2 1.678(2), P1-N3 1.694(2), P2-C1 1.606(2), P2-N4 1.699(2), P2-N5 1.669(2), P2-N6 1.671(2), P1-C1-P2 155.9(2), C1-P1-N1 110.2(1), C1-P1-N2 115.1(1), C1-P1-N3 121.8(1), C1-P2-N4 118.4(1), C1-P2-N5 111.3(1), C1-P2-N6 117.1(1), N1-P1-C1-P2 168.0(4), N4-P2-C1-P1 130.6(4). 1 P1-C1 1.619(1), P1-N4 1.680(1), P1-N5 1.714(1), P1-N1 1.665(1), N1-C2 1.298(2), N2-C2 1.377(2), N3-C2 1.382(2), P2-C1 1.617(1), P2-N9 1.719(1), P2-N10 1.680(1), P2-N6 1.664(1), N6-C11 1.299(2), N7-C11 1.376(2), N8-C11 1.379(2), P2-C1-P1 147.30(9), C1-P1-N4 109.52(6), C1-P1-N5 121.56(6), C1-P1-N1 119.85(6), C2-N1-P1 128.1(1), C1-P2-N9 120.76(6), C1-P2-N10 110.08(6), C1-P2-N6 119.47(6), C11-N6-P2 127.3(1), N4-P1-C1-P2 162.2(2), N10-P2-C1-P1 155.8(2).






Fig. 2 Molecular structure of $4 \cdot 2\text{HCl}$ with pyrrolidinium chloride as cocrystallite as well as of $1 \cdot 2\text{HBF}_4$ and $2 \cdot 2\text{HBF}_4$ (only one of the two independent molecules depicted, structure factors given for both). Peripheral hydrogen atoms and BF_4^- anions omitted for clarity, ellipsoids at 50% probability. # marked atoms generated via a 2-fold axis through C1. Selected bond length/Å and angles/ $^\circ$: $4 \cdot 2\text{HCl}$ P1–C1 1.799(1), P1–N1 1.612(2), P1–N2 1.630(2), P1–N3 1.616(2), P1–C1–P1# 119.5(1), N1–P1–C1 103.26(9), N2–P1–C1 109.07(7), N3–P1–C1 115.21(8), N1–P1–C1–P1# 177.03(7), C1–H1A...Cl2 3.600(2), 173.8; C1–H1B...Cl2# 3.600(2), 173.8; N4–H18A...Cl2 3.048(2), 174(3); N4–H19A...Cl1 3.018(2), 172(3). $1 \cdot 2\text{HBF}_4$ P1–C19 1.820(2), P1–N4 1.644(2), P1–N5 1.639(2), P1–N1 1.580(2), N1–C1 1.330(3), N2–C1 1.351(3), N3–C1 1.346(3), P2–C19 1.822(2), P2–N9 1.640(2), P2–N10 1.643(2), P2–N6 1.586(2), N6–C10 1.335(3), N7–C10 1.332(3), N8–C10 1.349(3), P1–C19–P2 113.4(1), N4–P1–C19 104.3(1), N5–P1–C19 109.7(1), N1–P1–C19 110.8(1), C1–N1–P1 136.1(2), N9–P2–C19 105.4(1), N10–P2–C19 108.8(1), N6–P2–C19 111.8(1), C10–N6–P2 132.6(2), N4–P1–C19–P2 169.1(1), N9–P2–C19–P1 165.2(1). $2 \cdot 2\text{HBF}_4$ P1–C1/P5–C22 1.820(4)/1.822(4), P1–N1/P5–N19 1.626(4)/1.630(4), P1–N2/P5–N20 1.642(4)/1.650(4), P1–N3/P5–N21 1.573(4)/1.571(4), P2–N3/P6–N21 1.582(4)/1.589(4), P2–N4/P6–N22 1.648(4)/1.639(4), P2–N5/P6–N23 1.639(4)/1.639(4), P2–N6/P6–N24 1.650(4)/1.655(4), P3–C1/P7–C22 1.819(5)/1.817(5), P3–N7/P7–N13 1.636(4)/1.645(4), P3–N8/P7–N14 1.647(4)/1.635(4), P3–N9/P7–N15 1.575(4)/1.567(4), P4–N9/P8–N15 1.577(4)/

3.048(2) Å, the slightly longer distance involving the bridging chlorine atom. Similar weak hydrogen bonds were described for $(\text{Ph})_6\text{-CDP} \cdot 2\text{H}^+$ with $[\text{InCl}_4]^-$ (3.60 Å and 4.03 Å),⁴⁸ $[\text{BeCl}_4]^{2-}$ (3.55 Å and 3.58 Å),⁴⁹ I^- (3.80 Å and 3.81 Å)⁵⁰ and Cl^- (3.38 Å)⁴⁹ anions. The difference between the latter and $4 \cdot 2\text{HCl}$ probably arise from a less polarized C–H bond due to the stronger electron pair donor **4**. Single crystals of the isolated $4 \cdot 2\text{HBF}_4$ were additionally obtained from chloroform and exhibits no significant differences in the structural properties (displayed in the ESI[†]). Fig. 2 shows the molecular structures of $1 \cdot 2\text{HBF}_4$ and $2 \cdot 2\text{HBF}_4$ as well. All three bisprotonated CDPs exhibit a strong influence of charge delocalization as the reason for their extraordinary basicity: upon protonation the P–C bonds elongate from 1.606 Å (**4**) and 1.618 Å (**1**) to 1.799 Å in $4 \cdot 2\text{HCl}$ and 1.821 Å in $1 \cdot 2\text{HBF}_4$ and $2 \cdot 2\text{HBF}_4$, whilst the P–N bonds become shorter to average 1.62 Å for pyrrolidine and 1.64 Å for dimethylamine substituents. This complies with distances found in protonated phosphazenes⁵¹ and phosphorus ylids⁵² and proves the electron donating effect of the amino substituents. The P–N bonds to the tmg groups in $1 \cdot 2\text{HBF}_4$ exhibits with 1.58 Å (1.66 Å in **1**) clearly double-bond character. The P–N=C angles are expanded from 127° and 128° to 132° and 136°. A diminishing difference of formal N–C single and double bonds in the tmg group indicates the conjugation within the CN_3 moiety. The formal P–N single and double bonds of the phosphazene substituents in $2 \cdot 2\text{HBF}_4$ equalize at 1.57–1.59 Å with P–N–P angles between 134° and 142°. Similar influence of negative hyperconjugation for charge delocalization was found in superbasic PAPS¹⁶ and protonated diphosphazenes.⁵³ The P–C–P angles in the bisprotonated forms (**4**: 120°, **1**: 113°, **2**: 121°) are more acute than in the free CDPs (**4**: 156°, **1**: 147°). The difference to ideal tetrahedral geometry presumably arise from the bulkiness of the PR_3 moieties.

NMR spectroscopic features

All six presented compounds were characterized by ^1H , ^{13}C , and ^{31}P NMR spectroscopy. Selected chemical shifts and couplings are collected in Table 1. Proton shifts of bis- and monoprotated CDPs lie around 3 ppm for CH_2 and below 1 ppm for CH groups, both decreasing with increasing basicity of the parent CDP indicating less polarized C–H bonds. This shielding trend is not observed in the ^{13}C NMR shifts of the carbon nuclei: the most basic CDP **1** exhibits a triplet at 9.5 ppm compared to –1.6 ppm (**4**) and –6.8 ppm ($(\text{dma})_6\text{-CDP}$).²⁹ Surprisingly the ^{13}C chemical shift for **1** is even higher than for its monoprotated form ($1 \cdot \text{HTFSI}$: 9.3 ppm) contrasting the typical trend

1.579(4), P4–N10/P8–N16 1.644(4)/1.652(4), P4–N11/P8–N17 1.636(4)/1.648(4), P4–N12/P8–N18 1.655(4)/1.637(4), P3–C1–P1/P5–C22–P7 120.9(2)/121.7(2), N1–P1–C1/N19–P5–C22 110.8(2)/109.8(2), N2–P1–C1/N20–P5–C22 103.8(2)/104.0(2), N3–P1–C1/N21–P5–C22 107.9(2)/108.4(2), P1–N3–P2/P5–N21–P6 138.2(3)/135.7(3), N7–P3–C1/N14–P7–C22 111.2(2)/112.4(2), N8–P3–C1/N13–P7–C22 105.0(2)/103.2(2), N9–P3–C1/N15–P7–C22 107.9(2)/107.4(2), P3–N9–P4/P7–N15–P8 133.6(3)/141.6(3), N2–P1–C1–P3/N20–P5–C22–P7 164.8(3)/166.8(3), N8–P3–C1–P1/N13–P7–C22–P5 165.6(3)/164.4(3).



Table 1 NMR shifts δ /ppm and couplings J /Hz of the presented compounds

	δ_{H} ($^2J_{\text{PH}}/^1J_{\text{PH}}$)	δ_{C} ($^1J_{\text{PC}}/^3J_{\text{PC}}$)	δ_{P}
4·2HBF ₄ ^a	3.43 (19)	26.4 (110)	32.7
4·HTFSI ^b	0.93 (7)	10.3 (192)	40.1
4 ^c	—	−1.6 (280)	11.5
1·2HBF ₄ ^a	3.16 (17)	25.2 (112)	20.8
1·HTFSI ^b	0.55 (4)	9.3 (185)	37.1
1 ^c	—	9.5 (209)	18.2
2·2HBF ₄ ^a	2.87 (19)	25.6 (122/7)	23.2–22.7, 20.6–20.3
2·HBF ₄ ^a	0.25 (6/3)	12.6 (194/4)	34.3–33.6, 16.5–15.8
2 ^d	—	—	7.7–7.0, 6.2–5.6
7 ^c	0.42 (3/2)	13.0 (187/186/2)	109.9, 39.9, 37.0, 15.1

^a In CD₃CN. ^b In THF-*d*₃. ^c In C₆D₆. ^d In C₆D₆, assigned from the isolated mixture of the reaction between 2·2HBF₄ and KH in THF (Fig. S29 in the ESI).

observed for other CDPs.^{31,54} The $^1J_{\text{PC}}$ couplings drastically increase with step by step deprotonation indicating larger *s*-character of the ylidic P–C bonds. In the ^{31}P NMR spectra signals for the monoprotonated forms lie between the bisprotonated at higher and the free CDPs at lower values and correlate with the group electronegativity of the phosphines ((dma)₆-CDP: 27.72 ppm; (dma)₆-CDP·HCl: 54.16 ppm).²⁹ This is not exactly the case for the bisprotonated and free CDPs. The ^{31}P NMR signals of all three forms of **2** are multiplets corresponding to an AA'XX' spin system with $^2J_{\text{PP}}$ and $^4J_{\text{PP}}$ coupling (Fig. S22, S25, and S29 in the ESI[†]). **7** exhibits four individual signals in shape of two doublets of doublets for bridging phosphorus atoms and two doublets for terminal phosphorus atoms with the P^{III} atom being characteristically deshielded.^{15,16} ^1H and ^{13}C NMR signals are slightly shifted to higher frequencies in comparison with 2·HBF₄, indicating that the mixed valent P^{III}/P^V phosphanylphosphazene substituent is a poorer donor than corresponding P₂ bisphosphazene.

NMR titration experiments were conducted for **4** against (tmg)P₁-*t*Bu ($\text{p}K_{\text{BH}}^+$ in THF: 29.1)⁶ and (dma)P₄-*t*Bu ($\text{p}K_{\text{BH}}^+$ in THF: 33.9).²⁰ The $\text{p}K_{\text{BH}}^+$ value for **4** therefore has to be in between 30.1 and 32.9, since only free (tmg)P₁-*t*Bu and protonated **4** or protonated (dma)P₄-*t*Bu and free **4** were detected, respectively. Basicity of **1** was determined *via* titration against (pyrr)P₄-*t*Bu ($\text{p}K_{\text{BH}}^+$ in THF: 35.3)²⁰ as reference. Protonated and base forms of both species were quantified by ^{31}P NMR integration and a $\text{p}K_{\text{BH}}^+$ value of 35.8 in THF was determined for **1**. To our knowledge this is the first report of an experimental $\text{p}K_{\text{BH}}^+$ value for a carbodiphosphorane. It approves **1** to be an exceptional strong non-ionic carbon base, 0.5 orders of magnitude more basic than the strongest uncharged Schwesinger-type nitrogen superbases measured in THF²⁰ and 2.3 orders of magnitude more basic than the so far strongest uncharged carbon superbases H₂C=P(2,4,6-(MeO)₃-C₆H₂)₂Ph ($\text{p}K_{\text{BH}}^+$ in THF: 33.5).²⁰ Singlet carbenes such as NHCs and CAACs are weak carbon bases in comparison, according to $\text{p}K_{\text{BH}}^+$ values around 23 in THF and DMSO⁵⁵ or calculated PAs.^{34,56} The exceptional C-basicity of the title compounds is

only surpassed by our PAP phosphorus superbases (pyrr)P₃P (36.7) and (dma)P₄P (37.2).¹⁶

Quantumchemical calculations

First and second proton affinity (PA) and gas-phase basicity (GB) of carbodiphosphoranes **1**, **2**, **4** and phosphine **7** are calculated utilizing M06-2X/6-11+G(2df,p)//M06-2X/6-31+G(d) theoretical model. $\text{p}K_{\text{BH}}^+$ values in THF are obtained using the same functional and basis set whereas solvent is treated as dielectric continuum utilizing the SMD solvation model. $\text{p}K_{\text{BH}}^+$ values are calculated as relative values using an isodesmic reaction approach⁵⁷ where Schwesingers (dma)P₄-*t*Bu phosphazene with $\text{p}K_{\text{BH}}^+$ of 33.9 (ref. 20) has served as a reference base. Calculated values for protonation at central carbon atom, and in case of **7** protonation at the P^{III} atom as well, are presented in Table 2. It appears that the first proton affinity as well as $\text{p}K_{\text{BH}}^+$ values of **1** and **2** are higher than in Schwesingers (dma)P₄-*t*Bu phosphazene which has PA of 293.3 kcal mol^{−1} calculated at the same level of theory. Interestingly first GB of **1** is slightly lower than the GB of (dma)P₄-*t*Bu (GB = 288.2 kcal mol^{−1}) implying that the higher $\text{p}K_{\text{BH}}^+$ value of **1** relative to (dma)P₄-*t*Bu is a result of a more pronounced solvation effect in the carbodiphosphorane. This is unexpected considering that the N–H bond in a protonated phosphazene has a higher polarity than the C–H bond in protonated CDP as a result of lower electronegativity of carbon relative to nitrogen. The calculated $\text{p}K_{\text{BH}}^+$ (THF) 39.1 of **2** would be far higher than the $\text{p}K_{\text{BH}}^+$ (THF) 33.9 of (dma)P₄-*t*Bu, the strongest commercially available superbases. As described isolation of neutral base **2** is not achieved experimentally as other C–H bonds in the precursor 2·H⁺ seemed to have a higher kinetic and thermodynamic acidity. In order to understand the deprotonation path of 2·H⁺ under the action of NaNH₂, the reaction profile is calculated and presented in Fig. S36 in the ESI[†]. It appears, that the deprotonation of peripheral NMe₂ group in combination with the irreversible elimination of *N*-methylmethanimine is thermodynamically feasible (exergonic), however, kinetically hindered by a high barrier ($\Delta G^\ddagger = 32.8$ kcal mol^{−1}). This explains, that deprotonation induced degradation is competitive to deprotonation of central carbon atom at elevated temperatures, though the central carbon atom in 2·H⁺ is the thermodynamically most acidic site. It appears that decomposition product – phosphine **7** – has a gas-phase

Table 2 Calculated first and second proton affinity (PA) and gas phase basicity (GB) together with $\text{p}K_{\text{BH}}^+$ values in THF

	PA/kcal mol ^{−1}	GB/kcal mol ^{−1}	$\text{p}K_{\text{BH}}^+$ in THF ^a	
4	1 st	291.1	282.2	32.8 (30.1–32.9)
	2 nd	191.6	184.0	—
1	1 st	294.4	287.2	34.9 (35.8 ± 1)
	2 nd	202.0	194.1	—
2	1 st	305.3	299.7	39.1
	2 nd	212.1	202.2	—
7	At carbon	275.9	268.7	24.4
	At phosphorus	276.2	268.8	21.1

^a Experimental values in parentheses.



basicity ($30.9 \text{ kcal mol}^{-1}$) much lower than CDP 2. Interestingly, GB value for protonation at central carbon and P^{III} phosphorus of 7 is almost the same, whereas $\text{p}K_{\text{BH}}^+$ in THF for protonation at P^{III} is by 3.3 orders of magnitude lower than $\text{p}K_{\text{BH}}^+$ for protonation at carbon, which again indicates a more pronounced solvation effect in C-protonated CDP.

Conclusions

In this work we presented the most basic uncharged carbon bases known so far. A convenient synthesis for first- and novel second-order carbodiphosphorane superbases was presented. The CDPs (pyrr)₆-CDP 4 and *sym*-(tmg)₂(dma)₄-CDP 1 were synthesized as free base as well as in their mono- and bisprotonated forms. In our attempt to synthesize the even more outstanding base *sym*-(dmaP₁)₂(dma)₄-CDP 2 an unexpected, but highly selective deprotonation at peripheral PNCH₃ bonds induced an irreversible elimination path towards phosphine 7. This reaction is indicating a potential basicity limit for phosphazene containing superbases. Structural as well as spectroscopic features were investigated and the basicity was quantified by theoretical and experimental means. Remarkable $\text{p}K_{\text{BH}}^+$ values for 4 and 1 confirm them as benchmark breakers for non-ionic carbon bases on the THF basicity scale. Compared to the top Schwesinger bases, this basicity is even more outstanding, if their molecular weight below 500 g mol^{-1} is considered. We expect, that such simply synthesized carbodiphosphoranes with water stable protonated forms will enter the field of organic superbase catalysis.¹

Experimental section

General

All Reactions with air or moisture sensitive substances were carried out under inert atmosphere using standard Schlenk techniques. Air or moisture sensitive substances were stored in a nitrogen-flushed glovebox. Solvents were purified according to common literature procedures and stored under an inert atmosphere over molsieve (3 Å or 4 Å).⁵⁸ Pyrrolidine and tetramethylguanidine were distilled from CaH₂, triflimidic acid was purified by sublimation under argon. Bis(dichlorophosphino) methane³⁸ (6), bis[bis(dimethylamino)phosphino]methane³⁸ (3), tris(dimethylamino)phosphazene⁴ and (pyrr)₄-tBu⁴ were prepared according to literature-known procedures. (dma)₄-tBu was purchased as 1 M solution in *n*-hexane and dried in high vacuum. All other reagents were used as provided.

¹H, ¹³C, and ³¹P NMR spectra were recorded on a Bruker Avance III HD 250, Avance II 300, Avance III HD 300 or Avance III HD 500 spectrometer. Chemical shift δ is denoted relatively to SiMe₄ (¹H, ¹³C) or 85% H₃PO₄ (³¹P). ¹H and ¹³C NMR spectra were referenced to the solvent signals.⁵⁹ Multiplicity is abbreviated as follows: s (singlet), d (doublet), t (triplet), q (quartet), m (multiplet), br. (broad signal). High resolution mass spectrometry were performed on a Thermo Fisher Scientific LTQ-FT Ultra (ESI(+)) or a Jeol AccuTOF GCv (LIFDI(+) = liquid injection field desorption ionization), elemental analysis on an Elementar Vario Micro Cube. IR spectra were recorded in a glovebox on

a Bruker Alpha ATR-FT-IR. CCDC 1903830 (4·2HCl + HpyrrCl), 1903833 (1·2HBF₄), 1903838 (2·2HBF₄), 1903840 (1), 1903841 (4·2HBF₄), and 1903843 (4) contain the supplementary crystallographic data for this paper.†

General procedure for the precipitation of BF₄-salts

The crude product was dissolved in a minimum amount of water and a concentrated aqueous sodium tetrafluoroborate solution (2.0 eq.) was added. The resulting precipitate was filtered off, rinsed three times with small portions of cold water, washed with THF and dried in high vacuum.

(pyrr)₆-CDP·2HBF₄ (4·2HBF₄)

6 (3.60 g, 16.5 mmol, 1.00 eq.) was dissolved in THF (60 mL), cooled to $-78 \text{ }^\circ\text{C}$ and pyrrolidine (17.7 mL, 216 mmol, 13.1 eq.) was added dropwise. Afterwards the cooling bath was removed and the mixture stirred for additional 6 h. Carbon tetrachloride (3.12 mL, 32.3 mmol, 1.96 eq.) was added at $-78 \text{ }^\circ\text{C}$ and the mixture allowed to warm to room temperature overnight. The suspension was filtered under air and the filter cake extracted with THF ($3 \times 60 \text{ mL}$). The solvent was removed under reduced pressure and the residue dried in high vacuum. The crude product was converted to its tetrafluoroborate salt as described in the general procedure and recrystallized from methanol/ethanol. 4·2HBF₄ (6.38 g, 9.52 mmol, 58%) was obtained as colourless solid.

[C₂₅H₅₀B₂F₈N₆P₂] (670.27 g mol⁻¹) ¹H NMR (500.2 MHz, CD₃CN): δ (ppm) = 3.43 (t, ²J_{PH} = 19 Hz, 2H, CH₂), 3.25–3.22 (m, 24H, H1), 1.97–1.95 (m, 24H, H2, (overlapped with the solvent signal)). ¹³C{¹H} NMR (125.8 MHz, CD₃CN): δ (ppm) = 48.7 (s, C1), 26.9–26.8 (m, C2), 26.4 (t, ¹J_{PC} = 110 Hz, CH₂). ³¹P{¹H} NMR (121.5 MHz, CD₃CN): δ (ppm) = 32.7. ESI(+) MS (MeOH): *m/z* (%) = 495.6 (100) [M – H – 2BF₄]⁺, 583.2 (5) [M – BF₄]⁺. ESI(+) HRMS: *m/z* [M – H – 2BF₄]⁺ calcd 495.3488, found 495.3505; [M – BF₄]⁺ calcd 583.3600, found 583.3611. Elemental analysis: calcd C 44.80%, H 7.52%, N 12.54%; found C 44.49%, H 7.50%, N 12.46%. IR (neat): $\tilde{\nu}$ (cm⁻¹) = 2970 (w), 2879 (w), 1458 (w), 1251 (w), 1210 (m), 1134 (m), 1047 (vs.), 1021 (vs.), 918 (m), 870 (m), 824 (m), 779 (m), 699 (m), 581 (w), 549 (w), 517 (m) 484 (s). XRD: for single crystal X-ray structure determination suitable single crystals were obtained by slow evaporation of a concentrated solution in chloroform.

sym-(tmg)₂(dma)₄-CDP·2HBF₄ (1·2HBF₄)

3 (831 mg, 3.29 mmol, 1.00 eq.) and tetramethylguanidine (1.14 g, 9.88 mmol, 3.00 eq.) were dissolved in THF (60 mL). Carbon tetrachloride (640 μL , 6.62 mmol, 2.01 eq.) was added at $-78 \text{ }^\circ\text{C}$ and the mixture allowed to warm to room temperature overnight. The suspension was filtered under air and the filter cake extracted with THF ($3 \times 20 \text{ mL}$). The solvent was removed under reduced pressure and the residue dried in high vacuum. The crude product was converted to its tetrafluoroborate salt as described in the general procedure and recrystallized from ethanol. 1·2HBF₄ (1.08 g, 1.66 mmol, 50%) was isolated as colourless solid.



[C₁₀H₅₀B₂F₈N₁₀P₂] (654.24 g mol⁻¹) ¹H NMR (500.2 MHz, CD₃CN): δ (ppm) = 3.16 (t, ²J_{PH} = 17 Hz, 2H, CH₂), 2.91 (s, 24H, H1), 25.3 (d, ³J_{PH} = 10 Hz, 24H, H2). ¹³C{¹H} NMR (125.8 MHz, CD₃CN): δ (ppm) = 161.6 (dd, 2 × ²J_{PC} = 2 Hz, CN₃), 40.9 (s, C1), 37.1 (dd, 2 × ²J_{PC} = 2 Hz, C2), 25.2 (t, ¹J_{PC} = 112 Hz, CH₂). ³¹P{¹H} NMR (202.5 MHz, CD₃CN): δ (ppm) = 20.8 (s, ¹J_{PC} = 113 Hz (satellites)). ESI(+)⁺ MS (MeOH): *m/z* (%) = 479.5 (100) [M - H - 2BF₄]⁺. ESI(+)⁺ HRMS: *m/z* [M - H - 2BF₄]⁺ calcd. 479.3622, found 479.3625. Elemental analysis: calcd C 34.88%, H 7.70%, N 21.41%; found C 34.98%, H 7.84%, N 21.39%. IR (neat): $\tilde{\nu}$ (cm⁻¹) = 2911 (br. w.), 1539 (s), 1486 (m), 1429 (m), 1401 (m), 1356 (m), 1289 (m), 1235 (w), 1186 (m), 1161 (m), 1046 (vs.), 1034 (vs.), 979 (vs.), 933 (vs.), 784 (s), 771 (s), 739 (m), 716 (m), 690 (w), 672 (w), 618 (w), 572 (m), 519 (m), 459 (m), 437 (m). XRD: for single crystal X-ray structure determination suitable single crystals were obtained from ethanol at -25 °C.

sym-(dmaP)₂(dma)₄-CDP · 2HBF₄ (2 · 2HBF₄)

3 (1.55 g, 6.14 mmol, 1.00 eq.) and tris(dimethylamino)phosphazene (3.28 g, 18.4 mmol, 3.00 eq.) were dissolved in THF (60 mL). Carbon tetrachloride (1.19 mL, 12.3 mmol, 2.00 eq.) was added at -78 °C and the mixture allowed to warm to room temperature overnight. The suspension was filtered under air and the filter cake extracted with THF (3 × 20 mL). The solvent was removed under reduced pressure and the residue dried in high vacuum. The crude product was converted to its tetrafluoroborate salt as described in the general procedure and recrystallized from ethanol/*n*-hexane. 2 · 2HBF₄ (2.58 g, 3.31 mmol, 54%) was isolated as colourless solid.

[C₂₁H₆₂B₂F₈N₁₂P₄] (780.31 g mol⁻¹) ¹H NMR (500.2 MHz, CD₃CN): δ (ppm) = 2.87 (t, ²J_{PH} = 19 Hz, 2H, CH₂), 2.68 (d, ³J_{PH} = 11 Hz, 24H, H2), 2.65 (d, ³J_{PH} = 10 Hz, 36H, H1). ¹³C{¹H} NMR (125.8 MHz, CD₃CN): δ (ppm) = 37.3 (m, C1, C2), 25.6 (tt, ¹J_{PC} = 122 Hz, ³J_{PC} = 7 Hz, CH₂). ³¹P{¹H} NMR (202.5 MHz, CD₃CN): δ (ppm) = 2.2–22.7 (m, P1), 20.6–20.3 (m, P2). ESI(+)⁺ MS (MeOH): *m/z* (%) = 303.5 (25) [M - 2BF₄]²⁺, 605.6 (60) [M - H - 2BF₄]⁺, 693.5 (100) [M - BF₄]⁺. ESI(+)⁺ HRMS: *m/z* [M - 2BF₄]²⁺ calcd 303.2080, found 303.2088; [M - H - 2BF₄]⁺ calcd 605.4087, found 605.4104; [M - BF₄]⁺ calcd 693.4195, found 693.4215. Elemental analysis: calcd C 32.32%, H 8.01%, N 21.54%; found C 31.94%, H 7.70%, N 21.18%. IR (neat): $\tilde{\nu}$ (cm⁻¹) = 2886 (w), 1539 (s), 1486 (m), 1429 (m), 1401 (m), 1356 (m), 1298 (m), 1234 (m), 1186 (w), 1161 (m), 1047 (vs.), 1035 (vs.), 979 (vs.), 933 (s), 784 (s), 771 (s), 739 (m), 715 (m), 690 (m), 672 (m), 572 (m), 519 (m), 459 (m), 439 (m). XRD: for single crystal X-ray structure determination suitable single crystals were obtained from ethanol/*n*-hexane at -25 °C.

(pyrr)₆-CDP (4)

A solution of potassium bis(trimethylsilyl)amide (558 mg, 2.80 mmol, 2.09 eq.) in THF (15 mL) was added to a suspension of 4 · 2HBF₄ (938 mg, 1.34 mmol, 1.00 eq.) in THF (40 mL) and stirred for 16 h at room temperature. All volatiles were removed *in vacuo*, the residue dissolved in *n*-hexane (20 mL) and filtered over Celite. The filter cake was extracted with *n*-hexane (2 × 15 mL) and the filtrate evaporated to dryness. 4 (481 mg,

973 μmol, 70%) was isolated as colourless solid. [C₂₅H₄₈N₆P₂] (494.65 g mol⁻¹) ¹H NMR (500.2 MHz, C₆D₆): δ (ppm) = 3.33–3.23 (m, 24H, H1), 1.75–1.64 (m, 24H, H2). ¹³C{¹H} NMR (125.8 MHz, C₆D₆): δ (ppm) = 47.4 (s, C1), 28.9 (s, C2), -1.6 (t, ¹J_{PC} = 280 Hz, PCP). ³¹P{¹H}-NMR (202.5 MHz, C₆D₆): δ (ppm) = 11.5. LIFDI(+)⁺ MS (*n*-hexane): *m/z* (%) = 495.4 (100) [M + H]⁺. LIFDI(+)⁺ HRMS: *m/z* [M + H]⁺ calcd 495.34939, found 495.35037. Elemental analysis: calcd C 60.70%, H 9.78%, N 16.99%; found C 60.39%, H 9.62%, N 17.42%. IR (neat): $\tilde{\nu}$ (cm⁻¹) = 2952 (m), 2836 (m), 1492 (w), 1435 (s), 1338 (m), 1319 (m), 1289 (w), 1191 (m), 1134 (m), 1046 (vs.), 1000 (vs.), 980 (vs.), 909 (s), 870 (m), 742 (m), 546 (vs.), 497 (vs.). XRD: for single crystal X-ray structure determination suitable single crystals were obtained from *n*-hexane at -25 °C.

sym-(tmg)₂(dma)₄-CDP (1)

A mixture of 1 · 2HBF₄ (190 mg, 290 μmol, 1.00 eq.) and sodium amide (113 mg, 2.90 mmol, 10.0 eq.) was stirred in THF (15 mL) for 16 h at room temperature. The suspension was filtered over Celite and the filter cake extracted with THF (3 × 5 mL). All volatiles were removed *in vacuo*, *n*-hexane (10 mL) added to the residue, filtered again over Celite and extracted with *n*-hexane (3 × 4 mL). Evaporation of the solvent and drying in high vacuum yielded 1 (86 mg, 0.17 mmol, 60%) as colourless solid. [C₁₉H₄₈N₁₀P₂] (478.61 g mol⁻¹) ¹H NMR (500.2 MHz, C₆D₆): δ (ppm) = 2.88 (dd, 2 × ^{3,5}J_{PH} = 5 Hz, 24H, H2), 2.73 (s, 24H, H1). ¹³C{¹H} NMR (125.8 MHz, C₆D₆): δ (ppm) = 156.0 (s, CN₃), 40.1 (s, C1), 38.3 (s, C2), 9.5 (t, ¹J_{PC} = 209 Hz, PCP). ³¹P{¹H} NMR (121.5 MHz, C₆D₆): δ (ppm) = 18.2. LIFDI(+)⁺ MS (*n*-hexane): *m/z* (%) = 479.4 (100) [M + H]⁺. LIFDI(+)⁺ HRMS: *m/z* [M + H]⁺ calcd 479.36169, found 479.36229. Elemental analysis: calcd C 47.68%, H 10.11%, N 29.27%; found C 47.54%, H 9.96%, N 29.47%. IR (neat): $\tilde{\nu}$ (cm⁻¹) = 3006 (w), 2847 (m), 2810 (m), 2778 (m), 1566 (vs.), 1496 (s), 1472 (m), 1453 (m), 1440 (m), 1421 (m), 1358 (vs.), 1281 (m), 1251 (m), 1235 (m), 1211 (m), 1173 (m), 1128 (s), 1052 (m), 971 (s), 949 (vs.), 917 (m), 860 (vs.), 796 (m), 748 (m), 685 (s), 652 (s), 629 (vs.), 568 (m), 527 (s), 452 (s). XRD: for single crystal X-ray structure determination suitable single crystals were obtained from *n*-hexane at -25 °C.

Attempted synthesis of *sym*-(dmaP)₂(dma)₄-CDP (2)

A mixture of 2 · 2HBF₄ (136 mg, 174 μmol, 1.0 eq.) and freshly ground sodium amide (75 mg, 1.9 mmol, 11 eq.) was suspended in THF (15 mL) and stirred for 72 h at 60 °C. The solid was removed by filtration over Celite and the filtrate evaporated to dryness. The residue was dissolved in *n*-pentane (20 mL), cleared *via* syringe filtration, the solvent removed and the residue dried in high vacuum to give 7 as pale yellow high viscous oil.

[C₁₉H₅₅N₁₀P₄] (561.62 g mol⁻¹) ¹H NMR (300.3 MHz, C₆D₆): δ (ppm) = 2.99 (d, ³J_{PH} = 9 Hz, 12H, H4), 2.88 (d, ³J_{PH} = 10 Hz, 12H, H3), 2.83 (d, ³J_{PH} = 11 Hz, 12H, H2), 2.32 (d, ³J_{PH} = 10 Hz, 18H, H1), 0.42 (dddd, 2 × ²J_{PH} = 3 Hz, 2 × ⁴J_{PH} = 2 Hz, 1H, CH). ¹³C{¹H} NMR (75.5 MHz, C₆D₆): δ (ppm) = 38.5 (dd, ²J_{PC} = 4 Hz, ⁴J_{PC} = 3 Hz, C3), 38.4 (d, ²J_{PC} = 16 Hz, C4), 38.1 (dd, ²J_{PC} = 4 Hz, ⁴J_{PC} = 1 Hz, C2) 37.1 (d, ²J_{PC} = 4 Hz, C1), 13.0 (ddd, ¹J_{PC} =



187 Hz, $^1J_{PC} = 186$ Hz, $^3J_{PC} = 2$ Hz, CH). $^{31}\text{P}\{^1\text{H}\}$ NMR (121.5 MHz, C_6D_6): δ (ppm) = 109.9 (d, $^2J_{PP} = 100$ Hz, P4), 39.9 (dd, $^2J_{PP} = 50$ Hz, $^2J_{PP} = 41$ Hz, P2), 37.0 (dd, $^2J_{PP} = 100$ Hz, $^2J_{PP} = 41$ Hz, P3), 15.1 (d, $^2J_{PP} = 50$ Hz, P1). LIFDI(+) MS (*n*-hexane): m/z (%) = 561.4 (100) $[\text{M}]^+$. LIFDI(+) HRMS: m/z $[\text{M}]^+$ calcd 561.35924, found 561.35562.

(pyrr)₆-CDP·HTFSI (4·HTFSI)

4 (8.954 mg, 18.10 μmol , 1.04 eq.) and triflimidic acid (4.911 mg, 17.46 μmol , 1.00 eq.) were mixed in THF-*d*₈ (0.5 mL) and used for analytics.

$[\text{C}_{27}\text{H}_{49}\text{F}_6\text{N}_7\text{O}_4\text{P}_2\text{S}_2]$ (775.79 g mol⁻¹) ^1H NMR (500.2 MHz, THF-*d*₈): δ (ppm) = 3.20–3.17 (m, 24H, H1), 1.88–1.85 (m, 24H, H2), 0.93 (t, $^2J_{PH} = 7$ Hz, 1H, CH). $^{13}\text{C}\{^1\text{H}\}$ NMR (125.8 MHz, THF-*d*₈): δ (ppm) = 121.1 (q, $^1J_{FC} = 323$ Hz, CF₃), 47.8 (s, C1), 26.9 (dd, $2 \times J_{PC} = 4$ Hz, C2), 10.3 (t, $^1J_{PC} = 192$ Hz, CH). $^{31}\text{P}\{^1\text{H}\}$ NMR (121.5 MHz, THF-*d*₈): δ (ppm) = 40.1. LIFDI(+) MS (THF): m/z (%) = 495.4 (100) $[\text{M} - \text{TFSI}]^+$. LIFDI(+) HRMS: m/z $[\text{M} - \text{TFSI}]^+$ calcd 495.34939, found 495.35146.

sym-(tmg)₂(dma)₄-CDP·HTFSI (1·HTFSI)

1 (9.273 mg, 19.38 μmol , 1.00 eq.) and triflimidic acid (5.517 mg, 19.62 μmol , 1.01 eq.) were mixed in THF-*d*₈ (0.5 mL) and used for analytics.

$[\text{C}_{21}\text{H}_{49}\text{F}_6\text{N}_{11}\text{O}_4\text{P}_2\text{S}_2]$ (759.75 g mol⁻¹) ^1H NMR (300.3 MHz, THF-*d*₈): δ (ppm) = 2.90 (s, 24H, H1), 2.67–2.64 (m, 24H, H2), 0.55 (t, $^2J_{PH} = 4$ Hz, 1H, CH). $^{13}\text{C}\{^1\text{H}\}$ NMR (75.5 MHz, THF-*d*₈): δ (ppm) = 161.1 (s, CN₃), 121.1 (q, $^1J_{FC} = 322$ Hz, CF₃), 40.3 (s, C1), 37.7 (dd, $2 \times ^2J_{PC} = 2$ Hz, C2), 9.3 (t, $^1J_{PC} = 185$ Hz, CH). $^{31}\text{P}\{^1\text{H}\}$ NMR (121.5 MHz, C_6D_6): δ (ppm) = 37.1. LIFDI(+) MS (THF): m/z (%) = 479.4 (100) $[\text{M} - \text{TFSI}]^+$. LIFDI(+) HRMS: m/z $[\text{M} - \text{TFSI}]^+$ calcd 479.36169, found 479.36232.

sym-(dmaP)₂(dma)₄-CDP·HBF₄ (2·HBF₄)

A mixture of 2·2HBF₄ (600 mg, 769 μmol , 1.00 eq.) and finely ground sodium amide (321 mg, 8.23 mmol, 10.7 eq.) was suspended in THF (20 mL), cooled to -78 °C and ammonia (ca. 40 mL) was condensed in. The mixture was allowed to warm to room temperature overnight, the solid removed by centrifugation and the supernatant evaporated to dryness. The residue was dissolved in dichloromethane (40 mL) and filtered over Celite. All volatiles were removed *in vacuo*, the residue washed with diethyl ether (2×40 mL) and dried in high vacuum. 2·HBF₄ (365 mg, 527 μmol , 69%) was isolated as colorless solid. $[\text{C}_{21}\text{H}_{61}\text{BF}_4\text{N}_{10}\text{P}_4]$ (692.50 g mol⁻¹) ^1H NMR (500.2 MHz, CD₃CN): δ (ppm) = 2.64 (d, $^3J_{PH} = 10$ Hz, 36H, H1), 2.60–2.57 (m, 24H, H2), 0.25 (tt, $^2J_{PH} = 6$ Hz, $^4J_{PH} = 3$ Hz, 1H, CH). $^{13}\text{C}\{^1\text{H}\}$ NMR (125.8 MHz, CD₃CN): δ (ppm) = 37.9 (d, $^2J_{PC} = 2$ Hz, C2), 37.4 (d, $^2J_{PC} = 5$ Hz, C1), 12.6 (tt, $^1J_{PC} = 194$ Hz, $^3J_{PC} = 4$ Hz, CH). $^{31}\text{P}\{^1\text{H}\}$ NMR (121.5 MHz, CD₃CN): δ (ppm) = 34.3–33.6 (m, P2), 16.5–15.8 (m, P1). LIFDI(+) MS (THF): m/z (%) = 605.4 (100) $[\text{M} - \text{BF}_4]^+$. LIFDI(+) HRMS: m/z $[\text{M} - \text{BF}_4]^+$ calcd 605.40926, found 605.41147. Elemental analysis: calcd C 36.42%, H 8.88%, N 24.27%; found C 36.25%, H 8.59%, N 24.21%. IR (neat): $\tilde{\nu}$ (cm⁻¹) = 3000 (w), 2883 (m), 2846 (m), 2804 (m), 1458 (m), 1288 (s), 1243 (m), 1183 (m), 1167 (m), 1092 (m), 1048 (s), 976 (vs.), 955

(vs.), 845 (m), 823 (m), 770 (m), 740 (s), 715 (s), 660 (s), 598 (m), 551 (w), 527 (m), 498 (s), 454 (m), 420 (w).

Conflicts of interest

The authors have no conflicts to declare.

Acknowledgements

This work was supported by Philipps-Universität Marburg.

Notes and references

- 1 T. Ishikawa, *Superbases for organic synthesis. Guanidines, amidines and phosphazenes and related organocatalysts*, Wiley, Chichester, West Sussex, U.K, 2009.
- 2 E. D. Raczyńska, J.-F. Gal and P.-C. Maria, *Chem. Rev.*, 2016, **116**, 13454.
- 3 R. Schwesinger and H. Schlemper, *Angew. Chem., Int. Ed. Engl.*, 1987, **26**, 1167; *Angew. Chem.*, 1987, **99**, 1212.
- 4 R. Schwesinger, H. Schlemper, C. Hasenfratz, J. Willaredt, T. Dambacher, T. Breuer, C. Ottaway, M. Fletschinger, J. Boele, H. Fritz, D. Putzas, H. W. Rotter, F. G. Bordwell, A. V. Satish, G.-Z. Ji, E.-M. Peters, K. Peters, H. G. von Schnering and L. Walz, *Liebigs Ann./Recl.*, 1996, **1996**, 1055.
- 5 B. Kovačević and Z. B. Maksić, *Org. Lett.*, 2001, **3**, 1523.
- 6 A. A. Kolomeitsev, I. A. Koppel, T. Rodima, J. Barten, E. Lork, G.-V. Rösenthaller, I. Kaljurand, A. Kütt, I. Koppel, V. Mäemets and I. Leito, *J. Am. Chem. Soc.*, 2005, **127**, 17656.
- 7 (a) R. A. Kunetskiy, S. M. Polyakova, J. Vavřík, I. Cisařová, J. Saame, E. R. Nerut, I. Koppel, I. A. Koppel, A. Kütt, I. Leito and I. M. Lyapkalo, *Chem.-Eur. J.*, 2012, **18**, 3621; (b) K. Vazdar, R. Kunetskiy, J. Saame, K. Kaupmees, I. Leito and U. Jahn, *Angew. Chem., Int. Ed.*, 2014, **53**, 1435; *Angew. Chem.*, 2014, **126**, 1459.
- 8 E. D. Nacsá and T. H. Lambert, *J. Am. Chem. Soc.*, 2015, **137**, 10246.
- 9 B. Kovačević, Z. B. Maksić and R. Vianello, *J. Chem. Soc., Perkin Trans. 2*, 2001, 886.
- 10 J. F. Kögel, B. Oelkers, B. Kovačević and J. Sundermeyer, *J. Am. Chem. Soc.*, 2013, **135**, 17768.
- 11 R. W. Alder, P. S. Bowman, W. R. S. Steele and D. R. Winterman, *Chem. Commun.*, 1968, 723.
- 12 (a) V. Raab, J. Kipke, R. M. Gschwind and J. Sundermeyer, *Chem.-Eur. J.*, 2002, **8**, 1682; (b) V. Raab, E. Gauchenova, A. Merkoulou, K. Harms, J. Sundermeyer, B. Kovačević and Z. B. Maksić, *J. Am. Chem. Soc.*, 2005, **127**, 15738; (c) L. Belding and T. Dudding, *Chem.-Eur. J.*, 2014, **20**, 1032; (d) L. Belding, P. Stoyanov and T. Dudding, *J. Org. Chem.*, 2016, **81**, 6.
- 13 (a) H. A. Staab, T. Saupe and C. Krieger, *Angew. Chem., Int. Ed. Engl.*, 1983, **22**, 731; *Angew. Chem.*, 1983, **95**, 748; (b) T. Saupe, C. Krieger and H. A. Staab, *Angew. Chem., Int. Ed. Engl.*, 1986, **25**, 451; *Angew. Chem.*, 1986, **98**, 460; (c) M. A. Zirnstein and H. A. Staab, *Angew. Chem., Int. Ed. Engl.*, 1987, **26**, 460; *Angew. Chem.*, 1987, **99**, 460; (d) R. Schwesinger, M. Mißfeldt, K. Peters and H. G. von



- Schnering, *Angew. Chem., Int. Ed. Engl.*, 1987, **26**, 1165; *Angew. Chem.*, 1987, **99**, 1210; (e) A. F. Pozharskii, O. V. Ryabtsova, V. A. Ozeryanskii, A. V. Degtyarev, O. N. Kazheva, G. G. Alexandrov and O. A. Dyachenko, *J. Org. Chem.*, 2003, **68**, 10109; (f) A. V. Degtyarev, O. V. Ryabtsova, A. F. Pozharskii, V. A. Ozeryanskii, Z. A. Starikova, L. Sobczyk and A. Filarowski, *Tetrahedron*, 2008, **64**, 6209; (g) J. F. Kögel, N.-J. Kneusels and J. Sundermeyer, *Chem. Commun.*, 2014, **50**, 4319; (h) A. F. Pozharskii, V. A. Ozeryanskii, V. Y. Mikshiev, A. S. Antonov, A. V. Chernyshev, A. V. Metelitsa, G. S. Borodkin, N. S. Fedik and O. V. Dyablo, *J. Org. Chem.*, 2016, **81**, 5574; (i) R. J. Schwamm, R. Vianello, A. Marsavelski, M. A. Garcia, R. M. Claramunt, I. Alkorta, J. Saame, I. Leito, C. M. Fitchett, A. J. Edwards and M. P. Coles, *J. Org. Chem.*, 2016, **81**, 7612; (j) J. F. Kögel, B. Kovačević, S. Ullrich, X. Xie and J. Sundermeyer, *Chem.-Eur. J.*, 2017, **23**, 2591.
- 14 (a) C. Lensink, S. K. Xi, L. M. Daniels and J. G. Verkade, *J. Am. Chem. Soc.*, 1989, **111**, 3478; (b) J. Münchenberg, H. Thönnessen, P. G. Jones and R. Schmutzler, *Phosphorus, Sulfur Silicon Relat. Elem.*, 1997, **123**, 57; (c) P. Mehlmann, C. Mück-Lichtenfeld, T. T. Y. Tan and F. Dielmann, *Chem.-Eur. J.*, 2017, **23**, 5929.
- 15 A. P. Marchenko, G. N. Koidan, A. M. Pinchuk and A. V. Kirsanov, *J. Gen. Chem. USSR*, 1984, **54**, 1581.
- 16 S. Ullrich, B. Kovačević, X. Xie and J. Sundermeyer, *Angew. Chem., Int. Ed.*, 2019, **58**, 10335; *Angew. Chem.*, 2019, **131**, 10443.
- 17 R. Schwesinger, C. Hasenfratz, H. Schlemper, L. Walz, E.-M. Peters, K. Peters and H. G. von Schnering, *Angew. Chem., Int. Ed. Engl.*, 1993, **32**, 1361; *Angew. Chem.*, 1993, **105**, 1420.
- 18 I. Leito, I. A. Koppel, I. Koppel, K. Kaupmees, S. Tshepelevitsh and J. Saame, *Angew. Chem., Int. Ed.*, 2015, **54**, 9262; *Angew. Chem.*, 2015, **127**, 9394.
- 19 (a) S. Goumri-Magnet, O. Guerret, H. Gornitzka, J. B. Cazaux, D. Bigg, F. Palacios and G. Bertrand, *J. Org. Chem.*, 1999, **64**, 3741; (b) I. A. Koppel, R. Schwesinger, T. Breuer, P. Burk, K. Herodes, I. Koppel, I. Leito and M. Mishima, *J. Phys. Chem. A*, 2001, **105**, 9575.
- 20 J. Saame, T. Rodima, S. Tshepelevitsh, A. Kütt, I. Kaljurand, T. Haljasorg, I. A. Koppel and I. Leito, *J. Org. Chem.*, 2016, **81**, 7349.
- 21 J. F. Kögel, D. Margetić, X. Xie, L. H. Finger and J. Sundermeyer, *Angew. Chem., Int. Ed.*, 2017, **56**, 3090; *Angew. Chem.*, 2017, **129**, 3136.
- 22 (a) A. J. Arduengo, R. L. Harlow and M. Kline, *J. Am. Chem. Soc.*, 1991, **113**, 361; (b) W. A. Herrmann, *Angew. Chem., Int. Ed.*, 2002, **41**, 1290; *Angew. Chem.*, 2002, **114**, 1342; (c) M. N. Hopkinson, C. Richter, M. Schedler and F. Glorius, *Nature*, 2014, **510**, 485; (d) V. Nesterov, D. Reiter, P. Bag, P. Frisch, R. Holzner, A. Porzelt and S. Inoue, *Chem. Rev.*, 2018, **118**, 9678; (e) A. Doddi, M. Peters and M. Tamm, *Chem. Rev.*, 2019, **119**, 6994.
- 23 (a) V. Lavallo, Y. Canac, C. Präsang, B. Donnadieu and G. Bertrand, *Angew. Chem., Int. Ed.*, 2005, **44**, 5705; *Angew. Chem.*, 2005, **117**, 5851; (b) M. Soleilhavoup and G. Bertrand, *Acc. Chem. Res.*, 2015, **48**, 256; (c) Z. R. Turner, *Chem.-Eur. J.*, 2016, **22**, 11461; (d) M. Melaimi, R. Jazzar, M. Soleilhavoup and G. Bertrand, *Angew. Chem., Int. Ed.*, 2017, **56**, 10046; *Angew. Chem.*, 2017, **129**, 10180; (e) C. M. Weinstein, G. P. Junor, D. R. Tolentino, R. Jazzar, M. Melaimi and G. Bertrand, *J. Am. Chem. Soc.*, 2018, **140**, 9255.
- 24 (a) C. A. Dyker, V. Lavallo, B. Donnadieu and G. Bertrand, *Angew. Chem., Int. Ed.*, 2008, **47**, 3206; *Angew. Chem.*, 2008, **120**, 3250; (b) A. Fürstner, M. Alcarazo, R. Goddard and C. W. Lehmann, *Angew. Chem., Int. Ed.*, 2008, **47**, 3210; *Angew. Chem.*, 2008, **120**, 3254; (c) O. Kaufhold and F. E. Hahn, *Angew. Chem., Int. Ed.*, 2008, **47**, 4057; *Angew. Chem.*, 2008, **120**, 4122; (d) C. Pranckevicius, L. Liu, G. Bertrand and D. W. Stephan, *Angew. Chem., Int. Ed.*, 2016, **55**, 5536; *Angew. Chem.*, 2016, **128**, 5626; (e) W.-C. Chen, W.-C. Shih, T. Jurca, L. Zhao, D. M. Andrada, C.-J. Peng, C.-C. Chang, S.-K. Liu, Y.-P. Wang, Y.-S. Wen, G. P. A. Yap, C.-P. Hsu, G. Frenking and T.-G. Ong, *J. Am. Chem. Soc.*, 2017, **139**, 12830; (f) S.-C. Chan, P. Gupta, X. Engelmann, Z. Z. Ang, R. Ganguly, E. Bill, K. Ray, S. Ye and J. England, *Angew. Chem., Int. Ed.*, 2018, **57**, 15717; *Angew. Chem.*, 2018, **130**, 15943.
- 25 (a) W. C. Kaska, D. K. Mitchell and R. F. Reichelderfer, *J. Organomet. Chem.*, 1973, **47**, 391; (b) W. Petz and G. Frenking, in *Transition Metal Complexes of Neutral eta¹-Carbon Ligands*, ed. R. Chauvin and Y. Canac, Springer Berlin Heidelberg, Berlin, Heidelberg, 2010, vol. 30, pp. 49–92; (c) W. Petz, *Coord. Chem. Rev.*, 2015, **291**, 1; (d) J. E. Münzer, N.-J. H. Kneusels, B. Weinert, B. Neumüller and I. Kuzu, *Dalton Trans.*, 2019, **48**, 11076.
- 26 (a) R. Tonner and G. Frenking, *Chem.-Eur. J.*, 2008, **14**, 3260; (b) R. Tonner and G. Frenking, *Chem.-Eur. J.*, 2008, **14**, 3273; (c) M. Alcarazo, C. W. Lehmann, A. Anoop, W. Thiel and A. Fürstner, *Nat. Chem.*, 2009, **1**, 295.
- 27 F. Ramirez, N. B. Desai, B. Hansen and N. McKelvie, *J. Am. Chem. Soc.*, 1961, **83**, 3539.
- 28 O. Gasser and H. Schmidbaur, *J. Am. Chem. Soc.*, 1975, **97**, 6281.
- 29 R. Appel, U. Baumeister and F. Knoch, *Chem. Ber.*, 1983, **116**, 2275.
- 30 H. Schmidbaur, O. Gasser and M. S. Hussain, *Chem. Ber.*, 1977, **110**, 3501.
- 31 R. Appel and K. Waid, *Z. Naturforsch. B Chem. Sci.*, 1981, **36**, 131.
- 32 A. P. Marchenko, G. N. Koidan, V. A. Oleinik, I. S. Zal'tsman and A. M. Pinchuk, *J. Gen. Chem. USSR*, 1988, **58**, 1485.
- 33 R. Tonner, F. Öxler, B. Neumüller, W. Petz and G. Frenking, *Angew. Chem., Int. Ed.*, 2006, **45**, 8038; *Angew. Chem.*, 2006, **118**, 8206.
- 34 R. Tonner, G. Heydenrych and G. Frenking, *ChemPhysChem*, 2008, **9**, 1474.
- 35 (a) R. Appel, F. Knoll, W. Mische, W. Morbach, H.-D. Wihler and H. Veltmann, *Chem. Ber.*, 1976, **109**, 58; (b) R. Appel, F. Knoll, H. Schöler and H.-D. Wihler, *Angew. Chem., Int. Ed. Engl.*, 1976, **15**, 702; *Angew. Chem.*, 1976, **88**, 769; (c)



- R. Appel and H. Veltmann, *Tetrahedron Lett.*, 1977, **18**, 399;
- (d) A. P. Marchenko, G. N. Koidan, V. A. Oleinik, A. A. Kudryavtsev and A. M. Pinchuk, *J. Gen. Chem. USSR*, 1987, **57**, 1916; (e) V. A. Oleinik, A. P. Marchenko, G. N. Koidan and A. M. Pinchuk, *J. Gen. Chem. USSR*, 1988, **58**, 625; (f) G. N. Koidan, A. P. Marchenko, V. A. Oleinik and A. M. Pinchuk, *J. Gen. Chem. USSR*, 1988, **58**, 1304.
- 36 F. Ramirez and N. McKelvie, *J. Am. Chem. Soc.*, 1957, **79**, 5829.
- 37 (a) J. Münchenberg, O. Böge, A. K. Fischer, P. G. Jones and R. Schmutzler, *Phosphorus, Sulfur Silicon Relat. Elem.*, 1994, **86**, 103; (b) J. F. Kögel, N. C. Abacılar, F. Weber, B. Oelkers, K. Harms, B. Kovačević and J. Sundermeyer, *Chem.–Eur. J.*, 2014, **20**, 5994.
- 38 Z. S. Novikova, A. A. Prishchenko and I. F. Lutsenko, *J. Gen. Chem. USSR*, 1977, **47**, 707.
- 39 T. Rodima, I. Kaljurand, A. Pihl, V. Mäemets, I. Leito and I. A. Koppel, *J. Org. Chem.*, 2002, **67**, 1873.
- 40 I. Kaljurand, T. Rodima, A. Pihl, V. Mäemets, I. Leito, I. A. Koppel and M. Mishima, *J. Org. Chem.*, 2003, **68**, 9988.
- 41 A. P. Marchenko, G. N. Koidan and A. M. Pinchuk, *J. Gen. Chem. USSR*, 1984, **54**, 2405.
- 42 K. Issleib and M. Lischewski, *J. Prakt. Chem.*, 1970, **312**, 135.
- 43 (a) T. Scherpf, C. Schwarz, L. T. Scharf, J.-A. Zur, A. Helbig and V. H. Gessner, *Angew. Chem., Int. Ed.*, 2018, **57**, 12859; *Angew. Chem.*, 2018, **130**, 13041; (b) P. Weber, T. Scherpf, I. Rodstein, D. Lichte, L. T. Scharf, L. J. Gooßen and V. H. Gessner, *Angew. Chem., Int. Ed.*, 2019, **58**, 3203; *Angew. Chem.*, 2019, **131**, 3235.
- 44 P. J. Quinlivan and G. Parkin, *Inorg. Chem.*, 2017, **56**, 5493.
- 45 H. Schmidbaur, G. Hasslberger, U. Deschler, U. Schubert, C. Kappenstein and A. Frank, *Angew. Chem., Int. Ed. Engl.*, 1979, **18**, 408; *Angew. Chem.*, 1979, **91**, 437.
- 46 E. A. V. Ebsworth, T. E. Fraser, D. W. H. Rankin, O. Gasser and H. Schmidbaur, *Chem. Ber.*, 1977, **110**, 3508.
- 47 (a) A. T. Vincent and P. J. Wheatley, *J. Chem. Soc., Dalton Trans.*, 1972, 617; (b) G. E. Hardy, J. I. Zink, W. C. Kaska and J. C. Baldwin, *J. Am. Chem. Soc.*, 1978, **100**, 8001.
- 48 W. Petz, M. Fahlbusch, E. Gromm and B. Neumüller, *Z. Anorg. Allg. Chem.*, 2008, **634**, 682.
- 49 W. Petz, K. Dehnicke and B. Neumüller, *Z. Anorg. Allg. Chem.*, 2011, **637**, 1761.
- 50 R. Ganguly and V. Jevtovic, *Acta Crystallogr., Sect. E: Crystallogr. Commun.*, 2017, **73**, 1259.
- 51 (a) S. Ullrich, CCDC 1903847: Experimental Crystal Structure Determination; (b) S. Ullrich, CCDC 1903849: Experimental Crystal Structure Determination.
- 52 H. Vogt, D. Wulff-Molder, F. Ritschl, M. Mücke, U. Skrabei and M. Meisel, *Z. Anorg. Allg. Chem.*, 1999, **625**, 1025.
- 53 S. Ullrich, CCDC 1903848: Experimental Crystal Structure Determination.
- 54 M. Gruber, W. Bauer, H. Maid, K. Schöll and R. R. Tykwinski, *Inorg. Chim. Acta*, 2017, **468**, 152.
- 55 (a) R. W. Alder, P. R. Allen and S. J. Williams, *Chem. Commun.*, 1995, 1267; (b) Y.-J. Kim and A. Streitwieser, *J. Am. Chem. Soc.*, 2002, **124**, 5757; (c) N. Wang, J. Xu and J. K. Lee, *Org. Biomol. Chem.*, 2018, **16**, 8230.
- 56 (a) R. W. Alder, M. E. Blake and J. M. Oliva, *J. Phys. Chem. A*, 1999, **103**, 11200; (b) R. Tonner and G. Frenking, *Angew. Chem., Int. Ed.*, 2007, **46**, 8695; *Angew. Chem.*, 2007, **119**, 8850; (c) A. A. Tukov, A. T. Normand and M. S. Nechaev, *Dalton Trans.*, 2009, 7015.
- 57 S. Sastre, R. Casasnovas, F. Muñoz and J. Frau, *Phys. Chem. Chem. Phys.*, 2016, **18**, 11202.
- 58 W. L. F. Armarego and D. D. Perrin, *Purification of laboratory chemicals*, Butterworth Heinemann, Oxford, Boston, 4th edn, 1996.
- 59 G. R. Fulmer, A. J. M. Miller, N. H. Sherden, H. E. Gottlieb, A. Nudelman, B. M. Stoltz, J. E. Bercaw and K. I. Goldberg, *Organometallics*, 2010, **29**, 2176.

