# Chemical Science

# **EDGE ARTICLE**



Cite this: Chem. Sci., 2019, 10, 8799

o All publication charges for this article have been paid for by the Royal Society of Chemistry

## Synthesis of $\gamma$ -substituted carbonyl compounds from DMSO-mediated oxidation of envnamides: mechanistic insights and carbon- and heterofunctionalizations\*

Quynh H. Nguyen, 问 Nguyen H. Nguyen, 问 Hanbyul Kim 问 and Seunghoon Shin

Oxidative coupling of 1,3-envnamides using DMSO as a terminal oxidant has been developed. Carbon as well as unmodified heteroatom nucleophiles, including aliphatic alcohols, thiols, and hydrazides, could be efficiently alkylated at the  $\gamma$ -position in a highly regioselective fashion. The kinetic analysis suggested a nucleophile-dependent mechanism ranging from a concerted  $S_N 2^{\prime\prime}$  to a carbocationic mechanism. Thus, the remote site-selectivity was ascribed to the partial positive charge developing at the terminal carbocationic center.

Received 24th July 2019 Accepted 4th August 2019

DOI: 10.1039/c9sc03663f

rsc.li/chemical-science

### Introduction

Inverting natural polarity (Umpolung) allows for an outstanding opportunity to streamline redox efficient processes and enables novel disconnections with complementary selectivity to the traditional methods.1 For example, under conditions of normal polarity, *α*-hetero-functionalization of carbonyl compounds often requires strongly basic conditions and/or elaborate, preoxidized heteroatom-electrophiles.2 In contrast, Umpolung protocols allow for the use of unmodified heteroatom donors, which greatly facilitates the synthesis.

Towards umpoled enolate (a<sup>2</sup>) synthons, oxidation of alkynes through transition metal catalysis (Au, Rh, Ir, Ru, and Zn)<sup>3</sup> and the hypervalent iodine chemistry of carbonyl compounds<sup>4</sup> have led the way (Scheme 1a). More recently, we<sup>5</sup> and others<sup>6,7</sup> have reported a non-metal protocol involving Brønsted acid that can catalyze the oxidation of ynamides 1 and induce subsequent addition of carbon nucleophiles, furnishing a-heteroaryl imides 3 (Scheme 1b). In this process, it was unambiguously proved that the key intermediate, adduct 2, underwent a concerted  $S_N 2'$ alkylation by the observation of chirality transfer from optically active N-oxides.5a

Compared to enolonium equivalents, direct access to umpoled  $a^4$  synthons is rare. Although synthesis of  $\gamma$ substituted-a, \beta-carbonyl compounds has been demonstrated using PhNO, DEAD, or oxygen gas,8 generally applicable

This journal is © The Royal Society of Chemistry 2019

methods for the introduction of O-, S-, N- as well as carbon nucleophiles either at the  $\gamma$ - or  $\varepsilon$ -position are highly desirable as they are found in the scaffolds of numerous biologically active

CHEMISTRY

View Article Online

View Journal | View Issue



γ-(or 1 Vinylogous enolonium Scheme chemistry: E-) functionalizations

Department of Chemistry, Research Institute for Natural Sciences and Center for New Directions in Organic Synthesis (CNOS), Hanyang University, 222 Wangsimni-ro, Seongdong-gu, Seoul 04763, Korea. E-mail: sshin@hanayng.ac.kr

<sup>†</sup> Electronic supplementary information (ESI) available. CCDC 1895889. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c9sc03663f

agents.<sup>9</sup> To this end, we envisioned that oxidation of 1,3-enynamides 4 would generate *dienolonium*, such as 6 that corresponds to an umpoled version of silyl dienol ethers in Mukaiyama reactions. It would be interesting to explore the unknown regioselectivity in the alkylation,<sup>10,11</sup> and to elucidate the underlying mechanism. Herein, we have developed an oxidative alkylation of enynamides 4 and dienynamides 5 that occurs with remarkable selectivity for the remote substitution furnishing  $\gamma$ -substituted 8 or  $\varepsilon$ -substituted 9, respectively, and disclosed that the reaction proceeded through a mechanism ranging from concerted  $S_N 2''$  or through a dienolonium carbocation intermediate 7, depending on the types of nucleophiles.

Meanwhile, one inherent disadvantage of the alkyne oxidation protocols is the necessity to use pyridine-N-oxide which liberates pyridine, complicating the acid catalysis. In this context, dimethyl sulfoxide (DMSO) is advantageous in that the byproduct (Me<sub>2</sub>S) is easily removable (bp =  $37 \degree C$ ) and not so basic (p $K_a = -5.4$  in water).<sup>12-14</sup> Initially, we were concerned that the nucleophiles may attack the O-dienoxysulfenium intermediate 6 (Scheme 1c) at the S-atom (S<sub>N</sub>2) rather than at C4 (S<sub>N</sub>2") as in interrupted Pummerer reactions.15 Notwithstanding, a selective oxygenative coupling at the remote position was obtained and the usefulness of this oxidation is further demonstrated by the introduction of exceptionally diverse nucleophiles, ranging from carbon (indoles, furans, and silyl enol ethers) to heteroatom donors (aliphatic alcohols, thiols, and hydrazides) with good to excellent remote regioselectivity.

#### Results and discussion

# Survey of reaction conditions and Umpolung alkylation with carbon nucleophiles

For the examination of site selectivity in the enynamide oxidation, we initially chose 1,3-enynamide **4a** bearing sterically encumbering *o*-MeO-C<sub>6</sub>H<sub>4</sub> at C4 as a substrate that presents a regiose-lectivity challenge (Table 1). For initial screening, 3 equiv. of *N*-Meindole along with 4 equiv. of *N*-oxides (**Ox1-4**) was employed. 2-Cl-Pyridine-*N*-oxide, **Ox1** mediated the formation of **8a** with a remarkable efficiency in only 5 min at rt, although the regiose-lectivity remained mediocre ( $\alpha : \gamma = 1 : 4.5$ , entry 1).<sup>16</sup> Interestingly, diphenyl sulfoxide **Ox5** was also found to be a suitable oxidant for enynamides, while previous studies on the ynamide oxidation documented that [3,3]-sigmatropic rearrangement occurred dominantly.<sup>14b</sup> To our delight, DMSO (**Ox7**) displayed an excellent reactivity with  $\alpha : \gamma = 1 : 5.6$  (entry 7).<sup>17</sup> Change of solvents (entries 8–10) to 1,4-dioxane slightly improved the ratio to  $\alpha : \gamma = 1 : 8.7$ , albeit at the expense of the reactivity (entry 10).

At this point, we turned to other derivatives of enynamides **4b–f** having different aryl groups at C4 (entries 11–15). It was revealed that both the  $\alpha$  :  $\gamma$  selectivity and the reactivity in the  $\gamma$ -arylation with indoles critically depended on the electrondensity of the Ar group at C4. For example, the reaction of 4-MeO-derivatives **4b** delivered **8ba** with a good ratio ( $\alpha$  :  $\gamma$  = 1 : 10) in an excellent yield (entry 11). However, *p*-tolyl (**4c**), Ph (**4d**), 4-Cl-C<sub>6</sub>H<sub>4</sub> (**4e**) and 4-CF<sub>3</sub>-C<sub>6</sub>H<sub>4</sub> (**4f**) derivatives showed

Table 1 A survey of reaction conditions and substrates<sup>4</sup>

$\begin{array}{c} \begin{array}{c} \text{AM-Me-indole (3 equiv.)} \\ \text{N}\\ \text{R}^{2}\\ \text{A}\\ A\\ \text{A}\\ $					
Entry	Substrate	Oxidant	Conditions	$2^{b}$ (%)	$\alpha:\gamma^c$
1	4a	Ox1	DCE, 5 min	<b>8aa</b> , 90	1:4.5
2	4a	Ox2	DCE, 15 min	8aa, 71	1:6.7
3	4a	Ox3	DCE, 6 h	8aa, 35	1:5.2
4	4a	Ox4	DCE, 10 h, 60 $^\circ \mathrm{C}$	8aa, 41	1:4.5
5	4a	Ox5	DCE, 5 min	<b>8aa,</b> 70	1:5.8
6	4a	Ox6	DCE, 5 min	<b>8aa,</b> 77	1:5.1
7	4a	Ox7	DCE, 5 min	8aa, 95	1:5.6
8	4a	Ox7	CH <sub>3</sub> CN, 15 min	8aa, 62	1:6.8
9	4a	Ox7	MTBE, 0.5 h	<b>8aa,</b> 42 <sup>d</sup>	1:7.8
10	4a	Ox7	1,4-Dioxane, 0.5 h	<b>8aa</b> , 66	1:8.7
11	4b	Ox7	DCE, 5 min	<b>8ba</b> , 96	1:10
12	4 <b>c</b>	Ox7	DCE, 10 min	8ca, 83	1:2.7
13	4 <b>d</b>	Ox7	DCE, 3 h	<b>8da</b> , 62	1:1
14	4e	Ox7	DCE, 7 h	<b>8ea</b> , 63	1:1
15	4 <b>f</b>	Ox7	DCE, 3 h	e	—
16 <sup>7</sup>	4b	Ox7	DCM, 3 h	<b>8ba</b> , 97	1:10
17 <sup>f,g</sup>	4b	Ox7	DCM, 3 h	<b>8ba</b> , 93	1:10

<sup>*a*</sup> **4** (0.1 mmol), oxidant **3** (4 equiv.), *N*-Me-indole (3 equiv.) and HNTf<sub>2</sub> (10 mol%) in solvents (0.1 M). <sup>*b*</sup> Isolated yields of **8** after chromatographic separation. <sup>*c*</sup> Determined from the crude <sup>1</sup>H NMR spectra. <sup>*d*</sup> Hydration byproduct was observed. <sup>*e*</sup> Hydration byproduct (74% yield). <sup>*f*</sup> *N*-Me-indole (1.2 equiv.) and DMSO (1.2 equiv.) were used in the presence of 2.5 mol% of HNTf<sub>2</sub>. <sup>*g*</sup> With 3.77 mmol of **4b**.

decreasing reactivity and regioselectivity in the order, suggesting that a carbocation stabilizing group is needed at the C4 position. Remarkably, under the optimized conditions (entry 16), the oxidative coupling of **4b** occurred with only 1.2 equiv. of *N*-Me-indole and 1.2 equiv. of DMSO at 2.5 mol% of HNTf<sub>2</sub>, without adversely affecting the yield (entry 16). Under these conditions, a gram-scale reaction (3.8 mmol of **4b**) proceeded smoothly as well (entry 17).

With this initial reaction profile and the optimized conditions, we then inspected the reaction scope with various carbon nucleophiles (Table 2). The variation of N-alkyl and N-sulfonyl groups as well as electron-rich Ar groups (8ga-8ka) was all well accommodated with good reactivity and regioselectivity. Various indoles could also be incorporated without any event as in 8bb-8bf. Notably, heterocyclic motifs such as 2-furyl (8bg), N-acyl indole (8la), or 2-thiophene (8ma) could be easily incorporated into the products. Notably, the formation of a quaternary center as in 8na and **80a** was very efficient with a useful level of  $\gamma$ -selectivity: even phenyl derivative 80a was obtained with higher selectivity compared to 8da (entry 13, Table 1).18 Subsequently, we went on extending the conjugation by preparing dienynamide 5a (n = 2;  $\mathbb{R}^{1}$ = Ts,  $R^2$  = Me;  $R^3$  = H; Ar = 4-MeO-C<sub>6</sub>H<sub>4</sub>). To our delight, its reaction went smoothly to furnish the corresponding  $\alpha, \beta, \gamma, \delta$ unsaturated imide 9aa with high ɛ-selectivity.



<sup>*a*</sup> **4** (0.2 mmol), DMSO (1.2 equiv.), Nu-H (1.2 equiv.) and HNTf<sub>2</sub> (2.5 mol%) in CH<sub>2</sub>Cl<sub>2</sub> (0.1 M) unless otherwise noted; isolated yields of **8** (or **9**) after chromatographic purification. <sup>*b*</sup> Determined from the crude <sup>1</sup>H NMR spectra. <sup>*c*</sup> 5 mol% of HNTf<sub>2</sub> was used; reaction time: 9 h.

On the other hand, the reaction of silyl enol ethers was much slower (Table 3). For example, the reaction of **4b** with disubstituted silyl enol ether **10a** in the presence of 20 mol% of HNTf<sub>2</sub> at 60 °C was incomplete even after 36 h, from which **11ba** was obtained in 53% yield. Tri-substituted silyl enol ethers **10b**, **10c** were more reactive, giving higher yields of **11bb** and **11bc**, respectively, in the presence of 5 mol% of HNTf<sub>2</sub>. However, tetra-substituted silyl enol ether **10d**, reacted sluggishly. These indicated that a balance between the electron-density and the steric hindrance is necessary for an efficient coupling.



<sup>*a*</sup> **4** (0.2 mmol), DMSO (2 equiv.), silyl enol ethers (3 equiv.) and  $HNTf_2$  (5–20 mol%) in  $CH_2Cl_2$  (0.1 M); isolated yields of **11** after chromatographic purification. <sup>*b*</sup> The  $\alpha/\gamma$  and diastereomeric ratio were determined from the crude <sup>1</sup>H NMR spectra.

Remarkably, all the reactions of silyl enol ethers were completely regioselective ( $\alpha$  :  $\gamma = 1$  : >25).

#### Umpolung alkylation with heteroatom nucleophiles

Heteroatom-functionalization at the  $a^4$  or  $a^6$  site may perhaps be the most important application of this Umpolung chemistry.<sup>2,4b</sup> Intermolecular Umpolung reaction leading to  $\alpha$ -heterofunctionalization has been studied with various nucleophiles, including (thio)phenols,<sup>19a</sup> oxyacids,<sup>19b,c</sup> halogen,<sup>19d</sup> thiocyanate,<sup>19d</sup> azides,<sup>19d,e</sup> amines,<sup>4f,19f,g</sup> and TEMPO.<sup>19h-j</sup> Although unmodified phenols/thiols<sup>19a</sup> and amines<sup>19f,g</sup> could be used as nucleophiles, use of oxidizable aliphatic (primary and secondary) alcohols/thiols is surprisingly rare.<sup>19</sup>

When benzyl alcohol (3 equiv.) was tested as a nucleophile in the presence of DMSO (4 equiv.) and HNTf<sub>2</sub> (2.5 mol%), we were delighted to observe the formation of 67% of 12ba with a complete  $\gamma$ -selectivity in less than 5 min at room temperature (Table 4). Previous hypervalent iodine chemistry allowed for the use of MeOH, EtOH and iPrOH, but required these alcohols as solvents.<sup>20</sup> In contrast, the current protocol can operate with only 3 equivalents of nucleophiles and is thus suitable for the more elaborate alcohol donors. Gratifyingly, the transformation was general not only for simple primary alcohols (12ba-12bd), but also for furfuryl, allyl, propargyl, and cinnamyl alcohols, furnishing 12be, 12bf, 12bg, and 12bh, respectively. Products with secondary alcohols 12bi, 12bj were also obtained in reasonable yields, but tert-butanol gave only a trace amount of the adduct (not shown). Notably, all the reactions of 4b to form **12ba–12bj** were complete in less than 1 h and an exclusive  $\gamma$ regioselectivity was obtained, in contrast to the carbon nucleophiles in Table 2.

The faster rates with alcohol nucleophiles allowed us to use a less electron-rich aryl group at C4. For example, the reaction with 4c (Ar = *p*-tol) with benzyl alcohol proceeded uneventfully. However, the reaction with 4d (Ar = Ph) was very slow and





 $^{a}$  4 (0.2 mmol), DMSO (4 equiv.), ROH (3 equiv.) and HNTf<sub>2</sub> (2.5 mol%) in CH<sub>2</sub>Cl<sub>2</sub> (0.1 M) unless otherwise noted; isolated yields of **12** (or **13**) after chromatographic purification.  $^{b}$  Determined from the crude <sup>1</sup>H NMR spectra.  $^{c}$  5 mol% of HNTf<sub>2</sub>; reaction time: 6 h.

required 5 mol% of HNTf<sub>2</sub> and a longer time (6 h) for completion, giving reasonable yields of **12dc** and **12dj** with an exclusive  $\gamma$ -selectivity. Alkoxylation products with a tetra-substituted center, **12nb** and **12oh** were also obtained in reasonable yields, employing 5 mol% HNTf<sub>2</sub>. Finally, the reaction could be extended to dienynamides **5a**, **5b**, affording completely  $\varepsilon$ selective alkoxylation products **13ab** and **13ba** in good yields.

Encouraged by the success of alcohol nucleophiles, the reaction scope was extended to *S*- and *N*-nucleophiles as well (Table 5). Gratifyingly, with these nucleophiles, only a slight excess of donors (1.2 equiv.) and DMSO (1.2 equiv.) in the presence of HNTf<sub>2</sub> (2.5 mol%) were sufficient. Aliphatic thiols were suitable nucleophiles, furnishing **14ba**, **14bb** and **14bc** with exclusive  $\gamma$ -selectivity, and the dienimide **15ab** with exclusive  $\varepsilon$ -selectivity. However, in contrast to Zn(OTf)<sub>2</sub>-catalyzed oxidative coupling,<sup>19a,d</sup> aromatic thiol derivatives (PhSH) failed to deliver the corresponding product **14** (not shown).

Towards *N*-functionalization, various *N*-nucleophile donors were tested, including amines, amides, and hydroxylamines. Among them, sulfonyl hydrazides were found to be effective nucleophiles. With *p*-toluenesulfonyl- or benzenesulfonyl hydrazide, the corresponding products **16** or **17** were obtained in a completely regioselective manner. The connectivity of a hydrazide adduct **16ma** was unambiguously confirmed by single crystal X-ray crystallography.<sup>21</sup>

#### Kinetic study and mechanistic insights

To gain a mechanistic insight into the remote Umpolung functionalizations and to probe the nature of the intermediates involved, kinetic analysis was conducted. To the best of our knowledge, kinetic analysis on the Brønsted acid-catalyzed oxidative coupling of ynamides that can be used to check the validity of the DFT theoretical studies<sup>13e,22</sup> is unavailable. We set



<sup>*a*</sup> 4 (0.2 mmol), DMSO (1.2 equiv.), Het-H (1.2 equiv.) and  $HNTf_2$  (2.5 mol%) in  $CH_2Cl_2$  (0.1 M); isolated yields after chromatographic purification. <sup>*b*</sup> Determined from the crude <sup>1</sup>H NMR spectra.



**Scheme 2** Hammett correlation with *N*-Me-indole,  $Ph(CH_2)_2OH$  and 2-Me-furan as nucleophiles according to eqn (1).

out to perform a Hammett study with a set of enynamides differing in *para*-substituents, **4b** (4-OMe), **4c** (4-Me), **4d** (4-H), **4e** (4-Cl), **4p** (4-Ph), and **4q** (4-Br) to probe the electronic effect of aryl groups at C4 according to eqn (1) (Scheme 2). In the substitution with *N*-Me-indole, the  $\log(k_R/k_H)$  displayed a linear correlation with  $\sigma^+$  parameters, with a negative  $\rho$  value (-1.3) (Scheme 2a). Upon changing the nucleophile to an alcohol, Ph(CH<sub>2</sub>)<sub>2</sub>OH, the rate became significantly more sensitive to  $\sigma^+$ with  $\rho = -3.7$  (Scheme 2b), while the reaction with 2-Me-furan nucleophile showed a decreased sensitivity ( $\rho = -0.8$ ) (Scheme 2c). Interestingly, the observed regioselectivity in the reaction of enynamides **4b** (R = 4-MeO) with nucleophiles was in good accordance with the reaction constant  $\rho$ . For example, increasing  $\alpha/\gamma$  ratio was observed in the order of 2-Me-indole (1/4.4 for **8bg**),



Scheme 3  $\,$  A continuum mechanism between  ${\sf S}_N 2^{\prime\prime}$  and via an enolonium cation.

*N*-Me-indole (1/10 for **8ba**), and Ph(CH<sub>2</sub>)<sub>2</sub>OH (0/100 for **12bd**), where the corresponding  $\rho$  values were -0.8, -1.3, and -3.7, in the order. The observed reaction constant  $\rho$  reflects the build-up of a positive charge at C4 in the turnover-limiting step (TLS) and based on this, we formulated a tentative mechanism as in Scheme 3. At one extreme, Ph(CH<sub>2</sub>)<sub>2</sub>OH ( $\rho = -3.7$ ) would favor a route where S–O cleavage occurs first, forming an enolonium carbocation **20**, which is then trapped to yield **21**. At the other end of the continuum, 2-Me-furan ( $\rho = -0.8$ ) will follow a route in which a more synchronous cleavage of the S–O bond and the formation of a new C–C bond (TS<sub>19-22</sub>) occur and as a result, less charge builds up at C4.<sup>23</sup> Here, the  $\gamma$ -selectivity in the alkoxylation may be due to the formation of the more stable conjugated product and this tendency weakens with the S<sub>N</sub>2" pathway.



Next, the temperature dependence of the reaction rate ( $k_{obs}$ ) for the reaction of **4b** with *N*-Me-indole was examined in the range of 298–328 K (Fig. 1).<sup>15</sup> From the Eyring equation, the activation parameters were determined to be  $\Delta H^{\ddagger} = 83.5 \text{ kJ mol}^{-1}$  and  $\Delta S^{\ddagger} = -37.6 \text{ J K}^{-1} \text{ mol}^{-1}$ . The large negative entropy of activation is consistent with an associative transition state, such as  $\text{TS}_{19-22}$ .<sup>24</sup> Other associative steps, such as protonation of the enynamides (**4b** to **18**) or addition of DMSO (**18** to **19**) were excluded, because if these earlier steps were turnover-limiting, different sensitivity ( $\rho$ ) depending on the nucleophiles (Scheme 2) should not have been observed.



Fig. 1 Eyring plot at 298–328 K in the reaction of 4b with *N*-Meindole to form 8ba.

Finally, the reaction order was examined with respect to various reactants. The rate  $(k_{obs})$  of the formation of **8ba** was found to be inversely proportional to the concentration of N-Me-indole (Fig. S5<sup>†</sup>), which, at first, seemed contradictory to the turnover-limiting trapping of 19 to form 22. However, this should be due to the depletion of HNTf<sub>2</sub> by the basicity of indoles and the indole moiety in the product 8ba, as  $k_{obs}$  is linearly related to the concentration of HNTf<sub>2</sub> (Fig. S7<sup>†</sup>). In fact, for PhCH<sub>2</sub>CH<sub>2</sub>OH,  $k_{obs}$  was apparently independent of the concentration of the alcohol (Fig. S6†), likely because the increased concentration of the nucleophile counter-balanced the buffering of the superacid (HNTf<sub>2</sub>). Meanwhile, the reaction rate  $(k_{obs})$  was zeroth order in DMSO, which suggested that once the keteniminium 18 is formed, trapping with DMSO to yield 19 is very rapid and much faster than the dissociation to 20 or attack by N-Me-indole. This accounts for the chemoselectivity of this Umpolung alkylation process that occurs efficiently in a situation of multiple possible nucleophiles.5b,d

#### Synthetic applications

The current Umpolung alkylation provides unique structures that are difficult to obtain otherwise. Therefore, we examined their synthetic utility which is summarized in Scheme 4. Initially, 13ba (from 1.91 mmol of 4b, 69%) and 16ba (1.13 mmol of 4b, 75%) were prepared on a larger scale without any event. Esterification of these occurred chemoselectively to furnish the corresponding  $\varepsilon$ -alkoxydienoates 23 or  $\gamma$ -hydrazidoenoates 24, respectively, in good yields (eqn (2)). The imide product 8ba could be converted into a Weinreb amide 25 under basic conditions (eqn (3)).<sup>25</sup> The  $\varepsilon$ -alkoxy- $\alpha$ , $\beta$ , $\gamma$ , $\delta$ -unsaturated imide 13ba underwent Diels-Alder cycloaddition with N-methyl maleimide to give 26 with an exclusive endo-selectivity and a good level of substrate-controlled diastereoselectivity (dr = 8:1, eqn (4)). Conjugate addition with thiolate gave 27 with dr = 2:1. Subsequently, attempts to cleave the N–N bond in this mixture with SmI2 failed and instead 28 was obtained as a side product, presumably because of the Lewis acidity of Sm(III). Interestingly, 28 was obtained as a single diastereomer,<sup>26</sup> from the major diastereomer of 27. The N-N bond cleavage could then be achieved through a non-reductive method comprising alkylation of 28 and the in situ E1cb-elimination to form the



Scheme 4 Synthetic applications of the products.

lactam **29** (86%) (eqn (5)).<sup>27</sup> Interestingly, the compound **16ba** was found to be quite sensitive to bases and, upon treatment with  $Et_3N$ , cyclic hydrazone **30** was obtained in an excellent yield. Presumably, this occurs through a series of events, comprising conjugate addition of  $Et_3N$ , proton transfer, ring closure, elimination of  $Et_3N$  and isomerization (eqn (6)).

### Conclusions

Herein, we reported a generally applicable synthesis of  $\gamma$ -substituted- $\alpha$ , $\beta$ -unsaturated carbonyl compounds. This DMSObased oxidation protocol can accommodate diverse carbonand heteronucleophiles, including oxidizable heteroatom donors, in a highly chemo- and regio-selective manner. The origin of the remote-selectivity was interrogated by kinetic analysis and this suggested that the reaction proceeds through an  $S_N 2''$  or *via* a dienolonium carbocation, depending on the nucleophiles. This may provide mechanistic insights which may be of use in devising strategies for the future enantioselective processes.

## Conflicts of interest

There are no conflicts to declare.

### Acknowledgements

The authors thank the National Research Foundation of Korea (grant numbers NRF-2012M3A7B4049653, NRF-2014-011165 and NRF-2017R1A2B4010888) for generous financial support.

### Notes and references

- 1 (*a*) D. Seebach, *Angew. Chem., Int. Ed.*, 1979, **16**, 239–336; (*b*) O. Miyata, T. Miyoshi and M. Ueda, *ARKIVOC*, 2013, 60–81.
- 2 A. de la Torre, V. Tona and N. Maulide, *Angew. Chem., Int. Ed.*, 2017, **56**, 12416–12423.
- 3 [Au]: (a) H.-S. Yeom and S. Shin, Acc. Chem. Res., 2014, 47, 966-977; (b) L. Zhang, Acc. Chem. Res., 2014, 47, 877-888; (c) D. B. Huple, S. Ghorpade and R.-S. Liu, Adv. Synth. 1348–1367; (*d*) A. Mukherjee, Catal. 2016, 358, R. B. Dateer, R. Chaudhuri, S. Bhunia, S. N. Karad and R.-S. Liu, J. Am. Chem. Soc., 2011, 133, 15372-15375; (e) D. Vasu, H.-H. Hung, S. Bhunia, S. A. Gawade, A. Das and R.-S. Liu, Angew. Chem., Int. Ed., 2011, 50, 6911-6914; (f) L. Li, C. Shu, B. Zhou, Y.-F. Yu, X.-Y. Xiao and L.-W. Ye, Chem. Sci., 2014, 5, 4057-4064; (g) C. Shu, Y.-H. Wang, B. Zhou, X.-L. Li, Y.-F. Ping, X. Lu and L.-W. Ye, J. Am. Chem. Soc., 2015, 137, 9567-9570; [Rh]: (h) R. Liu, G. N. Winston-McPherson, Z.-Y. Yang, X. Zhou, W. Song, I. A. Guzei, X. Xu and W. Tang, J. Am. Chem. Soc., 2013, 135, 8201–8204; [Ir]: (i) G. Song, D. Chen, Y. Su, K. Han, C.-L. Pan, A. Jia and X. Li, Angew. Chem., Int. Ed., 2011, 50, 7791–7796; [Ru]: (j) K. Patil and R.-S. Liu, Chem. Commun. 2009, 5233-5235; [Zn]: (k) L. Li, B. Zhou, Y.-H. Wang, C. Shu, Y.-F. Pan, X. Lu and L.-W. Ye, Angew. Chem., Int. Ed., 2015, 54, 8245-8249.
- 4 For reviews: (a) A. Yoshimura and V. V. Zhdankin, Chem. Rev., 2016, 116, 3328-3435; (b) E. A. Merritt and B. Olofsson, Synthesis, 2011, 517-538; for selected examples: (c) S. Arava, J. N. Kumar, S. Maksymenko, M. A. Iron, K. N. Parida, P. Fristrup and A. M. Szpilman, Chem., Int. Ed., 2017, 56, 2599-2603; (d) Angew. О. Shneider, E. Pisarevsky, P. Fristrup S. and Szpilman, Org. Lett., 2015, 17, 282–285; (e) A. M. S. Maksymenko, K. N. Parida, G. K. Pathe, A. A. More, Y. B. Lipisa and A. M. Szpilman, Org. Lett., 2017, 19, 6312-6315; (f) P. Mizar and T. Wirth, Angew. Chem., Int. Ed., 2014, 53, 5993-5997.
- 5 (a) D. V. Patil, S. W. Kim, Q. H. Nguyen, H. Kim, S. Wang, T. Hoang and S. Shin, Angew. Chem., Int. Ed., 2017, 56, 3670-3674; (b) S. W. Kim, T.-W. Um and S. Shin, Chem. Commun., 2017, 53, 2733-2736; (c) S. W. Kim, T.-W. Um and S. Shin, J. Org. Chem., 2018, 83, 4703-4711; (d) D. V. Patil and S. Shin, Asian J. Org. Chem., 2019, 8, 63-73.
- 6 (a) D. Kaldre, B. Maryasin, D. Kaiser, O. Gajsek, L. González and N. Maulide, *Angew. Chem., Int. Ed.*, 2017, 56, 2212–2215;
  (b) Y. Zhao, Y. Hu, C. Wang, X. Li and B. Wan, *J. Org. Chem.*,

2017, **82**, 3935–3942; (c) Y. Zhao, Y. Hu, X. Li and B. Wan, Org. Biomol. Chem., 2017, **15**, 3413–3417.

- 7 For a related enolonium chemistry from dehydration of amides: (a) D. Kaiser, A. de la Torre, S. Shaaban and N. Maulide, Angew. Chem., Int. Ed., 2017, 56, 5921-5925; (b) S. Shaaban, V. Tona, B. Peng and N. Maulide, Angew. Chem., Int. Ed., 2017, 56, 10938-10941; (c) N. Takeda, E. Futaki, Y. Kobori, M. Ueda and O. Miyata, Angew. Chem., Int. Ed., 2017, 56, 16342-16346; (d) Z. Zhang, Y. Luo, H. Du, J. Xu and P. Li, Chem. Sci., 2019, 10, 5156-5161.
- 8 (a) S. Bertelsen, M. Marigo, S. Brandes, P. Dinér and K. A. Jørgensen, J. Am. Chem. Soc., 2006, 128, 12973-12980;
  (b) A. J. Fraboni and S. E. Brenner-Moyer, Org. Lett., 2016, 18, 2146-2149; (c) H. J. Zhang, A. W. Schuppe, S.-T. Pan, J.-X. Chen, B. R. Wang, T. R. Newhouse and L. Yin, J. Am. Chem. Soc., 2018, 140, 5300-5310; (d) X. Chen, X. Liu and J. T. Mohr, Org. Lett., 2016, 18, 716-719.
- 9 (a) S. Kölker, J. Inherited Metab. Dis., 2018, 41, 1055-1063; (b)
  M. Chebib and G. A. R. Johnston, Clin. Exp. Pharmacol. Physiol., 1999, 26, 937-940; (c) E. Szabadi, J. Psychopharmacol., 2015, 29, 744-749; (d) J. D. Griffin and L. Ellman, Semin. Thromb. Hemostasis, 1978, 5, 27-40.
- 10 Lithium dienolates are known to react preferentially at the α position, while alkylation of silyl dienol ethers occurs at the γ carbon: (a) S. E. Denmark, J. R. Heemstra Jr and G. L. Beutner, Angew. Chem., Int. Ed., 2005, 44, 4682-4698; (b) M. Frías, W. Cieślik, A. Fraile, A. Rosado-Abón, A. F. Garrido-Castro, F. Yuste and J. Alemán, Chem.-Eur. J., 2018, 24, 10906-10933.
- 11 For γ-functionalization of enynamides: (a) A. M. Jadhav,
  S. A. Gawade, D. Vasu, R. B. Dateer and R.-S. Liu, *Chem.-Eur. J.*, 2014, 20, 1813–1817; (b) A. M. Jadhav, D. B. Huple,
  R. R. Singh and R.-S. Liu, *Adv. Synth. Catal.*, 2016, 358, 1017–1022; (c) R. R. Singh and R.-S. Liu, *Adv. Synth. Catal.*, 2016, 358, 1421–1427; (d) S. S. Giri, L.-H. Lin, P. D. Jadhav and R.-S. Liu, *Adv. Synth. Catal.*, 2017, 359, 590–596.
- 12 For an excellent review on the redox chemistry of sulfoxides: A. P. Pulis and D. J. Procter, *Angew. Chem., Int. Ed.*, 2016, 55, 9842–9860.
- 13 For oxygen atom transfer from sulfoxides: (a) C.-W. Li, K. Pati, G.-Y. Lin, S. M. A. Sohel, H.-H. Hung and R.-S. Liu, Angew. Chem., Int. Ed., 2010, 49, 9891–9894; (b) S. Mori, M. Takubo, T. Yanase, T. Maegawa, Y. Monguchi and H. Sajiki, Adv. Synth. Catal., 2010, 352, 1630–1634; (c) C.-F. Xu, M. Xu, Y.-X. Jia and C.-Y. Li, Org. Lett., 2011, 13, 1556–1559; (d) H.-S. Yeom and S. Shin, Org. Biomol. Chem., 2013, 11, 1089–1092; (e) T. Stopka, M. Niggemann and N. Maulide, Angew. Chem., Int. Ed., 2017, 56, 13270–13274.
- 14 Use of Ph<sub>2</sub>S=O often resulted in [3,3]-sigmatropic rearrangement rather than O-atom transfer: (a)
  A. B. Cuenca, S. Montserrat, K. M. Hossain, G. Mancha,
  A. Lledós, M. Medio-Simón, G. Ujaque and G. Asensio, Org. Lett., 2009, 11, 4906-4909; (b)
  B. Peng, X. Huang, L.-G. Xie and N. Maulide, Angew. Chem., Int. Ed., 2014, 53, 8718-8721.
- 15 These reactions would produce a by-product apparently identical to the one *via* hydration of **4**. For interrupted

Pummerer reactions, see: Z. He, A. P. Pulis and D. J. Procter, *Angew. Chem., Int. Ed.*, 2019, **58**, 7813–7817 and references therein.

- 16 Assignment of the  $\alpha$  and  $\gamma$ -isomers was based on the analysis of <sup>1</sup>H NMR, COSY, and the nOe experiments (8ga, 8ha, 8na, 12bd, and 16ma). See the ESI.†
- 17 In contrast, the corresponding oxygenative alkylation of ynamides (according to ref. 5*a*) did not proceed when the oxidant changed from **Ox1** · (2-Cl-pyridine-*N*-oxide) to **Ox7** (DMSO).
- 18 However, the absence of an aryl group at C4 (for example, n = 1;  $R^1 = Ms$ ,  $R^2 = Me$ ,  $R^3/Ar = (CH_2)_4$ ) did not provide the corresponding product, indicating that an aryl group at C4 is necessary.
- 19 (a) P.-P. Ruan, C.-H. Shen, L. Li, C.-Y. Liu and L.-W. Ye, Org. Chem. Front., 2016, 3, 989-993; (b) G. F. Koser, A. G. Relenvi, A. N. Kalos, L. Rebrovic and R. H. Wettach, J. Org. Chem., 1982, 47, 2487-2489; (c) K. Ji, Y. Zhao and L. Zhang, Angew. Chem., Int. Ed., 2013, 52, 6508-6512; (d) F. Pan, X.-L. Li, X.-M. Chen, C. Shu, P.-P. Ruan, C.-H. Shen, X. Lu and L.-W. Ye, ACS Catal., 2016, 6, 6055-6062; (e) V. Tona, A. de la Torre, M. Padmanaban, S. Ruider, L. Gonzalez and N. Maulide, J. Am. Chem. Soc., 2016, 138, 8348-8351; (f) D. H. Miles, J. Guasch and F. D. Toste, J. Am. Chem. Soc., 2015, 137, 7632-7635; (g) X. Xia, B. Chen, X. Zeng and B. Xu, Org. Biomol. Chem., 2018, 16, 6918-6922; (h) A. de la Torre, D. Kaiser and N. Maulide, J. Am. Chem. Soc., 2017, 139, 6578-6581; (i) S. Taninokuchi, R. Yazaki and T. Ohshima, Org. Lett., 2017, 19, 3187-3190; (j) X. Li, F. Lin, K. Huang, J. Wei, X. Li, X. Wang, X. Geng and N. Jiao, Angew. Chem., Int. Ed., 2017, 56, 12307-12311.
- 20 J. Yu, J. Tian and C. Zhang, *Adv. Synth. Catal.*, 2010, **352**, 531–546.
- 21 CCDC 1895889 (16ma).†
- 22 V. Tona, S. A. Ruider, M. Berger, S. Shaaban, M. Padmanaban, L.-X. Xie, L. González and N. Maulide, *Chem. Sci.*, 2016, 7, 6032–6040.
- 23 However, both the reactions of *N*-Me-indole and BnOH had some degree of 'concertedness', as suggested by the asymmetric induction test (14% ee and 29% ee, respectively), employing chiral *N*,*N*'-dioxide (ref. 5*a*; eqn (S4)<sup>†</sup>).
- 24 A reviewer suggested a possibility that the reaction with indoles may follow a concerted  $S_N 2''$  substitution and such a reaction may display a large negative entropy of activation. For a similarly large negative entropy of activation in the  $S_N 2$  reaction: K. Okamoto, S. Fukui, I. Nitta and H. Shingu, *Bull. Chem. Soc. Jpn.*, 1967, **40**, 2350–2353.
- 25 F. Sirindi, S. Miaskiewicz, N. Kern, A. Lalaut, A.-S. Felten, J. –
  M. Weibel, P. Pale and A. Blanc, *Tetrahedron*, 2017, 73, 5096–5106.
- 26 An nOe experiment supported the relative stereochemistry in 27.
- 27 J. Ferreira, S. C. M. Rees-Jones, V. Ramaotsoa, A. Msutu and R. Hunter, *Org. Biomol. Chem.*, 2016, **14**, 1545–1549.